

TRADESCANTIA YELLOW LEAF SPOT FUNGUS

Kordyana brasiliensis

The history of tradescantia yellow leaf spot in New Zealand

The tradescantia yellow leaf spot fungus is native to Brazil. Permission to release this fungus was granted by the EPA in 2013. However, the fungus was not imported and released immediately to allow some time to see how well the three species of tradescantia beetles establish and to determine if this additional agent would be needed. The beetles have established well and are showing great promise. However, some release sites have been flooded and, since tradescantia often occurs in riparian areas, flooding will likely be a regular occurrence in some tradescantia-infested areas. It is likely that the beetles will struggle in areas that are subject to regular flooding, and that the fungus will be better suited to these conditions. Releases of the tradescantia yellow leaf spot will therefore get underway early in 2018.

The fungus was imported from a CSIRO containment facility in Australia by Landcare Research in 2017. Australia is also considering the suitability of releasing the fungus against *Tradescantia fluminensis* there, but New Zealand will be the first country to carry out field releases.

See *Tradescantia leaf beetle*, *Tradescantia stem beetle*, *Tradescantia tip beetle*.



Yellow leaf spot

How would I find tradescantia yellow leaf spot?

Infection begins when spores germinate on the surface of tradescantia leaves. This requires some moisture to be present. The fungus penetrates leaves through stomata present on the undersides, and about 10 days later, diffuse leaf spots appear on the upper leaf surface. These spots develop, expand and turn yellow to brown as the infected leaves die, but generally remain attached to the stem. Look for these distinctive yellow spots on the leaves, especially after wet, humid weather during the warmer months. Within 15 days of infection the spot centres turn white underneath, and if humidity is high enough, new basidiospores form there. Basidiospores are very small and lightweight, and are easily spread through air turbulence. The whole cycle starts again as soon as they land on a new tradescantia leaf.

Occasionally other pathogens are found damaging tradescantia leaves in New Zealand. Some pathogens, such as species from the genus *Colletotrichum*, produce brown to black lesions with sometimes a yellow border. Other pathogens exhibit brown to black lesions spreading from the leaf tip towards the base. However, the characteristic white spots on the



Typical symptoms

TRADESCANTIA



lower leaf surface and the yellow spots on the upper leaf surface caused by the tradescantia yellow leaf spot fungus should prevent any confusion with other pathogens.

You are also unlikely to confuse damage from the tradescantia beetles with the yellow leaf spot fungus. The tradescantia leaf beetle (*Neolema ogloblini*) causes notching and skeletonisation of the leaves, and the tip beetle (*Neolema abbreviata*) damages the leaf tips. Adult tip beetles and adult stem beetles (*Lema basicostata*) make small holes in the leaves, and the stem beetle larvae cause the stems to turn brown, shrivel and collapse.

How does tradescantia yellow leaf spot damage tradescantia?

The infection process damages the epidermis of the leaves causing the leaves to shrivel and die prematurely.

Will tradescantia yellow leaf spot attack other plants?

The yellow leaf spot fungus is highly host-specific and highly unlikely to attack anything other than tradescantia (*Tradescantia fluminensis*).

How effective is tradescantia yellow leaf spot?

It is too soon to know what impact the tradescantia yellow leaf spot fungus will have, on its own or in conjunction with the tradescantia beetles. However, the four agents are expected to complement each other and a monitoring programme has been set up to measure their effectiveness. Laboratory studies suggest that the tradescantia smut grows best at temperatures around 20°C.

How can I get the most out of tradescantia yellow leaf spot?

The yellow leaf spot fungus will disperse naturally via spores on air currents. However, if help is needed to establish the fungus, especially in more remote areas, then it will be necessary to put potted, infected plants out at these sites. First of all put healthy plants out at sites where the fungal infection is present until symptoms appear on them. Infection should occur readily in this manner (note that it is not possible to work with the tiny fragile spores as they cannot be grown on artificial media). The fungus is able to infect tradescantia plants most effectively at temperatures between 16 and 22°C, and it requires a long period of high humidity. A release during high rainfall will also likely assist with the successful establishment of this pathogen. Note that if you wish to propagate and move tradescantia plants around in this manner you will need an exemption from the Ministry for Primary Industries.

How do I choose a release site?

Read *Guidelines for selecting release sites for biocontrol agents*.

Tradescantia is usually present in damp shaded areas but choose release sites that are sheltered from wind and sun where the leaves are more likely to remain sufficiently moist. Sites should also be safe from frost, fire and human interference such as spraying and slashing.

How do I manage the release sites?

Avoid any activities that will interfere with the yellow leaf spot, such as herbicide or fungicide application. If you need to undertake control measures then avoid the release site.

For further information contact:

Chantal Probst
Landcare Research
Private Bag 92170
Auckland 1142
NEW ZEALAND
Email: probstc@landcareresearch.co.nz
Ph (09) 574 4162
Fax (09) 574 4101