

Fauna of New Zealand

Ko te Aitanga Pepeke o Aotearoa

Vink, C. J.; Dupérré, N.; Malumbres-Olarte, J. 2013. Periegopidae (Arachnida: Araneae). *Fauna of New Zealand* 70: 41 pp.

The Copyright notice printed on page 4 applies to the use of this PDF.

This PDF is not to be posted on websites.

Links should be made to:

FNZ.LandcareResearch.co.nz

EDITORIAL BOARD

Dr R. M. Emberson, *c/- Department of Ecology, P.O. Box 84, Lincoln University,
New Zealand*

Dr M. J. Fletcher, *NSW Agricultural Scientific Collections Unit, Forest Road,
Orange, NSW 2800, Australia*

Dr R. J. B. Hoare, *Landcare Research, Private Bag 92170, Auckland, New
Zealand*

Dr M.-C. Larivière, *Landcare Research, Private Bag 92170, Auckland, New
Zealand*

Mr R. L. Palma, *Natural Environment Department, Museum of New Zealand
Te Papa Tongarewa, P.O. Box 467, Wellington, New Zealand*

SERIES EDITOR

Dr T. K. Crosby, *Landcare Research, Private Bag 92170, Auckland, New
Zealand*

**Fauna of New Zealand
Ko te Aitanga Pepeke o Aotearoa**

Number / Nama 70

**Periegopidae
(Arachnida: Araneae)**

C. J. Vink

**AgResearch, Biocontrol & Biosecurity
Private Bag 4749, Christchurch 8140, New Zealand
(present address: Canterbury Museum, Rolleston Avenue,
Christchurch 8013, New Zealand
cvink@canterburymuseum.com)
Entomology Research Museum
PO Box 84, Lincoln University, Lincoln 7647, New Zealand**

N. Dupérré

**341 15ème Rue, Laval, Québec, H7N 1L5, Canada
nadineduperre@gmail.com**

J. Malumbres-Olarte

**Department of Ecology
PO Box 84, Lincoln University, Lincoln 7647, New Zealand
jagoba.malumbres.olarte@gmail.com**



**Manaaki
Whenua
P R E S S**

**Lincoln, Canterbury, New Zealand
2013**

Copyright © Landcare Research New Zealand Ltd 2013

No part of this work covered by copyright may be reproduced or copied in any form or by any means (graphic, electronic, or mechanical, including photocopying, recording, taping information retrieval systems, or otherwise) without the written permission of the publisher.

Cataloguing in publication

Vink, C. J. (Cornelis Jacob)

Periegopidae (Arachnida: Araneae) / C. J. Vink, N. Dupérré and J. Malumbres-Olarte. – Lincoln, Canterbury, N.Z. : Manaaki Whenua Press, 2013.

(Fauna of New Zealand, ISSN 0111-5383 (print), ISSN 1179-7193 (online) ; no. 70).

ISBN 978-0-478-34740-1 (print)

ISBN 978-0-478-34741-8 (online)

I. Dupérré, N. (Nadine) II. Malumbres-Olarte, J. (Jagoba) III. Title. IV. Series.

UDC 595.44

Suggested citation:

Vink, C. J.; Dupérré, N.; Malumbres-Olarte, J. 2013. *Periegopidae* (Arachnida: Araneae). *Fauna of New Zealand* 70: 41 pp.

Prepared for publication by the series editor and the authors using computer-based text processing, layout, and printing at Landcare Research, Private Bag 92170, Auckland, New Zealand.

To access on-line extracts from this series visit:

<http://fnz.landcareresearch.co.nz/>

Māori text by Piripi Walker, Tokomāpuna Māori Language Services, Upper Hutt/Whakatiki.

Published by Manaaki Whenua Press, Landcare Research, P.O. Box 40, Lincoln, Canterbury, N.Z.

Website: <http://www.mwpress.co.nz/>

Printed by PrintLink Ltd, Wellington

Date of publication 7 March 2013

Front cover: Female *Periegops suterii* Urquhart from Kennedys Bush Reserve, MC (LUNZ 00012718)

(Photographer: Bryce McQuillan).

Publication of the *Fauna of New Zealand* series is supported by Core funding for Crown Research Institutes from the Ministry of Business, Innovation and Employment's Science and Innovation Group, for the Defining New Zealand's Land Biota research programme.

POPULAR SUMMARY

HE WHAKARĀPOPOTOTANGA

Class **Periegopidae**Order **Arachnida**Family **Araneae****Periegopid spiders**

There are only three species known in the family Periegopidae and all are in one genus, *Periegops*. These rare spiders have only ever been found in relict forests at limited locations in New Zealand (Banks Peninsula, Riccarton Bush, the Aldermen Islands, and East Cape) and Queensland, Australia. Periegopids are only found in forest with a deep leaf litter layer and well-drained soil. They do not build a web, but hunt on the forest floor. Periegopids can be most readily distinguished from other spiders found in New Zealand by having six eyes arranged in three widely spaced diads.

Contributor **Cor Vink** was born and educated in Christchurch, New Zealand. He completed a Ph.D. at Lincoln University on the taxonomy and systematics of New Zealand Lycosidae, a major part of which was published as a revision in *Fauna of New Zealand 44*. After completing his thesis he spent nine months at AgResearch as a postdoctoral research fellow investigating the genetics of hymenopteran parasitoids of weevil pests. From 2003 to 2005, Cor was a postdoctoral associate at San Diego State University, U.S.A., where he worked on developing new molecular markers for inferring deep phylogenetic relationships in spiders. At the end of 2005 he returned to New Zealand and joined the Biosecurity Group at AgResearch, Lincoln as a scientist. Cor is especially interested in the systematics of New Zealand spiders and is the adjunct curator of spiders at the Entomology Research Museum at Lincoln University. In 2013, Cor was appointed as Curator Natural History at Canterbury Museum, Christchurch.

(continued overleaf)




Illustration / Whakaahua: *Periegops suterii*, female
(Illustrator / Kaiwhakaahua: N. Dupérré).

Ngā pūngāwere Periegopid

E toru anake ngā momo e mōhiohia ana o te whānau Periegopidae, ā, nō te huinga kotahi, arā, a *Periegops*. Kua kitea ēnei pūngāwere onge tonu i ngā toenga ngahere i ngā wāhi ruarua nei i Aotearoa (i Horomaka, i te ngahere o Riccarton, i ngā moutere Aldermen, me Te Koringa-a-Paoa) me Queensland, i Ahitereiria. Kitea ai ēnei Periegopid i te ngahere he hōhonu nei ōna pūranga rau-rākau, i ngā one āhua pai te mimititanga wai. Kāore e hanga tukutuku mō rātou, engari ka whakangau i ā rātou kai i te papa tonu o te ngahere. Ka mōhiohia wawetia te Periegopid, ina whakaritea ki ētahi katoa o ngā pūngāwere o Aotearoa, nā te noho mai o ngā karu e ono, he tawhiti tonu te takoto o ia pūruatanga karu i ētahi atu.

Translation by **Piripi Walker**
Whakatiki

Contributor **Nadine Dupérré** is from Québec, Canada. She completed a Bachelor of Science at Université de Montréal in 1997. She began illustrating spiders in 1998 and to date has produced over 5000 illustrations. Nadine is probably best known amongst arachnologists for her illustrations in the book “Spiders of North America: an Identification Manual”. Nadine has published on the taxonomy of spiders and has also produced illustrations for publications on beetles and harvestmen. Nadine was a research assistant at the American Museum of Natural History, New York from 2008–2012 where she worked on Oonopidae with Norman Platnick. She is now living in Quito, Ecuador, working on a revision of the genus *Agyneta* (Linyphiidae) from North America, and is planning to continue her work on Linyphiidae in South America.

Contributor **Jagoba Malumbres-Olarte** is from the Basque Country, Spain. He completed a Ph.D. at Lincoln University on the ecology and diversity of spider communities in New Zealand native tussock grasslands in 2011. After he completed his thesis, he worked as a teaching assistant at Lincoln University and Imperial College London, UK. In the last six years Jagoba has been involved in numerous research projects on diverse topics, including the ecology of spider communities in sand dunes, conservation genetics of the Chatham Islands coxella weevil, the systematics of the New Zealand *Clubiona*, and the invasibility of spiders and plants in New Zealand. Jagoba is particularly interested in ecological and environmental processes that drive speciation and shape the diversity and composition of communities, and is currently developing new projects in this field.



Nadine Dupérré



Cor Vink



Jagoba Malumbres-Olarte

ABSTRACT

Two species of Periegopidae, both in the genus *Periegops* Simon, 1893, are found in New Zealand; *P. suterii* (Urquhart, 1892) and *P. keani* sp. nov. The genus and both species are described or redescribed, with information on synonymy, type data, material examined, and geographical distribution. Habitus images of adults, illustrations of important morphological features, and distribution maps are provided. A key is given.

The mitochondrial gene cytochrome *c* oxidase subunit 1 (COI) and the nuclear gene 28S ribosomal RNA were sequenced for both species. COI divergence between some specimens of *P. suterii* was unexpectedly high (8.6%, uncorrected distance), but there was no morphological differentiation or 28S sequence divergence between those specimens. A molecular phylogenetic analysis examining the relationships between eight specimens of *P. suterii* and three specimens of *P. keani* using COI data is presented.

Keywords. Arachnida, Araneae, Periegopidae, New Zealand, Haplogynae, classification, distribution, ecology, biology, new species, key, phylogeny.

Vink, C. J.; Dupérré, N.; Malumbres-Olarte, J. 2013. Periegopidae (Arachnida: Araneae). *Fauna of New Zealand* 70: 41 pp.

Received: 5 July 2012. Accepted: 19 November 2012.

CHECKLIST OF TAXA

Genus <i>Periegops</i> Simon, 1893.....	15
<i>suterii</i> (Urquhart, 1892).....	15
<i>keani</i> new species.....	17

CONTENTS

Acknowledgments	8	Phylogenetic analysis	13
Introduction	8	Methods	13
Morphology and terminology	8	Results	13
Methods and conventions	10	Discussion	14
Collecting	10	Key to Periegopidae known from New Zealand	15
Preservation	10	Biosystematics	15
Preparation	10	References	19
Measurements	10	Appendices	
Descriptions	10	A: Glossary of technical terms	23
Illustrations	11	B: Collection details of specimens examined	24
Text conventions	11	Illustrations	25
Molecular biology	11	Distribution maps	34
		Taxonomic index	35

ACKNOWLEDGMENTS

We are indebted to Thomas Buckley (New Zealand Arthropod Collection, Auckland) for his support of this project.

We thank the following for the loan of specimens in their care; Phil Sirvid (Museum of New Zealand Te Papa Tongarewa, Wellington), Grace Hall (New Zealand Arthropod Collection, Auckland), John Marris (Entomology Research Museum, Lincoln University), Cody Fraser (Otago Museum, Dunedin), Robert Raven (Queensland Museum, Brisbane), and Norman Platnick (American Museum of Natural History, New York). Thanks to Christine Rollard (Muséum national d'Histoire naturelle, Paris) for checking for the presence of the type of *Periegops hirsutus*.

Mike Bowie (Lincoln University) kindly supplied specimens that have been valuable in this study. Mike Fitzgerald and Grace Hall shared information on the locations where *Periegops* specimens had been collected. Thanks to John Kean (AgResearch) for collecting assistance on Ruamahuanui Island, to Rob Chappell (Department of Conservation) for logistics support and to Kristine Grayson (Victoria University of Wellington) for sharing her charter to the Aldermen Islands. Henry Hudson Vink and Lucy Hudson Vink helped collect specimens at Kennedys Bush Reserve. We thank Bryce McQuillan for his excellent photographs of *Periegops suterii*.

Facundo Labarque (California Academy of Sciences), Phil Sirvid, and Thomas Buckley provided helpful comments on an earlier draft of the manuscript.

Collecting permits were supplied by the New Zealand Department of Conservation (permits WK-30010-FAU and CA-28815-FAU). The Ruamaahua Island Trust granted cultural permission to collect on Ruamahuanui Island. This research was supported by Core funding for Crown Research Institutes from the Ministry of Business, Innovation and Employment's Science and Innovation Group through the Defining New Zealand's Land Biota Portfolio.

INTRODUCTION

Periegopidae Simon, 1893 is a family of six-eyed spiders comprising only three species, all in the genus *Periegops* Simon, 1893. Two species are endemic to New Zealand and the other is endemic to Queensland, Australia (Forster 1995). *Periegopids* can be distinguished from other families found in New Zealand by a combination of characters. The six eyes are arranged in three, widely spaced diads. The chelicerae have a lamina on the ventral surface and the maxillae are slender and are more than twice as long as wide and directed across the labium. The anterior two pairs of legs have asymmetrical superior claws; the proclaw of legs I and II has a double row of teeth whereas the retroclaw has a single row (Forster 1995: fig. 11–13). There is also a field of spicules on the median surface of the posterior median spinnerets (Labarque & Ramírez 2012; fig. 21F, 22E) and the venom outlet is anterior on the fang (Labarque & Ramírez 2012; fig. 18A); both of these characters require high microscope magnification to see them.

Periegops suterii (Urquhart, 1892) was first described from a female and two immature males collected by the early New Zealand zoologist, Henry Suter, at Dyer's Pass on the Port Hills near Christchurch (Urquhart 1892; Forster 1995). Urquhart placed his new species in a Northern Hemisphere genus and family; *Segestria* Latreille, 1804, and *Dysderidae* C.L. Koch, 1837, respectively. Shortly after, Simon (1893) erected the genus *Periegops* for his new species *P. hirsutus*, which he described from a female specimen sent to him at the Muséum national d'Histoire naturelle, Paris, from New Zealand. Although Simon (1893) did not specify the location where the specimen was collected from, it was likely that it was from Banks Peninsula (Forster 1995). The generic description was of a female and Simon (1893: 267) noted that the male was unknown; however, the description of *P. hirsutus* was listed as male although no male characteristics appear to be described (Simon 1893: 268). Bryant (1935a: 54) and Forster (1995: 92) noted that the listing as a male appears to be in error. Dalmas (1917: 338) also noted the specimen was female and this was the specimen seen by Chamberlain (1946). It is unlikely that Simon realised he had described the same species as Urquhart and *P. hirsutus* was later made a synonym of *P. suterii* by Chamberlain (1946). Simon (1893) was first to recognise that the genus *Periegops* was distinct from other genera and placed it in its own subfamily, *Periegopinae* Simon, 1893. Forster (1995) elevated the subfamily to family status when he described *Periegops australia* Forster, 1995 and redescribed *P. suterii*. Forster (1995: 96) also noted that he had examined a single

female specimen collected at East Cape, but there were “no clear characters by which the species could be satisfactorily defined”; a male would need to be collected and the palpal bulb examined to determine if it was a separate species. Forster & Forster (1999) mentioned an additional specimen from the Aldermen Islands, which they believed to be a female, and Vink (2006) reported that he had sequence data from a fragment of the mitochondrial gene cytochrome *c* oxidase subunit 1 (COI) [specimen Pk1] that suggested that the North Island specimens were a distinct species.

Periegopidae is part of a group of spiders called the Haplogynae Simon, 1893, which are araneomorph spiders that lack separate fertilisation ducts, do not have a sclerotised epigyne and the male pedipalp is relatively simple. Haplogynae are a monophyletic group (Platnick *et al.* 1991; Ramírez 2000) that includes 17 families. Periegopids have a number of characteristics that are shared with the haplogyne families Diguettidae F. O. Pickard-Cambridge, 1899, Drymusidae Simon, 1893, Plectreuridae Simon, 1893, Scytodidae Blackwall, 1864, and Sicariidae Keyserling, 1880. These include a lamina on the ventral surface of the chelicerae, slender maxillary lobes directed across the labium, and a limited posterior respiratory system with fused apodemes (Forster 1995). Forster (1995) grouped this set of families, along with Periegopidae, into the superfamily Sicarioidea, but did not provide a formal definition of the superfamily. The name Sicarioidea was first used by Berland (1932) and Forster (1995) appears to be the only other arachnologist that has used it.

A more commonly used superfamily name that includes this set of taxa is Scytodoidea (Bristowe 1938; Caporiacco 1938; Brignoli 1978; Lehtinen 1986); however, Scytodoidea has been used to refer to a larger set of haplogyne families that has also included Caponiidae Simon, 1890, Leptonetidae Simon, 1890, Ochyroceratidae Fage, 1912, Pholcidae C. L. Koch, 1850, and Tetrablemmidae O. Pickard-Cambridge, 1873. A morphological phylogenetic analysis by Platnick *et al.* (1991) supports a clade informally named “scytodoids” (Coddington & Levi 1991) that includes Diguettidae, Drymusidae, Leptonetidae, Ochyroceratidae, Pholcidae, Plectreuridae, Scytodidae, Sicariidae, Telemidae Fage, 1913, and Tetrablemmidae. The monophyly of this clade is supported by the reduction of the posterior spiracles to one and the tetrahedral posterior median spinnerets (Platnick *et al.* 1991), although these characters have been reversed in some species in the clade (Platnick *et al.* 1991).

Ramírez (2000) examined the morphology of haplogyne respiratory systems and added characters to the data matrix of Platnick *et al.* (1991) to produce a phylogeny

of the Haplogynae. Periegopidae was not included in the analyses but Ramírez (2000) stated that Periegopidae was sister to the Scytodidae. Periegopidae was also included in the scytodoid clade by Coddington (2005); its inclusion was presumably based on Forster (1995) and Ramírez (2000). Labarque & Ramírez (2007a, b) considered that among the scytodoids, Periegopidae was closest to Drymusidae and Scytodidae as they share two characters; asymmetrical superior claws on the anterior two pairs of legs and a field of spicules on the median surface of the posterior median spinnerets. More recently, Labarque & Ramírez (2012) expanded the morphological dataset of Platnick *et al.* (1991) and Ramírez (2000) to include Periegopidae along with a restricted Scytodoidea, which included Drymusidae, Scytodidae, and Sicariidae. *Periegops*, *Drymusa*, and Scytodidae formed a clade, which all shared the synapomorphies of asymmetrical superior claws and a field of spicules on the median surface of the posterior median spinnerets. Labarque & Ramírez (2012) concluded that Periegopidae was sister to Drymusidae, but they did not find any autapomorphies for *Periegops* and suggested that the two families might be synonymised.

Very little is known of the biology of periegopids. They are only found in forest with a deep leaf litter layer and well-drained soil. Specimens have been collected from under logs and rocks, in leaf litter, in pitfall traps, in rotted logs and root cavities, and in a mygalomorph burrow (Forster 1995; Vink 2006). Forster (1995) suggested that periegopids were rare as their habitat has been severely reduced due to human deforestation and only 21 *Periegops* specimens had been collected in New Zealand and Australia between 1891 and 1995. In surveys between October 2002 and May 2003, Vink (2006) found a further 24 specimens of *P. suterii* at ten different locations on the Banks Peninsula. Sirvid *et al.* (2012) classified *P. suterii* as having a relict distribution.

Periegops do not appear to build a web for prey capture, but do build silken retreats. It seems probable that periegopids are cursorial, night-time hunters as they have only been found on the forest floor, have been collected in pitfall traps and lack webs. On two occasions a single female has been found together with two or three males under logs and rocks (Forster 1995; Vink 2006), which implies that the female might possess some method of attracting males.

MORPHOLOGY AND TERMINOLOGY

The morphological structures required for the identification of New Zealand and Australian Periegopidae are

referred to in Fig. 1–16 and explained in the glossary of technical terms (Appendix A), and Paquin *et al.* (2010). As with almost all spiders, the male pedipalp is important when identifying periegopids to species. The female epigyne, which is usually useful for identifying spider species (Paquin *et al.* 2010), is just a lightly sclerotised area that covers the internal genitalia in periegopids, and many other haplogyne spiders, and therefore is not likely to be diagnostic. The internal genitalia, however, are diagnostic, as is the case with most spider species. The morphological nomenclature of the pedipalp and the internal genitalia follows Forster (1995).

A phylogenetic species concept (Wheeler & Platnick 2000) has been implemented in this study. It defines a species as the smallest aggregation of populations diagnosable by a unique combination of character states.

METHODS AND CONVENTIONS

Collecting. Periegopids can be collected by a variety of methods. The best method for collecting periegopids is by searching under rocks and logs that are sitting on or partially embedded in soil on open ground within forest in areas with a good litter layer over well drained soil (Vink 2006). Periegopids have also been found under wooden discs (M. H. Bowie, pers. comm.) that are facsimiles for natural fallen logs (Bowie & Frampton 2004). Periegopids have also been collected by sieving and searching leaf litter. Specimens of *P. suterii* and *P. australia* have been caught in pitfall traps, but unless specimens are collected and preserved within a couple of days of being caught they can start to decay, which can make identification difficult. Decay of specimens caught in pitfall traps can be mitigated by the use of a good preservative such as propylene glycol, which also preserves DNA but can shrivel soft tissue (Vink *et al.* 2005). Malumbres-Olarte *et al.* (in press) used propylene glycol in their pitfall traps and found that useful DNA was preserved after two weeks in the field.

Preservation. Periegopids are best preserved long-term in 70–75% ethanol. To ensure adequate DNA preservation, specimens should also be stored at $\leq -20^{\circ}\text{C}$ (Vink *et al.* 2005). Spiders can be stored in 95–100% ethanol to preserve DNA but as with lower ethanol concentrations, it is still best to combine this with storage temperatures $\leq -20^{\circ}\text{C}$ (Vink *et al.* 2005). Preservation in 95–100% ethanol makes specimens brittle, potentially rendering them unsuitable for morphological examination.

Preparation. Specimens should be labelled with the locality, including area code (Crosby *et al.* 1976, 1998), latitude and longitude, collection date, collector's name, habitat data, and collection method.

Most morphological features used in identification can be seen under an ordinary dissecting microscope. When examining a spider in ethanol it should be rested in washed quartz sand or glass beads to provide support for the specimen. This also allows the specimen to be positioned at any desired viewing angle. The features of the male pedipalp are best viewed by removing the left pedipalp at the junction between the trochanter and the femur, and viewed ventrally. In periegopids, as with other haplogyne spiders, there is no sclerotised external genitalia, only a slightly sclerotised plate, which makes distinguishing females from immature specimens difficult. Female internal genitalia were excised using a sharp entomological needle and were prepared for examination by placing the dissected genitalia in either lactic acid or 10% KOH solution for 1–3 hours at 37°C to dissolve soft tissue. Internal genitalia were only illustrated for *P. suterii* and *P. australia*.

Measurements. All measurements are in millimetres (mm). Where the measurements are expressed as a fraction, the numerator refers to the length of the structure and the denominator refers to its width. Measurements outside parentheses are for males and inside parentheses are for females. The size ranges given for the body and carapace length of each species represent the smallest and largest individuals of each sex that we examined. The mean body length and carapace length were calculated and the number of specimens measured given.

Descriptions. For the new species, illustrations, measurements and colour pattern descriptions were made from the type specimen. For the existing species, illustrations, measurements, and colour pattern descriptions were prepared from a non-type representative male and female specimen (with collection information shown).

Descriptions of colours are for ethanol-preserved specimens. It should be noted that colours and colour patterns can fade in older specimens, particularly those that have not been stored away from light.

Measurements were made with a micrometer ruler fitted to the eyepiece of a Leica M125 stereo microscope.

Characters diagnostic in other spider families (e.g., eye size and position, leg spination) are not diagnostic for Periegopidae species and have not been included in the descriptions.

Illustrations. Specimens to be illustrated were first photographed with a Nikon Coolpix 950 digital camera attached to a SMZ-U Nikon dissection microscope. The digital photos were then used to establish proportions and the illustrations were detailed and shaded by referring back to the structure under the microscope.

Map images were created using the geographic information system software ArcMap 10 (ESRI).

Text conventions. The area codes of Crosby *et al.* (1976, 1998) are used in collection records.

The following acronyms for repositories are used:

AMNH	American Museum of Natural History, New York, U.S.A.
LUNZ	Entomology Research Museum, Lincoln University, New Zealand
MONZ	Museum of New Zealand Te Papa Tongarewa, Wellington, New Zealand
NZAC	New Zealand Arthropod Collection, Auckland, New Zealand
OMNZ	Otago Museum, Dunedin, New Zealand
QMB	Queensland Museum, Brisbane, Australia

Molecular biology. To construct a molecular phylogeny of New Zealand Periegopidae and to facilitate the identification of immature specimens, we used the mitochondrial gene cytochrome *c* oxidase subunit I (COI) and the nuclear ribosomal RNA gene 28S. COI is one of the fastest evolving mtDNA genes and has been used to examine genetic differences between and among haplogyne spiders (e.g., Arnedo *et al.* 2001, 2009; Astrin *et al.* 2006; Starrett & Waters 2007; Binford *et al.* 2008; Dimitrov *et al.* 2008; Huber & Astrin 2009; Duncan *et al.* 2010) and New Zealand spider species (Vink & Paterson 2003; Vink *et al.* 2008, 2011a, b; Framenau *et al.* 2010; Vink & Dupérré 2010; Lattimore *et al.* 2011; Malumbres-Olarte & Vink 2012). 28S was selected because it is a slow evolving gene (Hedin & Maddison 2001) and has been used in phylogenetic analyses of haplogynes (Bruvo-Madaric *et al.* 2005; Binford *et al.* 2008) as well as other spiders (e.g., Hedin & Bond 2006; Rix *et al.* 2008; Wang *et al.* 2008; Spagna *et al.* 2010). Eight specimens of *P. suterii* and three specimens of *P. keani* were sequenced for COI. Each specimen was arbitrarily assigned a specimen code (Table 1). A subset of four specimens, two from each species, was sequenced for 28S.

DNA was extracted non-destructively (see Paquin & Vink 2009) from either two to three legs using a ZR Genomic DNA™ Tissue Minipreps (Zymo Research). The primers used to PCR amplify and sequence COI fragments were either LCO1490 (5'-GGT-

CAACAAATCATAAAGATATTGG-3') (Folmer *et al.* 1994) plus C1-N-2776-spider (5'-GGATAATCAGAATANCGNCGAGG-3') (Vink *et al.* 2005) or C1-J-1718-spider (5'-GGNGGATTTGGAAATTGRT-TRGTTCC-3') (Vink *et al.* 2005) plus C1-N-2776-spider. The two COI fragments were 1260 base pairs (bp) and 1055 bp long, respectively. Two different primer pairs were used to amplify and sequence two overlapping 28S fragments; 28S-B1 (5'-GACCGATAGCAAACAAGTACCG-3') (Bruvo-Madaric *et al.* 2005) plus 28S-B2 (5'-GATTAGTCTTTCGCCCTATA-3') (Bruvo-Madaric *et al.* 2005) and 28Sa (5'-GACCCGTCTTGAAACACGGA-3') (Nunn *et al.* 1996) plus LSUR (5'-GCTACTACCACCAAGATCTGCA-3') (Rix *et al.* 2008). The two 28S fragments were 819–821 bp and 830–833 bp long, respectively. PCR amplification was performed using *i*-StarTaq™ DNA Polymerase (iNtRON Biotechnology) in a Mastercycler® (Eppendorf) thermocycler with a cycling profile of 35 cycles of 94°C denaturation (30 s), 48°C (COI) or 60°C (28S) annealing (30 s), 72°C extension (1 min) with an initial denaturation of 3 min and a final extension of 5 min. Excess primers and salts were removed from the resulting double-stranded DNA using a DNA Clean & Concentrator™ Kit (Zymo Research). Double bands were observed when 28S PCR products of Pk1 were visualised via gel electrophoresis; both bands were excised from the gel and prepared for sequencing using a Zymoclean™ Gel DNA Recovery kit. For Pk1, the primer pair 28S-B1 plus 28S-B2 amplified a fragment of 28S DNA that BLAST database searching indicated may have come from a ciliate protozoan. For Pk2, the primer pair 28Sa plus LSUR amplified a fragment of 28S DNA from a soil nematode in the family Cephalobidae. Amplification of 28S was attempted for specimens Ps5 and Ps6, but was not possible owing to contamination by a fungus that the primers preferentially annealed to. Contaminants were identified by BLAST searching their sequences (Altschul *et al.* 1997). Purified PCR fragments were sequenced in both directions at the Core Instrumentation Facility (University of California, Riverside, USA), the Massey Genome Service (Massey University, New Zealand), or Macrogen (Korea). Sequence data were deposited in GenBank (www.ncbi.nlm.nih.gov/GenBank/ – see Table 1 for accession numbers). Sequences were edited using Sequencher 4.6 (Gene Codes Corporation).

Table 1. Specimens used for molecular analysis.

Species	Specimen code	Sex	Collection information	GenBank accession numbers
<i>Periegops suterii</i>	Ps1	♂	MC, Hinewai Reserve, 43°48.59'S, 173°01.28'E, 11 October 2002, CJV, AMNH (ARAMR-000615)	JX174281
<i>Periegops suterii</i>	Ps2	♀	MC, Hinewai Reserve, 43°48.59'S, 173°01.28'E, 11 October 2002, CJV, AMNH (ARAMR-000616)	JX174282
<i>Periegops suterii</i>	Ps3	♀	MC, Hay Reserve, 43°42.18'S, 172°53.87'E, 13 March 2003, CJV, LUNZ (00012720)	JX174283
<i>Periegops suterii</i>	Ps4	♂	MC, Kennedys Bush Reserve, 43°37.9'S, 172°37.3'E, 22 April 2011, CJV, H.P. & L.J. Hudson Vink, LUNZ (00012715)	JX017357, JX017353
<i>Periegops suterii</i>	Ps5	♂	MC, Montgomery Park Reserve, 43°44.7'S, 172°52.2'E, 5 November 2010, CJV & JM-O, LUNZ (00012722)	JX174284
<i>Periegops suterii</i>	Ps6	♂	MC, Hinewai Reserve, 43°48.69'S, 173°00.95'E, 11 January 2011, M.H. Bowie, LUNZ (00012735)	JX174285
<i>Periegops suterii</i>	Ps7	subadult ♀	MC, Montgomery Park Reserve, 43°44.7'S, 172°52.2'E, 5 November 2010, JM-O & CJV, LUNZ (00012725)	JX174286, JX174291
<i>Periegops suterii</i>	Ps8	♀	MC, Kennedys Bush Reserve, 43°37.9'S, 172°37.3'E, 13 August 2011, CJV & H.P. Hudson Vink, LUNZ (00012718)	JX174287
<i>Periegops keani</i>	Pk1	immature	CL, Ruamahuanui Island, 36°57.25'S, 176°05.52'E, 19 November 2003, B.M. Fitzgerald, MONZ (AS.002310)	JX174288, JX174292
<i>Periegops keani</i>	Pk2	♂	CL, Ruamahuanui Island, 36°57.25'S, 176°05.52'E, 23 March 2012, CJV & J.M. Kean, LUNZ (00012716)	JX174289, JX174293
<i>Periegops keani</i>	Pk3	immature	CL, Ruamahuanui Island, 36°57.25'S, 176°05.52'E, 23 March 2012, CJV & J.M. Kean, LUNZ (00012717)	JX174290

Table 2. Uncorrected distance matrix for cytochrome c oxidase subunit 1 (COI).

	Pk1	Pk2	Pk3	Ps1	Ps2	Ps3	Ps4	Ps5	Ps6	Ps7
Pk2	0.006									
Pk3	0.007	0.007								
Ps1	0.135	0.131	0.135							
Ps2	0.135	0.131	0.135	0.000						
Ps3	0.139	0.135	0.139	0.007	0.007					
Ps4	0.141	0.138	0.138	0.084	0.084	0.086				
Ps5	0.132	0.130	0.132	0.012	0.012	0.013	0.082			
Ps6	0.135	0.131	0.135	0.000	0.000	0.007	0.084	0.012		
Ps7	0.132	0.130	0.132	0.012	0.012	0.013	0.082	0.000	0.012	
Ps8	0.141	0.138	0.138	0.084	0.084	0.086	0.000	0.082	0.084	0.082

PHYLOGENETIC ANALYSIS

Methods. Sequences were aligned using Sequencher 4.6. There was no evidence of insertions/deletions or stop codons in the COI sequences and alignment was straightforward.

Uncorrected COI pairwise distances were calculated using PAUP* version 4.0b10 (Swofford 2002). Distances using the Kimura-2-Parameter (K2P) model (Kimura 1980) were also calculated for comparison with previously reported intraspecific distances in spiders (Robinson *et al.* 2009). Although K2P has been commonly used in most DNA barcoding studies (e.g., Robinson *et al.* 2009), there is no evidence that this model is better for species identification than simpler metrics such as uncorrected pairwise distances (Astrin *et al.* 2006; Collins *et al.* 2012).


Partitioned Bayesian analysis implemented in MrBayes version 3.1.2 (Ronquist & Huelsenbeck 2003) was used to estimate the COI phylogenetic tree topology. MrModeltest version 2.3 (Nylander 2008) implemented in PAUP* version 4.0b10 (Swofford 2002) was used to select the optimal model and model parameters. Within MrModeltest the Akaike Information Criterion was used for model selection (Posada & Buckley 2004). Based on the results of Brandley *et al.* (2005), the COI data were partitioned by codon with models selected for each codon; GTR+I (Lanave *et al.* 1984; Tavaré 1986) for the 1st and 3rd codon positions and HKY+I (Hasegawa *et al.* 1985) for the 2nd codon positions. Bayesian analysis was conducted by running two simultaneous, completely independent analyses each with four heated chains, sampling every 1000th tree. The analysis was run for 20 million generations, by which time the average standard

deviation of split frequencies had dropped below 0.002, which indicated that the two tree samples had converged. Tracer version 1.5 (Rambaut & Drummond 2009) was also used to determine if the analyses had sufficient effective sample sizes. MrBayes was used to construct majority rule consensus trees, discarding the first 25% of trees generated as burn-in. TreeView 1.6.6 (Page 1996) was used to view and save trees in graphic format.

Results. A 1031 base pair (bp) COI fragment was sequenced from eight specimens of *P. suterii* and three specimens of *P. keani*. Four COI haplotypes occurred among the eight specimens of *P. suterii*. Each of the three specimens of *P. keani* had a different COI haplotype. Inter- and intraspecific uncorrected pairwise distances between COI sequences of *P. suterii* and *P. keani* are shown in Table 2. The minimum divergence between the two species was 13.0% (uncorrected) and the mean divergence was 13.5% (uncorrected). The mean intraspecific divergences in *P. suterii* and *P. keani*, were 4.0% (uncorrected) and 0.6% (uncorrected), respectively. The maximum intraspecific divergence in *P. suterii* was 8.6% (uncorrected, 9.2% K2P-distance) and was 0.7% (uncorrected and K2P-distance) in *P. keani*.

The 1360 bp fragments of 28S from two specimens of *P. suterii* were identical. The amplification of 28S from two specimens of *P. keani* (Pk1 and Pk2) was only partially successful with an 821 bp fragment from Pk1 and an 833 bp fragment from Pk2; however, the 281 bp overlap was identical. There were 12 nucleotides that varied between the two species.

The phylogenetic analysis of the COI data (Text-fig. 1) showed that *P. suterii* and *P. keani* are monophyletic with *P. suterii* divided into two clades.



Text-fig. 1 Unrooted Bayesian consensus tree based on cytochrome c oxidase subunit I (COI) sequence data. Values on branches are posterior probabilities. Specimen codes are listed in Table 1. Branch lengths are proportional to the expected number of substitutions per site (see scale bar).

Discussion. *Periegops* COI sequences were very similar or identical (Table 2) for specimens collected at the same sites; however, specimens of *P. suterii* collected at Kennedys Bush Reserve, in the west of the Banks Peninsula (Map 1), were 8.2–8.6% (8.7–9.2% K2P-distance) divergent from *P. suterii* specimens collected from three localities in the eastern Bank Peninsula (Map 1). This high divergence was not found in the slower evolving nuclear marker; 28S sequences were identical from specimens from Kennedys Bush Reserve and Montgomery Park Reserve (Map 1). We also did not observe any morphological differences between Kennedys Bush Reserve specimens and specimens collected from other locations on the Banks Peninsula; therefore we are confident that all specimens are one species, *P. suterii*. A difference of 8.7–9.2% (K2P-distance) in COI sequence divergence is very high in spiders; much more than the average maximum intraspecific divergence of 3.16% (K2P-distance) observed in other spiders (Robinson *et al.* 2009). However, higher intraspecific divergences (>10%) have been observed in other haplogynes (Astrin *et al.* 2006; Binford *et al.* 2008), Hypochilidae (Hedin 2001), and Mygalomorphae (Bond *et al.* 2001), so perhaps COI divergence is higher in ancestral spiders than it is in the Entelegynae. Nevertheless, it is curious that high (8.2–8.6%) COI divergence was observed between specimens from localities only just over 23 km apart and at similar altitudes, while there was much less divergence (0.7–1.3%) between specimens from eastern Bank Peninsula localities, which were 5–15 km apart. Specimens from localities between Kennedys Bush Reserve and Montgomery Park Reserve, such as Kaituna Valley, could reveal if COI divergence is correlated with distance or whether there has been some sort of genetic isolation between the east and west of Banks Peninsula, similar to the geographic distribution of two closely related weta species in the genus *Hemideina* (Townsend *et al.* 1997).

KEY TO NEW ZEALAND PERIEGOPIDAE

- 1 Tibia of leg I less than $0.7\times$ length of carapace. When viewed retrolaterally, dorsal surface of cymbium of male pedipalp projects dorsally (Fig. 9). Flared flattened flanges near embolus tip (Fig. 6, 7). Found only on Banks Peninsula and Riccarton Bush (Map 1). ... (p. 15)... *Periegops suterii* (Urquhart)
- Tibia of leg I more than $0.75\times$ length of carapace. When viewed retrolaterally, dorsal surface of cymbium of male only slightly projects dorsally (Fig. 9). Flattened flanges near embolus tip not flared (Fig. 9a). Found only on Aldermen Islands and East Cape (Map 2). ... (p. 17)... *Periegops keani* sp. nov.

BIOSYSTEMATICS

Family PERIEGOPIDAE

Medium-sized, 3-clawed haplogyne spiders. 6 eyes in 3 widely separated diads (Fig. 1, 14, 15). Proclaw of legs I and II with double row of teeth, retroclaw with single row. Chelicerae with lamina on the ventral surface (Fig. 2) and slender maxillary lobes directed across the labium (Fig. 3). Male pedipalp simple, consisting of a bulb with a short embolus (Fig. 6–9). Slightly sclerotised area anterior to the epigastric furrow (Fig. 10, 11) and simple internal genitalia (Fig. 12, 13).

Genus *Periegops* Simon

Periegops Simon, 1893: 267. —Forster 1995: 93–94. Type species *Periegops hirsutus* Simon, 1893.

Description. Body length 5.9–10.0 mm. Carapace longer than wide, flat, without fovea, orange or orange-brown, darker anteriorly. 6 eyes in 3 widely separated diads; anterior median eyes absent. Chelicerae with lamina on ventral surface and membranous promarginal lobe with 5–7 setae (Fig. 2); 3 promarginal teeth and 1 reduced retromarginal tooth; fang length less than width of paturon; venom outlet anterior on fang (Labarque & Ramírez 2012; fig. 18A). Lateral surface of female chelicerae with stridulatory ridges (Labarque & Ramírez 2012; fig. 16A), with associated palpal femoral thorns (Labarque & Ramírez 2012; fig. 11A). Maxillae slender and more than $2\times$ as long as wide, directed across the labium (Fig. 3). Legs in descending order of length usually I, IV, II, III, but legs I and IV sometimes subequal in females. Male tibiae

of leg I with a row of strong setae on retrolateral surface (Fig. 4, 5). Proclaw of legs I and II with double row of teeth, retroclaw with single row. Both superior claws of legs II and IV a single row. Inferior claw of all tarsi with a single tooth. Male pedipalp simple with a bulb and a short embolus (Fig. 6–9); coiled sperm duct often visible within the bulb. Abdomen ovoid, dun or dirty cream usually with a distinct chevron pattern (Fig. 1, 14–16) and black setae. Slightly sclerotised area anterior to the epigastric furrow (Fig. 10, 11). Female internal genitalia simple, haplogyne, consisting of a rounded poreplate with many small invaginated cups (Fig. 12, 13). 6 spinnerets and small colulus; median surface of posterior median spinnerets with field of spicules (Labarque & Ramírez 2012; fig. 21F, 22E). Forster (1995) provided details of the respiratory system of *Periegops*.

Periegops suterii (Urquhart)

Fig. 1, 4, 6, 7, 10, 12, 14, 16; Map 1
Segestria suterii Urquhart, 1892: 230.

Periegops hirsutus Simon, 1893: 268, figs 223–224. —Synonymy by Chamberlain 1946: 88.

Periegops suteri (Urquhart) —Bryant 1935a: 53. —Bryant 1935b: 81, figs 4, 13, 25. —Chamberlain 1946: 88. —Forster 1967: 72, figs 81, 134. —Forster & Forster 1973: 151, figs 87b, 90. —Jocqué & Dippenaar-Schoeman 2006: 202, figs 78a–e. —Labarque & Ramírez 2012: 1, figs 1, 2A, 3A–B, 6A–B, 9A, 10A, 11A, 12A, 13A, 14A, 15A, 16A, 17A, 20A, 21, 22, 29A–B, 31A.

Periegops suterii (Urquhart) —Forster 1995: 94, figs 1–3, 6–14, 23, 24, 27, 31. —Forster & Forster 1999: 143, figs 1.14i, 10.9. —Paquin, Vink & Dupérré 2010: 37, figs 14.1–4.

Diagnosis. Distinguished from all other *Periegops* species by the shape of the embolus and cymbium of the male pedipalp. Flattened flanges near embolus tip more pronounced than those of *P. keani*, and embolus is shorter than that of *P. australia*. Dorsal surface of cymbium projects slightly more than that of *P. keani*, but much less than that of *P. australia*. Poreplate of internal genitalia not as rounded as that of *P. australia*. Leg I of males, females, and immatures shorter than in *P. keani*; tibia I less than $0.7\times$ length of carapace. Found only on Banks Peninsula and Riccarton Bush in the South Island.

Description. Carapace: red-orange anteriorly, orange posteriorly; with sparse black setae; fovea absent. Sternum: orange. Leg I orange-brown, patella, distal ends of femur and tibia light orange; legs II, III, and IV yellow-brown, darker at proximal end of each leg segment. Abdomen: dun with black setae; black-brown chevron pattern (Fig. 1, 14, 16). Chelicerae red-brown. Male pedipalp simple with a bulb and a short embolus, which is flattened and

twisted, flaring out to 2 flanges just before the tip (Fig. 6, 7); coiled sperm duct often visible within the bulb. Slightly sclerotised area anterior to epigastric furrow (Fig. 10). Female internal genitalia simple, haplogyne, consisting of a rounded poreplate with many small invaginated cups (Fig. 12).

Dimensions (mm). Male MC, Kennedys Bush Reserve LUNZ 00012715 (female MC, Kennedys Bush Reserve LUNZ 00012718): total length 6.13 (7.92); carapace 3.06/2.05 (3.82/2.60), height 1.15 (1.71); abdomen 2.87/1.80 (4.12/2.60); sternum 1.64/1.05 (1.93/1.25).

	I	II	III	IV	Palp
Femur	2.43 (2.72)	2.13 (2.41)	1.84 (2.1)	2.35 (2.56)	0.65 (0.95)
Patella	0.85 (0.98)	0.8 (0.98)	0.73 (0.9)	0.8 (0.97)	0.27 (0.38)
Tibia	2.08 (2.15)	1.75 (1.93)	1.45 (1.65)	2.03 (2.3)	0.6 (0.63)
Metatarsus	2.16 (2.17)	1.82 (1.97)	1.62 (1.78)	2.07 (2.28)	
Tarsus	0.88 (0.85)	0.76 (0.83)	0.68 (0.73)	0.68 (0.73)	0.22 (0.78)
Total	8.4 (8.87)	7.26 (8.12)	6.32 (7.16)	7.93 (8.84)	1.74 (2.74)

Variation. Size range, mean. Male body length 5.9–7.4, 6.6, $n = 12$. Female body length 6.8–9.3, 8.1, $n = 11$. Male carapace length 3.0–3.7, 3.4, $n = 12$. Female carapace length 3.4–4.8, 3.9, $n = 11$.

The ratios of the length of tibia I to the length of the carapace (measured for all specimens, including immatures) were between 0.54 and 0.70. The ratio of leg I segment length to carapace length has also been shown to be a reliable character for the separation of other closely related spider species (Vink *et al.* 2008).

The chevron pattern on the abdomen (Fig. 1, 14, 16) was visible in all but one of the specimens examined, a male (LUNZ 00012715). Forster (1995) noted that a female collected at Riccarton Bush did not have abdominal markings.

Type data. Type of *Segestria suterii*: Not examined. Original description based on a female specimen from “Dyer’s Pass, Canterbury”, which was collected with 2 immature males by H. Suter (Urquhart 1892). Forster (1995) stated the type was in poor condition in the Canterbury Museum, Christchurch, New Zealand, where all of Urquhart’s existing type specimens are housed (Court & Forster 1988; Nicholls *et al.* 2000; Paquin *et al.* 2008). However, the type of *Segestria suterii* was not included in the list of arachnid

primary types held in Canterbury Museum (Nicholls *et al.* 2000). It was not possible to check whether the type specimen was present at Canterbury Museum, as the collections have not been accessible since the Christchurch earthquake in February 2011 (S. D. Pollard, pers. comm.). The type is not in OMNZ (C. Fraser, pers. comm.), which houses New Zealand’s largest collection of spiders. It is possible that the type was amongst specimens at Ray Forster’s house at the time of his death and has as yet not been located. We consider the type of *Segestria suterii* missing, but not lost.

Type of *Periegops hirsutus*: Not examined. Original description based on a female specimen from an undisclosed location in New Zealand (Simon 1893). Although many of Simon’s type specimens are housed in the Muséum national d’Histoire naturelle, Paris, France, the type of *Periegops hirsutus* could not be found there (C. Rollard, pers. comm.).

Material examined. 32 non-type specimens (15 males, 15 females, 2 subadult females) — see Appendix B for collection details of specimens examined.

Distribution (Map 1). Found only on Banks Peninsula and Riccarton Bush (MC). Forster (1995) and Vink (2006) listed localities where *P. suterii* specimens have been found. Riccarton Bush is the only site outside of the Banks Peninsula where *P. suterii* has been found (in 1994), but subsequent searches there (Chinn 2006; Vink 2006) have failed to find it.

Biology. *Periegops suterii* has been found in forest with a deep leaf litter layer and well-drained soil. It has been found under logs and rocks, and in leaf litter, in both beech and podocarp forest. Within suitable locations, *P. suterii* has a patchy distribution, but specimens are often found years apart in the same specific locations. *Periegops suterii* does not appear to build a web for prey capture, but it does produce dragline silk and builds silken retreats. The lack of webs and their presence in pitfall traps suggest that it is a cursorial, night-time hunter. On two occasions a single female has been found together with two or three males under logs and rocks (Forster 1995; Vink 2006), which implies that the female might possess some method of attracting males. Forster (1995) suggested pheromones, but it could also be stridulation; Forster (1995) did not notice the stridulatory ridges on the chelicerae. Adults have been found from August to May.

DNA. Nine cytochrome *c* oxidase subunit 1 and two 28S rRNA sequences for this species are listed in GenBank (www.ncbi.nlm.nih.gov/GenBank/) under accession numbers EF537066, JX017353, JX017357, JX174281–JX174287, and JX174291.

Remarks. Although the types of *Segestria suterii* and *Periegops hirsutus* were not examined, we have no reason to doubt that these species are synonymous, as Chamberlain (1946) examined the type specimens of both species. Forster (1995: 92) stated that it can “be reliably assumed” that the type of *P. hirsutus* “came from the same locality as Urquhart’s *Segestria suterii*.” Also, Simon (1893) noted that the type specimen of *P. hirsutus* was 8 mm long, which is within the range of female body lengths we observed for *P. suterii*.

Despite the common usage of the name *Periegops suteri*, the correct original spelling of the specific epithet is *suterii* (as *Segestria suterii* Urquhart, 1892). Article 33.4 in the fourth edition of the *International Code of Zoological Nomenclature* (International Commission on Zoological Nomenclature 1999) states the change from *-i* to *-ii* is an incorrect subsequent spelling. There is provision in the *International Code of Zoological Nomenclature* for prevailing usage of an unjustified emendation (article 33.2.3.1) or incorrect subsequent spelling (article 33.3.1); however, *suteri* is not an emendation but an incorrect subsequent spelling and article 33.4 is an exception to articles 33.2.3.1 and 33.3.1. Both spellings have been used; *P. suterii* by Forster (1995), Forster & Forster (1999), Paquin *et al.* (2010), and Sirvid *et al.* (2011, 2012), and *P. suteri* by Bryant (1935a, b), Chamberlain (1946), Forster (1967), Forster & Forster (1973), Vink (2006), Jocqué & Dippenaar-Schoeman (2006), Brocknerhoff *et al.* (2008), Platnick (2012), and Labarque & Ramírez (2012). As this is not a case of prevailing usage of an incorrect subsequent spelling (article 33.3.1) and article 33.4 is quite clear on the matter, we advocate for and use the correct original spelling.

Periegops keani new species

Fig. 2, 3, 5, 9, 15; Map 2

Diagnosis. Distinguished from all other *Periegops* species by shape of embolus and cymbium of the male pedipalp. Flattened flanges near embolus tip not as pronounced as those of *P. suterii* and embolus shorter than that of *P. australis*. Dorsal surface of cymbium projects only slightly, less than that of *P. suterii* and much less than that of *P. australis*. Leg I of male and immatures longer than in *P. suterii*; tibia I more than 0.75× length of carapace. Found only on the Aldermen Islands and East Cape in the North Island.

Description. Carapace: dark orange-brown anteriorly, orange-brown posteriorly; with sparse black setae; fovea absent. Sternum: light orange brown. Leg I dark

orange-brown, patella, distal ends of femur, and tibia orange-brown; legs II, III, and IV orange-brown, darker at proximal end of each leg segment. Abdomen: dirty cream with black setae; chevron pattern barely visible (Fig. 15). Chelicerae red-brown. Male pedipalp simple with a bulb and a short embolus, which is flattened and twisted, flaring out slightly to 2 flanges just before the tip (Fig. 9); coiled sperm duct visible within the bulb. Female unknown.

Dimensions (mm). Male holotype: total length 10.03; carapace 4.99/3.48, height 1.66; abdomen 5.25/3.75; sternum 2.40/1.90.

	I	II	III	IV	Palp
Femur	4.99	4.26	3.73	4.41	1.01
Patella	1.41	1.08	1.21	1.26	0.45
Tibia	4.39	3.83	2.97	4.13	0.96
Metatarsus	4.44	3.88	3.23	3.98	
Tarsus	1.31	1.26	1.01	1.26*	0.45
Total	16.54	14.31	12.15	15.04	2.87

*Both tarsi IV were damaged so the length was estimated based on the proportions observed in immature species of *P. keani*.

Variation. The holotype male was the only specimen that did not have abdominal markings; all of the other subadult and immature specimens had a chevron pattern on the abdomen, like that of *P. suterii* (Fig. 1, 14, 16).

The ratios of the length of tibia I to the length of the carapace (measured for all specimens, including immatures) were between 0.77 and 0.88. Based on what we observed for *P. suterii*, it seems likely that female *P. keani* will have a tibia/carapace ratio within that range.

Type data. Holotype: male (LUNZ 00012716) labelled “NEW ZEALAND, CL, Ruamahuanui Island; 36°57.25'S, 176°05.52'E; NW campsite, under rock; 23 Mar 2012, C.J. Vink & J.M. Kean” (LUNZ).

Material examined. Type specimen plus 7 non-type specimens (1 subadult male, 1 subadult female, 5 immatures) — see Appendix B for collection details of specimens examined.

Distribution (Map 2). Found only on the Aldermen Islands and East Cape (CL, GB).

Biology. *Periegops keani* has been found in forest with a deep leaf litter layer and well-drained soil. On the forest floor it has been found under rocks and in a *Stanwellia* sp. (Araneae: Nemesiidae) burrow (Forster 1995). *Periegops keani* does not appear to build a web for prey capture and

probably builds silken retreats, like *P. suterii*. The lack of webs suggests that it is a cursorial, night-time hunter. Adults have been found in March and September, but given that the climates it is found in are milder than those of *P. suterii*, it seems likely that adults occur throughout the year.

DNA. Three cytochrome *c* oxidase subunit 1 and two 28S rRNA sequences for this species are listed in GenBank (www.ncbi.nlm.nih.gov/GenBank/) under accession numbers JX174288–JX174290, JX174292, and JX174293.

Etymology. The specific name is in honour of the senior author's friend and colleague, John Kean, who helped collect the type specimen.

Remarks. The bulb of the holotype was not oriented to the same angle with respect to the cymbium (Fig. 9) as the specimens illustrated of *P. suterii* (Fig. 6, 7) and *P. australia* (Fig. 8); however, when oriented in a slight dorsal direction (Fig. 9a), it looks the same as those of *P. suterii*, other than the flattened flanges near the embolus tip, which are not as pronounced in *P. keani*.

We consider that a female specimen found in forest below the East Cape Lighthouse is *P. keani*. Forster (1995) remarked that the genitalia of this specimen was indistinguishable from that of *P. suterii*. The female specimen from East Cape was collected by Grace Hall (NZAC) on 30 September 1993 and was loaned to the late Ray Forster, but has not been returned to NZAC (G. Hall pers. comm.) and it has not been located at OMNZ (C. Fraser, pers. comm.). Unsuccessful searches for further specimens have been conducted in the small (~10 hectare) patch of forest below the East Cape Lighthouse in October 1994 (G. Hall), February 1995 (G. Hall), September 1995 (G. Hall & P. J. Sirvid), November 1995 (G. Hall & L. J. Boutin), December 2009 (G. Hall), November 2010 (J. Malumbres-Olarte) and February 2012 (C. J. Vink). Searches of the forest floor (including rock and log turning) were conducted in all trips, *Stanwellia* sp. burrows were searched in most trips, and pitfall trapping and litter searches were also conducted in the former and later trips. *Periegops keani* may conceivably now be extinct from that site.

It is possible that the population at East Cape could be a separate species to *P. keani*, as *Periegops* probably has limited dispersal ability. Although the Aldermen Islands and East Cape are 233 km apart, the Aldermen Islands would have been connected to the mainland from the Last Glacial Maximum (McKinnon 1997; Trewick & Bland 2012) up until 12000 years ago (Schofield & Thompson 1964; McKinnon 1997) and there would have been continuous forest between the two sites (McKinnon 1997; Trewick & Bland 2012). If a fresh specimen could

be located at East Cape, 28S sequence data could confirm whether they were the same species.

Forster & Forster (1999) stated that they had examined a female specimen from the Aldermen Islands. This specimen was collected from Ruamahua Island in November 1972 and loaned to Ray Forster from NZAC (G. Hall, pers. comm.). We have examined this specimen and determined it to be a subadult male, as it has swollen palpal tibiae, which are characteristic of males in their final moult before adulthood. The specimen did not have an epigyne or internal genitalia and we believe Forster & Forster (1999) made a typographical error in stating the specimen was female. We consider that this specimen is *P. keani*. Ruamahua Island is only 2 km from Ruamahuanui Island and the land below the two islands is no more than 20 m under the sea (Hayward & Moore 1973), therefore, the two populations would have been connected up until 9000 years ago (Schofield & Thompson 1964; McKinnon 1997).

Size may help differentiate *P. keani* from *P. suterii* even though we have only examined a single adult specimen of the former species. The subadult male from Ruamahua Island is larger (body length = 8.8 mm, carapace length = 4.3 mm) than any of the 11 adult male *P. suterii* that we have examined. Also, a *P. keani* specimen (MONZ AS.002301) that appeared to be one or two moults from adulthood was larger (body length = 9.3, carapace length = 3.9) than five of the 11 adult specimens of *P. suterii* examined; based on this and the size of the male, it would be reasonable to expect an adult female *P. keani* to be larger than any *P. suterii* female.

REFERENCES

- Altschul, S. F.; Madden, T. L.; Schäffer, A. A.; Zhang, J.; Zhang, Z.; Miller, W.; Lipman, D. J. 1997: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25: 3389–3402.
- Amedo, M. A.; Gasparo, F.; Opatova, V. 2009: Systematics and phylogeography of the *Dysdera erythrina* species complex (Araneae, Dysderidae) in Sardinia. *ZooKeys* 16: 319–345.
- ; Oromí, P.; Ribera, C. 2001: Radiation of the spider genus *Dysdera* (Araneae, Dysderidae) in the Canary Islands: cladistic assessment based on multiple data sets. *Cladistics* 17: 313–353.
- Astrin, J. J.; Huber, B. A.; Misof, B.; Klütsch, C. F. C. 2006: Molecular taxonomy in pholcid spiders (Pholcidae, Araneae): evaluation of species identification methods using CO1 and 16S rRNA. *Zoologica Scripta* 35: 441–457.
- Berland, L. 1932: Les arachnides (scorpions, araignées, etc.): biologie systematique. Pp. 1–485 in *Encyclopédie entomologique (A)*, Vol. XVI. Paris, Paul Lechavalier and fils.
- Binford, G. J.; Callahan, M. S.; Bodner, M. R.; Rynerson, M. R.; Nuñez, P. B.; Ellison, C. E.; Duncan, R. P. 2008: Phylogenetic relationships of *Loxosceles* and *Sicarius* spiders are consistent with Western Gondwanan vicariance. *Molecular Phylogenetics and Evolution* 49: 538–553.
- Bond, J. E.; Hedin, M. C.; Ramirez, M. G.; Opell, B. D. 2001: Deep molecular divergence in the absence of morphological and ecological change in the Californian coastal dune endemic trapdoor spider *Aptostichus simus*. *Molecular Ecology* 10: 899–910.
- Bowie, M. H.; Frampton, C. M. 2004: A practical technique for non-destructive monitoring of soil surface invertebrates for ecological restoration programmes. *Ecological Management & Restoration* 5: 34–42.
- Brandley, M. C.; Schmitz, A.; Reeder, T. W. 2005: Partitioned Bayesian analyses, partition choice, and the phylogenetic relationships of scincid lizards. *Systematic Biology* 54: 373–390.
- Brignoli, P. M. 1978: Some remarks on the relationships between the Haplogynae, the semi-Entelegynae and the Cribellatae (Araneae). *Symposia of the Zoological Society of London* 42: 285–292.
- Bristowe, W. S. 1938: The classification of spiders. *Proceedings of the Zoological Society of London* 108: 285–322.
- Brockerhoff, E. G.; Pawson, S. M.; Dungan, R. J.; Sarfati, M. S.; Bowie, M. H.; Wratten, S. D. 2008: Terrestrial invertebrates. Pp. 493–548 in M. J. Winterbourn, G. A. Knox, C. J. Burrows and I. D. Marsden (eds), *The Natural History of Canterbury*. Christchurch, Canterbury University Press.
- Bruvo-Madaric, B.; Huber, B. A.; Steinacher, A.; Pass, G. 2005: Phylogeny of pholcid spiders (Araneae: Pholcidae): Combined analysis using morphology and molecules. *Molecular Phylogenetics and Evolution* 37: 661–673.
- Bryant, E. B. 1935a: Notes on some of Urquhart's species of spiders. *Records of the Canterbury Museum* 4: 53–70.
- 1935b: Some new and little known species of New Zealand spiders. *Records of the Canterbury Museum* 4: 71–94.
- Caporiacco, L. di 1938: Il sistema degli Araneidi. *Archivio Zoologico Italiano* 25 (Suppl. 4): 35–155.
- Chamberlain, G. 1946: Revision of Araneae of New Zealand. Part 2. *Records of the Auckland Institute and Museum* 3: 85–97.
- Chinn, W. G. 2006: The Invertebrates of Riccarton Bush / Putaringamotu. A qualitative and quantitative survey. Unpublished report for the Riccarton Bush Trust. 1–28 pp.
- Coddington, J. A. 2005: Phylogeny and classification of spiders. Pp. 18–24 in D. Ubick, P. Paquin, P. E. Cushing and V. D. Roth (eds), *Spiders of North America: An Identification Manual*. Keene, American Arachnological Society.
- ; Levi, H. W. 1991: Systematics and evolution of spiders (Araneae). *Annual Review of Ecology and Systematics* 22: 565–592.
- Collins, R. A.; Boykin, L. M.; Cruickshank, R. H.; Armstrong, K. F. 2012: Barcoding's next top model: an evaluation of nucleotide substitution models for specimen identification. *Methods in Ecology and Evolution* 3: 457–465.
- Court, D. J.; Forster, R. R. 1988: The spiders of New Zealand. Part VI. Araneidae-Araneinae. *Otago Museum Bulletin* 6: 68–124.
- Crosby, T. K.; Dugdale, J. S.; Watt, J. C. 1976: Recording specimen localities in New Zealand: an arbitrary system of areas and codes defined. *New Zealand Journal of Zoology* 3: 69.
- ; ——; —— 1998: Area codes for recording specimen localities in the New Zealand subregion. *New Zealand Journal of Zoology* 25: 175–183.

- Dimitrov, D.; Arnedo, M. A.; Ribera, C. 2008: Colonization and diversification of the spider genus *Pholcus* Walckenaer, 1805 (Araneae, Pholcidae) in the Macaronesian archipelagos: Evidence for long-term occupancy yet rapid recent speciation. *Molecular Phylogenetics and Evolution* 48: 596–614.
- Duncan, R. P.; Rynerson, M. R.; Ribera, C.; Binford, G. J. 2010: Diversity of *Loxosceles* spiders in Northwestern Africa and molecular support for cryptic species in the *Loxosceles rufescens* lineage. *Molecular Phylogenetics and Evolution* 55: 234–248.
- Folmer, O.; Black, M.; Hoeh, W.; Lutz, R.; Vrijenhoek, R. 1994: DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.
- Forster, R. R. 1967: The spiders of New Zealand. Part I. *Otago Museum Bulletin* 1: 1–124.
- 1995: The Australasian spider family Periegopidae Simon, 1893 (Araneae: Sicarioidea). *Records of the Western Australian Museum Supplement* 52: 91–105.
- ; Forster, L. M. 1973: New Zealand Spiders: an Introduction. Auckland, Collins. 254 pp.
- ; ——— 1999: Spiders of New Zealand and their Worldwide Kin. Dunedin, Otago University Press. 270 pp.
- Framenau, V. W.; Dupérré, N.; Blackledge, T. A.; Vink, C. J. 2010: Systematics of the new Australasian orb-weaving spider genus *Backbourkia* (Araneae: Araneidae: Araneinae). *Arthropod Systematics & Phylogeny* 68: 79–111.
- Hasegawa, M.; Kishino, K.; Yano, T. 1985: Dating the human-ape splitting by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution* 22: 160–174.
- Hayward, B. W.; Moore, P. R. 1973: Geology of the Aldermen Islands. *Tane* 19: 69–85.
- Hedin, M. C. 2001: Molecular insights into species phylogeny, biogeography, and morphological stasis in the ancient spider genus *Hypochilus* (Araneae: Hypochilidae). *Molecular Phylogenetics and Evolution* 18: 238–251.
- ; Bond, J. E. 2006: Molecular phylogenetics of the spider infraorder Mygalomorphae using nuclear rRNA genes (18S and 28S): Conflict and agreement with the current system of classification. *Molecular Phylogenetics and Evolution* 41: 454–471.
- ; Maddison, W. P. 2001: A combined molecular approach to phylogeny of the jumping spider subfamily Dendryphantinae (Araneae: Salticidae). *Molecular Phylogenetics and Evolution* 18: 386–403.
- Huber, B. A.; Astrin, J. J. 2009: Increased sampling blurs morphological and molecular species limits: revision of the Hispaniolan endemic spider genus *Tainonia* (Araneae: Pholcidae). *Invertebrate Systematics* 23: 281–300.
- International Commission on Zoological Nomenclature 1999: International Code of Zoological Nomenclature. London, The International Trust for Zoological Nomenclature. 306 pp.
- Jocqué, R.; Dippenaar-Schoeman, A. S. 2006: Spider Families of the World. Tervuren, Musée Royal de l’Afrique Centrale. 336 pp.
- Kimura, M. 1980: A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16: 111–120.
- Labarque, F. M.; Ramírez, M. J. 2007a: A new species of *Drymusa* Simon, 1891 (Araneae: Drymusidae) from Chile. *Studies on Neotropical Fauna and Environment* 42: 241–246.
- ; ——— 2007b Description of the female of *Drymusa serrana* Goloboff & Ramírez, 1991 (Araneae: Drymusidae) with notes on its biology. *Zootaxa* 1580: 27–33.
- ; ——— 2012: The placement of the spider genus *Periegops* and the phylogeny of Scytodoidea (Araneae: Araneomorphae). *Zootaxa* 3312: 1–44.
- Lanave, C.; Preparata, G.; Sacone, C.; Serio, G. 1984: A new method for calculating evolutionary substitution rates. *Journal of Molecular Evolution* 20: 86–93.
- Lattimore, V. L.; Vink, C. J.; Paterson, A. M.; Cruickshank, R. H. 2011: Unidirectional introgression within the genus *Dolomedes* (Araneae: Pisauridae) in southern New Zealand. *Invertebrate Systematics* 25: 70–79.
- Lehtinen, P. T. 1986: Evolution of the Scytodoidea. Pp. 149–159 in W. G. Eberhard, Y. D. Lubin and B. C. Robinson (eds), Proceedings of the Ninth International Congress of Arachnology, Panama, 1983. Washington, D.C., Smithsonian Institution Press.
- Malumbres-Olarte, J.; Vink, C. J. 2012: Redescription of *Clubiona blesti* Forster, 1979 (Araneae: Clubionidae) with a preliminary molecular phylogeny of New Zealand *Clubiona*. *Zootaxa* 3277: 27–42.

- ; ——; Ross, J. G.; Cruickshank, R. H.; Paterson, A. M. in press: The role of habitat complexity on spider communities in native alpine grasslands of New Zealand. *Insect Conservation and Diversity*.
- McKinnon, M. (ed.) 1997: Bateman New Zealand Historical Atlas. Auckland, David Bateman. 290 pp.
- Nicholls, D. C.; Sirvid, P. J.; Pollard, S. D.; Walker, M. 2000: A list of arachnid primary types held in Canterbury Museum. *Records of the Canterbury Museum 14*: 37–48.
- Nunn, G. B.; Theisen, B. F.; Christensen, B.; Arctander, P. 1996: Simplicity-correlated size growth of the nuclear 28S ribosomal RNA D3 expansion segment in the crustacean order isopoda. *Journal of Molecular Evolution 42*: 211–223.
- Nylander, J. A. A. 2008: MrModeltest 2.3. Uppsala, Department of Systematic Zoology, Uppsala University. Available from <http://www.abc.se/~nylander/>
- Page, R. D. M. 1996: TREEVIEW: An application to display phylogenetic trees on personal computers. *Computer Applications in the Biological Sciences 12*: 357–358.
- Paquin, P.; Vink, C. J. 2009: Testing compatibility between molecular and morphological techniques for arthropod systematics: a minimally destructive DNA extraction method that preserves morphological integrity, and the effect of lactic acid on DNA quality. *Journal of Insect Conservation 13*: 453–457.
- ; ——; Dupérré, N. 2010: Spiders of New Zealand: Annotated Family Key & Species List. Lincoln, Manaaki Whenua Press. vii +118 pp.
- ; ——; ——; Sirvid, P. J.; Court, D. J. 2008: *Nomina dubia* and faunistic issues with New Zealand spiders (Araneae). *Insecta Mundi 46*: 1–6.
- Platnick, N. I. 2012: The World Spider Catalog, version 13.0. The American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/INTRO1.html>
- ; Coddington, J. A.; Forster, R. R.; Griswold, C. E. 1991: Spinneret morphology and the phylogeny of haplogyne spiders (Araneae, Araneomorphae). *American Museum Novitates 3016*: 1–73.
- Posada, D.; Buckley, T. R. 2004: Model selection and model averaging in phylogenetics: advantages of Akaike Information Criterion and Bayesian approaches over likelihood ratio tests. *Systematic Biology 53*: 793–808.
- Rambaut, A.; Drummond, A. J. 2009: Tracer v1.5. Available from <http://beast.bio.ed.ac.uk/Tracer>.
- Ramírez, M. J. 2000: Respiratory system morphology and the phylogeny of haplogyne spiders (Araneae, Araneomorphae). *Journal of Arachnology 28*: 149–157.
- Rix, M. G.; Harvey, M. S.; Roberts, J. D. 2008: Molecular phylogenetics of the spider family Micropholcommatidae (Arachnida: Araneae) using nuclear rRNA genes (18S and 28S). *Molecular Phylogenetics and Evolution 46*: 1031–1048.
- Robinson, E. A.; Blagoev, G. A.; Hebert, P. D. N.; Adamowicz, S. J. 2009: Prospects for using DNA barcoding to identify spiders in species-rich genera. *ZooKeys 16*: 27–46.
- Ronquist, F.; Huelsenbeck, J. P. 2003: MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics 19*: 1572–1574.
- Schofield, J. C.; Thompson, H. R. 1964: Postglacial sea levels and isostatic uplift. *New Zealand Journal of Geology and Geophysics 7*: 359–370.
- Simon, E. 1893: Histoire naturelle des Araignées, seconde édition. Tome 1, Fascicule 2. Paris, Librairie encyclopédique de Roret. 257–488 pp.
- Sirvid, P. J.; Zhang, Z.-Q.; Harvey, M. S.; Rhode, B. E.; Cook, D. R.; Bartsch, I.; Staples, D. A. 2011: Chelicerata: horseshoe crabs, arachnids, sea spiders. Pp. 50–89 in D. P. Gordon (ed.), New Zealand Inventory of Biodiversity. Volume Two: Kingdom Animalia. Chaetognatha, Ecdysozoa, Ichnofossils. Christchurch, Canterbury University Press.
- ; Vink, C. J.; Wakelin, M. D.; Fitzgerald, B. M.; Hitchmough, R. A.; Stringer, I. A. N. 2012: The conservation status of New Zealand Araneae. *New Zealand Entomologist 35*: 85–90.
- Spagna, J. C.; Crews, S. C.; Gillespie, R. G. 2010: Patterns of habitat affinity and Austral/Holarctic parallelism in dictynoid spiders (Araneae: Entelegynae). *Invertebrate Systematics 24*: 238–257.
- Starrett, J.; Waters, E. R. 2007: Positive natural selection has driven the evolution of the Hsp70s in *Diguettia* spiders. *Biology Letters 3*: 439–444.
- Swofford, D. L. 2002: PAUP*: Phylogenetic Analysis Using Parsimony (*and Other Methods), Version 4.0b10. Sunderland, Massachusetts, Sinauer Associates.
- Tavaré, S. 1986: Some probabilistic and statistical problems in the analysis of DNA sequences. *Lectures on Mathematics in the Life Sciences 17*: 57–86.

- Townsend, J. A.; Brown, B.; Stringer, I. A. N.; Potter, M. A. 1997: Distribution, habitat and conservation status of *Hemideina ricta* and *H. femorata* on Banks Peninsula, New Zealand. *New Zealand Journal of Ecology* 21: 43–49.
- Trewick, S. A.; Bland, K. J. 2012: Fire and slice: palaeogeography for biogeography at New Zealand's North Island/South Island juncture. *Journal of the Royal Society of New Zealand* 42: 153–183.
- Urquhart, A. T. 1892: Descriptions of new species of Araneae. *Transactions of the New Zealand Institute* 24: 230–253.
- Vink, C. J. 2006: The spider *Periegops suteri* (Araneae: Periegopidae): description, ecology, localities and management recommendations. *Weta* 31: 34–39.
- ; Dupérré, N. 2010: Pisauridae (Arachnida: Araneae). *Fauna of New Zealand* 64: 1–60.
- ; ———; McQuillan, B. N. 2011a: The black headed jumping spider, *Trite planiceps* Simon, 1899 (Araneae: Salticidae): redescription including COI and paralogous 28S sequences. *New Zealand Journal of Zoology* 38: 317–331.
- ; Fitzgerald, B. M.; Sirvid, P. J.; Dupérré, N. 2011b: Reuniting males and females: redescriptions of *Nuisiana arboris* (Marples 1959) and *Cambridgea reinga* Forster & Wilton 1973 (Araneae: Desidae, Stiphidiidae). *Zootaxa* 2739: 41–50.
- ; Paterson, A. M. 2003: Combined molecular and morphological phylogenetic analyses of the New Zealand wolf spider genus *Anoteropsis* (Araneae: Lycosidae). *Molecular Phylogenetics and Evolution* 28: 576–587.
- ; Sirvid, P. J.; Malumbres-Olarte, J.; Griffiths, J. W.; Paquin, P.; Paterson, A. M. 2008: Species status and conservation issues of New Zealand's endemic *Latrodectus* spider species (Araneae : Theridiidae). *Invertebrate Systematics* 22: 589–604.
- ; Thomas, S. M.; Paquin, P.; Hayashi, C. Y.; Hedin, M. 2005: The effects of preservatives and temperatures on arachnid DNA. *Invertebrate Systematics* 19: 99–104.
- Wang, Q.; Li, S.; Wang, R.; Paquin, P. 2008: Phylogeographic analysis of Pimoidae (Arachnida: Araneae) inferred from mitochondrial cytochrome *c* oxidase subunit I and nuclear 28S rRNA gene regions. *Journal of Zoological Systematics and Evolutionary Research* 46: 96–104.
- Wheeler, Q. D.; Platnick, N. I. 2000: The phylogenetic species concept (*sensu* Wheeler and Platnick). Pp. 55–69 in Q. D. Wheeler and R. Meier (eds), *Species Concepts and Phylogenetic Theory: A Debate*. New York, Columbia University Press.

APPENDIX A: Glossary of technical terms

abdomen — posterior division of the spider body; sometimes referred to as the opisthosoma

bulb — refers to the male pedipalpal organ as a whole

carapace — the hard dorsal covering of the cephalothorax

cephalothorax — anterior division of the spider body; sometimes referred to as the prosoma

chelicerae — first pair of appendages of the cephalothorax, consisting of two segments (the distal segment is called the fang, basal segment is called the paturon)

coxa — first or basal segment of the legs

cymbium — tarsus of the adult male pedipalp

distal — near the apex

dorsal — upper (surface)

epigastric furrow — a transverse groove across the anterior ventral part of the abdomen

epigyne — the sclerotised region of female spiders covering the internal genitalia and located between the book lungs and anterior of the epigastric furrow

embolus — the intromittent structure of the bulb containing the terminal portion of the ejaculatory duct

femur — third segment of the legs and pedipalps

fovea — depression on the thoracic region of the carapace where muscles of sucking stomach are attached internally

haplogyne — the primitive form of spider genitalia where the female has the copulatory openings internally, within the gonopore, and typically lacks a sclerotised epigyne, and the male has relatively simple pedipalps. This form is found in the basal spiders, including Mygalomorphae, Gradungulidae and Haplogynae.

maxilla — (= endite) the expanded lobe of the palpal coxa situated laterally of the labium (plural **maxillae**)

metatarsus — sixth segment of the legs; absent in the pedipalps

paturon — the basal segment of a chelicerae

pedipalp — six-segmented second appendage of the cephalothorax, anterior to legs I

patella — fourth segment of the legs and pedipalp

promarginal — anterior margin

proximal — near the base

retrolateral — on the outer side i.e., the surface nearer to the posterior end of the body

retromarginal — posterior margin

sclerotised — hardened by sclerotin or other substances in the cuticle

seta — a sclerotised hair-like projection arising from the cuticle (plural **setae**)

sternum — plate on the ventral surface of the cephalothorax between the coxae of the legs

tarsus — last segment of the legs and pedipalp

tibia — fifth segment of legs and pedipalp

trochanter — second segment of the leg and pedipalp

ventral — lower (surface)

APPENDIX B: Collection details of specimens examined. Localities (including coordinates) and dates collected, collectors, and institutions of specimens examined.

Periegops suterii

MC. 3 ♂, 1 ♀, Riccarton Bush, 43°31.7'S, 172°35.7'E, 9 Apr 1994, A. D. Blest, OMNZ IV35925; 1 ♂, Kennedys Bush Reserve, 43°37.9'S, 172°37.3'E, 22 Apr 2011, C. J. Vink, H. P. & L. J. Hudson Vink, LUNZ 00012715; 1 ♀, Kennedys Bush Reserve, 43°37.9'S, 172°37.3'E, 13 Aug 2011, C. J. Vink & H. P. Hudson Vink, LUNZ 00012718; 1 ♀, Kennedys Bush Reserve, 43°37.9'S, 172°37.3'E, 13 Aug 2011, C. J. Vink & H. P. Hudson Vink, LUNZ 00012719; 1 ♂, Rhodes Bush, 43°40'S, 172°37'E, 13 Nov 1915, G. Archey, OMNZ IV35927; 1 ♀, Hay Reserve, 43°42.18'S, 172°53.87'E, 13 Mar 2003, C. J. Vink, LUNZ 00012720; 2 ♀, Kaituna Valley, 43°43'S, 172°45'E, 11 Sep 1949, R. R. Forster, OMNZ IV35923; 1 ♂, Montgomery Park Reserve, 43°44.7'S, 172°52.2'E, 5 Nov 2010, C. J. Vink & J. Malumbres-Olarte, LUNZ 00012721; 1 ♂, Montgomery Park Reserve, 43°44.7'S, 172°52.2'E, 5 Nov 2010, C. J. Vink & J. Malumbres-Olarte, LUNZ 00012722; 1 ♂, Montgomery Park Reserve, 43°44.7'S, 172°52.2'E, 5 Nov 2010, C. J. Vink & J. Malumbres-Olarte, LUNZ 00012723; 1 ♂, Montgomery Park Reserve, 43°44.7'S, 172°52.2'E, 5 Nov 2010, J. Malumbres-Olarte & C. J. Vink, LUNZ 00012724; 1 ♀, 1 subadult ♀, Montgomery Park Reserve, 43°44.7'S, 172°52.2'E, 5 Nov 2010, J. Malumbres-Olarte & C. J. Vink, LUNZ 00012725; 1 ♀, Panama Rock, 43°44.77'S, 173°02.49'E, 12 Jan 2011, M. H. Bowie, LUNZ 00012726; 1 ♀, Panama Rock, 43°44.77'S, 173°02.49'E, 12 Jan 2011, M. H. Bowie, LUNZ 00012727; 1 ♂, Otepatotu Reserve, 43°44.90'S, 173°00.95'E, 4 Jan 2011, M. H. Bowie, LUNZ 00012728; 1 ♂, Otepatotu Reserve, 43°44.90'S, 173°00.95'E, 12 Jan 2011, M. H. Bowie, LUNZ 00012729; 1 ♀, Otepatotu Reserve, 43°44.90'S, 173°00.95'E, 4 Jan 2011, M. H. Bowie, LUNZ 00012730; 1 ♀, Otepatotu Reserve, 43°44.93'S, 173°00.95'E, 12 Jan 2011, M. H. Bowie, LUNZ 00012731; 1 ♂, Tititipounamu, 43°45.47'S, 173°01.44'E, 4 Jan 2011, M. H. Bowie, LUNZ 00012732; 1 ♀, Little River, 43°46'S, 172°47'E, 10 Jan 1985, A. C. Harris, OMNZ IV35924; 1 ♂, Ellangowan Reserve, 43°47.89'S, 173°02.08'E, 12 Jan 2011, M. H. Bowie, LUNZ 00012733; 1 ♂, Hinewai Reserve, 43°48.59'S, 173°01.28'E, 11 Oct 2002, C. J. Vink, AMNH ARAMR-000615; 1 ♀, Hinewai Re-

serve, 43°48.59'S, 173°01.28'E, 11 Oct 2002, C. J. Vink, AMNH ARAMR-000616; 1 ♀, Hinewai Reserve, 43°48.6'S, 173°01.3'E, 27 Aug 1996, C. J. Vink, LUNZ 00012734; 1 ♂, Hinewai Reserve, 43°48.69'S, 173°00.95'E, 11 Jan 2011, M. H. Bowie, LUNZ 00012735; 1 ♀, Akaroa, 43°48.8'S, 172°57.5'E, 16 Oct 1920, G. Archey, OMNZ IV35928; 1 subadult ♀, Fishermans Bay, 43°49.54'S, 173°03.85'E, 21 Dec 2010, M. H. Bowie, LUNZ 00012736.

Periegops keani

CL. 1 immature, Ruamahuanui Island, 36°57.3'S, 176°05.5'E, 7 Nov 2002, M. D. Wakeelin, MONZ AS.002302; 1 immature, Ruamahuanui Island, 36°57.25'S, 176°05.52'E, 19 Nov 2003, B. M. Fitzgerald, MONZ AS.002310; 1 immature, Ruamahuanui Island, 36°57.25'S, 176°05.52'E, 19 Nov 2003, B. M. Fitzgerald, MONZ AS.002300; holotype ♂, Ruamahuanui Island, 36°57.25'S, 176°05.52'E, 23 Mar 2012, C. J. Vink & J. M. Kean, LUNZ 00012716; 1 immature, Ruamahuanui Island, 36°57.25'S, 176°05.52'E, 23 Mar 2012, C. J. Vink & J. M. Kean, LUNZ 00012717; 1 subadult ♀, 1 immature, Ruamahuanui Island, 36°57.25'S, 176°05.52'E, 18 Apr 2005, B. M. Fitzgerald & P. MacDonald, MONZ AS.002301; 1 subadult ♂, Ruamahuaui Island, 36°58.5'S, 176°05'E, 8-12 November 1972, G. W. Ramsay, NZAC.

ILLUSTRATIONS






Fig. 1 Schematic dorsal view, *Periegops suterii*, female (LUNZ 00012720). Actual size on the right.




(2) Anterior view of chelicerae




(3) Ventral view of mouthparts

Fig. 2, 3 *Periegops keani* (holotype): (2) anterior view of chelicerae; (3) ventral view of mouthparts.




(4) *Periegops suterii*




(5) *Periegops keani* (holotype)


Fig. 4, 5 Retrolateral view of leg I: (4) *Periegops suterii* (LUNZ 00012722); (5) *Periegops keani* (holotype).




(6) *Periegops suterii*
Kennedys Bush Reserve



(7) *Periegops suterii*
Montgomery Park Reserve




(8) *Periegops australia*
Mt Goonaneman




(9) *Periegops keani*
Ruamahuanui Island (holotype)

Fig. 6–9 Retrolateral view of left male pedipalps (b – bulb; c – cymbium; e – embolus), arrow indicates dorsal projection on the cymbium: (6) *Periegops suterii*, Kennedys Bush Reserve (LUNZ 00012715); (7) *Periegops suterii*, Montgomery Park Reserve (LUNZ 00012721); (8) *Periegops australia*, Mt Goonaneman (QMB S20421); (9) *Periegops keani*, Ruamahuanui Island (holotype) (9a = laterodorsal view of bulb and cymbium).




(10) *Periegops suterii*
Hay Reserve




(11) *Periegops australia*
Mt Goonaneman

Fig. 10, 11 Ventral view of female epigastric area: (10) *Periegops suterii*, Hay Reserve (LUNZ 00012720); (11) *Periegops australia*, Mt Goonaneman (QMB S20419).



(12) *Periegops suterii*
Hay Reserve



(13) *Periegops australia*
Mt Goonaneman

Fig. 12, 13 Dorsal view of female internal genitalia (p – poreplate; ic – invaginated cup): (12) *Periegops suterii*, Hay Reserve (LUNZ 00012720); (13) *Periegops australia*, Mt Goonaneman (QMB S20419).




Fig. 14 Dorsal view of *Periegops suterii*, male (LUNZ 00012721). Actual size on the right.






Fig. 15 Dorsal view of *Periegops keani*, male (holotype). Actual size on the right.




Fig. 16 Photograph of *Periegops suterii*, female (LUNZ 00012718). Photographer: Bryce McQuillan.

DISTRIBUTION MAPS



Map 1 Collection localities, *Periegops suterii*. Inset shows localities on and near Banks Peninsula, and labelled locations indicate collection sites for DNA specimens.



Map 2 Collection localities, *Periegops keani*.

TAXONOMIC INDEX

Page numbers in **bold** type denote a description, and in italic type illustrations. A suffixed letter 'k' indicates a key, and 'm' a map.

australia, *Periegops* 8, 10, 15, 17, 18, 28, 29, 30

Caponiidae 9

Cephalobidae 11

Diguetidae 9

Drymusa 9

Drymusidae 9

Dysderidae 8

Entelegynae 14

Haplogynae 9

Hemideina 14

hirsutus, *Periegops* 8, 15, 16, 17

Hypochoilidae 14

keani*, *Periegops 7, 11, 12, 13, 15k, **17–18**, 24, 26, 27, 28, 32, 34m

Leptonetidae 9

Nemesiidae 17

Ochyroceratidae 9

Periegopidae this contribution

Periegops this contribution

Periegops australia 8, 10, 15, 17, 18, 28, 29, 30

Periegops hirsutus 8, 15, 16, 17

Periegops keani 7, 11, 12, 13, 15k, **17–18**, 24, 26, 27, 28, 32, 34m

Periegops suteri 15, 17

Periegops suterii 5, 7, 8, 9, 10, 11, 12, 13, 14, 15k, **15–17**, 18, 24, 25, 27, 28, 29, 30, 31, 33, 34m

Pholcidae 9

Plectreuridae 9

Segestria 8

Segestria suterii 15, 16, 17

Scytodidae 9

Scytodoidea 9

Sicariidae 9

Sicarioidea 9

Stanwellia sp. 17, 18


suteri, *Periegops* 15, 17





suterii*, *Periegops 5, 7, 8, 9, 10, 11, 12, 13, 14, 15k, **15–17**, 18, 24, 25, 27, 28, 29, 30, 31, 33, 34m

suterii, *Segestria* 15, 16, 17

Telemidae 9

Tetrablemmidae 9





Area codes and boundaries used to categorise specimen locality data (after Crosby *et al.* 1976)

Base-map for plotting collection localities; this may be photocopied without copyright release

FAUNA OF NEW ZEALAND PUBLICATIONS

- 1 **Terebrantia** (Insecta: Thysanoptera). *Laurence A. Mound & Annette K. Walker*. ISBN 0-477-06687-9, 23 December 1982, 120 pp. \$29.95
- 2 **Osoriinae** (Insecta: Coleoptera: Staphylinidae). *H. Pauline McColl*. ISBN 0-477-06688-7, 23 December 1982, 96 pp. \$18.60
- 3 **Anthribidae** (Insecta: Coleoptera). *B. A. Holloway*. ISBN 0-477-06703-4, 23 December 1982, 272 pp. \$41.00
- 4 **Eriophyoidea except Eriophyinae** (Arachnida: Acari). *D. C. M. Manson*. ISBN 0-477-06745-X, 12 November 1984, 144 pp. \$29.95
- 5 **Eriophyinae** (Arachnida: Acari: Eriophyoidea). *D. C. M. Manson*. ISBN 0-477-06746-8, 14 November 1984, 128 pp. \$29.95
- 6 **Hydraenidae** (Insecta: Coleoptera). *R. G. Ordish*. ISBN 0-477-06747-6, 12 November 1984, 64 pp. \$18.60
- 7 **Cryptostigmata** (Arachnida: Acari)—a concise review. *M. Luxton*. ISBN 0-477-06762-X, 8 December 1985, 112 pp. \$29.95
- 8 **Calliphoridae** (Insecta: Diptera). *James P. Dear*. ISBN 0-477-06764-6, 24 February 1986, 88 pp. \$18.60
- 9 **Protura** (Insecta). *S. L. Tuxen*. ISBN 0-477-06765-4, 24 February 1986, 52 pp. \$18.60
- 10 **Tubulifera** (Insecta: Thysanoptera). *Laurence A. Mound & Annette K. Walker*. ISBN 0-477-06784-0, 22 September 1986, 144 pp. \$34.65
- 11 **Pseudococcidae** (Insecta: Hemiptera). *J. M. Cox*. ISBN 0-477-06791-3, 7 April 1987, 232 pp. \$49.95
- 12 **Pompilidae** (Insecta: Hymenoptera). *A. C. Harris*. ISBN 0-477-02501-3, 13 November 1987, 160 pp. \$39.95
- 13 **Encyrtidae** (Insecta: Hymenoptera). *J. S. Noyes*. ISBN 0-477-02517-X, 9 May 1988, 192 pp. \$44.95
- 14 **Lepidoptera**—annotated catalogue, and keys to family-group taxa. *J. S. Dugdale*. ISBN 0-477-02518-8, 23 September 1988, 264 pp. \$49.95
- 15 **Ambositrinae** (Insecta: Hymenoptera: Diapriidae). *I. D. Naumann*. ISBN 0-477-02535-8, 30 December 1988, 168 pp. \$39.95
- 16 **Nepticulidae** (Insecta: Lepidoptera). *Hans Donner & Christopher Wilkinson*. ISBN 0-477-02538-2, 28 April 1989, 92 pp. \$22.95
- 17 **Mymaridae** (Insecta: Hymenoptera)—introduction, and review of genera. *J. S. Noyes & E. W. Valentine*. ISBN 0-477-02542-0, 28 April 1989, 100 pp. \$24.95
- 18 **Chalcidoidea** (Insecta: Hymenoptera)—introduction, and review of genera in smaller families. *J. S. Noyes & E. W. Valentine*. ISBN 0-477-02545-5, 2 August 1989, 96 pp. \$24.95
- 19 **Mantodea** (Insecta), with a review of aspects of functional morphology and biology. *G. W. Ramsay*. ISBN 0-477-02581-1, 13 June 1990, 96 pp. \$24.95
- 20 **Bibionidae** (Insecta: Diptera). *Roy A. Harrison*. ISBN 0-477-02595-1, 13 November 1990, 28 pp. \$14.95
- 21 **Margarodidae** (Insecta: Hemiptera). *C. F. Morales*. ISBN 0-477-02607-9, 27 May 1991, 124 pp. \$34.95
- 22 **Notonemouridae** (Insecta: Plecoptera). *I. D. McLellan*. ISBN 0-477-02518-8, 27 May 1991, 64 pp. \$24.95
- 23 **Sciapodinae, Medeterinae** (Insecta: Diptera) with a generic review of the Dolichopodidae. *D. J. Bickel*. ISBN 0-477-02627-3, 13 January 1992, 74 pp. \$27.95
- 24 **Therevidae** (Insecta: Diptera). *L. Lyneborg*. ISBN 0-477-02632-X, 4 March 1992, 140 pp. \$34.95
- 25 **Cercopidae** (Insecta: Homoptera). *K. G. A. Hamilton & C. F. Morales*. ISBN 0-477-02636-2, 25 May 1992, 40 pp. \$17.95
- 26 **Tenebrionidae** (Insecta: Coleoptera): catalogue of types and keys to taxa. *J. C. Watt*. ISBN 0-477-02639-7, 13 July 1992, 70 pp. \$27.95
- 27 **Antarctoperlinae** (Insecta: Plecoptera). *I. D. McLellan*. ISBN 0-477-01644-8, 18 February 1993, 70 pp. \$27.95
- 28 **Larvae of Curculionoidea** (Insecta: Coleoptera): a systematic overview. *Brenda M. May*. ISBN 0-478-04505-0, 14 June 1993, 226 pp. \$55.00
- 29 **Cryptorhynchinae** (Insecta: Coleoptera: Curculionidae). *C. H. C. Lyal*. ISBN 0-478-04518-2, 2 December 1993, 308 pp. \$65.00
- 30 **Hepialidae** (Insecta: Lepidoptera). *J. S. Dugdale*. ISBN 0-478-04524-7, 1 March 1994, 164 pp. \$42.50
- 31 **Talitridae** (Crustacea: Amphipoda). *K. W. Duncan*. ISBN 0-478-04533-6, 7 October 1994, 128 pp. \$36.00
- 32 **Sphecidae** (Insecta: Hymenoptera). *A. C. Harris*. ISBN 0-478-04534-4, 7 October 1994, 112 pp. \$33.50
- 33 **Moranilini** (Insecta: Hymenoptera). *J. A. Berry*. ISBN 0-478-04538-7, 8 May 1995, 82 pp. \$29.95
- 34 **Anthicidae** (Insecta: Coleoptera). *F. G. Werner & D. S. Chandler*. ISBN 0-478-04547-6, 21 June 1995, 64 pp. \$26.50
- 35 **Cydnidae, Acanthosomatidae, and Pentatomidae** (Insecta: Heteroptera): systematics, geographical distribution, and bioecology. *M.-C. Larivière*. ISBN 0-478-09301-2, 23 November 1995, 112 pp. \$42.50
- 36 **Leptophebiidae** (Insecta: Ephemeroptera). *D. R. Towns & W. L. Peters*. ISBN 0-478-09303-9, 19 August 1996, 144 pp. \$39.50
- 37 **Coleoptera**: family-group review and keys to identification. *J. Klimaszewski & J. C. Watt*. ISBN 0-478-09312-8, 13 August 1997, 199 pp. \$49.50
- 38 **Naturalised terrestrial Stylommatophora** (Mollusca: Gastropoda). *G. M. Barker*. ISBN 0-478-09322-5, 25 January 1999, 253 pp. \$72.50
- 39 **Molytini** (Insecta: Coleoptera: Curculionidae: Molytinae). *R. C. Craw*. ISBN 0-478-09325-X, 4 February 1999, 68 pp. \$29.50
- 40 **Cixiidae** (Insecta: Hemiptera: Auchenorrhyncha). *M.-C. Larivière*. ISBN 0-478-09334-9, 12 November 1999, 93 pp. \$37.50

- 41 Coccidae** (Insecta: Hemiptera: Coccoidea). C. J. Hodgson & R. C. Henderson. ISBN 0-478-09335-7, 23 February 2000, 264 pp. \$72.50
- 42 Aphodiinae** (Insecta: Coleoptera: Scarabaeidae). Z. T. Stebnicka. ISBN 0-478-09341-1, 15 June 2001, 64 pp. \$29.50
- 43 Carabidae** (Insecta: Coleoptera): catalogue. A. Laroche & M.-C. Larivière. ISBN 0-478-09342-X, 15 June 2001, 285 pp. \$72.50
- 44 Lycosidae** (Arachnida: Araneae). C. J. Vink. ISBN 0-478-09347-0, 23 December 2002, 94 pp. \$37.50
- 45 Nemonychidae, Belidae, Brentidae** (Insecta: Coleoptera: Curculionoidea). G. Kuschel. ISBN 0-478-09348-9, 28 April 2003, 100 pp. \$40.00
- 46 Nesameletidae** (Insecta: Ephemeroptera). Terry R. Hitchings & Arnold H. Staniczek. ISBN 0-478-09349-7, 14 May 2003, 72 pp. \$32.50
- 47 Erotylidae** (Insecta: Coleoptera: Cucujoidea): phylogeny and review. R. A. B. Leschen. ISBN 0-478-09350-0, 5 June 2003, 108 pp. \$42.50
- 48 Scaphidiinae** (Insecta: Coleoptera: Staphylinidae). I. Löbl & R. A. B. Leschen. ISBN 0-478-09353-5, 18 November 2003, 94 pp. \$37.50
- 49 Lithinini** (Insecta: Lepidoptera: Geometridae: Ennominae). J. D. Weintraub & M. J. Scoble. ISBN 0-478-09357-8, 29 April 2004, 48 pp. \$24.50
- 50 Heteroptera** (Insecta: Hemiptera): catalogue. M.-C. Larivière & A. Laroche. ISBN 0-478-09358-6, 14 May 2004, 330 pp. \$89.00
- 51 Coccidae** (Insecta: Hemiptera: Coccoidea): adult males, pupae and prepupae of indigenous species. C. J. Hodgson & R. C. Henderson. ISBN 0-478-09360-8, 22 June 2004, 228 pp. \$65.00
- 52 Raphignathoidea** (Acari: Prostigmata). Qing-Hai Fan & Zhi-Qiang Zhang. ISBN 0-478-09371-3, 20 May 2005, 400 pp. \$89.00
- 53 Harpalini** (Insecta: Coleoptera: Carabidae: Harpalinae). A. Laroche & M.-C. Larivière. ISBN 0-478-09369-1, 4 July 2005, 160 pp. \$55.00
- 54 Hierodoris** (Insecta: Lepidoptera: Gelechoidea: Oecophoridae), and overview of Oecophoridae. Robert J. B. Hoare. ISBN 0-478-09378-0, 24 December 2005, 100 pp. \$40.00
- 55 Criconematina** (Nematoda: Tylenchida). W. M. Wouts. ISBN 0-478-09381-0, 24 March 2006, 232 pp. ... \$65.00
- 56 Tyrophagus** (Acari: Astigmata: Acaridae). Qing-Hai Fan & Zhi-Qiang Zhang. ISBN 978-0-478-09386-5, 4 June 2007, 291 pp. \$80.00
- 57 Apoidea** (Insecta: Hymenoptera). B. J. Donovan. ISBN 978-0-478-09389-6, 7 September 2007, 295 pp. \$89.00
- 58 Alysiinae** (Insecta: Hymenoptera: Braconidae). J. A. Berry. ISBN 978-0-478-09390-2, 7 September 2007, 95 pp. \$45.00
- 59 Erotylinae** (Insecta: Coleoptera: Cucujoidea: Erotylidae): taxonomy and biogeography. Paul E. Skelley & Richard A. B. Leschen. ISBN 978-0-478-09391-9, 7 September 2007, 59 pp. \$30.00
- 60 Carabidae** (Insecta: Coleoptera): synopsis of supraspecific taxa. A. Laroche & M.-C. Larivière. ISBN 978-0-478-09394-0, 21 November 2007, 188 pp. \$54.00
- 61 Lucanidae** (Insecta: Coleoptera). B. A. Holloway. ISBN 978-0-478-09395-7, 21 November 2007, 254 pp. \$75.00
- 62 Trechini** (Insecta: Coleoptera: Carabidae: Trechinae). J. I. Townsend. ISBN 978-0-478-34717-9 (print), 978-0-478-34716-6 (online), 16 June 2010, 101 pp. ... \$49.50
- 63 Auchenorrhyncha** (Insecta: Hemiptera): catalogue. M.-C. Larivière, M. J. Fletcher & A. Laroche. ISBN 978-0-478-34720-3 (print), 978-0-478-34721-0 (online), 16 June 2010, 232 pp. \$75.00
- 64 Pisauridae** (Arachnida: Araneae). C. J. Vink & N. Dupérré. ISBN 978-0-478-34722-7 (print), 978-0-478-34723-4 (online), 13 July 2010, 60 pp. \$37.50
- 65 Izatha** (Insecta: Lepidoptera: Gelechoidea: Oecophoridae). Robert J. B. Hoare. ISBN 978-0-478-34724-1 (print), 978-0-478-34725-8 (online), 2 September 2010, 201 pp. \$75.00
- 66 Diaspididae** (Insecta: Hemiptera: Coccoidea). R. C. Henderson. ISBN 978-0-478-34726-5 (print), 978-0-478-34727-2 (online), 23 May 2011, 275 pp. \$89.00
- 67 Peloridiidae** (Insecta: Hemiptera: Coleorrhyncha). M.-C. Larivière, D. Burckhardt & A. Laroche. ISBN 978-0-478-34730-2 (print), 978-0-478-34731-9 (online), 14 November 2011, 78 pp. \$48.00
- 68 Simuliidae** (Insecta: Diptera). Douglas A. Craig, Ruth E. G. Craig & Trevor K. Crosby. ISBN 978-0-478-34734-0 (print), 978-0-478-34735-7 (online), 29 June 2012, 336 pp. \$95.00
- 69 Carabidae** (Insecta: Coleoptera): synopsis of species, Cicindelinae to Trechinae (in part). A. Laroche & M.-C. Larivière. ISBN 978-0-478-34738-8 (print), 978-0-478-34739-5 (online), 7 March 2013, 193 pp. \$75.00
- 70 Periegopidae** (Arachnida: Araneae). C. J. Vink, N. Dupérré & J. Malumbres-Olarte. ISBN 978-0-478-34740-1 (print), 978-0-478-34741-8 (online), 7 March 2013, 41 pp. \$29.00

Visit the Manaaki Whenua Press Website at:

<http://www.mwpress.co.nz/>

for further information.

To access on-line extracts and PDFs from this series

visit: <http://fnz.landcareresearch.co.nz/>

**Taxonomic groups covered in the
Fauna of New Zealand series**

Insecta

Coleoptera

- Family-group review and keys to identification (*J. Klimaszewski & J. C. Watt*, FNZ 37, 1997)
- Anthribidae (*B. A. Holloway*, FNZ 3, 1982)
- Anthicidae (*F. G. Werner & D. S. Chandler*, FNZ 34, 1995)
- Carabidae: catalogue (*A. Laroche & M.-C. Larivière*, FNZ 43, 2001); synopsis of supraspecific taxa (*A. Laroche & M.-C. Larivière*, FNZ 60, 2007); synopsis of species, Cicindelinae to Trechinae (in part) (*A. Laroche & M.-C. Larivière*, FNZ 69, 2013)
- Carabidae: Harpalinae: Harpalini (*A. Laroche & M.-C. Larivière*, FNZ 53, 2005)
- Carabidae: Trechinae: Trechini (*J. I. Townsend*, FNZ 62, 2010)
- Curculionidae: Cryptorhynchinae (*C. H. C. Lyal*, FNZ 29, 1993)
- Curculionidae: Molytinae: Molytini (*R. C. Craw*, FNZ 39, 1999)
- Curculionoidea: Nemonychidae, Belidae, Brentidae (*G. Kuschel*, FNZ 45, 2003)
- Curculionoidea larvae: a systematic overview (*Brenda M. May*, FNZ 28, 1993)
- Erotylidae: phylogeny and review (*Richard A. B. Leschen*, FNZ 47, 2003); Erotylinae: taxonomy and biogeography (*Paul E. Skelley & Richard A. B. Leschen*, FNZ 59, 2007)
- Hydraenidae (*R. G. Ordish*, FNZ 6, 1984)
- Lucanidae (*B. A. Holloway*, FNZ 61, 2007)
- Scarabaeidae: Aphodiinae (*Z. T. Stebnicka*, FNZ 42, 2001)
- Staphylinidae: Osoriinae (*H. Pauline McColl*, FNZ 2, 1982)
- Staphylinidae: Scaphidiinae (*I. Löbl & Richard A. B. Leschen*, FNZ 48, 2003)
- Tenebrionidae: catalogue of types and keys to taxa (*J. C. Watt*, FNZ 26, 1992)

Diptera

- Bibionidae (*Roy A. Harrison*, FNZ 20, 1990)
- Calliphoridae (*James P. Dear*, FNZ 8, 1986)
- Dolichopodidae: Sciapodinae, Medeterinae with a generic review (*D. J. Bickel*, FNZ 23, 1992)
- Simuliidae (*Douglas A. Craig, Ruth E. G. Craig, Trevor K. Crosby*, FNZ 68, 2012)
- Therevidae (*L. Lyneborg*, FNZ 24, 1992)

Ephemeroptera

- Leptophlebiidae (*D. R. Towns & W. L. Peters*, FNZ 36, 1996)
- Nesameletidae (*Terry R. Hitchings & Arnold H. Staniczek*, FNZ 46, 2003)

Hemiptera

- Auchenorrhyncha: catalogue (*M.-C. Larivière, M. J. Fletcher & A. Laroche*, FNZ 63, 2010)
- Cercopidae (*K. G. A. Hamilton & C. F. Morales*, FNZ 25, 1992)
- Cixiidae (*M.-C. Larivière*, FNZ 40, 1999)
- Coccidae (*C. J. Hodgson & R. C. Henderson*, FNZ 41, 2000); adult males, pupae and prepupae of indigenous species (*C. J. Hodgson & R. C. Henderson*, FNZ 51, 2004)
- Cydnidae, Acanthosomatidae, and Pentatomidae (*M.-C. Larivière*, FNZ 35, 1995)
- Diaspididae (*R. C. Henderson*, FNZ 66, 2011)
- Heteroptera: catalogue (*M.-C. Larivière & A. Laroche*, FNZ 50, 2004)
- Margarodidae (*C. F. Morales*, FNZ 21, 1991)

- Pseudococcidae (*J. M. Cox*, FNZ 11, 1987)
- Peloriidae (*M.-C. Larivière, D. Burckhardt & A. Laroche*, FNZ 67, 2011).

Hymenoptera

- Apoidea (*B. J. Donovan*, FNZ 57, 2007)
- Braconidae: Alysini (*J. A. Berry*, FNZ 58, 2007)
- Chalcidoidea: introduction, and review of smaller families (*J. S. Noyes & E. W. Valentine*, FNZ 18, 1989)
- Diapriidae: Ambositrinae (*I. D. Naumann*, FNZ 15, 1988)
- Encyrtidae (*J. S. Noyes*, FNZ 13, 1988)
- Mymaridae (*J. S. Noyes & E. W. Valentine*, FNZ 17, 1989)
- Pompilidae (*A. C. Harris*, FNZ 12, 1987)
- Pteromalidae: Eunotinae: Moranilini (*J. A. Berry*, FNZ 33, 1995)
- Sphecidae (*A. C. Harris*, FNZ 32, 1994)

Lepidoptera

- Annotated catalogue, and keys to family-group taxa (*J. S. Dugdale*, FNZ 14, 1988)
- Geometridae: Ennominae: Lithinini (*Jason D. Weintraub & Malcolm J. Scoble*, FNZ 49, 2004)
- Hepialidae (*J. S. Dugdale*, FNZ 30, 1994)
- Nepticulidae (*Hans Donner & Christopher Wilkinson*, FNZ 16, 1989)
- Oecophoridae: *Hierodoris* (*Robert J. B. Hoare*, FNZ 54, 2005); *Izatha* (*Robert J. B. Hoare*, FNZ 65, 2010).

- Mantodea**, with a review of aspects of functional morphology and biology (*G. W. Ramsay*, FNZ 19, 1990)

Plecoptera

- Antarctoperlinae (*I. D. McLellan*, FNZ 27, 1993)
- Notonemouridae (*I. D. McLellan*, FNZ 22, 1991)

- Protura** (*S. L. Tuxen*, FNZ 9, 1986)

Thysanoptera

- Terebrantia (*Laurence A. Mound & Annette K. Walker*, FNZ 1, 1982)
- Tubulifera (*Laurence A. Mound & Annette K. Walker*, FNZ 10, 1986)

Arachnida

Acari

- Acaridae: *Tyrophagus* (*Qing-Hai Fan & Zhi-Qiang Zhang*, FNZ 56, 2007)
- Cryptostigmata—a concise review (*M. Luxton*, FNZ 7, 1985)
- Eriophyoidea except Eriophyinae (*D. C. M. Manson*, FNZ 4, 1984)
- Eriophyinae (*D. C. M. Manson*, FNZ 5, 1984)
- Raphignathoidea (*Qing-Hai Fan & Zhi-Qiang Zhang*, FNZ 52, 2005)

Araneae

- Lycosidae (*C. J. Vink*, FNZ 44, 2002)
- Periegopidae (*C. J. Vink, N. Dupérré & Malumbres-Olarte*, FNZ 70, 2013)
- Pisauridae (*C. J. Vink & N. Dupérré*, FNZ 64, 2010)

Crustacea

- Amphipoda**
- Talitridae (*K. W. Duncan*, FNZ 31, 1994)

Mollusca

Gastropoda

- Naturalised terrestrial Stylommatophora (*G. M. Barker*, FNZ 38, 1999)

Nematoda

- Tylenchida: Criconematina (*W. M. Wouts*, FNZ 55, 2006)

NOTICES

This series of refereed publications has been established to encourage those with expert knowledge to publish concise yet comprehensive accounts of elements in the New Zealand fauna. The series is professional in its conception and presentation, yet every effort is made to provide resources for identification and information that are accessible to the non-specialist.

Fauna of N.Z. deals with non-marine invertebrates only, since the vertebrates are well documented, and marine forms are covered by the series *NIWA Biodiversity Memoirs*.

Contributions are invited from any person with the requisite specialist skills and resources. Material from the N.Z. Arthropod Collection is available for study.

Contributors should discuss their intentions with a member of the Editorial Board or with the Series Editor before commencing work; all necessary guidance will be given.

Subscribers should address enquiries to *Fauna of N.Z.*, Manaaki Whenua Press, Landcare Research, P.O. Box 40, Lincoln 7640, New Zealand.

NGĀ PĀNUI

Kua whakatūria tēnei huinga pukapuka hei whakahauhau i ngā tohunga whai mātauranga kia whakaputa i ngā kōrero poto, engari he whaikiko tonu, e pā ana ki ngā aitanga pepeke o Aotearoa. He tōtika tonu te āhua o ngā tuhituhi, engari ko te tino whāinga, kia mārama te marea ki ngā tohu tautuhi o ia ngārara, o ia ngārara, me te roanga atu o ngā kōrero mō tēnā, mō tēnā.

He titiro whāiti tā tēnei pukapuka ki ngā mea noho whenua, kāore he tuarā; i pēnei ai i te mea kei te mōhio whānuitia ngā mea whai tuarā, ā, ko ngā mea noho moana, koirā te tino kaupapa o te huinga pukapuka *NIWA Biodiversity Memoirs*.

Ka āhei te tangata ki te **whakauru tuhituhinga** mehemea kei a ia ngā tohungatanga me ngā rauemi e tutuki pai ai tana mahi. Heoi anō, e wātea ana te Kohinga Angawaho o Aotearoa hei āta tiroiro mā te tangata mehemea he āwhina kei reira.

Me whāki te kaituhi i ōna whakaaro ki tētahi o te Kāhui Ārahi Whakarōpūtanga Tuarā-Kore, ki te Ētita rānei i mua i te tīmatanga, ā, mā rātou a ia e ārahi mō te wāhi ki tana tuhinga.

Ko te hunga pīrangi **hoko pukapuka**, me tuhi ki *Fauna of N.Z.*, Manaaki Whenua Press, Manaaki Whenua, Pouaka Poutāpeta 40, Lincoln 7640, Aotearoa.