

**Fauna of  
New Zealand**  
Ko te Aitanga Pepeke  
o Aotearoa

**EDITORIAL BOARD**

**Dr R. M. Emberson, L. c/-** *Department of Ecology, P.O. Box 84, Lincoln University, New Zealand*

**Dr M. J. Fletcher,** *NSW Agricultural Scientific Collections Unit, Forest Road, Orange, NSW 2800, Australia*

**Prof. G. Giribet,** *Museum of Comparative Zoology, Harvard University, 26 Oxford Street, Cambridge, MA 02138, U.S.A.*

**Dr R. J. B. Hoare,** *Landcare Research, Private Bag 92170, Auckland, New Zealand*

**Dr M.-C. Larivière,** *Landcare Research, Private Bag 92170, Auckland, New Zealand*

**Mr R. L. Palma,** *Museum of New Zealand Te Papa Tongarewa, P.O. Box 467, Wellington, New Zealand*

**Dr C. J. Vink,** *Canterbury Museum, Rolleston Ave, Christchurch, New Zealand*

**CHIEF EDITOR**

**Prof Z.-Q. Zhang,** *Landcare Research, Private Bag 92170, Auckland, New Zealand*

**Associate Editors**

**Dr T. R. Buckley, Dr R. J. B. Hoare, Dr M.-C. Larivière, Dr R. A. B. Leschen, Dr D. F. Ward, Dr Z. Q. Zhao,**  
*Landcare Research, Private Bag 92170, Auckland, New Zealand*

**Honorary Editor**

**Dr T. K. Crosby,** *Landcare Research, Private Bag 92170, Auckland, New Zealand*

---

**Fauna of New Zealand**  
**Ko te Aitanga Pepeke o Aotearoa**

**Number / Nama 76**

**Phthiraptera**

**(Insecta)**

**A catalogue of parasitic lice from New Zealand**

by  
**Ricardo L. Palma**

**Museum of New Zealand Te Papa Tongarewa, PO Box 467, Wellington, New Zealand**  
**ricardop@tepapa.govt.nz**

**Auckland, New Zealand**  
**2017**

**Copyright © Landcare Research New Zealand Ltd 2017**

No part of this work covered by copyright may be reproduced or copied in any form or by any means (graphic, electronic, or mechanical, including photocopying, recording, taping information retrieval systems, or otherwise) without the written permission of the publisher.

**Cataloguing in publication**

Palma, Ricardo L.

Phthiraptera (Insecta) : a catalogue of parasitic lice from New Zealand / by Ricardo L. Palma. -- Lincoln, N.Z. : Landcare Research NZ, 2017.

1 online resource

(Fauna of New Zealand, ISSN 1179-7193 (online) ; no. 76 = Ko te Aitanga Pepeke o Aotearoa ; nama 76)

ISBN 978-0-947525-18-7 (online)

ISBN 978-0-947525-19-4 (print)

1.Lice -- New Zealand. I. Title. II. Landcare Research New Zealand Ltd.

UDC 595.751(931)

**Suggested citation:**

Palma, RL. 2017. Phthiraptera (Insecta): a catalogue of parasitic lice from New Zealand. Lincoln, N.Z. : Landcare Research.

<https://doi.org/10.7931/J2/FNZ.76>

Prepared for publication by Zhi-Qiang Zhang and the authors using computer-based text processing, layout, and printing at Landcare Research, Private Bag 92170, Auckland, New Zealand.

To access on-line extracts from this series visit:

<http://fnz.landcareresearch.co.nz/>

Published by Landcare Research, P.O. Box 40, Lincoln, Canterbury, N.Z.

Website: <http://www.landcareresearch.co.nz/>

Printed by 3A

Cover design: Birgit Rhode

Date of publication 30 September 2017

**Publication of the Fauna of New Zealand series is supported by Core funding for Crown Research Institutes from the Ministry of Business, Innovation and Employment's Science and Innovation Group, for the Defining New Zealand's Land Biota research programme.**

## POPULAR SUMMARY

Class **Insecta**

Order **Phthiraptera**

### Lice

The simple mention of words like “lice” or “nits” is enough to bring strong feelings of revulsion from most people, reminding them of unpleasant childhood memories when they were subjected to uncomfortable treatments to get rid of the infestation. However, those feelings were caused mainly by only one kind of louse: head lice, which— together with its relatives: body and pubic lice—live exclusively on human beings. Unfortunately, the majority of the human population is unaware of the great number of other lice living permanently on almost all birds and most mammals, which cause no harm to humans and very little to their animal hosts. At present, there are about 5,500 species of parasitic lice described and named from many birds and mammals, which are as relevant to the world biodiversity as their hosts. Also, most people are unaware of the vast diversity of body shapes, sizes and colours exhibited by lice. Examples of this morphological diversity can be appreciated from the photographs included in this book. Parasitic lice are wingless, small, flat-bodied insects, grouped in what scientists call an “Order”, and are subdivided in about 20 “families” depending on their body features. According to their feeding habits—either eating feathers and other solid matter or taking liquid blood—two major groups have been named as “chewing” and “sucking” lice, respectively. However, these two groups are not recognised as scientifically accurate because the chewing lice are represented by different morphological types which are not genetically closely related.

The many birds and the few native and introduced mammals which live in New Zealand are hosts to over 400 species of lice, which are listed and annotated in this catalogue. Most of these species parasitise seabirds and shorebirds with wide geographic distributions over the world oceans. There is also a number of lice living exclusively on New Zealand terrestrial birds, in particular kiwis, parrots, pigeons and some perching birds. The fact that most louse species live only on one host species, or on a small number of closely related hosts, allows the identification of the host from the identity of its lice, without even examining the host. This characteristic has been successfully used to identify partial remains of bird bodies found in beaches, on roads, or in the forest. Also, some bird species that are difficult to distinguish from each other by their own features, kiwis for example, have been shown to be different by studying their distinct lice.

The plight for the survival of endangered bird species is well publicised and known in New Zealand, but people do not realise that the unique lice living on those birds are also endangered and need to be conserved as much as their hosts, especially when birds are managed and handled for their reproduction in captivity. There are overseas examples, such as the case of the Californian condor, where the management of an endangered bird to ensure its survival has caused the extinction of its unique lice, a situation that should be avoided because those lice are also part of world biodiversity.

### Kutu

Mā te whakahuatia noa o te kupu "kutu" o te kupu "riha" rānei e pupū ake ai te anuanu i te nuinga o te tangata, me te hoki anō o ngā mahara ki te tamarikitanga i wetiweti ai rātou ki ngā mahi patu i te hanga e mui ana i te māhunga. Heoi rā, kotahi anō te momo kutu nāna i pupū ake ai te anuanu i te tangata: ko te kutu i te māhunga. Engari he hoa anō o te kutu nei ko te tinana tangata anō tōna kāinga, ko te kutu piri tinana, me tērā momo ka kitea i ngā huruhuru o raro. Ko te mate kē ia, kāore te nuinga o te tāngata e mōhio arā noa atu ngā momo kutu e rarau ana te noho i te nuinga o te manu me te nuinga o te whāngote, ka mutu kāore o rātou pānga kino ki te tangata, he iti rānei te pānga kino ki te kararehe e noho ana hei piringa mō rātou. I tēnei wā kei tōna 5,500 ngā momo kutu pirinoa e mōhioia ana, kua whakaingoatia hoki, e noho ana i te manu me te whāngote, ka mutu e rite ana tō rātou hāngaitanga ki te kanorau koiora o te ao pērā i o rātou kaiatawhai. Waihoki e noho kūare ana te nuinga o te tangata ki te rerekē o te āhua, o te rahi me ngā tae o ngā momo kutu katoa. Ka kitea he tauira o ngā rerekētanga nei i ngā whakaahua e takoto mai ana ki te pukapuka nei. Kāore he parirau o te kutu pirinoa, he hanga iti, he papatahi te tinana, ka mutu kua whakarōpūhia e ngā kaupūtaiao ki te "Pūtoi", kua wehewehea hoki ki ētahi whānau e 20 nei. Kei te āhua tonu o ngā tinana. Kei te āhua anō o tā rātou i kai ai—he kai hurumanu rānei me ētahi atu kai mārō, he kai toto rānei—e rua ngā rōpū matua kua kīia he kutu "ngaungau", he kutu "momi" rānei. Heoi anō, kāore e kīia ngā rōpū e rua nei he

rōpū tika ā-pūtaiao nei i te mea he āhua kē, he āhua kē tō ngā kutu ngaungau, ka mutu kāore e noho whanaunga tata ana ā-ira nei.

E noho ana te tini manu me ngā whāngote taketake o Āotearoa, tae atu ki ērā i haria mai i whenua kē hei kaiatawhai mō ētahi momo kutu e 400 neke atu, ka mutu kua whakarārangihia, kua tautuhia ēnei ki te pukapuka nei. Ko te nuinga e piri ana ki te manu moana, ki te manu tahatika, ngā manu e tāwhai ana ki ngā moana o te ao. Arā anō ētahi momo kutu ka piri noa ki ngā manu whenua o Aotearoa, ngā manu pērā i te kiwi, i te kākā, i te kūkū me ētahi manu noho pae. I te mea ka kitea noatia te nuinga o ngā momo kutu i tētahi momo kaiatawhai kotahi, i ētahi kaiatawhai rānei e noho whanaunga tata ana, ka taea te tautuhi te kaiatawhai mā te tautuhi i te kutu. Ehara i te mea me āta mātai te kaiatawhai. E whakamahia ana tēnei āhuatanga ki te tautuhi i ngā toenga tinana manu ka kitea i tātahi, i ngā huarahi, i te ngahere rānei. Ka mutu ko ētahi momo manu e uaua ana te wehe tētahi i tētahi nā runga i ngā huruhuru, pērā i te kiwi, ka kitea te motuhaketanga o tētahi momo kiwi i tētahi atu mā te mātai i ō rātou kutu.

Kua puta whānui te rongō mō ngā raruraru e pā ana ki te whakarauora i ngā manu e noho mōrearea ana i Aotearoa nei, engari kāore te tangata e mōhio mō te noho mōrearea anō o ngā kutu motuhake e atawhaitia ana e aua manu. Me āta tiaki anō ngā kutu nei pērā i ō rātou kaiatawhai. E hāngai ana tēnei kōrero ki ngā manu e tiakina ana, e atawhaitia ana i ngā whare manu kia puta ai he hua e ora tonu ai tōna momo. Kei tāwāhi ngā tauira pērā i te condor o Karapōnia, e tiakina ana te manu noho mōrearea kia ora tonu ai tōna momo, ko te hua o tēnei ko te wharengaro o ana kutu motuhake. Kia kaha te karo i tēnei āhuatanga, inā rā, he wāhanga nui tonu aua kutu rā o te kanorau koiora o te ao.



Contributor Ricardo L. Palma was born in Argentina. He attended the University of Buenos Aires, where he completed a Master's Degree in Biological Sciences, majoring in Entomology, in January 1971. After finishing compulsory military service, he left his native country to work at the University of Concepción, Chile, lecturing on entomology and researching the taxonomy of the family Mydidae (Diptera). Following the September 11 military coup in 1973, he left Chile as a political refugee and was given permanent residence in New Zealand, where he obtained a research assistant position under the direction of Professor Robert L.C. Pilgrim at the University of Canterbury, Christchurch, in May 1974. After two years in Canterbury, he was successful in attaining the position of Scientist (Entomology) in the then National Museum of New Zealand, now Te Papa, where he was promoted to Curator of Entomology in 1991—position he held until his retirement in May 2016 after working at the museum for 40 years. He has published over 130 peer-reviewed papers, several book chapters and two books dealing mainly with parasitic lice (Phthiraptera), together with a great numbers of co-authors from many countries. Other animal groups covered in his publications are flies, beetles, feather mites, ticks and birds. His study of lice from kiwis indicated the existence of a previously unrecognised kiwi species, of which he became one of the authors of its original description. He is currently the editor of *Zootaxa* for Phthiraptera papers, member of the *Fauna of New Zealand* Series Editorial board and of several overseas editorial boards, as well as member of the Checklist Committee of the Ornithological Society of New Zealand, and is the New Zealand representative to the international Working Group on Avian Nomenclature. Although not officially employed, Te Papa has given him the position of Honorary Research Associate, which entitles him to pursue his research and editorial work at the museum two to three days per week, when he is not travelling overseas.

I whānau mai te kaituhi, a Ricardo L. Palma ki Āketina. I haere ia ki te Whare Wānanga o Buenos Aires, i reira nei ia e whakatutuki ana i tana Tohu Paerua Pūtaiao Koiora i te Kohitātea o te tau 1971. Ka mutu tana haere hei hoia i raro i te ture, ka wehe atu ia i tana kāinga taketake ake ki te mahi i te Whare Wānanga o Concepción i Hiri, ki te kauhau mō te mātaītanga pepeke me te rangahau i te pūnaha whakarōpūtanga o te whānau Mydidae (Diptera). Whai muri mai i te tukipoto taua o te 11 o Mahuru i te tau 1973, ka rere ia i Hiri hei rerenga tōrangapū, ka whakaaehia tana noho tūturu ki Aotearoa. Ka noho ia hei kaiāwhina rangahau i raro i ngā tohutohu a Ahorangi Robert L.C. Pilgrim i te Whare Wānanga o Ōtautahi i te Haratua 1974. Ka pau te rua tau ki Ōtautahi, ka whiwhi ia i te tūranga Kaipūtaiao (Mātai Pepeke) i te Whare Taonga o Aotearoa o taua wā, e mōhiotia nei ko Te Papa. I reira ka whakapikia ia ki te tūranga o te Kaitiaki Mātai Pepeke i te tau 1991—he tūranga mau tonu ā eke rawa ia ki te ahungarua i te Haratua 2016, ka 40 tau ia e mahi ana i te Te Papa. Neke atu i te 130 ngā pepa, i āta arotakea e ōna anō tāngata pūtaiao, kua tāia, ka hia nei ngā upoko pukapuka me ētahi pukapuka e rua e pā ana ki te kutu pirinoa (Phthiraptera) i tuhia nei e rātou ko ētahi atu o ētahi whenua maha tonu. Kua whai wāhi atu anō ia ki ētahi atu tānga mō ētahi atu kāhui pepeke pērā i te rango, i te pītara, i ngā kutu hurumanu i te kutu wae waru me te manu. I ana mātaītanga i ngā kutu i te kiwi, ka kitea e ia tētahi momo kiwi hou, ā, ko ia tētahi o ngā kaituhi e whakamārama ana i tōna āhua ake. I tēnei wā ko ia te ētita o te *Zootaxa* mō ngā pepa e pā ana ki te Phthiraptera, he mema anō ia o te poari mō te Raupapa Tānga mō Ngāi Kīrehe o Aotearoa me ētahi atu poari ētita o te ao, me tana noho anō hei mema o te Komiti Rārangi Hihira o te Hunga Mātai Manu o Aotearoa, ko ia anō te māngai o Aotearoa ki te Rōpū Mahi mō te Whakaingoa Manu o te ao. Ehara ia i te kaimahi tūturu, heoi anō kua tohua ia e Te Papa ki tana tūnga hei Hoa Rangahau Hōnore, ka mutu nā konei ia i āhei ai ki te rangahau tonu i āna kaupapa, ki te whai tonu i āna mahi ētita i te whare taonga mō te rua ki te toru rā i te wiki, mēnā kāore ia e haere ana i tāwāhi.

Māori translation by Te Haumihiata Mason

## ABSTRACT

The parasitic lice (Insecta: Phthiraptera) from the New Zealand Subregion are all listed and annotated with data from both literature records and collections. The current scientific name, its taxonomic history, data on type material, type host, other hosts, geographic distribution within New Zealand and elsewhere, New Zealand literature references and other significant references are given for each species or subspecies. The louse fauna comprises 424 species/subspecies distributed in 101 genera, 14 families and three suborders. Among them, 381 species/subspecies (90%) are from birds, of which 58 taxa are introduced by human agency, and 43 species (10%) are from mammals, of which 37 have been introduced by humans. The total number of species/subspecies includes 22 new records of louse taxa for this country. Six species/subspecies deleted from the New Zealand louse fauna are listed and discussed. A host-parasite list of all the hosts known to harbour lice in the New Zealand Subregion, including 18 new host-lice associations, and a list of bird species which breed in the Subregion but with no lice collected from them yet, are given. Citations to literature references for every publication known to include a record or a mention of a louse from the New Zealand Subregion, and other significant publications dealing with various aspects of louse taxonomy, biology, ecology, phylogeny and evolution are also given.

The New Zealand louse fauna is characterised by (1) low endemism, with only 11% at species level and 2% at generic level, (2) a high proportion (22.5%) of species introduced by human agency, (3) a very low number of species from native mammals (six or 1.8% of the total fauna), and (4) a greater number of species from seabirds and shorebirds (213 or 56%) than from terrestrial birds (168 or 44%).

**Keywords:** Insecta, Phthiraptera, Amblycera, Ischnocera, Anoplura, parasitic lice, catalogue, New Zealand Subregion, synonymies, primary types, type hosts, New Zealand localities, geographic distribution, references, endemic, native, introduced, new records, new host-lice associations, hosts, birds, mammals, Aves, Mammalia.

urn:lsid:zoobank.org:pub:A2DB72BC-A954-47BF-86F2-74655475BD4C

Accepted by Zhi-Qiang Zhang: 1 April 2017; published: 30 September 2017

## CHECKLIST OF TAXA

Order PHTHIRAPTERA Haeckel, 1896 .....	29
Suborder AMBLYCERA Kellogg, 1896 .....	29
Family BOOPIIDAE Mjöberg, 1910 .....	29
Genus <i>Boopia</i> Piaget, 1880 .....	29
<i>Boopia notafusca</i> Le Souëf, 1902 .....	29
Genus <i>Heterodoxus</i> Le Souëf & Bullen, 1902 .....	29
<i>Heterodoxus ampullatus</i> Kéler, 1971 .....	29
Family GYROPIDAE Kellogg, 1896 .....	30
Genus <i>Gliricola</i> Mjöberg, 1910 .....	30
Subgenus <i>Gliricola</i> Mjöberg, 1910 .....	30
<i>Gliricola (Gliricola) porcelli</i> (Schrank, 1781) .....	30
Genus <i>Gyropus</i> Nitzsch, 1818 .....	30
<i>Gyropus ovalis</i> Burmeister, 1838 .....	30
Family LAEMOBOTHRIIDAE Mjöberg, 1910 .....	31
Genus <i>Laemobothrion</i> Nitzsch, 1818 .....	31
Subgenus <i>Laemobothrion</i> Nitzsch, 1818 .....	31
<i>Laemobothrion (Laemobothrion) timunculi</i> (Linnaeus, 1758) .....	31
Family MENOPONIDAE Mjöberg, 1910 .....	31
Genus <i>Actornithophilus</i> Ferris, 1916 .....	32
<i>Actornithophilus bicolor</i> (Piaget, 1880) .....	32
<i>Actornithophilus ceruleus</i> (Timmermann, 1954) .....	32
<i>Actornithophilus grandiceps</i> (Piaget, 1880) .....	33



<i>Actornithophilus hoplopteri</i> (Mjöberg, 1910) sensu lato .....	33
<i>Actornithophilus limosae</i> (Kellogg, 1908) .....	33
<i>Actornithophilus ochraceus</i> (Nitzsch, 1818) sensu lato .....	34
<i>Actornithophilus pediculoides</i> (Mjöberg, 1910) .....	34
<i>Actornithophilus piceus lari</i> (Packard, 1870) .....	35
<i>Actornithophilus sedes</i> Eichler, 1944 New Record .....	35
<i>Actornithophilus spinulosus</i> (Piaget, 1880) .....	36
<i>Actornithophilus umbrinus</i> (Burmeister, 1838) .....	36
Genus <i>Amyrsidea</i> Ewing, 1927 .....	37
Subgenus <i>Argimenopon</i> Eichler, 1947 .....	37
<i>Amyrsidea</i> ( <i>Argimenopon</i> ) <i>minuta</i> Emerson, 1961 .....	37
<i>Amyrsidea</i> ( <i>Argimenopon</i> ) <i>perdicis</i> (Denny, 1842) .....	37
Genus <i>Ancistrona</i> Westwood, 1874 .....	38
<i>Ancistrona vagelli</i> (J.C. Fabricius, 1787) .....	38
Genus <i>Apterygon</i> Clay, 1961 .....	39
<i>Apterygon dumosum</i> Tandan, 1972 .....	39
<i>Apterygon hintoni</i> Clay, 1966 .....	40
<i>Apterygon mirum</i> Clay, 1961 .....	40
<i>Apterygon okarito</i> Palma & Price, 2004 .....	41
Genus <i>Austromenopon</i> Bedford, 1939 .....	41
<i>Austromenopon aegialitidis</i> (Durrant, 1906) sensu lato .....	41
<i>Austromenopon affine</i> (Piaget, 1890) .....	41
<i>Austromenopon atrofulvum</i> (Piaget, 1880) .....	42
<i>Austromenopon beckii</i> (Kellogg, 1906) .....	43
<i>Austromenopon brevifimbriatum</i> (Piaget, 1880) .....	43
<i>Austromenopon bulweriae</i> Timmermann, 1963 .....	44
<i>Austromenopon cursorium</i> (Giebel, 1874) New Record .....	44
<i>Austromenopon elliotti</i> Timmermann, 1954 .....	45
<i>Austromenopon enigki</i> Timmermann, 1963 .....	45
<i>Austromenopon fuscofasciatum</i> (Piaget, 1880) .....	46
<i>Austromenopon haematopi</i> Timmermann, 1954 .....	46
<i>Austromenopon himantopi</i> Timmermann, 1954 .....	46
<i>Austromenopon limosae</i> Timmermann, 1954 .....	47
<i>Austromenopon longithoracicum</i> (Piaget, 1880) .....	47
<i>Austromenopon lutescens lutescens</i> (Burmeister, 1838) .....	47
<i>Austromenopon meyeri</i> (Giebel, 1874) .....	48
<i>Austromenopon navigans</i> (Kellogg, 1896) .....	48
<i>Austromenopon ossifragae</i> (Eichler, 1949) .....	49
<i>Austromenopon paululum</i> (Kellogg & Chapman, 1899) .....	49
<i>Austromenopon phaeopodis</i> (Schrank, 1802) .....	50
<i>Austromenopon pinguis</i> (Kellogg, 1896) .....	51
<i>Austromenopon popellus</i> (Piaget, 1890) .....	51
<i>Austromenopon stammeri</i> Timmermann, 1963 .....	52
<i>Austromenopon transversum</i> (Denny, 1842) .....	53
<i>Austromenopon</i> species 1 .....	53
<i>Austromenopon</i> species 2 .....	53
<i>Austromenopon</i> species 3 .....	54
Genus <i>Bonomiella</i> Conci, 1942 .....	54
<i>Bonomiella columbae</i> Emerson, 1957 .....	54
Genus <i>Ciconiphilus</i> Bedford, 1939 .....	55
<i>Ciconiphilus decimfasciatus</i> (Boisduval & Lacordaire, 1835) .....	55
<i>Ciconiphilus pectiniventris</i> (Harrison, 1916) .....	55
Genus <i>Colpocephalum</i> Nitzsch, 1818 .....	56
<i>Colpocephalum eucarenum</i> Burmeister, 1838 .....	56
<i>Colpocephalum fregili</i> Denny, 1842 .....	57
<i>Colpocephalum leptopygos</i> Nitzsch [in Giebel], 1874 .....	57
<i>Colpocephalum pilgrimi</i> Price, 1967 .....	57

<i>Colpocephalum subzerafae</i> Tendeiro, 1988 .....	58
<i>Colpocephalum tausi</i> (Ansari, 1951) .....	58
<i>Colpocephalum turbinatum</i> Denny, 1842 .....	59
Genus <i>Cuculiphilus</i> Uchida, 1926 .....	59
Subgenus <i>Cuculiphilus</i> Uchida, 1926 .....	59
<i>Cuculiphilus (Cuculiphilus) fasciiventris</i> Carriker, 1955 .....	59
<i>Cuculiphilus (Cuculiphilus) platygaster</i> (Giebel, 1874) .....	59
Genus <i>Dennyus</i> Neumann, 1906 .....	60
Subgenus <i>Takamatsuia</i> Uchida, 1926 .....	60
<i>Dennyus (Takamatsuia) species</i> New Record .....	60
Genus <i>Eidmanniella</i> Kéler, 1938 .....	60
<i>Eidmanniella albescens</i> (Piaget, 1880) .....	60
<i>Eidmanniella eurygaster</i> (Nitzsch [in Giebel], 1866) .....	61
<i>Eidmanniella pellucida</i> (Rudow, 1869) .....	61
<i>Eidmanniella pustulosa</i> (Nitzsch [in Giebel], 1866) .....	62
Genus <i>Eucolpocephalum</i> Bedford, 1930 .....	62
<i>Eucolpocephalum femorale</i> (Piaget, 1880) .....	63
Genus <i>Franciscoloa</i> Conci, 1942 .....	63
Subgenus <i>Franciscoloa</i> Conci, 1942 .....	63
<i>Franciscoloa (Franciscoloa) pallida</i> (Piaget, 1880) .....	63
Genus <i>Heteromenopon</i> Carriker, 1954 .....	64
Subgenus <i>Keamenopon</i> Price & Beer, 1967 .....	64
<i>Heteromenopon (Keamenopon) kea</i> (Kellogg, 1907) .....	64
<i>Heteromenopon (Keamenopon) species</i> .....	64
Genus <i>Hohorstiella</i> Eichler, 1940 .....	64
<i>Hohorstiella lata</i> (Piaget, 1880) .....	65
<i>Hohorstiella timorensis</i> Tendeiro, 1980 New Record .....	65
Genus <i>Holomenopon</i> Eichler, 1941 .....	65
<i>Holomenopon leucoxanthum</i> (Burmeister, 1838) .....	65
<i>Holomenopon tadornae</i> (Gervais, 1844) .....	66
<i>Holomenopon species 1</i> .....	66
<i>Holomenopon species 2</i> .....	67
Genus <i>Kurodaia</i> Uchida, 1926 .....	67
Subgenus <i>Conciella</i> Eichler, 1949 .....	67
<i>Kurodaia (Conciella) cryptostigmatia</i> (Nitzsch [in Giebel], 1861) .....	67
Genus <i>Longimenopon</i> Thompson, 1948 .....	68
<i>Longimenopon galeatum</i> Timmermann, 1957 .....	68
<i>Longimenopon species</i> .....	68
Genus <i>Menacanthus</i> Neumann 1912 .....	69
<i>Menacanthus eurysternus</i> (Burmeister, 1838) .....	69
<i>Menacanthus pallidulus</i> (Neumann, 1912) .....	70
<i>Menacanthus rhipidurae</i> Palma & Price, 2005 .....	70
<i>Menacanthus stramineus</i> (Nitzsch, 1818) .....	71
<i>Menacanthus species</i> .....	71
Genus <i>Menopon</i> Nitzsch, 1818 .....	72
<i>Menopon gallinae</i> (Linnaeus, 1758) .....	72
Genus <i>Myrsidea</i> Waterston, 1915 .....	72
<i>Myrsidea hihi</i> Sychra, Kolencik & Palma, 2016 .....	72
<i>Myrsidea ivanliteraki</i> Sychra, Kolencik & Palma, 2016 .....	73
<i>Myrsidea novaeseelandiae</i> Sychra, Kolencik & Palma, 2016 .....	73
<i>Myrsidea serini</i> (Séguy, 1944) .....	73
<i>Myrsidea thoracica</i> (Giebel, 1874).....	74
<i>Myrsidea species</i> .....	74
Genus <i>Nosopon</i> Hopkins, 1950 .....	75
<i>Nosopon lucidum</i> (Rudow, 1869) .....	75
Genus <i>Plegadiphilus</i> Bedford, 1939 .....	75
<i>Plegadiphilus plegadis</i> (Dubinin, 1938) .....	75

<i>Plegadiphilus threskiornis</i> Bedford, 1939 .....	76
Genus <i>Pseudomenopon</i> Mjöberg, 1910 .....	76
<i>Pseudomenopon concretum</i> (Piaget, 1880) .....	76
<i>Pseudomenopon pilgrimi</i> Price, 1974 .....	77
<i>Pseudomenopon pilosum</i> (Scopoli, 1763) .....	77
<i>Pseudomenopon scopulacorne</i> (Denny, 1842) .....	77
Genus <i>Trinoton</i> Nitzsch, 1818 .....	78
<i>Trinoton nigrum</i> Le Souëf, 1902 .....	78
<i>Trinoton querquedulae</i> (Linnaeus, 1758) .....	78
Family RICINIDAE Neumann, 1890 .....	79
Genus <i>Ricinus</i> De Geer, 1778 .....	79
<i>Ricinus</i> species .....	79
Suborder ISCHNOCERA Kellogg, 1896 .....	80
Family PHILOPTERIDAE Burmeister, 1838 .....	80
Genus <i>Acidoproctus</i> Piaget, 1878 .....	80
<i>Acidoproctus emersoni</i> Timmermann, 1962 New Record .....	80
<i>Acidoproctus gottwaldhirschi</i> Eichler, 1958 .....	80
Genus <i>Alcedoecus</i> Clay & Meinertzhagen, 1939 .....	80
<i>Alcedoecus alato-clypeatus</i> (Piaget, 1885) .....	81
<i>Alcedoecus delphax</i> (Nitzsch [in Giebel], 1866) New Record .....	81
Genus <i>Anaticola</i> Clay, 1936 .....	82
<i>Anaticola anseris</i> (Linnaeus, 1758) .....	82
<i>Anaticola crassicornis</i> (Scopoli, 1763) .....	82
<i>Anaticola magnificus</i> Ansari, 1955 .....	83
<i>Anaticola</i> species .....	83
Genus <i>Anatoecus</i> Cummings, 1916 .....	83
<i>Anatoecus dentatus</i> (Scopoli, 1763) sensu lato .....	83
<i>Anatoecus dentatus magnicornutus</i> Złotorzycka, 1970 .....	84
<i>Anatoecus icterodes</i> (Nitzsch, 1818) sensu lato .....	85
<i>Anatoecus icterodes oloris</i> Złotorzycka, 1970 .....	85
<i>Anatoecus singhi</i> Palma, 2015 .....	86
Genus <i>Aquanirmus</i> Clay & Meinertzhagen, 1939 .....	86
<i>Aquanirmus australis</i> Kettle, 1974 .....	86
<i>Aquanirmus podiceps</i> (Denny, 1842) New Record .....	87
Genus <i>Ardeicola</i> Clay, 1936 .....	87
<i>Ardeicola expallidus</i> Blagoveshtchensky, 1940 .....	87
<i>Ardeicola neopallidus</i> Price, Hellenthal & Palma, 2003 .....	88
<i>Ardeicola pilgrimi</i> Tandan, 1972 .....	88
<i>Ardeicola plataleae</i> (Linnaeus, 1758) .....	88
<i>Ardeicola raphidius</i> (Nitzsch [in Giebel], 1866) .....	89
<i>Ardeicola stellaris</i> (Denny, 1842) .....	89
Genus <i>Austrogoniodes</i> Harrison, 1915 .....	90
<i>Austrogoniodes antarcticus</i> Harrison, 1937 .....	90
<i>Austrogoniodes concii</i> (Kéler, 1952) .....	90
<i>Austrogoniodes cristati</i> Kéler, 1952 .....	91
<i>Austrogoniodes hamiltoni</i> Harrison, 1937 .....	91
<i>Austrogoniodes macquariensis</i> Harrison, 1937 .....	92
<i>Austrogoniodes macquariensis</i> Harrison, 1937 sensu lato .....	93
<i>Austrogoniodes mawsoni</i> Harrison, 1937 .....	93
<i>Austrogoniodes struthes</i> Harrison, 1915 nomen dubium .....	94
<i>Austrogoniodes vanalphenae</i> Banks & Palma, 2003 .....	94
<i>Austrogoniodes waterstoni</i> (Cummings, 1914) .....	94
Genus <i>Bedfordiella</i> Thompson, 1937 .....	95
<i>Bedfordiella unica</i> Thompson, 1937 .....	95
Genus <i>Brueelia</i> Kéler, 1936 .....	96
<i>Brueelia amsel</i> (Eichler, 1951) .....	96
<i>Brueelia breueri</i> Balát, 1955 New Record .....	96

<i>Brueelia callaeincola</i> Valim & Palma, 2015 .....	97
<i>Brueelia cyclothorax</i> (Burmeister, 1838) .....	97
<i>Brueelia delicata</i> (Nitzsch [in Giebel], 1866) .....	97
<i>Brueelia densilimba</i> (Nitzsch [in Giebel], 1866) New Record .....	98
<i>Brueelia merulensis</i> (Denny, 1842) .....	98
<i>Brueelia nebulosa</i> (Burmeister, 1838) .....	99
<i>Brueelia parviguttata</i> (Blagoveshtchensky, 1940) New record .....	99
<i>Brueelia semiannulata</i> (Piaget, 1883) .....	100
<i>Brueelia turdinulae</i> Ansari, 1956 .....	100
<i>Brueelia</i> species .....	101
Genus <i>Campanulotes</i> Kéler, 1939 .....	101
<i>Campanulotes bidentatus compar</i> (Burmeister, 1838) .....	101
Genus <i>Capraiella</i> Conci, 1941 .....	102
<i>Capraiella</i> species .....	102
Genus <i>Carduiceps</i> Clay & Meinertzhagen, 1939 .....	102
<i>Carduiceps cingulatus cingulatus</i> (Denny, 1842) .....	102
<i>Carduiceps cingulatus lapponicus</i> Emerson, 1953 .....	102
<i>Carduiceps zonarius</i> (Nitzsch [in Giebel], 1866) .....	103
Genus <i>Chelopistes</i> Kéler, 1939 .....	103
<i>Chelopistes meleagridis</i> (Linnaeus, 1758) .....	104
Genus <i>Coloceras</i> Taschenberg, 1882 .....	104
<i>Coloceras chinense</i> (Kellogg & Chapman, 1902) New Record .....	104
<i>Coloceras harrisoni</i> (Tendeiro, 1972) .....	105
<i>Coloceras hemiphagae</i> (Tendeiro, 1972) .....	105
<i>Coloceras novaeseelandiae</i> (Tendeiro, 1972) .....	105
<i>Coloceras restinctum</i> (Tendeiro, 1972) .....	106
Genus <i>Columbicola</i> Ewing, 1929 .....	106
<i>Columbicola columbae columbae</i> (Linnaeus, 1758) .....	106
Genus <i>Cuclotogaster</i> Carriker, 1936 .....	107
<i>Cuclotogaster heterographus</i> (Nitzsch [in Giebel], 1866) .....	107
<i>Cuclotogaster synoicus</i> (Clay, 1938) .....	108
Genus <i>Cuculicola</i> Clay & Meinertzhagen, 1939 .....	108
<i>Cuculicola kui</i> Kettle, 1980 .....	108
<i>Cuculicola latirostris</i> (Burmeister, 1838) .....	108
Genus <i>Degeeriella</i> Neumann, 1906 .....	109
<i>Degeeriella fusca</i> (Denny, 1842) .....	109
<i>Degeeriella rufa rufa</i> (Burmeister, 1838) .....	109
Genus <i>Docophoroides</i> Giglioli, 1864 .....	110
<i>Docophoroides brevis</i> (Dufour, 1835) .....	110
<i>Docophoroides harrisoni</i> Waterston, 1917 .....	110
<i>Docophoroides murphyi</i> (Kellogg, 1914) .....	111
<i>Docophoroides simplex</i> (Waterston, 1914) .....	112
Genus <i>Emersoniella</i> Tendeiro, 1965 .....	112
<i>Emersoniella bracteata</i> (Nitzsch [in Giebel], 1866) New Record .....	112
Genus <i>Episbates</i> Thompson, 1935 .....	113
<i>Episbates pederiformis</i> (Dufour, 1835) .....	113
Genus <i>Forficuloecus</i> Conci, 1941 .....	113
<i>Forficuloecus meinertzhageni</i> Guimarães, 1974 .....	113
<i>Forficuloecus pilgrimi</i> Guimarães, 1985 .....	114
<i>Forficuloecus</i> species New Record .....	114
Genus <i>Fulicoffula</i> Clay & Meinertzhagen, 1938 .....	115
<i>Fulicoffula lurida</i> (Nitzsch, 1818) .....	115
<i>Fulicoffula stammeri</i> Eichler, 1958 New Record .....	115
Genus <i>Goniocotes</i> Burmeister, 1838 .....	116
<i>Goniocotes chrysocephalus</i> Giebel, 1874 .....	116
<i>Goniocotes gallinae</i> (De Geer, 1778) .....	116
<i>Goniocotes pusillus</i> (Nitzsch [in Giebel], 1866) .....	116

Genus <i>Goniodes</i> Nitzsch, 1818 .....	117
<i>Goniodes colchici</i> (Denny, 1842) .....	117
<i>Goniodes dispar</i> Burmeister, 1838 .....	118
<i>Goniodes dissimilis</i> (Denny, 1842) .....	118
<i>Goniodes ortygis</i> (Denny, 1842) .....	118
<i>Goniodes pavonis</i> (Linnaeus, 1758) .....	119
<i>Goniodes retractus</i> Le Souëf, 1902 .....	119
<i>Goniodes stefani</i> Clay & Hopkins, 1955 .....	120
Genus <i>Haffneria</i> Timmermann, 1966 .....	120
<i>Haffneria grandis</i> (Piaget, 1880) .....	120
Genus <i>Halipeurus</i> Thompson, 1936 .....	121
<i>Halipeurus angusticeps</i> (Piaget, 1880) .....	121
<i>Halipeurus bulweriae</i> Timmermann, 1960 .....	122
<i>Halipeurus confusus</i> Palma, 2011 .....	122
<i>Halipeurus consimilis</i> Timmermann, 1960 .....	123
<i>Halipeurus diversus</i> (Kellogg, 1896) .....	123
<i>Halipeurus falsus pacificus</i> Edwards, 1961 .....	124
<i>Halipeurus gravis priapululus</i> Timmermann, 1961 .....	124
<i>Halipeurus kermadecensis</i> (Johnston & Harrison, 1912) .....	125
<i>Halipeurus leucophryna</i> Timmermann, 1960 .....	125
<i>Halipeurus marquesanus</i> (Ferris, 1932) .....	126
<i>Halipeurus mirabilis</i> Thompson, 1940 .....	126
<i>Halipeurus mundaе</i> Edwards, 1961 .....	127
<i>Halipeurus noctivagus</i> Timmermann, 1960 .....	127
<i>Halipeurus pelagicus</i> (Denny, 1842) .....	128
<i>Halipeurus pelagodromae</i> Palma, 2011 .....	128
<i>Halipeurus placodus</i> Edwards, 1961 .....	129
<i>Halipeurus pricei</i> Palma, 2011 .....	129
<i>Halipeurus procellariae</i> (J.C. Fabricius, 1775) .....	130
<i>Halipeurus spadix spadix</i> Timmermann, 1961 .....	130
<i>Halipeurus spadix subclavus</i> Timmermann, 1961 .....	131
<i>Halipeurus theresae</i> Timmermann, 1969 .....	131
<i>Halipeurus thompsoni</i> Edwards, 1961 .....	132
<i>Halipeurus turtur</i> Edwards, 1961 .....	132
Genus <i>Harrisoniella</i> Bedford, 1929 .....	133
<i>Harrisoniella ferox</i> (Giebel, 1867) .....	133
<i>Harrisoniella hopkinsi</i> Eichler, 1952 .....	133
Genus <i>Ibidoecus</i> Cummings, 1916 .....	134
<i>Ibidoecus bisignatus</i> (Nitzsch [in Giebel], 1866) .....	134
<i>Ibidoecus dianaе</i> Tandan, 1958 .....	135
<i>Ibidoecus platalae</i> (Denny, 1842) .....	135
Genus <i>Incidifrons</i> Ewing, 1929 .....	136
<i>Incidifrons fulicae</i> (Linnaeus, 1758) .....	136
<i>Incidifrons porzanae</i> Blagoveshtchensky, 1951 New Record .....	136
Genus <i>Lagopoecus</i> Waterston, 1922 .....	137
<i>Lagopoecus docophoroides</i> (Piaget, 1880) .....	137
Genus <i>Lipeurus</i> Nitzsch, 1818 .....	137
<i>Lipeurus caponis</i> (Linnaeus, 1758) .....	137
<i>Lipeurus maculosus maculosus</i> Clay, 1938 .....	138
Genus <i>Luniceps</i> Clay & Meinertzhagen, 1939 .....	138
<i>Luniceps actophilus</i> (Kellogg & Chapman, 1899) .....	138
<i>Luniceps drosti</i> Timmermann, 1954 .....	139
<i>Luniceps falcinellus</i> Timmermann, 1954 .....	139
<i>Luniceps limosae</i> Bechet, 1968 .....	140
<i>Luniceps limosella</i> Timmermann, 1954 .....	140
<i>Luniceps numenii madagascariensis</i> Gustafsson & Olsson, 2012 .....	140
<i>Luniceps numenii oliveri</i> (Johnston & Harrison, 1912) .....	141

<i>Luniceps superciliosus</i> Gustafsson & Olsson, 2012 .....	141
Genus <i>Melibrueelia</i> Valim & Palma, 2015 .....	142
<i>Melibrueelia novaeseelandiae</i> Valim & Palma, 2015 .....	142
Genus <i>Naubates</i> Bedford, 1930 .....	142
Subgenus <i>Guenterion</i> Palma & Pilgrim, 2002 .....	142
<i>Naubates (Guenterion) clypeatus</i> (Giebel, 1874) .....	142
<i>Naubates (Guenterion) damma</i> Timmermann, 1961 .....	143
<i>Naubates (Guenterion) heteroproctus</i> Harrison, 1937 .....	143
<i>Naubates (Guenterion) lessonii</i> Palma & Pilgrim, 2002 .....	144
<i>Naubates (Guenterion) prioni</i> (Enderlein, 1908) .....	144
<i>Naubates (Guenterion) pterodromi</i> Bedford, 1930 .....	145
Subgenus <i>Naubates</i> Bedford, 1930 .....	146
<i>Naubates (Naubates) fuliginosus</i> (Taschenberg, 1882) .....	146
<i>Naubates (Naubates) harrisoni</i> Bedford, 1930 .....	146
<i>Naubates (Naubates) thieli</i> Timmermann, 1965 .....	147
Genus <i>Neopsittaconirmus</i> Conci, 1942 .....	147
<i>Neopsittaconirmus albus</i> (Le Souëf & Bullen, 1902).....	147
<i>Neopsittaconirmus kea</i> (Kellogg, 1907) .....	148
Genus <i>Nesiotinus</i> Kellogg, 1903 .....	148
<i>Nesiotinus demersus</i> Kellogg, 1903 .....	148
Genus <i>Ornithobius</i> Denny, 1842 .....	149
<i>Ornithobius bucephalus</i> (Giebel, 1874) .....	149
<i>Ornithobius fuscus</i> Le Souëf, 1902.....	149
<i>Ornithobius goniopleurus</i> (Denny, 1842).....	150
Genus <i>Oxylipeurus</i> Mjöberg, 1910 .....	150
<i>Oxylipeurus clavatus</i> (McGregor, 1917) .....	150
<i>Oxylipeurus ellipticus</i> (Kéler, 1958) .....	151
<i>Oxylipeurus mesopelios colchicus</i> Clay, 1938 .....	151
<i>Oxylipeurus polytrapezius polytrapezius</i> (Burmeister, 1838) .....	152
Genus <i>Paraclisis</i> Timmermann, 1965 .....	152
<i>Paraclisis diomedeeae</i> (J.C. Fabricius, 1775) .....	152
<i>Paraclisis hyalina</i> (Neumann, 1911) .....	153
<i>Paraclisis miriceps</i> (Kellogg & Kuwana, 1902) .....	154
<i>Paraclisis obscura</i> (Rudow, 1869) .....	154
Genus <i>Pectinopygus</i> Mjöberg, 1910 .....	155
<i>Pectinopygus annulatus</i> (Piaget, 1880) .....	155
<i>Pectinopygus australis</i> Thompson, 1948 .....	155
<i>Pectinopygus bassani</i> (O. Fabricius, 1780) .....	156
<i>Pectinopygus carunculatus</i> Timmermann, 1964 .....	156
<i>Pectinopygus carunculatus</i> Timmermann, 1964 sensu lato .....	157
<i>Pectinopygus dispar</i> (Piaget, 1880) .....	157
<i>Pectinopygus garbei</i> (Pessôa & Guimarães, 1935) .....	157
<i>Pectinopygus gyricornis</i> (Denny, 1842) .....	158
<i>Pectinopygus punctatus</i> Timmermann, 1964 .....	158
<i>Pectinopygus setosus</i> (Piaget, 1880) .....	159
<i>Pectinopygus turbinatus</i> (Piaget, 1890) .....	159
<i>Pectinopygus varius</i> Timmermann, 1964 .....	160
<i>Pectinopygus</i> species .....	160
Genus <i>Pelmatocerandra</i> Enderlein, 1908 .....	160
<i>Pelmatocerandra setosa</i> (Giebel, 1876).....	160
Genus <i>Penenirmus</i> Clay & Meinertzhagen, 1938 .....	161
<i>Penenirmus</i> species .....	161
Genus <i>Perineus</i> Thompson, 1936 .....	161
<i>Perineus circumfasciatus</i> Kéler, 1957 .....	161
<i>Perineus concinnoides</i> Kéler, 1957 .....	162
<i>Perineus macronecti</i> Palma & Pilgrim, 1988 .....	163
<i>Perineus nigrolimbatus</i> (Giebel, 1874) .....	163

Genus <i>Philoceanus</i> Kellogg, 1903 .....	164
<i>Philoceanus fasciatus</i> (Carriker, 1958) .....	164
<i>Philoceanus garrodiae</i> (Clay, 1940) .....	164
<i>Philoceanus robertsi</i> (Clay, 1940) .....	165
Genus <i>Philopteroides</i> Mey, 2004 .....	165
<i>Philopteroides fuliginosus</i> Valim & Palma, 2013 .....	165
<i>Philopteroides macrocephalus</i> Valim & Palma, 2013 .....	166
<i>Philopteroides novaezealandiae</i> Mey, 2004 .....	166
<i>Philopteroides pilgrimi</i> Valim & Palma, 2013 .....	166
<i>Philopteroides xenicus</i> Mey, 2004 .....	167
Genus <i>Philopterus</i> Nitzsch, 1818 .....	167
<i>Philopterus irkutensis</i> Fedorenko, 1985 New Record .....	167
<i>Philopterus modularis</i> Denny, 1842 New Record .....	167
<i>Philopterus novaezealandiae</i> Palma & Price, 2000 .....	168
<i>Philopterus stadleri</i> (Eichler, 1959) New record .....	168
<i>Philopterus turdi</i> Denny, 1842 .....	169
<i>Philopterus</i> species 1 .....	169
<i>Philopterus</i> species 2 .....	170
Genus <i>Pseudonirmus</i> Mjöberg, 1910 .....	170
<i>Pseudonirmus charcoti</i> (Neumann, 1907) .....	170
<i>Pseudonirmus gurlti</i> (Taschenberg, 1882) .....	170
<i>Pseudonirmus lugubris</i> (Taschenberg, 1882) .....	171
Genus <i>Psittoecus</i> Conci, 1942 .....	171
<i>Psittoecus vanzolinii</i> Guimarães, 1974 .....	171
Genus <i>Quadriceps</i> Clay & Meinertzhagen, 1939 .....	172
<i>Quadriceps assimilis</i> (Piaget, 1890) .....	172
<i>Quadriceps auratus</i> (Haan, 1829) .....	173
<i>Quadriceps birostris</i> (Giebel, 1874) .....	173
<i>Quadriceps caspius</i> (Giebel, 1874) .....	174
<i>Quadriceps cedemajori</i> Timmermann, 1969 .....	174
<i>Quadriceps charadrii orarius</i> (Kellogg, 1896) .....	175
<i>Quadriceps coenocoryphae</i> Timmermann, 1955 .....	175
<i>Quadriceps dominella</i> Timmermann, 1953 .....	176
<i>Quadriceps ellipticus</i> (Nitzsch [in Giebel], 1866) sensu lato .....	176
<i>Quadriceps hemichrous</i> (Nitzsch [in Giebel], 1866) .....	176
<i>Quadriceps hopkinsi apophoretus</i> Timmermann, 1969 .....	177
<i>Quadriceps hopkinsi hopkinsi</i> Timmermann, 1952 .....	177
<i>Quadriceps normifer alpha</i> (Kellogg, 1914) .....	178
<i>Quadriceps normifer normifer</i> (Grube, 1851) .....	178
<i>Quadriceps normifer parvopallidus</i> (Eichler, 1951) .....	179
<i>Quadriceps novaeseelandiae</i> Timmermann, 1953 .....	179
<i>Quadriceps nycthemerus</i> (Burmeister, 1838) .....	180
<i>Quadriceps ornatus fuscolaminulatus</i> (Enderlein, 1908) .....	180
<i>Quadriceps punctatus lingulatus</i> (Waterston, 1914) .....	181
<i>Quadriceps punctatus sublingulatus</i> Timmermann, 1952 New Record .....	181
<i>Quadriceps renschi</i> Timmermann, 1954 .....	182
<i>Quadriceps ridgwayi</i> (Kellogg, 1906) .....	182
<i>Quadriceps sellatus houri</i> Hopkins, 1949 .....	183
<i>Quadriceps sellatus sellatus</i> (Burmeister, 1838) .....	183
<i>Quadriceps semifissus</i> (Nitzsch [in Giebel], 1866) .....	184
<i>Quadriceps separatus</i> (Kellogg & Kuwana, 1902) .....	184
<i>Quadriceps strepsilaris</i> (Denny, 1842) .....	185
<i>Quadriceps</i> species .....	185
Genus <i>Rallicola</i> Johnston & Harrison, 1911 .....	186
Subgenus <i>Aptericola</i> Harrison, 1915 .....	186
<i>Rallicola (Aptericola) gadowi</i> Harrison, 1915 .....	186
<i>Rallicola (Aptericola) gadowi</i> Harrison, 1915 sensu lato .....	186

<i>Rallicola (Apterocola) gracilentus</i> Clay, 1953 .....	187
<i>Rallicola (Apterocola) pilgrimi</i> Clay, 1972 .....	187
<i>Rallicola (Apterocola) rodericki</i> Palma, 1991 .....	188
Subgenus <i>Huiacola</i> Mey, 1990 .....	188
<i>Rallicola (Huiacola) extinctus</i> (Mey, 1990) .....	188
Subgenus <i>Rallicola</i> Johnston & Harrison, 1911 .....	189
<i>Rallicola (Rallicola) fulicae</i> (Denny, 1842) .....	189
<i>Rallicola (Rallicola) harrisoni</i> Emerson, 1955 .....	189
<i>Rallicola (Rallicola) lugens</i> (Giebel, 1874) .....	190
<i>Rallicola (Rallicola) ortyometrae philippensis</i> Emerson, 1966 .....	190
<i>Rallicola (Rallicola) tabuensis</i> Emerson, 1966 .....	191
<i>Rallicola (Rallicola) takahe</i> Holloway, 1956 .....	191
Genus <i>Saemundssonina</i> Timmermann, 1936 .....	191
Subgenus <i>Puffinoecus</i> Eichler, 1949 .....	192
<i>Saemundssonina (Puffinoecus) enderleini</i> (Eichler, 1949) .....	192
<i>Saemundssonina (Puffinoecus) jamaicensis</i> Timmermann, 1962 New Record .....	192
<i>Saemundssonina (Puffinoecus) orientalis</i> (Uchida, 1949) .....	193
<i>Saemundssonina (Puffinoecus) puellula</i> Timmermann, 1965 .....	193
<i>Saemundssonina (Puffinoecus) valida</i> (Kellogg & Chapman, 1899) .....	194
<i>Saemundssonina (Puffinoecus) species</i> .....	194
Subgenus <i>Saemundssonina</i> Timmermann, 1936 .....	194
<i>Saemundssonina (Saemundssonina) albatrossa</i> Palma, 2012 .....	194
<i>Saemundssonina (Saemundssonina) albemarlensis</i> (Kellogg & Kuwana, 1902) .....	195
<i>Saemundssonina (Saemundssonina) antarctica</i> (Wood, 1937) .....	195
<i>Saemundssonina (Saemundssonina) bicolor</i> (Rudow, 1870) .....	196
<i>Saemundssonina (Saemundssonina) cephalus</i> (Denny, 1842) .....	196
<i>Saemundssonina (Saemundssonina) chathamensis</i> Timmermann, 1977 .....	197
<i>Saemundssonina (Saemundssonina) conica conica</i> (Denny, 1842) .....	197
<i>Saemundssonina (Saemundssonina) desolata</i> Timmermann, 1959 .....	198
<i>Saemundssonina (Saemundssonina) euryrhynga</i> (Giebel, 1874) .....	198
<i>Saemundssonina (Saemundssonina) gaini</i> (Neumann, 1913) .....	199
<i>Saemundssonina (Saemundssonina) gygisa</i> Palma, 2012 .....	200
<i>Saemundssonina (Saemundssonina) haematopi</i> (Linnaeus, 1758) .....	200
<i>Saemundssonina (Saemundssonina) hexagona</i> (Giebel, 1874) .....	200
<i>Saemundssonina (Saemundssonina) inexpectata</i> Timmermann, 1951 .....	201
<i>Saemundssonina (Saemundssonina) lari</i> (O. Fabricius, 1780) sensu lato .....	201
<i>Saemundssonina (Saemundssonina) limosae</i> (Denny, 1842) .....	202
<i>Saemundssonina (Saemundssonina) lockleyi</i> Clay, 1949 .....	203
<i>Saemundssonina (Saemundssonina) marina</i> Timmermann, 1956 .....	203
<i>Saemundssonina (Saemundssonina) melanocephalus</i> (Burmeister, 1838) .....	204
<i>Saemundssonina (Saemundssonina) nereis</i> Timmermann, 1956 .....	204
<i>Saemundssonina (Saemundssonina) platycephalus</i> (Kellogg & Kuwana, 1902) New Record .....	204
<i>Saemundssonina (Saemundssonina) platygaster</i> (Denny, 1842) sensu lato .....	205
<i>Saemundssonina (Saemundssonina) platygaster balati</i> Timmermann, 1969 .....	206
<i>Saemundssonina (Saemundssonina) pterodromae</i> Timmermann, 1959 .....	206
<i>Saemundssonina (Saemundssonina) remota</i> Timmermann, 1951 .....	207
<i>Saemundssonina (Saemundssonina) scolopacisphaeopodis scolopacisphaeopodis</i> (Schrank, 1803) .....	207
<i>Saemundssonina (Saemundssonina) stammeri</i> Timmermann, 1959 .....	208
<i>Saemundssonina (Saemundssonina) sternaes</i> (Linnaeus, 1758) .....	208
<i>Saemundssonina (Saemundssonina) thalassidromae incisa</i> Timmermann, 1950 .....	208
<i>Saemundssonina (Saemundssonina) thompsoni</i> Timmermann, 1951 .....	209
<i>Saemundssonina (Saemundssonina) tringae</i> (O. Fabricius, 1780) sensu lato .....	209
<i>Saemundssonina (Saemundssonina) uppalensis</i> (Rudow, 1870) .....	210
<i>Saemundssonina (Saemundssonina) species</i> New Record .....	211
Genus <i>Strigiphilus</i> Mjöberg, 1910 .....	211
<i>Strigiphilus aitkeni</i> Clay, 1966 .....	211
<i>Strigiphilus cursitans</i> (Nitzsch [in Giebel], 1861) .....	211



<i>Strigiphilus vapidus</i> Clay, 1977 .....	212
<i>Strigiphilus</i> species .....	212
Genus <i>Sturnidoecus</i> Eichler, 1944 .....	212
<i>Sturnidoecus sturni</i> (Schrank, 1776) .....	213
<i>Sturnidoecus</i> species .....	213
Genus <i>Trabeculus</i> Rudow, 1866 .....	213
<i>Trabeculus aviator</i> (Evans, 1912) .....	214
<i>Trabeculus flemingi</i> Timmermann, 1959 .....	214
<i>Trabeculus fuscoclypeatus</i> (Johnston & Harrison, 1912) .....	214
<i>Trabeculus hexakon</i> (Waterston, 1914) .....	215
<i>Trabeculus hexakon</i> (Waterston, 1914) sensu lato .....	216
<i>Trabeculus mirabilis</i> (Kellogg, 1896) .....	216
<i>Trabeculus schillingi</i> Rudow, 1866 .....	217
<i>Trabeculus schillingi</i> Rudow, 1866 sensu lato .....	218
<i>Trabeculus</i> species 1 .....	218
<i>Trabeculus</i> species 2 .....	218
Family TRICHODECTIDAE Kellogg, 1896 .....	219
Genus <i>Bovicola</i> Ewing, 1929 .....	219
Subgenus <i>Bovicola</i> Ewing, 1929 .....	219
<i>Bovicola (Bovicola) bovis</i> (Linnaeus, 1758) .....	219
<i>Bovicola (Bovicola) caprae</i> (Gurlt, 1843) .....	220
<i>Bovicola (Bovicola) limbatus</i> (Gervais, 1844) .....	220
<i>Bovicola (Bovicola) longicornis</i> (Nitzsch, 1818) .....	221
<i>Bovicola (Bovicola) ovis</i> (Schrank, 1781) .....	222
Subgenus <i>Lepikentron</i> Kéler, 1938 .....	222
<i>Bovicola (Lepikentron) breviceps</i> (Rudow, 1866) .....	222
Subgenus <i>Spinibovicola</i> Lyal, 1985 .....	223
<i>Bovicola (Spinibovicola) hemitrangi</i> (Cummings, 1916) .....	223
Genus <i>Felicola</i> Ewing, 1929 .....	223
Subgenus <i>Felicola</i> Ewing, 1929 .....	223
<i>Felicola (Felicola) subrostratus</i> (Burmeister, 1838) .....	224
Genus <i>Trichodectes</i> Nitzsch, 1818 .....	224
Subgenus <i>Trichodectes</i> Nitzsch, 1818 .....	224
<i>Trichodectes (Trichodectes) canis</i> (De Geer, 1778) .....	224
Subgenus <i>Stachiella</i> Kéler, 1938 .....	225
<i>Trichodectes (Stachiella) ermineae</i> (Hopkins, 1941) .....	225
<i>Trichodectes (Stachiella) mustelae</i> (Schrank, 1803) .....	225
Genus <i>Tricholipeurus</i> Bedford, 1929 .....	226
<i>Tricholipeurus lipeuroides</i> (Méglin, 1884) .....	226
<i>Tricholipeurus parallelus</i> (Osborn, 1896) .....	227
Genus <i>Werneckiella</i> Eichler, 1940 .....	227
<i>Werneckiella equi</i> (Denny, 1842) .....	227
<i>Werneckiella ocellata</i> (Piaget, 1880) .....	228
Suborder ANOPLURA Leach, 1815 .....	228
Family ECHINOPHTHIRIIDAE Enderlein, 1904 .....	229
Genus <i>Antarctophthirus</i> Enderlein, 1906 .....	229
<i>Antarctophthirus carlinii</i> Leonardi et al., 2014 .....	229
<i>Antarctophthirus lobodontis</i> Enderlein, 1909 .....	229
<i>Antarctophthirus mawsoni</i> Harrison, 1937 .....	230
<i>Antarctophthirus microchir</i> (Trouessart & Neumann, 1888) .....	230
<i>Antarctophthirus ogmorhini</i> Enderlein, 1906 .....	231
Genus <i>Lepidophthirus</i> Enderlein, 1904 .....	231
<i>Lepidophthirus macrorhini</i> Enderlein, 1904 .....	231
Family HAEMATOPINIDAE Enderlein, 1904 .....	232
Genus <i>Haematopinus</i> Leach, 1815 .....	232
<i>Haematopinus asini</i> (Linnaeus, 1758) .....	232
<i>Haematopinus eurysternus</i> Denny, 1842 .....	233

<i>Haematopinus suis</i> (Linnaeus, 1758) .....	234
Family HOPLOPLEURIDAE Ewing, 1929 .....	234
Genus <i>Hoplopleura</i> Enderlein, 1904 .....	234
<i>Hoplopleura pacifica</i> Ewing, 1924 .....	234
Family LINOGNATHIDAE Webb, 1946 .....	235
Genus <i>Linognathus</i> Enderlein, 1905 .....	235
<i>Linognathus ovillus</i> (Neumann, 1907) .....	235
<i>Linognathus pedalis</i> (Osborn, 1896) .....	236
<i>Linognathus setosus</i> (von Olfers, 1816) .....	236
<i>Linognathus stenopsis</i> (Burmeister, 1838) .....	237
<i>Linognathus vituli</i> (Linnaeus, 1758) .....	237
Genus <i>Solenopotes</i> Enderlein, 1904 .....	238
<i>Solenopotes burmeisteri</i> (Fahrenheit, 1919) .....	238
<i>Solenopotes capillatus</i> Enderlein, 1904 .....	238
Family PEDICULIDAE Leach, 1817 .....	239
Genus <i>Pediculus</i> Linnaeus, 1758 .....	239
<i>Pediculus humanus capitis</i> De Geer, 1778 .....	239
<i>Pediculus humanus humanus</i> Linnaeus, 1758 .....	240
Family PTHIRIDAE Ewing, 1929 .....	240
Genus <i>Pthirus</i> Leach, 1815 .....	240
<i>Pthirus pubis</i> (Linnaeus, 1758) .....	241
Family POLYPLACIDAE Fahrenheit, 1912 .....	241
Genus <i>Haemodipsus</i> Enderlein, 1904 .....	241
<i>Haemodipsus lyriocephalus</i> (Burmeister, 1838) .....	241
<i>Haemodipsus ventricosus</i> (Denny, 1842) .....	242
Genus <i>Polyplax</i> Enderlein, 1904 .....	243
<i>Polyplax serrata</i> (Burmeister, 1838) .....	243
<i>Polyplax spinulosa</i> (Burmeister, 1838) .....	243

## CONTENTS

POPULAR SUMMARY .....	5
ABSTRACT .....	8
CHECKLIST OF TAXA .....	8
CONTENTS .....	18
INTRODUCTION .....	19
History of collections, research and publications on New Zealand lice .....	24
METHODS AND CONVENTIONS .....	26
Primary type specimens .....	27
Abbreviations for institutions holding primary types .....	28
THE CHECKLIST .....	29
SPECIES & SUBSPECIES DELETED FROM THE NEW ZEALAND LOUSE FAUNA .....	244
HOST-PARASITE LIST .....	245
LIST OF BIRD SPECIES WHICH BREED IN THE NEW ZEALAND SUBREGION BUT WITH NO LICE .....	272
ACKNOWLEDGEMENTS .....	272
REFERENCES .....	273
TAXONOMIC INDEX .....	360

## INTRODUCTION

The insect order Phthiraptera includes all the lice exclusively parasitic on birds and mammals. Historically, these lice were placed in two orders, the Mallophaga or chewing lice and the Anoplura or sucking lice. However, phylogenetic research based on morphology (Clay 1970; Lyal 1985b) showed that the “Mallophaga” was not a monophyletic unit. Therefore, this group was rejected, the Anoplura was demoted to a suborder, and the single order Phthiraptera was adopted and subdivided in four suborders: Amblycera, Ischnocera, Rhynchophthirina and Anoplura. More recently, phylogenetic research based on molecular evidence (Yoshizawa & Johnson 2003, 2010) and morphology of the male genitalia (Yoshizawa & Johnson 2006) has shown the Amblycera to be more closely related to the suborder of the Psocoptera containing the “booklice” than to the other three suborders of parasitic lice. Hence, Yoshizawa & Johnson (2006: 358) proposed that the single order Psocodea should be recognised to include all parasitic, bark and book lice, rejecting the order Phthiraptera as a valid group. Notwithstanding that position, the practical advantages of grouping together all parasitic lice in a single unit cannot be ignored and, therefore, I recognise the Phthiraptera as a useful group for the purpose of this faunistic catalogue.

Parasitic lice are wingless, small (adult length range 1–12 mm), dorso-ventrally compressed insects living permanently on warm-blooded vertebrates (Marshall 1981). They evolved from psocodean ancestors, probably living in nests, during the early to mid Cretaceous, 115–130 million years ago (Rózsa & Vas 2015a: 215). Most species of birds and mammals have lice (Price *et al.* 2003; Durden & Musser 1994b) living among their plumage or pelage, where they shelter, feed, reproduce and die. Most species of lice are highly host-specific (Clayton *et al.* 2015: 25), with one louse species per host species but, more frequently, one louse species parasitises two or more closely related hosts and, in a few exceptional cases, one species may be found on a wide range of host species belonging to several families (see *Menacanthus eurysternus* below). The geographical distribution of lice is, with some exceptions, that of their hosts (Clay 1964b, 1976b). Parasitic lice have developed morphological, behavioural and physiological adaptations to survive on their hosts. Similarly, because lice are detrimental to host health and fitness, hosts have developed adaptations to control their lice populations. This reciprocal natural selection pressure has led to the coevolution of hosts and lice (Johnson & Clayton 2003; Clayton *et al.* 2015). Thus, the phylogenetic relationships of lice often parallel those of their hosts and may help both to elucidate the relationships of the latter (e.g. Hughes *et al.* 2007; Hammer *et al.* 2010) and to distinguish closely related host taxa, which are otherwise poorly defined (e.g. Melville 1985; Palma 1991a; Whiteman *et al.* 2009).

Wise (1977: 55) was the first to compile a list of the species and subspecies of Phthiraptera recorded in the literature from New Zealand birds and mammals, including them in the suborders “Mallophaga” and Anoplura. Pilgrim & Palma (1982) augmented that list considerably but only for the chewing lice from birds, and in the form of a host-parasite list, which was amended and added to by Palma (1999). Due to the relatively high number of records listed at the generic level in those latter lists—i.e. without species identifications—it was then decided not to prepare a proper systematic catalogue of the louse species. However, subsequent taxonomic publications and further identification work on available collections have reduced considerably those ‘Genus only’ entries, to the extent that a more comprehensive catalogue is now a feasible option.

The sucking lice (suborder Anoplura) recorded in the literature from New Zealand were first listed by Wise (1977: 66), who included 19 species and subspecies from both native and introduced mammal hosts. In the revision of their 1981 checklist of ectoparasites of terrestrial mammals from New Zealand, Tenquist & Charleston (2001) included 35 introduced species of chewing and sucking lice. For a summary of species and subspecies of lice listed from New Zealand since Wise (1977), see Table 1.

This catalogue includes all the species and subspecies of parasitic lice recorded in the literature from New Zealand as well as 22 new records, of which 15 are the result of identifications of material previously published as “Genus sp.”, and seven are the result of identifications of newly collected samples, although three of the latter are identified to genus level only due to a lack of adequate material (see below). Also, this catalogue includes 18 new host-lice associations for New Zealand.

According to the status of their hosts, I have subdivided the 424 species/subspecies of lice included in this catalogue in four categories as defined in Tables 2 and 3, as follows: “Endemic”, “Native”, “Introduced” and “Uncertain”. The 27 species of uncertain status are determined according to three criteria:

1. Lack of adequate material to obtain a species identification. This condition applies to 19 species, listed as “*Genus species*”, e.g. *Forficuloecus* species.

2. The origins of their hosts are uncertain because they are regarded as both introduced by humans and self-introduced. This condition applies to seven louse species: four from the black swan (*Cygnus atratus*) and three from the sulphur-crested cockatoo (*Cacatua galerita*) (see details below).

3. One species is regarded as a “*nomen dubium*”: *Austrogoniodes strutheus* Harrison, 1915 (see below).

**TABLE 1.** Numbers of species and subspecies of lice from New Zealand in published lists.

	Amblycera		Ischnocera		Anoplura	TOTALS
	ex birds	ex mammals	ex birds	ex mammals	ex mammals	
<b>Wise 1977</b>	29	0	95	11	19	154
<b>Pilgrim &amp; Palma 1982*</b>	72	–	195	–	–	267
<b>Palma 1999*</b>	20	–	39	–	–	59
<b>Tenquist &amp; Charleston 2001</b>	–	4	–	14	17	35
<b>Palma 2010*</b>	78	4	230	14	22	348
<b>This catalogue*</b>	87	4	268	15	24	398

\* Numbers do not include ‘*Genus only*’ entries

In addition to the 19 species of uncertain status listed as “*Genus species*”, there are a further nine endemic or native species also listed as “*Genus species*”. However, these latter species are represented by extensive samples of both sexes which need to be formally described and named (e.g. *Trabeculus* species 1) or which need revisional studies to identify them (e.g. *Longimenopon* species).

Phthiraptera are generally the least endemic group among insects, due to the fact that most genera are represented in all continents. The New Zealand louse fauna is not an exception, with no endemic family, and only two genera (*Apterygon*, *Melibrueelia*) and two subgenera of *Rallicola* (*Aptericola*, *Huiacola*) endemic to this country. These endemic genera and subgenera are from endemic terrestrial hosts: kiwis and passerines. There is also a number of endemic species of lice parasitic on other endemic hosts —such as the takahe, the blue duck, several parrots, plovers, and several passerines (see below under “Host-parasite list” and Table 3)—but they belong to non-endemic louse genera.

Total numbers and percentages of the four categories of lice according to the status of their hosts are given in Table 2. Several louse species parasitise hosts that breed in New Zealand only but have extensive geographic ranges reaching other countries, and they cannot be considered as truly endemic to New Zealand. There are many examples of this category among lice from petrels and albatrosses.

**TABLE 2.** Numbers of species and subspecies of lice recorded from New Zealand in this catalogue.

	Amblycera		Ischnocera		Anoplura	TOTALS	%
	ex birds	ex mammals	ex birds	ex mammals	ex mammals		
Endemic	11	–	35	–	–	46	11
Native	65	–	185	–	6	256	60.5
Introduced	13	4	45	15	18	95	22.5
Uncertain	9	–	18	–	–	27	6
<b>TOTALS</b>	<b>98</b>	<b>4</b>	<b>283</b>	<b>15</b>	<b>24</b>	<b>424</b>	<b>100</b>

**Endemic:** Louse species/subspecies recorded exclusively from the New Zealand Subregion

**Native:** Louse species/subspecies recorded from the New Zealand Subregion and other parts of the world, from host species present in the Subregion by their own means.

**Introduced:** Louse species/subspecies recorded from the New Zealand Subregion and other parts of the world, from host species introduced to the Subregion by human agency.

**Uncertain:** Louse species/subspecies which either are unidentified due to lack of adequate material, or have been recorded from hosts of unconfirmed origins.

**TABLE 3.** List of louse genera with numbers of species/subspecies recorded from New Zealand under various categories.

Genus	TOTAL number of species & subspecies	ENDEMIC species & subspecies	NATIVE species & subspecies	INTRODUCED species & subspecies	Species & subspecies of UNCERTAIN status	NEW RECORDS
<i>Acidoproctus</i>	2	1	1	-	-	1
<i>Actornithophilus</i>	11	-	11	-	-	1
<i>Alcedoecus</i>	2	-	2	-	-	1
<i>Amyrsidea</i>	2	-	-	2	-	-
<i>Anaticola</i>	5	-	3	1	1	-
<i>Anatoecus</i>	5	-	2	1	2	-
<i>Ancistrona</i>	1	-	1	-	-	-
<i>Antarctophthirus</i>	5	-	5	-	-	-
<i>Apterygon</i>	4	4	-	-	-	-
<i>Aquanirmus</i>	2	1	1	-	-	1
<i>Ardeicola</i>	6	-	6	-	-	-
<i>Austrogoniodes</i>	10	1	8	-	1	-
<i>Austromenopon</i>	27	-	26	-	1	1
<i>Bedfordiella</i>	1	-	1	-	-	-
<i>Bonomiella</i>	1	-	-	1	-	-
<i>Boopia</i>	1	-	-	1	-	-
<i>Bovicola</i>	7	-	-	7	-	-
<i>Brueelia</i>	12	1	-	10	1	3
<i>Campanulotes</i>	1	-	-	1	-	-
<i>Capraiella</i>	1	-	-	-	1	-
<i>Carduiceps</i>	3	-	3	-	-	-
<i>Chelopistes</i>	1	-	-	1	-	-
<i>Ciconiphilus</i>	2	-	1	1	-	-
<i>Coloceras</i>	5	4	-	1	-	1
<i>Colpocephalum</i>	7	1	4	2	-	-
<i>Columbicola</i>	1	-	-	1	-	-
<i>Cuclotogaster</i>	2	-	-	2	-	-
<i>Cuculicola</i>	2	1	1	-	-	-
<i>Cuculiphilus</i>	2	-	2	-	-	-
<i>Degeeriella</i>	2	-	2	-	-	-
<i>Dennyus</i>	1	-	-	-	1	1
<i>Docophoroides</i>	4	-	4	-	-	-
<i>Eidmanniella</i>	4	-	4	-	-	-
<i>Emersoniella</i>	1	-	-	1	-	1
<i>Episbates</i>	1	-	1	-	-	-
<i>Eucolpocephalum</i>	1	-	1	-	-	-
<i>Felicola</i>	1	-	-	1	-	-
<i>Forficuloecus</i>	3	2	-	-	1	1
<i>Franciscocola</i>	1	-	-	-	1	-

..... continued on the next page

TABLE 3 (continued)

Genus	TOTAL number of species & subspecies	ENDEMIC species & subspecies	NATIVE species & subspecies	INTRODUCED species & subspecies	Species & subspecies of UNCERTAIN status	NEW RECORDS
<i>Fulicoffula</i>	2	-	2	-	-	1
<i>Gliricola</i>	1	-	-	1	-	-
<i>Goniocotes</i>	3	-	-	3	-	-
<i>Goniodes</i>	7	-	-	7	-	-
<i>Gyropus</i>	1	-	-	1	-	-
<i>Haematopinus</i>	3	-	-	3	-	-
<i>Haemodipsus</i>	2	-	-	2	-	-
<i>Haffneria</i>	1	-	1	-	-	-
<i>Halipeurus</i>	23	-	23	-	-	-
<i>Harrisoniella</i>	2	-	2	-	-	-
<i>Heterodoxus</i>	1	-	-	1	-	-
<i>Heteromenopon</i>	2	1	-	-	1	-
<i>Hohorstiella</i>	2	-	1	1	-	1
<i>Holomenopon</i>	4	1	2	-	1	-
<i>Hoplopleura</i>	1	-	-	1	-	-
<i>Ibidoecus</i>	3	-	3	-	-	-
<i>Incidifrons</i>	2	-	2	-	-	1
<i>Kurodaia</i>	1	-	1	-	-	-
<i>Laemobothrion</i>	1	-	1	-	-	-
<i>Lagopoecus</i>	1	-	-	1	-	-
<i>Lepidophthirus</i>	1	-	1	-	-	-
<i>Linognathus</i>	5	-	-	5	-	-
<i>Lipeurus</i>	2	-	-	2	-	-
<i>Longimenopon</i>	2	-	2	-	-	-
<i>Luniceps</i>	8	-	8	-	-	-
<i>Melibrueelia</i>	1	1	-	-	-	-
<i>Menacanthus</i>	5	1	1	2	1	-
<i>Menopon</i>	1	-	-	1	-	-
<i>Myrsidea</i>	6	2	-	3	1	-
<i>Naubates</i>	9	-	9	-	-	-
<i>Neopsittaconirmus</i>	2	1	-	-	1	-
<i>Nesiotinus</i>	1	-	1	-	-	-
<i>Nosopon</i>	1	-	1	-	-	-
<i>Ornithobius</i>	3	-	-	2	1	-
<i>Oxylipeurus</i>	4	-	-	4	-	-
<i>Paraclisis</i>	4	-	4	-	-	-
<i>Pectinopygus</i>	13	3	9	-	1	-
<i>Pediculus</i>	2	-	-	2	-	-
<i>Pelmatocerandra</i>	1	-	1	-	-	-
<i>Penenirmus</i>	1	-	-	-	1	-

..... continued on the next page

TABLE 3 (continued)

Genus	TOTAL number of species & subspecies	ENDEMIC species & subspecies	NATIVE species & subspecies	INTRODUCED species & subspecies	Species & subspecies of UNCERTAIN status	NEW RECORDS
<i>Perineus</i>	4	-	4	-	-	-
<i>Philoceanus</i>	3	-	3	-	-	-
<i>Philopteroides</i>	5	5	-	-	-	-
<i>Philopterus</i>	7	1	-	5	1	3
<i>Plegadiphilus</i>	2	-	2	-	-	-
<i>Polyplax</i>	2	-	-	2	-	-
<i>Pseudomenopon</i>	4	1	3	-	-	-
<i>Pseudonirmus</i>	3	-	3	-	-	-
<i>Psittoecus</i>	1	-	-	-	1	-
<i>Pthirus</i>	1	-	-	1	-	-
<i>Quadriceps</i>	28	3	24	-	1	1
<i>Rallicola</i>	12	8	4	-	-	-
<i>Ricinus</i>	1	-	-	-	1	-
<i>Saemundssonina</i>	39	1	36	-	2	3
<i>Solenopotes</i>	2	-	-	2	-	-
<i>Strigiphilus</i>	4	-	2	1	1	-
<i>Sturnidoecus</i>	2	-	-	1	1	-
<i>Trabeculus</i>	10	1	9	-	-	-
<i>Trichodectes</i>	3	-	-	3	-	-
<i>Tricholipeurus</i>	2	-	-	2	-	-
<i>Trinoton</i>	2	-	1	-	1	-
<i>Werneckiella</i>	2	-	-	2	-	-
<b>101 genera</b>	<b>424</b>	<b>46</b>	<b>256</b>	<b>95</b>	<b>27</b>	<b>22</b>

**Endemic:** Louse species/subspecies recorded exclusively from the New Zealand Subregion

**Native:** Louse species/subspecies recorded from the New Zealand Subregion and other parts of the world, from host species present in the Subregion by their own means.

**Introduced:** Louse species/subspecies recorded from the New Zealand Subregion and other parts of the world, from host species introduced to the Subregion by human agency.

**Uncertain:** Louse species/subspecies which either are unidentified due to lack of adequate material, or have been recorded from hosts of unconfirmed origins.

**New records:** Louse species/subspecies recorded in this Catalogue for the first time from New Zealand.

At the species level, according to the statistics given in Price *et al.* (2003: 3), the chewing louse fauna of New Zealand comprises about 9% of the world fauna. The sucking lice represent only about 4.5% of the world fauna (see Durden & Musser 1994a: 4 for world data) but, if the introduced species are excluded, that percentage is just above 1%. This is a direct reflection of New Zealand's lack of native terrestrial mammals (King 2005). The paucity of New Zealand mammal lice is also shown by the statistics within this country, with 381 (90%) species from birds against 43 (10%) from mammals, including introduced species (see Table 2). However, if introduced species are excluded from both host groups, the difference is even greater, with an overwhelming 323 (98.2%) species of bird lice against six (1.8%) of mammal lice from native seals and sea lions.

Another feature of the New Zealand bird louse fauna is the greater number of species from seabirds and shorebirds (213 or 56%) than those from terrestrial birds (168 or 44%). Again, that difference is much greater if the 58 species of lice from introduced terrestrial birds are excluded, with 213 species (66%) from seabirds and shorebirds against 110 (34%) from terrestrial birds. This difference is a characteristic shared with louse faunas from

other oceanic archipelagos, such as the Galápagos (Palma & Peck 2013), the Faroes (Palma & Jensen 2005), and Tristan da Cunha (Hänel & Palma 2007).

Considering the taxonomy of the hosts, the New Zealand louse fauna is characterised by all the species parasitic on some endemic bird families (e.g. Apterygidae: kiwis; Strigopidae: parrots; Callaeidae: wattlebirds; Notiomystidae: stichbird), and an exceptionally high proportion of species/subspecies from the avian orders Procellariiformes (albatrosses and petrels), Sphenisciformes (penguins), Pelecaniformes (shags, gannets and pelicans), and Charadriiformes (plovers, shorebirds, waders, gulls, etc.). Numbers of lice from these eight host groups combined add up to 229 species/subspecies, or 60% of the total louse fauna parasitising New Zealand birds.

The total New Zealand louse fauna is much greater than the figures given in Table 2. In addition to the species of uncertain status and the need of some generic revisions discussed above, there are still 18 bird species which breed in New Zealand without lice recorded from them in this country: six are rare endemics, six are native, and six are introduced by humans, but all of them are seldom available to search for lice (see list below). Also, there is a great number of regular and occasional migratory birds as well as rare stragglers (see Checklist Committee 2010) known to harbour lice outside this country, but with no lice collected from them in New Zealand yet.

Considering the high number of species of birds and mammals introduced to New Zealand by human agency (Checklist Committee 2010; King 2005), there could have been an even greater number of louse taxa in this country, if the so-called sorting events, such as “missing the boat”, “drowning on arrival” and others (Paterson *et al.* 2003; Johnson & Clayton 2004; MacLeod *et al.* 2010) had not prevented a considerable number of lice from becoming established in New Zealand. Paterson *et al.* (1999: 219) list 18 species of birds introduced by humans with the lice recorded from them in their home countries and the lice recorded from them in New Zealand. With one exception, all hosts harbour fewer louse species in New Zealand than in their home ranges. Extreme examples are: the peafowl (*Pavo cristatus*) with only three louse species established in New Zealand out of nine in its home range, and the dunnoek or hedge sparrow (*Prunella modularis*) plus the rook (*Corvus frugilegus*) each with only one species established in New Zealand out of five in their home ranges (Paterson *et al.* 1999: 220–221; but six according to Price *et al.* 2003: 338, 354).

A number of bird species have gone extinct in New Zealand since Europeans settled this country in 1840 (Checklist Committee 2010), and their host-specific lice also went extinct before any could be collected. However, preserved specimens of some of those extinct birds have been kept in museum collections around the world, and they have been the source of louse samples that have been described and named as different species. At least five extinct louse species have been found, described and named from New Zealand endemic birds of the genera *Apteryx* (kiwis), *Hemiphaga* (endemic pigeons), *Heteralocha* (huia) and *Xenicus* (endemic wrens) (Rózsa & Vas 2015b; see below). A louse species from *Sceloglaux albifacies*, an extinct endemic owl, has not been described yet because of lack of adequate material (see below under *Strigiphilus* sp.).

Lice together with all other parasites constitute a significant, perhaps the largest, part of the world biodiversity (Price 1980; Zimmer 2000), but they are becoming extinct together with their hosts in a process known as “coextinction” (Stork & Lyal 1993; Koh *et al.* 2004; Dunn *et al.* 2009). Therefore, it has been argued that parasites have the same rights as their hosts to be conserved and protected (Windsor 1995), but this concept has created a great dilemma with complex consequences which are difficult to resolve (Pérez *et al.* 2013).

### History of collections, research and publications on New Zealand lice

The first record of lice from New Zealand was made by Banks (1769: 186), referring to the lice he observed on the hair of some Maori people in Poverty Bay. Subsequently, there were several reports of both human head and body lice from this country during pre-European times, summarised and discussed by Andrews (1976a,b). The first citation of a louse by its scientific name was by Polack (1838: 320) who named human lice as “*pediculus humanus*”. This record was also included by White & Doubleday (1843: 283) in their list of New Zealand “Annulose animals”, inserted in Dieffenbach’s *Travels in New Zealand*. Since this book, there appears to be no more published records of lice from New Zealand until Hutton (1904) who, in his classic *Index Faunae Novae Zealandiae*, listed four species of sucking lice: one native from the New Zealand sea lion and three “naturalised” species, the short-nosed cattle louse, and head and body lice from humans.

The first papers recording and naming actual specimens of parasitic lice collected from hosts in New Zealand were published by Neumann (1907b) and by Kellogg (1907). Henry C. Wilkie, a New Zealand government



veterinarian, collected an undisclosed number of specimens from a sheep and sent them to L.-G. Neumann, a French parasitologist resident in Toulouse, France. Neumann (1907b) described and named those lice as "*Haematopinus ovillus*", together with a sample collected from Scottish sheep by William Evans in Edinburgh. This louse species is now placed in the genus *Linognathus* and is commonly known as the "sheep face louse" (Murray 1955a). In the same year, Vernon L. Kellogg, a professor at Stanford University, California, recorded three different feather lice—two as new "varieties" and one as a described species—from a sample collected on a kea and sent to him by Mr D.L. van Dine of Hawaii (Kellogg 1907). Those lice are now regarded as three full species exclusive to the kea (see below).

A few years later, Johnston & Harrison (1912) published a report on the lice collected from birds and goats on the Kermadec Islands by W.L. Wallace, as member of an expedition of New Zealand naturalists guided by Tom Iredale and W.R.B. Oliver, during 1907–1908. The lice were sent to Australia, where the authors resided, by Augustus Hamilton, the curator of the Dominion Museum (now Museum of New Zealand Te Papa Tongarewa = MONZ), who received the samples from W.R.B. Oliver. This was the first paper on lice published in a New Zealand scientific journal—the *Transactions of the New Zealand Institute*—and deals with 16 species of chewing lice of which seven are described and named as new. However, only three of the seven new species are still regarded as valid, although the type specimens of all seven are extant and kept in the MONZ collection (Palma *et al.* 1989).

Harrison (1915b) published the first paper dealing with kiwi lice, describing the new subgenus *Aptericola* to include the species of *Rallicola* exclusive to kiwis, and three new species: *R. (A.) gadowi* from the South Island brown kiwi, *R. (A.) novaezealandiae* from the Stewart Island brown kiwi and *R. (A.) gracilis* from the great spotted kiwi. The latter species was later renamed as *R. (A.) gracilentus* by Clay (1953) as the name "*gracilis*" proved to be preoccupied.

From the late 1920s to the 1960s, several genera and many species of lice which parasitise New Zealand seabirds were described and/or revised by Bedford (1929, 1930), Thompson (1935c, 1936, 1937b, 1940b, 1948c), Edwards (1961), and especially Timmermann (1936, 1951a–c, 1952a–d, 1953a–b, 1954a–f, 1957a–b, 1959a–c, 1960, 1961a–d, 1963, 1965, 1966). However, the original descriptions were based on material from other countries, with only a few exceptions: e.g. *Trabeculus flemingi* Timmermann, 1959; *Halipeurus consimilis* Timmermann, 1960; *Halipeurus falsus pacificus* Edwards, 1961; *Austromenopon stammeri* Timmermann, 1963.

The first louse species described from a New Zealand bird by an entomologist resident in this country was *Rallicola (Rallicola) takahe* Holloway, 1956. The original type series was collected from the South Island takahe, a flightless rail that had been regarded as extinct until rediscovered by G.B. Orbell in November 1948 (Checklist Committee 2010: 189). The author of the takahe louse, Beverley A. Holloway, was then a member of the Dominion Museum (now MONZ) staff. However, that was her only publication on lice as she then changed her research interest to other insect groups.

During the mid-1950s, Robert L.C. Pilgrim, Professor of Zoology at the University of Canterbury, (Christchurch, New Zealand) began collecting New Zealand lice and established contact with louse specialists of that time, in particular Dr Theresa Clay at the British Museum of Natural history (now Natural History Museum), Dr Kary C. Emerson in Oklahoma State University (Stillwater, Oklahoma, U.S.A.) and Professor Roger D. Price at the University of Minnesota (Minneapolis, U.S.A.). These useful connections provided Pilgrim with a reference collection properly prepared and identified, as well as several new louse species from endemic New Zealand birds (e.g. *Rallicola (Rallicola) harrisoni* Emerson, 1955 and *Pseudomenopon pilgrimi* Price 1974 from wekas; *Rallicola (Aptericola) pilgrimi* Clay, 1972 from the little spotted kiwi; *Colpocephalum pilgrimi* Price 1967 from the kea; and *Austromenopon bulleri* Price & Clay 1972 from Buller's albatross).

Other new species of lice from New Zealand birds described by foreign entomologists during the 1960s and 1970s were: *Pectinopygus punctatus* Timmermann, 1964, *Pectinopygus varius* Timmermann, 1964, *Ardeicola pilgrimi* Tandan, 1972, and *Patellinirmus novaeseelandiae* Tendeiro, 1972. The primary types of these species as well as those of *Austromenopon bulleri*, *Colpocephalum pilgrimi*, *Pseudomenopon pilgrimi*, and *Trabeculus flemingi* are held in the collection of the Canterbury Museum (Nicholls *et al.* 1998: 30).

Also, during the 1970s, a group of scientists based at the then New Zealand Ministry of Agriculture began research on the epidemiology and control of the lice introduced with domestic mammals into New Zealand, in particular those from sheep and cattle. Under the leadership of Peter R. Kettle and Allen C.G. Heath, they published

over 25 papers on mammal lice (see References below), as well as descriptions of two new species of lice from native birds: *Aquanirmus australis* Kettle, 1974 from the New Zealand dabchick, and *Cuculicola kui* Kettle, 1980 from the shining cuckoo.

Professor Pilgrim continued to build his collection and, from 1974 to 1976, I worked with him on bird lice, which had been the subject of my MSc degree thesis at the Universidad de Buenos Aires in Argentina (Palma 2011a). In April 1976, I was appointed as a research entomologist at the National Museum (now MONZ) and was able to continue with research on bird lice. In 1985, the “R.L.C. Pilgrim Collection”, then the largest and most comprehensive collection of New Zealand lice, was transferred to the National Museum of New Zealand (now MONZ). It was combined with the museum’s louse collection, which had been growing steadily since my appointment. Other institutions holding considerable numbers of New Zealand lice in their collections are: the Natural History Museum (NHML, London, England), the Auckland Museum (Auckland, New Zealand), Landcare Research Ltd (NZAC, Auckland, New Zealand) and the Canterbury Museum (CMNZ, Christchurch, New Zealand).

Thus, from 1974 until the present, I have been working on the louse fauna of New Zealand, building up a national collection at MONZ, and publishing papers on various genera, especially those from seabirds, as well as faunistic accounts from several oceanic islands. My work has been both facilitated and enriched by having collaborated with a great number of colleagues whose expertise greatly increased the quality of the resulting publications. All their names are listed as my coauthors in the References section below, but some deserve an especial mention. They are: the late Robert L.C. Pilgrim, the late M. Durno Murray, Roger D. Price (Arkansas, U.S.A.), Donald S. Horning (New South Wales, Australia), Terry D. Galloway (Winnipeg, Manitoba, Canada), Oldřich Sychra (Brno, Czech Republic), Michel P. Valim (Brazil), Jens-Kjeld Jensen (Nólsoy, Faroe Islands), and Adrian M. Paterson (Lincoln, Canterbury, New Zealand).

Although morphological descriptions of new species and generic revisions of New Zealand lice have continued into the 3<sup>rd</sup> Millennium (e.g. Palma & Pilgrim 2002; Palma & Price 2000, 2004, 2005; Valim & Palma 2013, 2015), the history of phthirapteran research is now characterised by molecular and phylogenetic studies of genera and generic complexes. Not surprisingly, several studies have dealt with lice from New Zealand seabirds, partly as a consequence of the comprehensive collections held in MONZ and also thanks to a group of young scientists who have taken the new techniques with great interest. Thus, several papers on the following genera have so far been published: Page *et al.* (2004) on the *Philoceanus* complex, Banks *et al.* (2006) on *Austrogoniodes*, Hughes *et al.* (2007) on the *Pectinopygus* complex, Hammer *et al.* (2010) on *Halipeurus*, and Bush *et al.* (2015, 2016) on the *Brueelia* complex.

Regarding the history of faunistic accounts of New Zealand lice, the first checklist was published by Wise (1977), who listed a total of 154 species and subspecies, but omitted the hosts. In 1982, Pilgrim & Palma published a list of lice from New Zealand birds in the format of a “host-lice list”, including 267 named species and subspecies; this list was updated by Palma (1999). Tenquist & Charleston (2001) listed all the lice recorded in New Zealand from introduced mammals, with a total of 35 species. The latest complete list of New Zealand lice from both birds and mammals—comprising 347 species and subspecies— was assembled by Palma (2010), and was included in a comprehensive inventory of the biodiversity of this country (Gordon 2010). A summary with statistics taken from the five abovementioned lists is shown in Table 1.

Although the main subject of this historical section is New Zealand, readers interested in information about louse researchers, collections and faunistic accounts in the rest of the world will find the following publications useful: Emerson (1967, 1972a,b, 1979), Lakshminarayana (1979), Ledger (1980), Złotorzycka & Modrzejewska (1988), Cocker (1989), Carriker (2001), Powell (2001), Mey (2003), Garfield (2007), Mey *et al.* (2007), and Martín-Mateo (2002, 2009).

## METHODS AND CONVENTIONS

In the Checklist, all taxa are listed in alphabetical order: families within each suborder, genera within each family, and species within each genus. Taxonomy and nomenclature of chewing lice follow Price *et al.* (2003), except where indicated; those of sucking lice follow Durden & Musser (1994a). Species qualified as “*sensu lato*” (14 in total) refer to populations either (1) with a range of measurements and morphological variation at present

considered different from the named species but not sufficiently distinct to warrant the erection of separate taxa (Pilgrim & Palma 1982: 2), or (2) without detailed systematic revisions available.

Synonymies and references for genera and species of lice are listed in chronological order but are not comprehensive. They include original citations, different generic combinations, and most of the references relevant to the New Zealand louse fauna. In the species synonymies, quotation marks “ ” around species or subspecies names indicate misidentifications made by the author(s) cited immediately after the names.

The geographic coverage of this Checklist is the same as that in the *Checklist of the Birds of New Zealand* (Checklist Committee 2010: 497, 500), with localities given as area codes taken from Crosby *et al.* (1976, 1998) and sequenced approximately from north to south, followed by outer island groups, and Antarctica. Since Crosby *et al.* (1976, 1998) do not include Norfolk and Macquarie Islands, these localities are given in full. Details of punctual localities are not relevant in the case of parasites living on highly mobile hosts. It should be noted that louse species recorded from Norfolk Island are now included in this catalogue following the inclusion of this island in the New Zealand Subregion (Checklist Committee 2010: 2).

This catalogue is primarily based on published articles and books, with additional data taken from collections held in the Museum of New Zealand Te Papa Tongarewa (MONZ) and in the New Zealand Arthropod Collection (NZAC). Several other institutions are included as repositories of primary types (see below). The material examined of new hitherto unpublished records, and published records from New Zealand localities given under each louse species refer to specimens collected from natural and regular hosts only. Specimens, localities and host records resulting from contaminations or straggling have been excluded (see Pilgrim & Palma 1982: 2). Voucher specimens of newly recorded species and of those identified to genus only have been slide-mounted in Canada balsam following the technique published by Palma (1978) and are held in the MONZ insect collection.

An attempt was made to include all literature citations to all references and records of all the lice known from New Zealand, with the exception of newspaper articles; however, involuntary omissions can be expected. Under “Significant references”, those publications that were deemed to be useful for various reasons are included.

Taxonomy, nomenclature, vernacular English names, and sequence of New Zealand bird taxa follow Checklist Committee (2010), those for other bird species and subspecies follow Dickinson (2003). In cases of nomenclatorial disagreement between these two latter sources, the Checklist Committee (2010) is followed. Taxonomy, nomenclature, vernacular English names, and sequence of New Zealand mammals follow King (2005).

The pattern used for *species/subspecies* entries is:

**Name of *species/subspecies* author/s, date of publication**

Original *genus* and *species* author/s, date: page number, figs.

Combinations of *species/subspecies* with other *genera*, date: page number, figs.

Synonymies relevant to New Zealand, date: page number, figs.

Current *genus* and *species/subspecies* date: page number, figs.

Primary type status, sex, and repository institution (Reference/s, if data differ from or were not given in the original description).

Type host/s:

Other host/s:

New Zealand host/s:

New Zealand locality/ies: [listed as area codes proposed by Crosby *et al.* 1976, 1998]

Geographic distribution: [given as those of the hosts, not from actual louse records]

New Zealand reference/s: [in chronological order]

Other significant reference/s: [in chronological order]

Material examined and repository/ies: [only given for new records]

Remarks: [additional data on: louse taxonomy, host status, host distribution, misidentifications, new records, ecology, etc.]

**Primary type specimens**

Repository institutions of primary type specimens (holotypes, lectotypes, syntypes, neotypes) are given for most species, except in a few cases where there is no information available. Several colleagues provided information

about primary types under their care, and their names are given as a “pers. comm.” where relevant. Dr Karla Schneider (MLUH) has kindly searched for many of the type specimens of species described by Nitzsch (1818), Burmeister (1838a), Nitzsch (in Giebel 1861; 1866), Rudow (1866; 1869; 1870), Giebel (1874), and Taschenberg (1882)—originally deposited in the Zoological Museum of Halle University—without success. With the exception of a few lectotypes and neotypes erected subsequently, all those types are listed here as “presumed lost” with references to relevant papers, in particular Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

In his eight parts comprising “*The Piaget Collection of Mallophaga*”, Thompson (1937–1939) does not specifically mention the words “holotype”, “lectotype” or “syntypes”. Therefore, with the exception of those taxa where Piaget had a single specimen for his description (the holotype) and those which had a lectotype subsequently designated, I regard the specimens listed by Thompson (1937–1939) are syntypes.

### Abbreviations for institutions holding primary types

AMNH	American Museum of Natural History, New York, U.S.A.
AMSA	Australian Museum, Sydney, Australia.
ANIC	Australian National Insect Collection, CSIRO, Canberra ACT, Australia.
BPBM	Bernice P. Bishop Museum, Honolulu, Hawaii, U.S.A.
CENPAT	Centro Nacional Patagónico, Puerto Madryn, Chubut, Argentina.
CZLP	Centro de Zoologia, Lisboa, Portugal.
CMNZ	Canterbury Museum, Christchurch, New Zealand.
CUIC	Cornell University Insect Collection, Department of Entomology, Cornell University, Ithaca, New York, U.S.A.
EMEC	Essig Museum, Division of Entomology, University of California, Berkeley, California, U.S.A.
FMLA	Fundación Miguel Lillo, Tucumán, Argentina.
GNHS	Göteborg Natural History Museum, Göteborg, Sweden.
MCZC	Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, U.S.A.
MLUH	Institut für Zoologie und Zoologische Sammlungen, Martin-Luther-Universität Halle-Wittenberg, Halle (Saale), Germany.
MNHN	Muséum National d’Histoire Naturelle, Paris, France.
MNHW	Museum of Natural History, University of Wrocław, Wrocław, Poland.
MONZ	Museum of New Zealand Te Papa Tongarewa, Wellington, New Zealand (formerly Dominion Museum 1907–1973, and National Museum of New Zealand 1973–1992).
MZUSP	Museu de Zoologia da Universidade de São Paulo, São Paulo, Brazil.
NHML	Natural History Museum, London, England.
NHMR	Naturhistorisches Museum im Thüringer, Rudolstadt, Germany.
NSMJ	National Science Museum, Tokyo, Japan.
NZAC	New Zealand Arthropod Collection, Landcare Research Ltd, Auckland, New Zealand (formerly Entomology Division Collection, DSIR).
OSUM	Ohio State University Museum, Columbus, Ohio, U.S.A.
SAIM	South African Institute for Medical Research, Johannesburg, South Africa.
SAMS	South African Museum, Cape Town, South Africa.
SDEI	Senckenberg Deutsches Entomologisches Institut, Muencheberg, Germany.
SMDV	Spencer Entomological Collection, Beaty Biodiversity Museum, Faculty of Science, The University of British Columbia, Vancouver, British Columbia, Canada.
USNM	United States National Museum of Natural History, Smithsonian Institution, Washington, D.C., U.S.A.
ZMAS	Zoological Institute of the Russian Academy of Sciences, Saint Petersburg, Russia.
ZMHG	Zoologisches Institut und Zoologisches Museum, Hamburg, Germany.
ZMHU	Museum für Naturkunde, Leibniz Institut, Humboldt-Universität zu Berlin, Germany.

## THE CHECKLIST

### Order PHTHIRAPTERA Haeckel, 1896

Phthiraptera Haeckel 1896. *Systemat. Phylog.*: 703.

### Suborder AMBLYCERA Kellogg, 1896

Amblycera Kellogg, 1896a. *Proc. Calif. Acad. Sci.* 6: 68.

### Family BOOPIIDAE Mjöberg, 1910

Boopiidae Mjöberg, 1910a. *Arkiv Zool.* 6(13): 21. Type genus: *Boopia* Piaget, 1880.

### Genus *Boopia* Piaget, 1880

*Boopia* Piaget, 1880. *Pédiculines*: 599. Type species: *Boopia tarsata* Piaget, 1880 (by monotypy).

### *Boopia notafusca* Le Souëf, 1902

Figs 1–2

*Boopia nota-fusca* Le Souëf, 1902a: 50, fig. 1.

*Boopia notafusca* Le Souëf, 1902; Kéler 1971: 31, fig. 90H.

*Boopia notafusca* Le Souëf, 1902; Palma 1996a: 162.

*Boopia notafusca* Le Souëf, 1902; Price *et al.* 2003: 73.

*Boopia notafusca* Le Souëf, 1902; Palma 2010: 407.

Lectotype ♂ in NHML (Barker 1996: 94).

Type host: *Wallabia bicolor* (Desmarest, 1804).

New Zealand host: *Petrogale penicillata* (J.E. Gray, 1825).

Other hosts: *Macropus robustus* Gould, 1841 and *Macropus giganteus* Shaw, 1790, but both need confirmation (Kéler 1971: 33).

New Zealand locality: AK (Kawau Island).

Geographic distribution: Australasia.

New Zealand references: Palma (1996a); Tenquist & Charleston (2001: 486); King (2005: 52); Palma (2010); Vermeulen *et al.* (2016: 71).

Other significant references: Tillyard (1926: 134, fig. O1); Kéler (1971); Murray & Calaby (1971: 83); Barker (1996: 94); Price *et al.* (2003).

Remarks: *Boopia notafusca* is native to Australia, and was introduced to New Zealand, probably with its type host, by human agency (King 2005: 50). The New Zealand record represents a host-association not yet recorded in Australia.

### Genus *Heterodoxus* Le Souëf & Bullen, 1902

*Heterodoxus* Le Souëf & Bullen, 1902b. *Vict. Naturalist* 18: 159. Type species: *Heterodoxus macropus* Le Souëf & Bullen, 1902b (by monotypy).

### *Heterodoxus ampullatus* Kéler, 1971

Figs 3–4

*Heterodoxus ampullatus* Kéler, 1971: 52, figs 34–38, 115G, 116P, 123.

*Heterodoxus ampullatus* Kéler, 1971; Palma 1996a: 161.

*Heterodoxus ampullatus* Kéler, 1971; Price *et al.* 2003: 74.

*Heterodoxus ampullatus* Kéler, 1971; Palma 2010: 407.

Holotype ♂ in ANIC (Barker 1996: 95).

Type host: *Petrogale penicillata* (J.E. Gray, 1825).

New Zealand host: *Petrogale penicillata* (J.E. Gray, 1825).

Other hosts: None.

New Zealand locality: AK (Kawau Island).

Geographic distribution: Australasia.

New Zealand references: Palma (1996a); Tenquist & Charleston (2001: 504); King (2005: 52); Palma (2010); Vermeulen *et al.* (2016: 71).

Other significant references: Murray & Calaby (1971: 84); Clay (1981a: 65, map 1); Barker & Close (1990: 1082, fig. 2); Barker (1996: 95); Price *et al.* (2003).

Remarks: *Heterodoxus ampullatus* is native to Australia, and was introduced to New Zealand with brush-tailed rock wallabies by human agency (King 2005: 50).

### Family GYROPIDAE Kellogg, 1896

Gyropidae Kellogg, 1896a. *Proc. Calif. Acad. Sci.* 6: 68. Type genus: *Gyropus* Nitzsch, 1818.

### Genus *Gliricola* Mjöberg, 1910

#### Subgenus *Gliricola* Mjöberg, 1910

*Gliricola* Mjöberg, 1910b. *Zool. Anz.* 35: 292. Type species: *Gyropus gracilis* (Nitzsch, 1818) = *Gliricola porcelli* (Schrank, 1781) (by monotypy).

#### *Gliricola (Gliricola) porcelli* (Schrank, 1781)

Figs 5–6

*Pediculus porcelli* Schrank, 1781: 500, pl. 1: fig. 1.

*Gyropus gracilis* Nitzsch, 1818: 304. Unnecessary *nomen novum* for *Pediculus porcelli* Schrank, 1781.

*Gyropus gracilis* Nitzsch, 1818; Mjöberg 1910b: 292, figs 7, 10–14.

*Gliricola porcelli* (Schrank, 1781); Hopkins & Clay 1952: 145.

*Gliricola porcelli* (Schrank, 1781); Tenquist & Charleston 2001: 500.

*Gliricola (Gliricola) porcelli* (Schrank, 1781); Price *et al.* 2003: 76.

*Gliricola porcelli* (Schrank, 1781); Palma 2010: 407.

Neotype ♂ in NHML (Clay & Hopkins 1954: 254).

Type host: *Cavia porcellus* (Linnaeus, 1758).

New Zealand host: *Cavia porcellus* (Linnaeus, 1758).

Other hosts: *Cavia aperea* Erxleben, 1777; *Cavia fulgida* Wagler, 1831; *Cavia tschudii* Fitzinger, 1857.

New Zealand localities: AK, NN, MC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Tenquist & Charleston (1981: 265); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Mjöberg (1910b); Werneck (1936: 397, figs 1–9); Séguy (1944: 53, figs 40–43); Webb (1946: 52); Symmons (1952: 379, fig. 25); Clay & Hopkins (1954: 254); Kéler (1957c: 97, figs 3b, 11, 13, 41, 42a); Emerson & Price (1975: 13, figs 23–26); Price (1987: 219); Barker (1996: 105); Martín-Mateo (2002: 140); Price *et al.* (2003); Palma & Jensen (2005: 50, 69).

Remarks: *Gliricola (Gliricola) porcelli* is native to South America and was introduced to New Zealand and other countries with guinea pigs by human agency (King 2005: 9).

### Genus *Gyropus* Nitzsch, 1818

*Gyropus* Nitzsch, 1818. *Germar's Mag. Entomol.* 3: 303. Type species: *Gyropus ovalis* Burmeister, 1838a (by subsequent designation).

#### *Gyropus ovalis* Burmeister, 1838

Figs 7–8

*Gyropus ovalis* Burmeister, 1838a: 443.

*Gyropus ovalis* Burmeister, 1838; Hopkins & Clay 1952: 161.

*Gyropus ovalis* Burmeister, 1838; Tenquist & Charleston 2001: 501.

*Gyropus ovalis* Burmeister, 1838; Price *et al.* 2003: 77.

*Gyropus ovalis* Burmeister, 1838; Palma 2010: 407.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Cavia porcellus* (Linnaeus, 1758).

New Zealand host: *Cavia porcellus* (Linnaeus, 1758).

Other hosts: *Cavia aperea* Erxleben, 1777; *Cavia fulgida* Wagler, 1831; *Cavia tschudii* Fitzinger, 1857.

New Zealand localities: AK, WN, NN, MC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Tenquist & Charleston (1981: 266); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Werneck (1936: 419, figs 34–39); Séguy (1944: 52, figs 37–39); Werneck (1948: 39); Webb (1946: 52); Symmons (1952: 379, figs 20–24); Kéler (1957c: 97, figs 3e, 20a, 42b,c); Emerson & Price (1975: 28, figs 71–74); Price (1987: 219); Barker (1996: 105); Martín-Mateo (2002: 139); Price *et al.* (2003); Bartlow *et al.* (2016: 222).

Remarks: *Gyropus ovalis* is native to South America and was introduced to New Zealand and other countries with guinea pigs by human agency (King 2005: 9).

### Family LAEMOBOTHRIDAE Mjöberg, 1910

Laemobothriidae Mjöberg, 1910a. *Arkiv Zool.* 6(13): 53. Type genus: *Laemobothrion* Nitzsch, 1818.

#### Genus *Laemobothrion* Nitzsch, 1818

##### Subgenus *Laemobothrion* Nitzsch, 1818

*Laemobothrion* Nitzsch, 1818. *German's Mag. Entomol.* 3: 301. Type species: *Laemobothrion maximum* (Scopoli, 1763) (by subsequent designation).

#### *Laemobothrion (Laemobothrion) tinnunculi* (Linnaeus, 1758)

Fig. 9

*Pediculus tinnunculi* Linnaeus, 1758: 612.

*Ricinus tinnunculi* (Linnaeus, 1758); Latreille 1804: 104.

*Laemobothrion tinnunculi* (Linnaeus, 1758); Hopkins & Clay 1952: 186.

*Laemobothrion tinnunculi* (Linnaeus, 1758); Nelson & Price 1965: 253, figs 4–5, 9, 11, 15.

*Laemobothrion tinnunculi* (Linnaeus, 1758); Pilgrim & Palma 1982: 17.

*Laemobothrion (Laemobothrion) tinnunculi* (Linnaeus, 1758); Murray *et al.* 1993: 960.

*Laemobothrion tinnunculi* (Linnaeus, 1758); Palma 2010: 407.

Neotype ♀ in NHML (Clay & Hopkins 1950: 230, pl. 1: fig. 1).

Type host: *Falco tinnunculus* Linnaeus, 1758.

New Zealand host: *Falco cenchroides cenchroides* Vigors & Horsfield, 1827.

Other hosts: At least 15 other species of *Falco* (see Price *et al.* 2003: 82).

New Zealand locality: WN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Clay & Hopkins (1950: 228, figs 1–5, pl. 1: fig. 1); Nelson & Price (1965); Lakshminarayana (1970: 132, figs 2c, 4d); Butler & O'Connor (1994: 451); Palma (1996b: 107); Martín-Mateo (2002: 136, fig. 43B,D,I); Price *et al.* (2003: 82).

Remarks: *Falco cenchroides cenchroides* breeds in Australia and is an infrequent straggler to New Zealand (Checklist Committee 2010: 174). Only one small sample of *Laemobothrion (Laemobothrion) tinnunculi* has been collected from one nankeen kestrel in New Zealand (voucher specimens in MONZ).

### Family MENOPONIDAE Mjöberg, 1910

Menoponidae Mjöberg, 1910a. *Arkiv Zool.* 6(13): 26. Type genus: *Menopon* Nitzsch, 1818.

**Genus *Actornithophilus* Ferris, 1916**

*Actornithophilus* Ferris, 1916a. *Canad. Entomol.* 48: 303. Type species: *Colpocephalum uniseriatum* Piaget, 1880 = *Actornithophilus uniseriatus* (Piaget, 1880) (by original designation).

*Clypeodon* Timmermann, 1954c. *Ann. Mag. Nat. Hist. (Ser. 12)* 7: 830. Type species: *Colpocephalum incisum* Piaget, 1880 = *Actornithophilus incisus* (Piaget, 1880) (by original designation).

***Actornithophilus bicolor* (Piaget, 1880)**

*Colpocephalum bicolor* Piaget, 1880: 561, pl. 47: fig. 1.

*Actornithophilus bicolor* (Piaget, 1880); Hopkins & Clay 1952: 21.

*Actornithophilus bicolor* (Piaget, 1880); Clay 1962: 239, 242, pl. 11: fig. 3.

*Actornithophilus bicolor* (Piaget, 1880); Pilgrim & Palma 1982: 21.

*Actornithophilus bicolor* (Piaget, 1880); Murray *et al.* 2006a: 1964.

*Actornithophilus bicolor* (Piaget, 1880); Palma 2010: 407.

Syntypes ♂♀, lost (Clay 1951a: 176).

Type host: *Arenaria interpres* (Linnaeus, 1758).

New Zealand host: *Arenaria interpres* (Linnaeus, 1758).

Other hosts: *Arenaria melanocephala* (Vigors, 1829).

New Zealand locality: MC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Emerson & Ward (1958: 50); Clay (1962); Hackman & Nyholm (1968: 75); Amerson & Emerson (1971: 14, 27); Butler & O'Connor (1994: 450); Price *et al.* (2003: 83); Martín-Mateo (2002: 71); Palma & Jensen (2005: 51, 64); Palma & Peck (2013: 9).

Remarks: Although the ruddy turnstone is a very common summer visitor to New Zealand (Checklist Committee 2010: 207), there are only two records of *Actornithophilus bicolor* from this country.

***Actornithophilus ceruleus* (Timmermann, 1954)**

Figs 10–11

*Clypeodon ceruleus* Timmermann, 1954c: 830, pl. 26: fig. c.

*Actornithophilus (Clypeodon) ceruleus* (Timmermann, 1954c): 832, figs 2cd.

*Clypeodon ceruleus* Timmermann, 1954; Timmermann 1957a: 111, fig. 84b, pl. 15: fig. b.

*Actornithophilus ceruleus* (Timmermann, 1954); Clay 1962: 201, 237.

*Actornithophilus ceruleus* (Timmermann, 1954); Nelson 1969: 199.

*Actornithophilus ceruleus* (Timmermann, 1954); Watt 1971: 233, 243, fig. 1.

*Actornithophilus ceruleus* (Timmermann, 1954); Wise 1977: 56.

*Actornithophilus ceruleus* (Timmermann, 1954); Pilgrim & Palma 1982: 23.

*Actornithophilus ceruleus* (Timmermann, 1954) *s. l.*; Pilgrim & Palma 1982: 23.

*Actornithophilus ceruleus* (Timmermann, 1954); Murray *et al.* 2006a: 1965.

*Actornithophilus ceruleus* (Timmermann, 1954); Palma 2010: 407.

Holotype ♂ in NHML (Timmermann 1954c: pl. 26: fig. c).

Type host: *Procelsterna cerulea cerulea* (Bennett, 1840).

New Zealand hosts: *Procelsterna cerulea albivitta* Bonaparte, 1856; *Anous minutus minutus* Boie, 1844.

Other hosts: *Anous tenuirostris* (Temminck, 1823); *Anous minutus melanogenys* Peters, 1934.

New Zealand localities: ND, BP, MC, KE, Norfolk Island.

Geographic distribution: Tropical and subtropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Nelson (1969); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1957a); Clay (1962); Amerson & Emerson (1971: 19, 27); Palma (1996b: 110); Price *et al.* (2003: 83); Silva *et al.* (2014: 942).

Remarks: The main New Zealand population of *Actornithophilus ceruleus* is in the Kermadec Islands, where the New Zealand hosts breed. Pilgrim & Palma (1982: 23) regarded the population of *Actornithophilus ceruleus* from *Anous*



*minutus minutus* as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted.

### ***Actornithophilus grandiceps* (Piaget, 1880)**

*Colpocephalum grandiceps* Piaget, 1880: 558, pl. 46: fig. 7.  
*Actornithophilus grandiceps* (Piaget, 1880); Hopkins & Clay 1952: 21.  
*Actornithophilus grandiceps* (Piaget, 1880); Clay 1962: 226, 239, fig. 66.  
*Actornithophilus grandiceps*; Baker 1974: 20.  
*Actornithophilus grandiceps* (Piaget, 1880); Pilgrim & Palma 1982: 19.  
*Actornithophilus grandiceps* (Piaget, 1880); Murray *et al.* 1993: 961.  
*Actornithophilus grandiceps* (Piaget, 1880); Palma 2010: 407.

Lectotype ♂ in NHML (Clay 1951a: 182).

Type host: *Haematopus ostralegus* Linnaeus, 1758.

New Zealand hosts: *Haematopus finschi* Martens, 1897; *Haematopus unicolor* J.R. Forster, 1844; *Haematopus chathamensis* Hartert, 1927.

Other hosts: *Haematopus ater* Vieillot & Oudart, 1825; *Haematopus bachmani* Audubon, 1838; *Haematopus fuliginosus* Gould, 1845; *Haematopus longirostris* Vieillot, 1817; *Ibidorhyncha struthersii* Vigors, 1832.

New Zealand localities: ND, AK, SD, MB, NN, NC, MC, SC, WD, CO, DN, SL, CH.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Baker (1974); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 223); Palma (2010).

Other significant references: Clay (1962); Green & Palma (1991: 4, 32); Forrester *et al.* (1995: 25); Palma (1996b: 110); Price *et al.* (2003: 83); Palma & Jensen (2005: 51, 63); Palma & Peck (2013: 10).

Remarks: *Actornithophilus grandiceps* is widespread on most species of oystercatchers, and found in large numbers per host.

### ***Actornithophilus hoplopteri* (Mjöberg, 1910) *sensu lato***

*Colpocephalum hoplopteri* Mjöberg, 1910a: 40, fig. 24, pl. 5: fig. 3.  
*Actornithophilus hoplopteri* (Mjöberg, 1910); Hopkins & Clay 1952: 21.  
*Actornithophilus hoplopteri* (Mjöberg, 1910) *s. l.*; Clay 1962: 202, 238, 240, fig. 1, pl. 7: fig. 1, pl. 9: fig. 4.  
*Actornithophilus hoplopteri* (Mjöberg, 1910) *s. l.*; Pilgrim & Palma 1982: 20.  
*Actornithophilus hoplopteri* (Mjöberg, 1910); Murray *et al.* 1993: 962.  
*Actornithophilus hoplopteri* (Mjöberg, 1910); Palma 2010: 407.

Syntypes ♀♀, probably lost (Daniel Gustafsson pers. comm. August 2012).

Type host: *Vanellus spinosus* (Linnaeus, 1758).

New Zealand host: *Vanellus miles novaehollandiae* Stephens, 1819.

Other hosts: *Charadrius vociferus* Linnaeus, 1758; and 10 other species of *Vanellus* (see Price *et al.* 2003: 83).

New Zealand localities: NN, NC, MC, SC, WD, CO, DN, SL.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 221); Palma (2010).

Other significant references: Clay (1962); Green & Palma (1991: 4, 32); Forrester *et al.* (1995: 25); Palma (1996b: 110); Price *et al.* (2003: 83).

Remarks: *Actornithophilus hoplopteri* is a widespread and frequently collected species, showing some morphological variability among populations from different hosts (Clay 1962: 203, fig. 1); hence, I regard the New Zealand population as “*sensu lato*”.

### ***Actornithophilus limosae* (Kellogg, 1908)**

“*Colpocephalum ochraceum*” Grube, 1851: 490 (not *Colpocephalum ochraceum* Nitzsch, 1818).  
*Colpocephalum limosae* Kellogg, 1908: 56. *Nomen novum* for *Colpocephalum ochraceum* Grube, 1951.  
*Actornithophilus limosae* (Kellogg, 1908); Hopkins & Clay 1952: 22.  
*Actornithophilus limosae* (Kellogg, 1908); Clay (1962: 224, 239, fig. 68, pl. 8: figs 1–2).

*Actornithophilus limosae* (Kellogg, 1908); Watt 1971: 233, 243.

*Actornithophilus limosae* (Kellogg, 1908); Wise 1977: 56.

*Actornithophilus limosae* (Kellogg, 1908); Pilgrim & Palma 1982: 21.

*Actornithophilus limosae* (Kellogg, 1908); Murray *et al.* 2006a: 1964.

*Actornithophilus limosae* (Kellogg, 1908); Palma 2010: 407.

Status, sex and repository of types unknown.

Type host: *Limosa lapponica lapponica* (Linnaeus, 1758).

New Zealand host: *Limosa lapponica baueri* Naumann, 1836.

Other hosts: *Limosa fedoa* (Linnaeus, 1758); *Limosa haemastica* (Linnaeus, 1758).

New Zealand localities: ND, SD, MB, WD, SL, KE, SN, CA.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay (1962); Amerson & Emerson (1971: 14, 27); Price *et al.* (2003: 84).

Remarks: *Actornithophilus limosae* is a relatively widespread species, but found in small numbers per host. The eastern bar-tailed godwit is the most numerous wader that visits New Zealand every year (Checklist Committee 2010: 203).

### ***Actornithophilus ochraceus* (Nitzsch, 1818) *sensu lato***

“*Pulex avis pluvialis*” Redi, 1668: pl. 2: upper fig.

*Liotheum (Colpocephalum) ochraceum* Nitzsch, 1818: 299. *Nomen novum* for “*Pulex avis pluvialis*” Redi, 1668.

*Colpocephalum timidum* Kellogg, 1896a: 145, pl. 12: fig. 6.

*Colpocephalum timidum* Kellogg, 1896; Johnston & Harrison 1912: 364.

*Actornithophilus timidus* (Kellogg, 1896); Thompson 1938a: 208.

*Actornithophilus ochraceus* (Nitzsch, 1818); Hopkins & Clay 1952: 22.

*Actornithophilus timidus* (Kellogg, 1896); Hopkins & Clay 1952: 24.

*Actornithophilus ochraceus* (Nitzsch, 1818); Clay 1962: 203, 240, figs 1–2, 25–28, 55–56, 63, pl. 9: fig. 2, pl. 10: fig. 3.

*Actornithophilus timidus* (Kellogg, 1896); Watt 1971: 233, 243.

*Actornithophilus timidus* (Kellogg, 1896); Wise 1977: 56.

*Actornithophilus ochraceus* (Nitzsch, 1818) *s. l.*; Pilgrim & Palma 1982: 20.

*Actornithophilus ochraceus* (Nitzsch, 1818); Murray *et al.* 1993: 962.

*Actornithophilus ochraceus* (Nitzsch, 1818); Palma 2010: 407.

Neotype ♂ in NHML (Clay 1962: 204).

Type host: *Pluvialis apricaria* (Linnaeus, 1758).

New Zealand hosts: *Pluvialis fulva* (J.F. Gmelin, 1789); *Charadrius obscurus obscurus* (J.F. Gmelin, 1789); *Charadrius obscurus aquilonius* Dowding, 1994.

Other hosts: *Pluvialis dominicus* (Statius Müller, 1776); *Pluvialis squatarola* (Linnaeus, 1758); at least 14 other species of *Charadrius* (see Price *et al.* 2003: 84).

New Zealand localities: ND, AK, SL, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Johnston & Harrison (1912: 363); Thompson (1938a); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Clay (1958c: 144); Clay & Hopkins (1960: 45); Clay (1962); Amerson & Emerson (1971: 13, 27); Moreby (1976: 93); Palma (1996b: 111); Price *et al.* (2003: 84); Palma & Jensen (2005: 51, 63).

Remarks: *Actornithophilus ochraceus* is a widespread and frequently collected species, showing some morphological variability among populations from different hosts (Clay 1962: 202, fig. 1); hence, I regard the New Zealand population as “*sensu lato*”. *Charadrius obscurus aquilonius* is a new host record for *Actornithophilus ochraceus* (voucher specimens in MONZ).

### ***Actornithophilus pediculoides* (Mjöberg, 1910)**

*Colpocephalum pediculoides* Mjöberg, 1910a: 44, pl. 2: fig. 6.

*Rediella pediculoides* (Mjöberg, 1910); Hopkins & Clay 1952: 322.

*Longimenopon pediculoides* (Mjöberg, 1910); Emerson 1956b: 296, figs 1–4.

*Actornithophilus pediculoides* (Mjöberg, 1910); Clay 1962: 191, 238.

*Actornithophilus pediculoides* (Mjöberg, 1910); Pilgrim & Palma 1982: 21.

*Actornithophilus pediculoides* (Mjöberg, 1910); Murray *et al.* 2006a: 1964.

*Actornithophilus pediculoides* (Mjöberg, 1910); Palma 2010: 407.

Syntypes ♂♀ in GNHS (Daniel Gustafsson pers. comm. August 2012).

Type host: *Arenaria interpres* (Linnaeus, 1758).

New Zealand host: *Arenaria interpres* (Linnaeus, 1758).

Other hosts: None.

New Zealand locality: MC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Emerson (1956b); Emerson & Ward (1958: 51); Clay (1962); Price *et al.* (2003: 84).

Remarks: *Actornithophilus pediculoides* lives mainly inside the rachis of primary wing feathers of the ruddy turnstone, its only host (see Clay 1962: 192). Although *Arenaria interpres* is a very common summer visitor to New Zealand (Checklist Committee 2010: 207), there is only a single record of *Actornithophilus pediculoides* from this country.

### ***Actornithophilus piceus lari* (Packard, 1870)**

*Colpocephalum lari* Packard, 1870: 96, pl. 1: fig. 1.

*Actornithophilus lari* (Packard, 1870); Hopkins & Clay 1952: 22.

*Actornithophilus piceus lari* (Packard, 1870); Timmermann 1954c: 839.

*Actornithophilus piceus* (Denny, 1842) *sens. lat.*; Clay 1962: 201, 237.

*Actornithophilus piceus* (Denny, 1842) *sens. lat.*; Clay & Moreby 1967: 158, 169, figs 53, 62.

*Actornithophilus piceus* (Denny, 1842) *s. l.*; Pilgrim & Palma 1982: 22.

*Actornithophilus piceus lari* (Packard, 1870); Palma 1996b: 112.

*Actornithophilus piceus lari* (Packard, 1870); Palma & Horning 2002: 5, 17.

*Actornithophilus piceus lari* (Packard, 1870); Price *et al.* 2003: 84.

*Actornithophilus piceus lari* (Packard, 1870); Murray *et al.* 2006a: 1965.

“*Actornithophilus piceus* (Packard, 1870)””; Palma 2010: 407. Error for *Actornithophilus piceus lari* (Packard, 1870).

Status, sex and repository of types unknown.

Type host: *Larus marinus* Linnaeus, 1758.

New Zealand hosts: *Larus dominicanus dominicanus* Lichtenstein, 1823; *Larus novaehollandiae scopulinus* J.R. Forster, 1843.

Other hosts: At least 23 other species of *Larus* (see Price *et al.* 2003: 84); *Pagophila eburnea* (Phipps, 1774); *Rhodostethia rosea* (Macgillivray, 1924); *Rissa brevirostris* (Bruch, 1853); *Rissa tridactyla* (Linnaeus, 1758); *Xema sabini* (Sabine, 1819).

New Zealand localities: AK, WN, NC, MC, SC, SL, Macquarie Island.

Geographic distribution: Cosmopolitan.

New Zealand references: Pilgrim & Palma (1982); Palma (1996b); Palma & Horning (2002); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1954c); Kéler (1957c: fig. 21a); Timmermann (1957a: 102, pl. 14: fig. d); Clay (1962); Clay & Moreby (1967); Green & Palma (1991: 4, 33); Forrester *et al.* (1995: 29); Palma (1996b: 112); Price *et al.* (2003); González-Acuña *et al.* (2011: 300).

Remarks: Although widespread on many host species, *Actornithophilus piceus lari* has a low prevalence of infestation on New Zealand gulls.

### ***Actornithophilus sedes* Eichler, 1944**

### **New Record**

*Actornithophilus sedes* Eichler, 1944a: 56.

*Actornithophilus sedes* Eichler, 1944; Clay 1962: 197, 240, fig. 64, pl. 4: fig. 4, pl. 11: fig. 2.

*Actornithophilus* sp.: Pilgrim & Palma 1982: 22.

*Actornithophilus sedes* Eichler, 1944; Price *et al.* 2003: 85.

*Actornithophilus* sp.: Murray *et al.* 2006a: 1964.

Holotype, probably ♀, repository unknown.

Type host: *Glareola pratincola* (Linnaeus, 1766).

New Zealand host: *Glareola maldivarum* J.R. Forster, 1795.

Other hosts: None.

New Zealand locality: NN.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a).

Other significant references: Clay (1962); Price *et al.* (2003).

Material examined and repository: 1 ♀ (1 sample, MONZ).

Remarks: This is the first record of *Actornithophilus sedes* for New Zealand, because the New Zealand references cited above reported this louse as “*Actornithophilus* sp.” only. Its host, the oriental pratincole is a rare visitor to New Zealand (Checklist Committee 2010: 223)

### ***Actornithophilus spinulosus* (Piaget, 1880)**

*Colpocephalum spinulosum* Piaget, 1880: 563, pl. 47: fig. 3.

*Actornithophilus spinulosus* (Piaget, 1880); Hopkins & Clay 1952: 23.

*Actornithophilus spinulosus* (Piaget, 1880); Clay 1962: 224, 243, fig. 67, pl. 8: fig. 3.

*Actornithophilus spinulosus* (Piaget, 1880); Pilgrim & Palma 1982: 20.

*Actornithophilus spinulosus* (Piaget, 1880); Murray *et al.* 2006a: 1964.

*Actornithophilus spinulosus* (Piaget, 1880); Palma 2010: 407.

Lectotype ♂ in NHML (Clay 1953b: 653).

Type host: *Limosa limosa limosa* (Linnaeus, 1758).

New Zealand host: *Limosa limosa melanuroides* Gould, 1846.

Other hosts: None.

New Zealand locality: AU.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay (1962); Price *et al.* (2003: 85); Adam (2007: 165).

Remarks: Despite the widespread distribution of its hosts, *Actornithophilus spinulosus* has been recorded from only a few localities around the world. The Asiatic black-tailed godwit is an uncommon annual visitor to New Zealand (Checklist Committee 2010: 203).

### ***Actornithophilus umbrinus* (Burmeister, 1838)**

*Colpocephalum umbrinum* Burmeister, 1838a: 438.

*Colpocephalum umbrinum* Piaget, 1880: 556, pl. 46: fig. 6. Preoccupied by *Colpocephalum umbrinum* Burmeister, 1838.

*Colpocephalum umbrosum* Harrison, 1916: 56. Unnecessary *nomen novum* for *Colpocephalum umbrinum* Piaget, 1880.

*Actornithophilus umbrinus* (Burmeister, 1838); Hopkins & Clay 1952: 24.

*Actornithophilus umbrosus* (Harrison, 1916); Hopkins & Clay 1952: 24.

*Actornithophilus umbrinus* (Burmeister, 1838); Clay 1962: 212, figs 6, 19, 57, 69, pl. 6: figs 4–6, pl. 10: fig. 1.

*Actornithophilus umbrinus* (Burmeister, 1838) *s. l.*; Pilgrim & Palma 1982: 21.

*Actornithophilus umbrinus* (Burmeister, 1838); Palma 1999: 380.

*Actornithophilus umbrinus* (Burmeister, 1838); Murray *et al.* 2006a: 1964.

*Actornithophilus umbrinus* (Burmeister, 1838); Palma 2010: 407.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Calidris ferruginea* (Pontoppidan, 1763).

New Zealand hosts: *Calidris canutus rogersi* (Mathews, 1913); *Calidris acuminata* (Horsfield, 1821).

Other hosts: At least 16 other species of *Calidris* (see Price *et al.* 2003: 85); *Tringa macularia* Linnaeus, 1766; *Phalaropus fulicarius* (Linnaeus, 1758); *Phalaropus lobatus* (Linnaeus, 1758); *Phalaropus tricolor* (Vieillot, 1819); *Tryngites subruficollis* (Vieillot, 1819).

New Zealand localities: NC, MC, SC, KE, CA.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Palma (1999); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay (1962); Hackman & Nyholm (1968: 75); Amerson & Emerson (1971: 15, 28); Green & Palma (1991: 4, 33); Hunter & Colwell (1994: 402); Forrester *et al.* (1995: 26); Palma (1996b: 112); Price *et al.* (2003: 85); Palma & Jensen (2005: 51, 63).

Remarks: *Actornithophilus umbrinus* is a frequently collected species widespread on many species of waders. Pilgrim & Palma (1982: 23) regarded the population of *Actornithophilus umbrinus* from *Calidris canutus* as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted.

### Genus *Amyrsidea* Ewing, 1927

*Amyrsidea* Ewing, 1927. *Jour. Wash. Acad. Sci.* 17: 90. Type species: *Menopon ventrale* Nitzsch [*in Giebel*], 1866 = *Amyrsidea* (*Amyrsidea*) *ventralis* (Nitzsch [*in Giebel*], 1866) (by original designation).

#### Subgenus *Argimenopon* Eichler, 1947

*Argimenopon* Eichler, 1947. *Arch. Zool.* 39A(2): 5. Type species: *Argimenopon polytrichum* Eichler, 1947 = *Amyrsidea* (*Argimenopon*) *polytrichum* Eichler, 1947 (by original designation).

### *Amyrsidea* (*Argimenopon*) *minuta* Emerson, 1961

Figs 12–13

*Amyrsidea minuta* Emerson, 1961a: 117, figs 1–3.

*Amyrsidea minuta* Emerson, 1961; Pilgrim & Palma 1982: 18.

*Amyrsidea* (*Argimenopon*) *minuta* Emerson, 1961; Scharf & Price 1983: 447, figs 18, 20.

*Amyrsidea minuta* Emerson, 1961; Murray *et al.* 1993: 960.

*Amyrsidea* (*Argimenopon*) *minuta* Emerson, 1961; Palma 2010: 407.

Holotype ♂ in SMDV (Karen Needham pers. comm. November 2014).

Type host: *Pavo cristatus* Linnaeus, 1758.

New Zealand host: *Pavo cristatus* Linnaeus, 1758.

Other host: *Pavo muticus* Linnaeus, 1766.

New Zealand localities: WA, NC, MC, SC.

Geographic distribution: Eurasia; North America; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Scharf & Price (1983); Palma (1996b: 113); Price *et al.* (2003: 87); Adam (2007: 157, figs 2a,b).

Remarks: *Amyrsidea* (*Argimenopon*) *minuta* is native to Asia, and was introduced to New Zealand and other countries with peafowl by human agency (Checklist Committee 2010: 28).

### *Amyrsidea* (*Argimenopon*) *perdicis* (Denny, 1842)

*Liotheum* (*Menopon*) *perdicis* Denny, 1842: 200, 225, pl. 21: fig. 9.

*Menopon megalosomum* Overgaard, 1943: 13, figs 5–6.

*Amyrsidea megalosoma* (Overgaard, 1943); Hopkins & Clay 1952: 28.

*Amyrsidea perdicis* (Denny, 1842); Hopkins & Clay 1952: 29.

*Amyrsidea perdicis* (Denny, 1842); Pilgrim & Palma 1982: 18.

*Amyrsidea* (*Argimenopon*) *perdicis* (Denny, 1842); Scharf & Price 1983: 445, figs 14, 16.

*Amyrsidea perdicis* (Denny, 1842); Murray *et al.* 1993: 960.

*Amyrsidea* (*A.*) *perdicis* (Denny, 1842); Palma 2010: 407.

Syntypes ♀♀ in NHML (Clay 1949b: 902).

Type host: *Perdix perdix perdix* (Linnaeus, 1758).

New Zealand hosts: *Perdix perdix perdix* (Linnaeus, 1758); *Phasianus colchicus* Linnaeus, 1758.

Other hosts: *Alectoris rufa* (Linnaeus, 1758); *Francolinus capensis* (J.F. Gmelin, 1789); *Syrnaticus reevesii* (J.E. Gray, 1829); *Bonasa umbellus* (Linnaeus, 1766); *Tympanuchus cupido* (Linnaeus, 1758); *Tympanuchus pallidicinctus* (Ridgway, 1873); *Tympanuchus phasianellus* (Linnaeus, 1758).

New Zealand localities: BP, HB, RI, WN, NC, MC, SC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Scharf & Price (1983); Modrzejewska & Złotorzycka (1987: 659, figs 1, 8); Kopociński *et al.* (1998: 81); Price *et al.* (2003: 87); Adam (2007: 158, figs 2c,d, 3a).

Remarks: *Amyrsidea (Argimenopon) perdicis* was introduced to New Zealand and other countries with grey partridges by human agency, but this bird species appears to have died out in New Zealand (Checklist Committee 2010: 348). However, *Amyrsidea (A.) perdicis* has spread onto a number of adventive hosts—including the common pheasant in New Zealand—due to the human practice of mixing various species of game birds in captivity (Scharf & Price 1983: 446).

### Genus *Ancistrona* Westwood, 1874

*Ancistrona* Westwood, 1874. *Thesaurus Entomol. Oxon.*: 197. Type species: *Ancistrona procellariae* Westwood, 1874 = *Ancistrona vagelli* (J.C. Fabricius, 1787) (by monotypy).

#### *Ancistrona vagelli* (J.C. Fabricius, 1787)

Figs 14–15

- Pediculus vagelli* J.C. Fabricius, 1787: 369.  
*Ancistrona procellariae* Westwood, 1874: 197, pl. 37: fig. 4.  
*Ancistrona gigas* Piaget, 1883: 152, pl. 9: fig. 1.  
*Ancistrona procellariae* Westwood, 1874; Johnston & Harrison 1912: 364.  
*Ancistrona vagelli* (J.C. Fabricius, 1787); Harrison 1937: 14.  
*Ancistrona vagelli* (J.C. Fabricius, 1787); Hopkins & Clay 1952: 36.  
*Ancistrona vagelli* (J.C. Fabricius, 1787); Timmermann 1965: 177, figs 114, 122.  
*Ancistrona* sp.; Watson 1967: 70.  
*Ancistrona* sp.?; Clay & Moreby 1967: 158, 168, fig. 59.  
*Ancistrona procellariae*; Clay & Moreby 1967: 177, fig. 52.  
*Ancistrona* sp.; Clay & Moreby 1970: 217, 218.  
*Ancistrona procellariae* Westwood, 1874; Watt 1971: 233, 243.  
*Ancistrona* sp.; Watt 1971: 233, 242.  
*Ancistrona procellariae* Westwood, 1874; Wise 1977: 56.  
*Ancistrona vagelli* (J.C. Fabricius, 1787); Wise 1977: 56.  
*Ancistrona* sp.; Horning *et al.* 1980: 4, 9.  
*Ancistrona* sp.; Pilgrim & Palma 1982: 7–13, 30, notes 17–18.  
*Ancistrona* sp.; Murray *et al.* 1990: 1369–1372.  
*Ancistrona vagelli* (J.C. Fabricius, 1787); Palma 1999: 375–378.  
*Ancistrona vagelli* (J.C. Fabricius, 1787); Palma & Horning 2002: 2 (figs), 5, 16.  
*Ancistrona vagelli* (J.C. Fabricius, 1787); Palma 2010: 407.

Neotype ♂ in NHML (Clay & Hopkins 1960: 6).

Type host: *Fulmarus glacialis glacialis* (Linnaeus, 1758).

New Zealand hosts: *Fulmarus glacialoides* (A. Smith, 1840); *Thalassoica antarctica* (J.F. Gmelin, 1789); *Daption capense capense* (Linnaeus, 1758); *Daption capense australe* Mathews, 1913; *Lugensa brevirostris* (Lesson, 1833); *Pterodroma macroptera gouldi* (Hutton, 1869); *Pterodroma lessonii* (Garnot, 1826); *Pterodroma magentae* (Giglioli & Salvadori, 1869); *Pterodroma inexpectata* (J.R. Forster, 1844); *Pterodroma cervicalis* (Salvin, 1891); *Pterodroma nigripennis* (Rothschild, 1893); *Pterodroma cookii* (G.R. Gray, 1843); *Pterodroma longirostris* (Stejneger, 1888); *Halobaena caerulea* (J.F. Gmelin, 1789); *Pachyptila vittata* (G. Forster, 1777); *Pachyptila salvini salvini* (Mathews, 1912); *Pachyptila desolata* (J.F. Gmelin, 1789); *Pachyptila belcheri* (Mathews, 1912); *Pachyptila turtur* (Kuhl, 1820); *Pachyptila crassirostris crassirostris* (Mathews, 1912); *Pachyptila crassirostris*

*pyramidalis* Fleming, 1939; *Procellaria aequinoctialis* Linnaeus, 1758; *Procellaria westlandica* Falla, 1946; *Procellaria parkinsoni* G.R. Gray, 1862; *Procellaria cinerea* J.F. Gmelin, 1789; *Puffinus pacificus pacificus* (J.F. Gmelin, 1789); *Puffinus bulleri* Salvin, 1888; *Puffinus carneipes* Gould, 1844; *Puffinus griseus* (J.F. Gmelin, 1789); *Puffinus tenuirostris* (Temminck, 1835); *Puffinus huttoni* Mathews, 1912; *Puffinus assimilis kermadecensis* Murphy, 1927; *Puffinus assimilis haurakiensis* Fleming & Serventy, 1943; *Pelagodroma albiclunis* Murphy & Irving, 1951.

Other hosts: *Pterodroma arminjoniana* (Giglioli & Salvadori, 1869); *Pterodroma externa* (Salvin, 1875); *Pterodroma hypoleuca* (Salvin, 1888); *Pterodroma incerta* (Schlegel, 1863); *Pterodroma mollis* (Gould, 1844); *Pterodroma phaeopygia* (Salvin, 1876); *Puffinus creatopus* Coues, 1864; *Puffinus gravis* (O'Reilly, 1818); *Puffinus opisthomelas* Coues, 1864; *Puffinus puffinus* (Brünnich, 1764); *Pseudobulweria rostrata* (Peale, 1848); *Pelagodroma marina* (Latham, 1790); *Oceanites oceanicus* (Kuhl, 1820).

New Zealand localities: ND, CL, BP, WO, HB, TK, WI, WN, SD, MB, NN, NC, MC, SC, WD, CO, DN, SL, KE, CH, SI, BO, SN, AN, Macquarie Island.

Geographic distribution: Cosmopolitan.

New Zealand references: Johnston & Harrison (1912: 363); Harrison (1937); Thompson (1939: 15); Watson (1967); Clay & Moreby (1970); Gressitt (1970: 326); Watt (1971); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Palma & Pilgrim (1983: 148); Murray *et al.* (1990); Palma (1996b: 114); Paterson *et al.* (1999: 222); Palma (1999); Marris (2000: 188); Palma & Imber (2000: 229); Palma & Horning (2002); Greenslade (2006: figs 20.1–20.2); Palma (2010).

Other significant references: Kellogg (1908: 75, fig. 24); Kellogg (1914: 89); Eichler (1941a: 362, fig. 27); Séguy (1944: 159, figs 212); Kéler (1952: 209, figs 4–5); Symmons (1952: 371, figs 12–13); Séguy (1953: 599, figs 57–58); Clay & Hopkins (1955: 51); Kéler (1957c: figs 10, 34b); Tendeiro (1958: 448, pl. 4: figs 7–8); Clay & Hopkins (1960: 4, figs 1–6); Timmermann (1965); Clay & Moreby (1967); Bourgeois & Threlfall (1979: 1356); Fowler & Shaw (1990: 15); Green & Palma (1991: 4, 25); Forrester *et al.* (1995: 5); Foster *et al.* (1996: 85); Palma (1996b: 114); Furness & Palma (1992: 35, 39); Price *et al.* (2003: 89); Palma & Jensen (2005: 51, 60); Jensen & Palma (2005: 228); Hänel & Palma (2007: 112, 121, 130); Palma & Peck (2013: 11).

Remarks: *Ancistrona vagelli* is a monotypic louse species, recorded from a great number of petrel species but not abundant on any particular host. Although populations from different hosts show some variation in size, there are no significant features that would justify subdividing this species into more than one taxon.

### Genus *Apterygon* Clay, 1961

*Apterygon* Clay, 1961. *Ann. Mag. Nat. Hist.* (13) 3(33): 571. Type species: *Apterygon mirum* Clay, 1961 (by monotypy).  
Endemic to New Zealand.

### *Apterygon dumosum* Tandan, 1972

*Apterygon dumosum* Tandan, 1972a: 54, figs 1, 4, 5, 11–15, 20–30.

*Apterygon* sp. *incertae sedis*; Tandan 1972: 68.

*Apterygon dumosum* Tandan, 1972; Pilgrim & Palma 1982: 3.

*Apterygon dumosum* Tandan, 1972 *s. l.*; Pilgrim & Palma 1982: 3.

*Apterygon dumosum* Tandan, 1972; Murray *et al.* 1990: 1367.

*Apterygon dumosum* Tandan, 1972; Palma & Price 2004: 70, 73.

*Apterygon dumosum* Tandan, 1972; Palma 2010: 407.

Holotype ♀ in NZAC.

Type host: *Apteryx australis lawryi* Rothschild, 1893.

New Zealand hosts: *Apteryx australis australis* Shaw, 1813; *Apteryx australis lawryi* Rothschild, 1893; *Apteryx owenii* Gould, 1847.

Other hosts: None.

New Zealand localities: WN, MB, FD, SI.

Geographic distribution: New Zealand.

New Zealand references: Tandan (1972a); Reid & Williams (1975: 324); Wise (1977: 56); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1991a: 318); Baker *et al.* (1995: 8256); Burbidge *et al.* (2003: 174); Palma & Price (2004); Sales (2005: 15); Heath (2010: 151); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003: 89).

Remarks: *Apterygon dumosum* is an endemic and “at risk” species (Buckley *et al.* 2012). Pilgrim & Palma (1982: 3) regarded the populations of *Apterygon dumosum* from *Apteryx australis australis* and *Ap. owenii* as somewhat different from that of the type host, and qualified them as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted. Furthermore, it is likely that the population of *Ap. dumosum* from *Apteryx owenii* was introduced to Kapiti Island from the South Island with its host by human agency (Checklist Committee 2010: 22), and may also be the result of a host switch from *Apteryx australis australis*.

### ***Apterygon hintoni* Clay, 1966**

*Apterygon hintoni* Clay, 1966a: 292, figs 1–3, pl. 5: fig. 2, pl. 6: figs 2, 4.

*Apterygon hintoni* Clay, 1966; Tandan 1972a: 59, figs 2, 6–10, 16–18.

*Apterygon hintoni* Clay, 1966; Pilgrim & Palma 1982: 3.

*Apterygon hintoni* Clay, 1966; Murray *et al.* 1990: 1367.

*Apterygon hintoni* Clay, 1966; Palma 2010: 407.

Holotype ♂ in NZAC (Tandan 1972a: 64).

Type host: *Apteryx haastii* Potts, 1872.

New Zealand host: *Apteryx haastii* Potts, 1872.

Other hosts: None.

New Zealand localities: NN, WD, BR.

Geographic distribution: South Island, New Zealand.

New Zealand references: Clay (1966a: 292); Pilgrim (1970: 75); Tandan (1972a); Reid & Williams (1975: 324); Wise (1977: 56); Pilgrim & Palma (1982); Murray *et al.* (1990); Baker *et al.* (1995: 8256); Burbidge *et al.* (2003: 174); Palma & Price (2004: 70, 73); Sales (2005: 15); Heath (2010: 151); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003: 89).

Remarks: *Apterygon hintoni* is an endemic and vulnerable species (Buckley *et al.* 2012), exclusively parasitic on great spotted kiwis.

### ***Apterygon mirum* Clay, 1961**

*Apterygon mirum* Clay, 1961: 574, figs 1–8, pl. 9: figs 1–3.

*Apterygon mirum* Clay, 1961; Clay 1966a: 292, pl. 5: fig. 1, pl. 6: figs 1, 3.

*Apterygon mirum* Clay, 1961; Tandan 1972a: 65, figs 3, 19, 31–42.

*Apterygon mirum* Clay, 1961; Pilgrim & Palma 1982: 3.

*Apterygon mirum* Clay, 1961; Murray *et al.* 1990: 1367.

*Apterygon mirum* Clay, 1961; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Apteryx mantelli* Bartlett, 1852.

New Zealand host: *Apteryx mantelli* Bartlett, 1852.

Other hosts: None

New Zealand localities: ND, CL, BP, GB, TK, WA (captive).

Geographic distribution: North Island, New Zealand.

New Zealand references: Clay (1961); Clay (1966a); Pilgrim (1970: 74); Tandan (1972a); Reid & Williams (1975: 324); Wise (1977: 56); Pilgrim & Palma (1982); Murray *et al.* (1990); Baker *et al.* (1995: 8256); Burbidge *et al.* (2003: 174); Palma & Price (2004: 70, 73); Sales (2005: 15); Heath (2010: 151); Palma (2010: 407); Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003: 89).

Remarks: *Apterygon mirum* is an endemic and vulnerable species (Buckley *et al.* 2012), exclusively parasitic on North Island brown kiwis.



***Apterygon okarito* Palma & Price, 2004**

Figs 16–17

*Apterygon* sp.; Pilgrim & Palma 1982: 3.*Apterygon* sp.; Murray *et al.* 1990: 1367.*Apterygon* new species; Baker *et al.* 1995: 8256.*Apterygon* new species; Burbidge *et al.* 2003: 172, 174.*Apterygon okarito* Palma & Price, 2004: 68, figs 1–4.*Apterygon okarito* Palma & Price, 2004; Palma 2010: lower fig. p. 295, 407.

Holotype ♂ in MONZ.

Type host: *Apteryx rowi* Tennyson, Palma, Robertson, Worthy & Gill, 2003.New Zealand host: *Apteryx rowi* Tennyson, Palma, Robertson, Worthy & Gill, 2003.

Other hosts: None.

New Zealand locality: WD (Okarito).

Geographic distribution: Westland, South Island, New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Baker *et al.* (1995); Burbidge *et al.* (2003); Palma & Price (2004: 73); Sales (2005: 15); Checklist Committee (2010: 20); Heath (2010: 151); Palma (2010); Buckley *et al.* (2012: 137, App. 2).

Other significant references: None.

Remarks: *Apterygon okarito* is an endemic and critically threatened species (Buckley *et al.* 2012), exclusively parasitic on Okarito brown kiwis.**Genus *Austromenopon* Bedford, 1939***Austromenopon* Bedford, 1939. *Onderstepoort Jour. Vet. Sci. Animal Ind.* 12(1): 122. Type species: *Menopon crocatum* Nitzsch [*in* Giebel], 1866 = *Austromenopon crocatum* (Nitzsch [*in* Giebel], 1866) (by original designation).*Procellariophaga* Eichler, 1949a. *Boll. Soc. Entomol. Italiana* 79: 12. Type species: *Procellariophaga ossifragae* Eichler, 1949a = *Austromenopon ossifragae* (Eichler, 1949) (by original designation).***Austromenopon aegialitidis* (Durrant, 1906) *sensu lato****Menopon aegialitidis* Durrant, 1906: 529, fig. 1c.*Austromenopon aegialitidis* (Durrant, 1906); Hopkins & Clay 1952: 46.*Austromenopon aegialitidis* (Durrant, 1906) *s. l.*; Pilgrim & Palma 1982: 20.*Austromenopon aegialitidis* (Durrant, 1906); Murray *et al.* 1993: 962.*Austromenopon aegialitidis* (Durrant, 1906); Palma 1999: 380.*Austromenopon aegialitidis* (Durrant, 1906); Palma 2010: 407.

Holotype ♀ in OSUM, probably (Palma 1996b: 114).

Type host: *Charadrius vociferus* Linnaeus, 1758.New Zealand hosts: *Vanellus miles novaehollandiae* Stephens, 1819; *Charadrius bicinctus exilis* Falla, 1978.Other hosts: *Vanellus vanellus* (Linnaeus, 1758); *Vanellus coronatus* (Boddaert, 1783); at least 9 other species of *Charadrius* (see Price *et al.* 2003: 90).

New Zealand localities: NC, MC, SC, WD, AU.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 221); Palma (1999); Palma (2010).Other significant references: Clay (1959: 164, fig. 7); Mester (1971: 115, fig. 4); Green & Palma (1991: 4, 32); Forrester *et al.* (1995: 25); Palma (1996b: 114); Price *et al.* (2003: 90).Remarks: *Austromenopon aegialitidis* is a morphologically variable species from a wide range of hosts and in need of a detailed systematic study. Therefore, I regard the New Zealand populations as “*sensu lato*”. *Charadrius bicinctus exilis* is a new host record for *A. aegialitidis* (voucher specimens in MONZ).***Austromenopon affine* (Piaget, 1890)***Menopon affine* Piaget, 1890a: 248, pl. 10: fig. 3.

- Procellariphaga affinis* (Piaget, 1890); Eichler 1949b: 346.  
*Procellariphaga affinis* (Piaget, 1890); Hopkins & Clay 1952: 298.  
*Austromenopon affine* (Piaget, 1890); Timmermann 1965: 166.  
*Austromenopon affine* (Piaget, 1890); Clay & Moreby 1967: 159, 168, figs 72, 74.  
*Austromenopon affine* (Piaget, 1890); Price & Clay 1972: 488, figs 1–9.  
*Austromenopon affine* (Piaget, 1890); Pilgrim & Palma 1982: 5.  
*Austromenopon affine* (Piaget, 1890); Murray *et al.* 1990: 1368.  
*Austromenopon affine* (Piaget, 1890); Palma 2010: 407.

Lectotype ♀ in NHML (Clay 1949b: 816).

Type host: *Diomedea exulans* Linnaeus, 1758.

New Zealand hosts: *Diomedea exulans* Linnaeus, 1758; *Diomedea antipodensis antipodensis* Robertson & Warham, 1992; *Diomedea epomophora* Lesson, 1825; *Diomedea sanfordi* Murphy, 1917.

Other hosts: None.

New Zealand localities: WA, WN, SD, MB, NC, MC, SC, WD, CO, DN, CH, AN, CA.

Geographic distribution: Southern Hemisphere.

New Zealand references: Clay (1964a: 230); Gressitt (1964: 538); Price & Clay (1972); Wise (1977: 56); Pilgrim & Palma (1982); Murray *et al.* (1990: 1368); Marris (2000: 188); Palma (2001: 66, fig. 3); Palma (2010).

Other significant references: Timmermann (1965); Clay & Moreby (1967); Palma (1996b: 114); Price *et al.* (2003: 90).

Remarks: *Austromenopon affine* is a small species, exclusively parasitic on the larger albatross of the genus *Diomedea*.

#### ***Austromenopon atrofulvum* (Piaget, 1880)**

- Menopon atrofulvum* Piaget, 1880: 483, pl. 39: fig. 2.  
*Austromenopon atrofulvum* (Piaget, 1880); Hopkins & Clay 1952: 47.  
*Austromenopon atrofulvum* (Piaget, 1880); Watt 1971: 233, 243.  
*Austromenopon* sp.; Watt 1971: 233, 243.  
*Austromenopon atrofulvum* (Piaget, 1880) *s. l.*; Horning *et al.* 1980: 4, 11.  
*Austromenopon atrofulvum* (Piaget, 1880) *s. l.*; Pilgrim & Palma 1982: 22–23.  
*Austromenopon atrofulvum* (Piaget, 1880); Murray *et al.* 2006a: 1965.  
*Austromenopon atrofulvum* (Piaget, 1880); Palma 2010: 407.

Lectotype ♂ in NHML (Clay 1949b: 819).

Type host: “*Platalea leucorodia*”, in error (see Hopkins & Clay 1952: 47).

New Zealand hosts: *Procelsterna cerulea albivitta* Bonaparte, 1856; *Anous minutus minutus* Boie, 1844; *Hydroprogne caspia* (Pallas, 1770); *Onychoprion fuscatus serratus* (J.R. Forster, 1830); *Sterna vittata bethunei* Buller, 1896; *Sterna paradisaea* Pontoppidan, 1763; *Sterna striata* J.F. Gmelin, 1789.

Other hosts: *Anous tenuirostris* (Temminck, 1823); *Anous stolidus* (Linnaeus, 1758); *Gelochelidon nilotica* (J.F. Gmelin, 1789); three species of *Chlidonias*; at least 10 other species of *Sterna*, and five species of *Thalasseus* (see Price *et al.* 2003: 90).

New Zealand localities: AK, WN, NC, MC, SC, WD, CO, DN, KE, CH, SN.

Geographic distribution: Cosmopolitan.

New Zealand references: Watt (1971); Wise (1977: 56); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay (1949b: 819); Clay (1959: 163, figs 6, 35); Clay & Moreby (1967: 158, fig. 67); Amerson & Emerson (1971: 17, 28); Moreby (1976: 93); Green & Palma (1991: 4, 33); Forrester *et al.* (1995: 30); Palma (1996b: 115); Price *et al.* (2003: 90); Palma & Peck (2013: 12); Silva *et al.* (2014: 942).

Remarks: *Austromenopon atrofulvum* is the only species of *Austromenopon* parasitic on members of the Sternidae. Horning *et al.* (1980) and Pilgrim & Palma (1982) regarded the populations of *Austromenopon atrofulvum* from all New Zealand hosts as *sensu lato*; however, my examination of more samples, including some from other host species, shows that such qualification is not warranted. *Sterna paradisaea* is a new host record for this louse species in New Zealand (voucher specimens in MONZ).

***Austromenopon beckii* (Kellogg, 1906)**

*Menopon beckii* Kellogg, 1906: 322.

*Menopon beckii* Kellogg, 1906; Thompson 1938c: 460, figs 2, 3a,c.

*Austromenopon becki* [sic] (Kellogg, 1906); Hopkins & Clay 1952: 47.

*Austromenopon becki* [sic] (Kellogg, 1906); Watt 1971: 233, 243, fig. 2.

*Austromenopon beckii* (Kellogg, 1906); Wise 1977: 56.

*Austromenopon becki* [sic] (Kellogg, 1906) *s. l.*; Pilgrim & Palma 1982: 14.

*Austromenopon becki* [sic] (Kellogg, 1906); Murray *et al.* 1990: 1373.

*Austromenopon beckii* (Kellogg, 1906); Price *et al.* 2003: 90.

*Austromenopon beckii* (Kellogg, 1906); Palma 2010: 407.

Holotype ♀ in EMEC.

Type host: *Phaethon aethereus mesonauta* Peters, 1930.

New Zealand hosts: *Phaethon rubricauda* Boddaert, 1783; *Phaethon lepturus dorotheae* Mathews, 1913.

Other hosts: *Phaethon aethereus aethereus* Linnaeus, 1758; *Phaethon lepturus lepturus* Daudin, 1802; *Phaethon lepturus fulvus* J.F. Brandt, 1838.

New Zealand localities: ND, KE.

Geographic distribution: Tropical and subtropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Thompson (1938c); Carriker (1949: 18); Timmermann (1954d: 198, fig. 4); Palma (1996b: 115); Price *et al.* (2003); Palma & Peck (2013: 12).

Remarks: *Austromenopon beckii* is the only species of *Austromenopon* parasitic on members of the Phaethontiformes.

Pilgrim & Palma (1982: 14) regarded the population of *A. beckii* from *Phaethon rubricauda* as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of samples from four additional hosts shows that making such difference is not warranted. A sample of *A. beckii* from *Phaethon lepturus dorotheae* from Northland is a new host record for this louse in New Zealand (voucher specimens in MONZ).

***Austromenopon brevifimbriatum* (Piaget, 1880)**

*Menopon brevifimbriatum* Piaget, 1880: 499, pl. 41: fig. 1.

*Menopon numerosum* Kellogg, 1896a: 159, pl. 15: fig. 5.

*Procellariphaga daptionis* Eichler, 1949b: 344, figs 21–23.

*Procellariphaga brevifimbriata* (Piaget, 1880); Eichler 1949b: 346.

*Procellariphaga numerosa* [sic] (Kellogg, 1896); Eichler 1949b: 346.

*Procellariphaga brevifimbriata* (Piaget, 1880); Hopkins & Clay 1952: 298.

*Procellariphaga numerosus* [sic] (Kellogg, 1896); Hopkins & Clay 1952: 299.

*Austromenopon brevifimbriatum* (Piaget, 1880); Timmermann 1963: 410, fig. 5.

*Austromenopon daptionis* (Eichler, 1949); Timmermann 1963: 411.

*Austromenopon oschei* Timmermann, 1963: 412, fig. 6.

*Austromenopon brevifimbriatum* (Piaget, 1880); Timmermann 1965: 165, 169, fig. 105.

*Austromenopon daptionis* (Eichler, 1949); Timmermann 1965: 170.

*Austromenopon oschei* Timmermann, 1963; Timmermann 1965: 170, fig. 106.

*Austromenopon ?daptionis* (Eichler, 1949); Clay & Moreby 1967: 159, 168.

*Austromenopon oschei* Timmermann, 1963; Clay & Moreby 1967: 159, 168, figs 69, 71.

*Austromenopon brevifimbriatum* (Piaget, 1880); Price & Clay 1972: 494, figs 25–26, 28–33.

*Austromenopon brevifimbriatum* (Piaget, 1880); Pilgrim & Palma 1982: 7.

*Austromenopon brevifimbriatum* (Piaget, 1880); Murray *et al.* 1990: 1369.

*Austromenopon brevifimbriatum* (Piaget, 1880); Palma 2010: 407.

Lectotype ♂ in NHML (Clay 1949b: 820).

Type host: *Fulmarus glacialis* (Linnaeus, 1758).

New Zealand hosts: *Fulmarus glacialis* (A. Smith, 1840); *Thalassoica antarctica* (J.F. Gmelin, 1789); *Daption capense capense* (Linnaeus, 1758).

Other host: *Pagodroma nivea* (G. Forster, 1777).

New Zealand localities: ND, AK, TK, WI, WN, SD, MB, NN, NC, MC, SC, CO, DN, SL, Macquarie Island.

Geographic distribution: All oceans at high latitudes. Subarctic Islands; north-west Europe; Greenland; Iceland; Australasia; Subantarctic Islands; Antarctica.

New Zealand references: Pilgrim & Palma (1982: 7); Murray *et al.* (1990); Palma (1996b: 115); Paterson *et al.* (1999: 222); Palma & Horning (2002: 5, 16); Palma (2010).

Other significant references: Timmermann (1963; 1965); Clay & Moreby (1967); Price & Clay (1972); Green & Palma (1991: 4, 26); Price *et al.* (2003: 90); Palma & Jensen (2005: 51, 60).

Remarks: *Austromenopon brevifimbriatum* is highly prevalent on all its hosts, except on *Pagodroma nivea*. Both *Fulmarus glacialis* and *Thalassoica antarctica* breed on the coast of Antarctica and are stragglers to New Zealand seas (Checklist Committee 2010: 80, 82).

### ***Austromenopon bulweriae* Timmermann, 1963**

*Austromenopon bulweriae* Timmermann, 1963: 420, fig. 10.

*Austromenopon bulweriae* Timmermann, 1963; Timmermann 1965: 174, fig. 109.

*Austromenopon bulweriae* Timmermann, 1963; Price & Clay 1972: 493, figs 22–23.

*Austromenopon bulweriae* Timmermann, 1963; Palma 1999: 376, 383, note 3.

*Austromenopon bulweriae* Timmermann, 1963; Price *et al.* 2003: 90.

*Austromenopon bulweriae* Timmermann, 1963; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Bulweria bulwerii* (Jardine & Selby, 1828).

New Zealand host: *Bulweria bulwerii* (Jardine & Selby, 1828).

Other hosts: None.

New Zealand locality: WN.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Timmermann (1965); Price & Clay (1972); Price *et al.* (2003).

Remarks: *Austromenopon bulweriae* is exclusively parasitic on Bulwer's petrels. This host, with a single record until 2013, is considered as a very rare straggler to New Zealand (Checklist Committee 2010: 107). A second record occurred in Canterbury in January 2014, but no *Austromenopon bulweriae*, was found on it.

### ***Austromenopon cursorium* (Giebel, 1874)**

### **New Record**

*Menopon cursorius* Giebel, 1874: 296.

*Austromenopon cursorius* [sic] (Giebel, 1874); Hopkins & Clay 1952: 47.

*Austromenopon cursorius* [sic] (Giebel, 1874); Clay 1959: 162, 167, fig. 4.

*Austromenopon* sp.: Pilgrim & Palma 1982: 22.

*Austromenopon cursorium* (Giebel, 1874); Price *et al.* 2003: 90.

*Austromenopon* sp.: Murray *et al.* 2006a: 1964.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Cursorius cursor* (Latham, 1787).

New Zealand host: *Glareola maldivarum* J.R. Forster, 1795.

Other host: *Cursorius coromandelicus* (J.F. Gmelin, 1789).

New Zealand locality: NN.

Geographic distribution: Asia; Australasia; Africa; Atlantic Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a).

Other significant references: Timmermann (1954d: 198, fig. 5, pl. 1: figs c, d); Clay (1959); Price *et al.* (2003).

Material examined and repository: 1♂, 3♀ (1 sample, MONZ).

Remarks: This is the first record of *Austromenopon cursorium* for New Zealand, because the New Zealand references cited above reported this louse as "*Austromenopon* sp." only. Its host, the oriental pratincole is a rare visitor to New Zealand (Checklist Committee 2010: 223)

***Austromenopon elliotti* Timmermann, 1954**

- Austromenopon elliotti* Timmermann, 1954d: 205, fig. 19, pl. 1: figs e,f.  
*Austromenopon elliotti* Timmermann, 1954; Clay 1964a: 230.  
*Austromenopon elliotti* Timmermann, 1954; Timmermann 1965: 177.  
*Austromenopon elliotti* Timmermann, 1954; Price & Clay 1972: 496, figs 39–41.  
*Austromenopon elliotti* Timmermann, 1954; Wise 1977: 56.  
*Austromenopon elliotti* Timmermann, 1954; Pilgrim & Palma 1982: 13.  
*Austromenopon elliotti* Timmermann, 1954; Murray *et al.* 1990: 1372.  
*Austromenopon elliotti* Timmermann, 1954; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Pelecanoides urinatrix* (J.F. Gmelin, 1789).

New Zealand hosts: *Pelecanoides urinatrix urinatrix* (J.F. Gmelin, 1789); *Pelecanoides urinatrix chathamensis* Murphy & Harper, 1916; *Pelecanoides urinatrix exsul* Salvin, 1896; *Pelecanoides georgicus* Murphy & Harper, 1916.

Other hosts: *Pelecanoides urinatrix dacunhae* Nicholl, 1906.

New Zealand localities: ND, CL, BP, GB, TO, WI, WA, WN, SD, MB, NC, MC, SC, CH, SI, SN, AN, CA.

Geographic distribution: Southern Hemisphere.

New Zealand references: Clay (1964a); Gressitt (1964: 538); Price & Clay (1972); Wise (1977); Horning *et al.* (1980: 4, 10); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Timmermann (1965); Payne & Prince (1979: 316); Green & Palma (1991: 4, 27); Furness & Palma (1992: 35, 42); Palma (1996b: 116); Price *et al.* (2003: 91); Hänel & Palma (2007: 112, 121, 131).

Remarks: *Austromenopon elliotti* is naturally and frequently found on diving petrels only. A record of this louse species from *Pterodroma inexpectata* in New Zealand (Price & Clay 1972: 497) is the result of natural straggling or contamination by human agency.

***Austromenopon enigki* Timmermann, 1963**

- Austromenopon enigki* Timmermann, 1963: 425, fig. 14.  
*Austromenopon enigki* Timmermann, 1963; Timmermann 1965: 177, fig. 113.  
*Austromenopon enigki* Timmermann, 1963; Price & Clay 1972: 493, fig. 27.  
*Austromenopon enigki* Timmermann, 1963; Pilgrim & Palma 1982: 13.  
*Austromenopon* sp.; Pilgrim & Palma 1982: 13.  
*Austromenopon enigki* Timmermann, 1963; Murray *et al.* 1990: 1372.  
*Austromenopon* sp.; Murray *et al.* 1990: 1372.  
*Austromenopon* species 1; Marris 2000: 188.  
*Austromenopon enigki* Timmermann, 1963; Price *et al.* 2003: 91.  
*Austromenopon enigki* Timmermann, 1963; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Pelagodroma marina* (Latham, 1790).

New Zealand hosts: *Pelagodroma marina dulciae* Mathews, 1912; *Pelagodroma marina maoriana* Mathews, 1912; *Fregetta tropica* (Gould, 1844); *Pealeornis maoriana* Mathews, 1932.

Other hosts: *Pelagodroma marina marina* (Latham, 1790); *Oceanites oceanicus exasperatus* Mathews, 1912; *Oceanites pincoyae* Harrison *et al.*, 2013; *Fregetta grallaria* (Vieillot, 1817).

New Zealand localities: ND, CL, WN, CH, AN.

Geographic distribution: Antarctica; Pacific, Atlantic and Indian Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 222); Marris (2000); Palma (2010).

Other significant references: Timmermann (1965); Price & Clay (1972); Green & Palma (1991: 5, 27); Palma (1996b: 116); Price *et al.* (2003); Harrison *et al.* (2013: 186).

Remarks: Records of *Austromenopon enigki* from *Fregetta tropica* (Antipodes Islands, New Zealand), *Fregetta grallaria* (Gough Island; Juan Fernández Islands), *Pealeornis maoriana* (Hauraki Gulf, New Zealand), and *Oceanites oceanicus exasperatus* (South Orkney Islands) represent new host-louse associations (voucher specimens in MONZ).

***Austromenopon fuscofasciatum* (Piaget, 1880)**

- Menopon fuscofasciatum* Piaget, 1880: 492, pl. 40: fig. 9.  
*Austromenopon fuscofasciatum* (Piaget, 1880); Hopkins & Clay 1952: 47.  
*Austromenopon fuscofasciatum* (Piaget, 1880); Clay 1959: 161, 164, 167, figs 16, 22.  
*Austromenopon fuscofasciatum* (Piaget, 1880); Pilgrim & Palma 1982: 22.  
*Austromenopon fuscofasciatum*; Crossland 1993: 305.  
*Austromenopon fuscofasciatum* (Piaget, 1880); Palma 1999: 380.  
*Austromenopon fuscofasciatum* (Piaget, 1880); Murray *et al.* 2006a: 1965.  
*Austromenopon fuscofasciatum* (Piaget, 1880); Palma 2010: 407.

Lectotype ♀ in NHML (Clay 1949b: 829).

Type host: *Coprotheres pomarinus* (Temminck, 1815).

New Zealand hosts: *Catharacta antarctica lonnbergi* Mathews, 1912; *Catharacta maccormicki* (Saunders, 1893); *Stercorarius parasiticus* (Linnaeus, 1758); *Stercorarius longicaudus* Vieillot, 1819.

Other hosts: *Catharacta antarctica hamiltoni* Hagen, 1952; *Catharacta chilensis* (Bonaparte, 1857).

New Zealand localities: ND, WO, WI, WN, SD, MB, NC, MC, SC, CH, Macquarie Island, RO.

Geographic distribution: Cosmopolitan.

New Zealand references: Pilgrim & Palma (1982); Crossland (1993); Palma (1996b: 116); Palma (1999); Palma & Horning (2002: 5, 17); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay (1949b: 829); Clay (1959); Clay & Moreby (1967: 158, fig. 66); Hackman & Nyholm (1968: 75); Furness & Palma (1992: 35, 42); Cohen *et al.* (1997: 186); Price *et al.* (2003: 91); Palma & Jensen (2005: 51, 64); Hänel & Palma (2007: 112, 121, 131).

Remarks: *Austromenopon fuscofasciatum* is exclusively parasitic on members of the family Stercorariidae.

***Austromenopon haematopi* Timmermann, 1954**

- Austromenopon haematopi* Timmermann, 1954d: 199, fig. 6.  
*Austromenopon haematopi* Timmermann, 1954; Clay 1959: 163, 165, 167, figs 10, 25, 34.  
*Austromenopon haematopi* Timmermann, 1954; Palma 1999: 380.  
*Austromenopon haematopi* Timmermann, 1954; Price *et al.* 2003: 91.  
*Austromenopon haematopi* Timmermann, 1954; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Haematopus moquini* Bonaparte, 1856.

New Zealand host: *Haematopus finschi* Martens, 1897.

Other hosts: *Haematopus ostralegus* Linnaeus, 1758; *Haematopus bachmani* Audubon, 1838.

New Zealand locality: Auckland.

Geographic distribution: Eurasia; North America; Africa; New Zealand.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Timmermann (1957a: 94, fig. 67a); Clay (1959); Price *et al.* (2003); Palma & Jensen (2005: 51, 63).

Remarks: *Austromenopon haematopi* is restricted to oystercatchers, but infrequently collected.

***Austromenopon himantopi* Timmermann, 1954**

- Austromenopon himantopi* Timmermann, 1954d: 199, fig. 6a.  
*Austromenopon himantopi* Timmermann, 1954; Clay 1959: 163, 167, figs 24, 28, 33.  
*Austromenopon himantopi* Timmermann, 1954; Pilgrim & Palma 1982: 21.  
*Austromenopon himantopi* Timmermann, 1954; Murray *et al.* 1993: 962.  
*Austromenopon himantopi* Timmermann, 1954; Palma 2010: 407.

Holotype ♂ in ZMHG.

Type host: *Himantopus himantopus himantopus* (Linnaeus, 1758).

New Zealand host: *Himantopus himantopus leucocephalus* Gould, 1837.

Other host: *Himantopus himantopus mexicanus* (Stadius Müller, 1776).

New Zealand localities: BP, SD, MB, NC, MC, SC, WD.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 221, 223); Palma (2010).

Other significant references: Timmermann (1957a: 94, fig. 67b); Clay (1959); Hinojos & Canaris (1988: 328); Palma (1996b: 116); Martín-Mateo (2002: 63, figs 18E,G); Price *et al.* (2003: 91).

Remarks: *Austromenopon himantopi* has been found only on stilts, but infrequently collected.

#### ***Austromenopon limosae* Timmermann, 1954**

*Austromenopon limosae* Timmermann, 1954d: 202, fig. 10.

*Austromenopon limosae* Timmermann, 1954; Clay 1959: 163, 167, figs 30, 44.

*Austromenopon limosae* Timmermann, 1954; Pilgrim & Palma 1982: 20.

*Austromenopon limosae* Timmermann, 1954; Murray *et al.* 2006a: 1964.

*Austromenopon limosae* Timmermann, 1954; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Limosa limosa limosa* (Linnaeus, 1758).

New Zealand host: *Limosa limosa melanuroides* Gould, 1846.

Other hosts: *Limosa fedoa* (Linnaeus, 1758).

New Zealand locality: AU.

Geographic distribution: Eurasia; Americas; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1957a: 95, fig. 69); Clay (1959); Price *et al.* (2003: 91).

Remarks: *Austromenopon limosae* appears to be restricted to two species of *Limosa*. Another *Austromenopon* species parasitises *Limosa lapponica* (see below). The Asiatic black-tailed godwit is an uncommon annual visitor to New Zealand (Checklist Committee 2010: 203).

#### ***Austromenopon longithoracicum* (Piaget, 1880)**

*Menopon longithoracicum* Piaget, 1880: 500, pl. 41: fig. 5.

*Procellariophaga longithoracica* (Piaget, 1880); Eichler 1949b: 346.

*Procellariophaga longithoracica* (Piaget, 1880); Hopkins & Clay 1952: 298.

*Austromenopon longithoracicum* (Piaget, 1880); Timmermann 1963: 418, figs 2–3.

*Austromenopon longithoracicum* (Piaget, 1880); Timmermann 1965: 165, 174, pl. 11: figs 1–2.

*Austromenopon longithoracicum* (Piaget, 1880); Price & Clay 1972: 497, figs 42–46.

*Austromenopon longithoracicum* (Piaget, 1880); Price *et al.* 2003: 91.

*Austromenopon longithoracicum* (Piaget, 1880); Scofield *et al.* 2011: 213.

Lectotype ♂ in NHML (Clay 1949b: 836).

Type host: “*Procellaria cinerea*”, in error (see Price & Clay 1972: 497).

New Zealand host: *Calonectris leucomelas* (Temminck, 1836).

Other hosts: None.

New Zealand locality: WO.

Geographic distribution: Japan; Korea; eastern China; Pacific Ocean.

New Zealand reference: Scofield *et al.* (2011).

Other significant references: Eichler (1949b); Timmermann (1963; 1965); Price & Clay (1972); Price *et al.* (2003); Palma (2011b: 21).

Remarks: *Austromenopon longithoracicum* is exclusively parasitic on the streaked shearwater, a host recorded only once in New Zealand (Checklist Committee 2010: 111; Scofield *et al.* 2011).

#### ***Austromenopon lutescens lutescens* (Burmeister, 1838)**

*Menopon lutescens* Burmeister, 1838a: 440.

*Austromenopon lutescens* (Burmeister, 1838); Hopkins & Clay 1952: 47.

*Austromenopon lutescens* (Burmeister, 1838); Clay 1959: 163, 167, fig. 43.

*Austromenopon lutescens* (Burmeister, 1838) *s. l.*; Pilgrim & Palma 1982: 21.

*Austromenopon lutescens* (Burmeister, 1838); Price *et al.* 2003: 91.

*Austromenopon lutescens lutescens* (Burmeister, 1838); Palma & Jensen 2005: 51, 63.

*Austromenopon lutescens* (Burmeister, 1838); Murray *et al.* 2006a: 1964.

*Austromenopon lutescens* (Burmeister, 1838); Palma 2010: 407.

Types presumed lost. See Clay (1949b: 837).

Type host: *Philomachus pugnax* (Linnaeus, 1758).

New Zealand hosts: *Arenaria interpres* (Linnaeus, 1758); *Calidris canutus rogersi* (Mathews, 1913).

Other hosts: *Calidris alpina* (Linnaeus, 1758); *Calidris alba* (Pallas, 1764); *Tringa totanus* (Linnaeus, 1758); *Tringa cinerea* (Güldenstaedt, 1774); *Tringa glareola* Linnaeus, 1758; *Tringa brevipes* (Vieillot, 1816).

New Zealand localities: WN, NC, MC, SC, CA.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1957a: 95); Clay (1959); Brelieh & Tovornik (1962: 93); Cabot (1975: 146); Butler & O'Connor (1994: 450); Forrester *et al.* (1995: 27); Martín-Mateo (2002: 63); Price *et al.* (2003); Palma & Jensen (2005).

Remarks: In agreement with Palma & Jensen (2005: 51) and contrary to Price *et al.* (2003: 91) and Palma (2010: 407), I find the subdivision of *Austromenopon lutescens* into subspecies to better fit the morphological variability exhibited by this taxon.

#### ***Austromenopon meyeri* (Giebel, 1874)**

*Menopon Meyeri* Giebel, 1874: 296.

*Austromenopon meyeri* (Giebel, 1874); Hopkins & Clay 1952: 48.

*Austromenopon meyeri* (Giebel, 1874); Clay 1959: 163, 167, fig. 42.

*Austromenopon meyeri* (Giebel, 1874); Watt 1971: 233, 243.

*Austromenopon meyeri* (Giebel, 1874); Wise 1977: 56.

*Austromenopon meyeri* (Giebel, 1874); Pilgrim & Palma 1982: 21.

*Austromenopon meyeri* (Giebel, 1874); Murray *et al.* 2006a: 1964.

*Austromenopon meyeri* (Giebel, 1874); Palma 2010: 407.

Holotype ♀, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Limosa lapponica lapponica* (Linnaeus, 1758).

New Zealand host: *Limosa lapponica baueri* Naumann, 1836.

Other hosts: None.

New Zealand localities: SD, MB, WD, KE, SN.

Geographic distribution: Eurasia; Australasia; Pacific Ocean.

New Zealand references: Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1954d: 202, fig. 9); Clay (1959); Price *et al.* (2003: 91).

Remarks: *Austromenopon meyeri* appears to be restricted to *Limosa lapponica*. Another *Austromenopon* species parasitises two other species of *Limosa* (see above). The eastern bar-tailed godwit is the most numerous wader that visits New Zealand every year (Checklist Committee 2010: 203).

#### ***Austromenopon navigans* (Kellogg, 1896)**

Figs 18–19

*Menopon navigans* Kellogg, 1896a: 156, pl. 14: figs 4–5.

*Procellariphaga navigans* (Kellogg, 1896); Hopkins & Clay 1952: 299.

*Austromenopon* sp.; Timmermann 1954d: fig. 20.

*Austromenopon navigans* (Kellogg, 1896); Timmermann 1965: 166.

*Austromenopon navigans* (Kellogg, 1896); Price & Clay 1972: 490, figs 13–15, 17.

*Austromenopon bulleri* Price & Clay 1972: 491, fig. 16.

*Austromenopon bulleri* Price & Clay 1972; Wise 1977: 56.

*Austromenopon bulleri* Price & Clay 1972; Horning *et al.* 1980: 4, 9.

*Austromenopon navigans* (Kellogg, 1896); Horning *et al.* 1980: 4, 9.

*Austromenopon bulleri* Price & Clay 1972; Pilgrim & Palma 1982: 6.



*Austromenopon navigans* (Kellogg, 1896); Pilgrim & Palma 1982: 6.

*Austromenopon bulleri* Price & Clay 1972; Murray *et al.* 1990: 1368.

*Austromenopon navigans* (Kellogg, 1896); Murray *et al.* 1990: 1368–1369.

*Austromenopon navigans* (Kellogg, 1896); Palma 1994a: 64.

*Austromenopon navigans* (Kellogg, 1896); Price *et al.* 2003: 91.

*Austromenopon navigans* (Kellogg, 1896); Palma 2010: 407.

Syntypes ♂♀, lost (Price & Clay 1972: 491). Holotype ♀ of *Austromenopon bulleri* in CMNZ (Nicholls *et al.* 1998: 30).

Type host: *Phoebastria albatrus* (Pallas, 1769).

New Zealand hosts: *Thalassarche chrysostoma* (J.R. Forster, 1785); *Thalassarche melanophris* (Temminck, 1828);

*Thalassarche bulleri bulleri* (Rothschild, 1893); *Thalassarche bulleri platei* (Reichenow, 1898); *Thalassarche cauta stadi* Falla, 1933; *Thalassarche eremita* Murphy, 1930; *Thalassarche salvini* (Rothschild, 1893).

Other hosts: *Thalassarche chlororhynchos* (J.F. Gmelin, 1789); *Thalassarche cauta cauta* (Gould, 1841).

New Zealand localities: ND, HB, WI, WN, SD, MB, NC, MC, SC, CH, SN.

Geographic distribution: Southern Hemisphere and North Pacific Ocean.

New Zealand references: Price & Clay (1972); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1994a); Nicholls *et al.* (1998: 30); Palma (1999: 375, 384, note B); Palma (2010).

Other significant references: Timmermann (1965); Green & Palma (1991: 5, 25); Palma (1996b: 117); Price *et al.* (2003); Palma & Peck (2013: 14).

Remarks: *Austromenopon navigans* is one of two species of *Austromenopon* parasitic on the smaller albatrosses. *Austromenopon bulleri* was described from the New Zealand endemic Buller's albatross, but later synonymised under *A. navigans* by Palma (1994a: 64).

#### ***Austromenopon ossifragae* (Eichler, 1949)**

*Procellariophaga ossifragae* Eichler, 1949a: 12.

*Procellariophaga ossifragae* Eichler, 1949b: 345, figs 24–27.

*Austromenopon ossifragae* (Eichler, 1949); Timmermann 1963: 409, fig. 4.

*Austromenopon ossifragae* (Eichler, 1949); Timmermann 1965: 167, fig. 104.

*Austromenopon ossifragae* (Eichler, 1949); Price & Clay 1972: 491, figs 18–20.

*Austromenopon ossifragae* (Eichler, 1949); Wise 1977: 56.

*Austromenopon ossifragae* (Eichler, 1949); Pilgrim & Palma 1982: 7.

*Austromenopon ossifragae* (Eichler, 1949); Murray *et al.* 1990: 1369.

*Austromenopon ossifragae* (Eichler, 1949); Palma 2010: 407.

Syntypes ♀♀ in ZMHG (Weidner 1966: 261).

Type host: *Macronectes giganteus* (J.F. Gmelin, 1789).

New Zealand hosts: *Macronectes giganteus* (J.F. Gmelin, 1789); *Macronectes halli* Mathews, 1912.

Other hosts: None.

New Zealand localities: ND, AK, BP, TK, WA, WN, MB.

Geographic distribution: Southern Hemisphere.

New Zealand references: Clay (1964a: 230); Gressitt (1964: 538); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1369); Paterson *et al.* (1999: 222); Palma (2010).

Other significant references: Timmermann (1963; 1965); Clay & Moreby (1967: 158, 168, figs 68, 70, 75); Price & Clay (1972); Green & Palma (1991: 5, 25); Palma (1996b: 117); Price *et al.* (2003: 92).

Remarks: *Austromenopon ossifragae* is exclusively parasitic on giant petrels. Eichler (1949a,b) described this louse as a new species twice.

#### ***Austromenopon paululum* (Kellogg & Chapman, 1899)**

*Menopon paululum* Kellogg & Chapman, 1899: 119, pl. 8: fig. 2.

*Procellariophaga paulula* (Kellogg & Chapman, 1899); Eichler 1949b: 346.

*Procellariophaga paulula* (Kellogg & Chapman, 1899); Hopkins & Clay 1952: 299.

*Austromenopon spec.*; Timmermann 1963: 414, fig. 8a.

*Austromenopon paululum* (Kellogg & Chapman, 1899); Timmermann 1963: 416.

*Austromenopon piekarskii* Timmermann, 1963: 417, fig. 9.  
*Austromenopon paululum* (Kellogg & Chapman, 1899); Timmermann 1965: 171.  
*Austromenopon piekarskii* Timmermann, 1963; Timmermann 1965: 173, fig. 108.  
*Austromenopon* sp.; Watt 1971: 233, 243.  
*Austromenopon paululum* (Kellogg & Chapman, 1899); Price & Clay 1972: 494, fig. 34.  
*Austromenopon paululum* (Kellogg & Chapman, 1899); Horning *et al.* 1980: 4, 10.  
*Austromenopon paululum* (Kellogg & Chapman, 1899); Pilgrim & Palma 1982: 11.  
*Austromenopon paululum* (Kellogg & Chapman, 1899); Palma 1999: 376, 383, note 2.  
*Austromenopon paululum* (Kellogg & Chapman, 1899); Palma 2010: 407.

Lectotype ♀ in EMEC (Price & Clay 1972: 496).

Type host: *Puffinus opisthomelas* Coues, 1864.

New Zealand hosts: *Puffinus pacificus pacificus* (J.F. Gmelin, 1789); *Puffinus pacificus chlororhynchus* Lesson, 1831; *Puffinus bulleri* Salvin, 1888; *Puffinus carneipes* Gould, 1844; *Puffinus griseus* (J.F. Gmelin, 1789); *Puffinus tenuirostris* (Temminck, 1835); *Puffinus newelli* Henshaw, 1900; *Puffinus puffinus* (Brünnich, 1764); *Puffinus gavia* (J.R. Forster, 1844); *Puffinus huttoni* Mathews, 1912; *Puffinus assimilis kermadecensis* Murphy, 1927; *Puffinus assimilis haurakiensis* Fleming & Serventy, 1943; *Puffinus elegans* Giglioli & Salvadori, 1869.

Other hosts: *Puffinus gravis* (O'Reilly, 1818); *Puffinus creatopus* Coues, 1864; *Puffinus yelkouan* (Acerbi, 1827); *Puffinus mauretanicus* Lowe, 1921; *Puffinus nativitatis* Streets, 1877; *Puffinus lherminieri* Lesson, 1839; *Puffinus auricularis* Townsend, 1890; *Puffinus assimilis boydi* Mathews, 1912; *Puffinus assimilis tunneyi* Mathews, 1912.

New Zealand localities: ND, CL, AK, BP, WI, WA, WN, SD, MB, NN, NC, MC, SC, WD, CO, DN, KE, CH, SI, SN, AN.

Geographic distribution: Pacific, Atlantic and Indian Oceans.

New Zealand references: Watt (1971); Price & Clay (1972: 496); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990: 1371); Paterson *et al.* (1999: 222); Palma (1999); Galloway (2005: 16); Palma (2010).

Other significant references: Eichler (1949b); Timmermann (1963); Timmermann (1965); Bourgeois & Threlfall (1979: 1356); Fowler & Shaw (1990: 15); Green & Palma (1991: 5, 27); Forrester *et al.* (1995: 5); Foster *et al.* (1996: 85); Palma (1996b: 117); Martín-Mateo (2002: 62, figs 18B,I, 19D); Price *et al.* (2003: 92); Palma & Jensen (2005: 51, 60); Hänel & Palma (2007: 112, 121, 130); Palma & Peck (2013: 14).

Remarks: *Austromenopon paululum* parasitises most species of the genus *Puffinus*. The Manx shearwater, *Puffinus puffinus*, is a new host record for *Austromenopon paululum* in New Zealand (voucher specimens in MONZ).

### ***Austromenopon phaeopodis* (Schrank, 1802)**

*Pediculus phaeopodis* Schrank, 1802: 361.  
*Austromenopon phaeopodis* (Schrank, 1802); Hopkins & Clay 1952: 48.  
*Austromenopon phaeopodis* (Schrank, 1802); Timmermann 1954d: 203, fig. 12.  
*Austromenopon phaeopodis* (Schrank, 1802); Clay 1959: 157, 162, 165–167, figs 1–2, 14, 18, 20, 29, 36–37, 40.  
*Austromenopon phaeopodis* (Schrank, 1802); Pilgrim & Palma 1982: 20.  
*Austromenopon phaeopodis* (Schrank, 1802); Murray *et al.* 2006a: 1964.  
*Austromenopon phaeopodis* (Schrank, 1802); Palma 2010: 407.

Neotype ♂ in NHML (Clay 1959: 166).

Type host: *Numenius phaeopus phaeopus* (Linnaeus, 1758).

New Zealand host: *Numenius phaeopus variegatus* (Scopoli, 1786).

Other host: *Numenius tahitiensis* (J.F. Gmelin, 1789).

New Zealand localities: TK, KE.

Geographic distribution: Eurasia; Americas; Australasia; Pacific Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1954d); Timmermann (1957a: 96, fig. 70); Emerson & Ward (1958: 50); Clay (1959); Clay & Hopkins (1960: 10); Moreby (1976: 93); Price *et al.* (2003: 92); Palma & Jensen (2005: 52, 64).

Remarks: *Austromenopon phaeopodis* appears to be restricted to species of *Numenius*. Although the Asiatic whimbrel is a frequent visitor to New Zealand (Checklist Committee 2010: 201), its lice have been infrequently collected in this country.

### ***Austromenopon pinguis* (Kellogg, 1896)**

- Colpocephalum pingue* Kellogg, 1896a: 144, pl. 12: fig. 5.  
*Procellariphaga pinguis* (Kellogg, 1896); Eichler 1949b: 346.  
*Procellariphaga pinguis* (Kellogg, 1896); Hopkins & Clay 1952: 299.  
*Austromenopon pingue* (Kellogg, 1896); Timmermann 1965: 166.  
*Austromenopon pinguis* (Kellogg, 1896); Price & Clay 1972: 490, figs 10–12.  
*Austromenopon pinguis* (Kellogg, 1896); Horning *et al.* 1980: 4, 9.  
*Austromenopon pinguis* (Kellogg, 1896); Pilgrim & Palma 1982: 6.  
*Austromenopon pinguis* (Kellogg, 1896); Murray *et al.* 1990: 1368–1369.  
*Austromenopon pinguis* (Kellogg, 1896); Palma 2010: 407.

Lectotype ♀ in EMEC (Price & Clay 1972: 490).

Type host: *Phoebastria albatrus* (Pallas, 1769).

New Zealand hosts: *Thalassarche chrysostoma* (J.R. Forster, 1785); *Thalassarche melanophris* (Temminck, 1828); *Thalassarche impavida* Mathews, 1912; *Thalassarche bulleri bulleri* (Rothschild, 1893); *Thalassarche bulleri platei* (Reichenow, 1898); *Thalassarche cauta steadi* Falla, 1933; *Thalassarche salvini* (Rothschild, 1893); *Phoebetria palpebrata* (J.R. Forster, 1785).

Other hosts: *Phoebastria nigripes* (Audubon, 1839); *Phoebastria immutabilis* (Rothschild, 1893); *Thalassarche chlororhynchos* (J.F. Gmelin, 1789); *Thalassarche cauta cauta* (Gould, 1841); *Phoebetria fusca* (Hilsenberg, 1822).

New Zealand localities: ND, AK, BP, HB, WI, WA, WN, NN, NC, MC, SC, CO, DN, SI, CH, BO, SN, AU, CA.

Geographic distribution: Southern Hemisphere and north Pacific Ocean.

New Zealand references: Price & Clay (1972); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Timmermann (1965); Green & Palma (1991: 5, 25); Palma (1996b: 118); Price *et al.* (2003: 92); Hänel & Palma (2007: 112, 121, 129); Palma & Peck (2013: 15).

Remarks: *Austromenopon pinguis* is the smallest of the two species of *Austromenopon* parasitic on the smaller albatrosses.

### ***Austromenopon popellus* (Piaget, 1890)**

- Menopon popellus* Piaget, 1890a: 251, pl. 10: fig. 5.  
*Procellariphaga popellus* (Piaget, 1890); Hopkins & Clay 1952: 299.  
*Austromenopon spec.*; Timmermann 1963: 402, 413, figs 1, 7.  
*Austromenopon popellus* (Piaget, 1890); Timmermann 1963: 413.  
*Austromenopon spec.*; Timmermann 1965: 169, fig. 107, pl. 11: fig. 3.  
*Austromenopon popellus* (Piaget, 1890); Timmermann 1965: 170.  
*Austromenopon popellus* (Piaget, 1890); Price & Clay 1972: 491, fig. 24.  
*Austromenopon popellus* (Piaget, 1890); Pilgrim & Palma 1982: 8, 11.  
*Austromenopon popellus* (Piaget, 1890) *s. l.*; Pilgrim & Palma 1982: 8.  
*Austromenopon popellus* (Piaget, 1890); Murray *et al.* 1990: 1369, 1371.  
*Austromenopon popellus* (Piaget, 1890); Palma 2010: 407.

Holotype ♂ in NHML (Clay 1949b: 908).

Type host: “*Podica senegalensis*” in error (see Hopkins & Clay 1952: 299).

New Zealand hosts: *Lugensa brevirostris* (Lesson, 1833); *Pterodroma macroptera gouldi* (Hutton, 1869); *Pterodroma lessonii* (Garnot, 1826); *Pterodroma magentae* (Giglioli & Salvadori, 1869); *Pterodroma neglecta neglecta* (Schlegel, 1863); *Pterodroma mollis* (Gould, 1844); *Pterodroma externa* (Salvin, 1875); *Pterodroma cervicalis* (Salvin, 1891); *Procellaria aequinoctialis* Linnaeus, 1758; *Procellaria westlandica* Falla, 1946; *Procellaria parkinsoni* G.R. Gray, 1862; *Procellaria cinerea* J.F. Gmelin, 1789.

Other hosts: *Pterodroma incerta* (Schlegel, 1863); *Pterodroma solandri* (Gould, 1844); *Pterodroma ultima* Murphy, 1949; *Pterodroma hasitata hasitata* (Kuhl, 1820); *Pterodroma occulta* Imber & Tennyson, 2001; *Pterodroma arminjoniana* (Giglioli & Salvadori, 1869); *Pterodroma heraldica* (Salvin, 1888); *Pterodroma atrata* (Mathews, 1912); *Pterodroma phaeopygia* (Salvin, 1876); *Pterodroma hypoleuca* (Salvin, 1888); *Pseudobulweria rostrata* (Peale, 1848).

New Zealand localities: ND, AK, CL, BP, WO, TK, WI, WA, WN, NN, NC, MC, SC, WD, SL, KE, Norfolk Island, CH, SI, AN.

Geographic distribution: Pacific, Atlantic and Indian Oceans.

New Zealand references: Price & Clay (1972: 493); Pilgrim & Palma (1982); Palma & Pilgrim (1983: 148); Murray *et al.* (1990: 1369, 1371); Marris (2000: 188); Palma & Imber (2000: 229); Palma (2010).

Other significant references: Timmermann (1963; 1965); Price & Clay (1972); Green & Palma (1991: 5, 26); Zonfrillo (1993: 327); Forrester *et al.* (1995: 5); Furness & Palma (1992: 35, 40); Palma (1996b: 118); Price *et al.* (2003: 92); Hänel & Palma (2007: 112, 121, 130); Palma & Peck (2013: 15).

Remarks: *Austromenopon popellus* is a widespread species on most species of *Pterodroma*. Pilgrim & Palma (1982: 8–9) regarded the population of *A. popellus* from *Lugensa brevisrostris* and *Pt. magentae* as somewhat different from those from several other hosts, and qualified them as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted.

Records of *A. popellus* from *Pterodroma cookii* (G.R. Gray, 1843) and *Pterodroma leucoptera* (Gould, 1844) in Price *et al.* (2003: 92), and from *Pterodroma pycrofti* Falla, 1933 in Pilgrim & Palma (1982: 9) are erroneous. The species of *Austromenopon* from these hosts and other related *Pterodroma* species is undescribed (see below under ***Austromenopon* species 1**).

Furthermore, only two specimens of *A. popellus*—one female reported by Price & Clay (1972: 493) and one male by Pilgrim & Palma (1982: 8)—have been recorded from *Pterodroma inexpectata* (J.R. Forster, 1844), and both are most likely stragglers because a different species represented by several samples from various localities has also been recorded from *Pt. inexpectata* (see below under ***Austromenopon* species 2**).

### ***Austromenopon stammeri* Timmermann, 1963**

*Austromenopon stammeri* Timmermann, 1963: 421, fig. 12.

*Austromenopon stammeri* Timmermann, 1963; Timmermann 1965: 175, fig. 111.

*Austromenopon stammeri* Timmermann, 1963; Price & Clay 1972: 496, figs 35–38.

*Austromenopon stammeri* Timmermann, 1963; Wise 1977: 56.

*Austromenopon stammeri* Timmermann, 1963; Horning *et al.* 1980: 4, 10.

*Austromenopon stammeri* Timmermann, 1963; Pilgrim & Palma 1982: 9.

*Austromenopon stammeri* Timmermann, 1963; Murray *et al.* 1990: 1370.

*Austromenopon stammeri* Timmermann, 1963; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Pachyptila turtur* (Kuhl, 1820).

New Zealand hosts: *Halobaena caerulea* (J.F. Gmelin, 1789); *Pachyptila vittata* (G. Forster, 1777); *Pachyptila salvini salvini* (Mathews, 1912); *Pachyptila desolata* (J.F. Gmelin, 1789); *Pachyptila belcheri* (Mathews, 1912); *Pachyptila turtur* (Kuhl, 1820); *Pachyptila crassirostris crassirostris* (Mathews, 1912); *Pachyptila crassirostris pyramidalis* C.A. Fleming, 1939.

Other hosts: None.

New Zealand localities: AK, TK, WI, WA, WN, NC, MC, SC, WD, CH, SI, BO, SN, AN.

Geographic distribution: Southern Hemisphere.

New Zealand references: Timmermann (1963); Pilgrim (1970: 74); Price & Clay (1972); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 222); Marris (2000: 188); Hänel & Palma (2007: 112, 121, 130); Palma (2010).

Other significant references: Timmermann (1965); Green & Palma (1991: 5, 26); Furness & Palma (1992: 35, 39); Palma (1996b: 118); Price *et al.* (2003: 92).

Remarks: *Austromenopon stammeri* is a prevalent and abundant species on all species and subspecies of prions.

**Austromenopon transversum (Denny, 1842)**

*Liotheum (Menopon) transversum* Denny, 1842: 201.

*Liotheum (Menopon) ridibundis* [sic] Denny, 1842: 201.

*Liotheum (Menopon) transversus* [sic] Denny, 1842: 226, pl. 21; fig. 7.

*Liotheum (Menopon) ridibundus* [sic] Denny, 1842: 227, pl. 20; fig. 3.

*Menopon transversum* Denny, 1842 [sic]; Harrison 1916: 46 (as junior synonym of *Menopon ridibundum* Denny, 1842 [sic]).

*Austromenopon ridibundus* [sic] (Denny, 1842); Hopkins & Clay 1952: 48.

*Austromenopon transversum* (Denny, 1842); Hopkins & Clay 1952: 48.

*Austromenopon transversum* (Denny, 1842); Clay 1959: 161, 164, 166, figs 15, 19, 21.

*Austromenopon transversum* (Denny, 1842); Clay & Moreby 1967: 158, 169, figs 63, 73.

*Austromenopon transversum* (Denny, 1842); Pilgrim & Palma 1982: 22.

*Austromenopon transversum* (Denny, 1842); Murray *et al.* 2006a: 1965.

*Austromenopon transversum* (Denny, 1842); Palma 2010: 407.

Lectotype ♀ in NHML (Clay 1959: 166).

Type host: *Rissa tridactyla* (Linnaeus, 1758).

New Zealand hosts: *Larus dominicanus* Lichtenstein, 1823; *Larus novaehollandiae scopulinus* J.R. Forster, 1844; *Larus bulleri* Hutton, 1871.

Other hosts: *Pagophila eburnea* (Phipps, 1774); *Rhodostethia rosea* (Macgillivray, 1924); *Rissa brevirostris* (Bruch, 1853); *Xema sabini* (Sabine, 1819); *Creagrus furcatus* (Nébox, 1842) and at least 19 other species of *Larus* (see Price *et al.* 2003: 93).

New Zealand localities: WN, SD, MB, NN, NC, MC, SC, WD, SL, CA.

Geographic distribution: Cosmopolitan.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1954d: 204, figs 15–16); Timmermann (1957a: 97, figs 73–74); Clay (1959); Clay & Moreby (1967); Choe & Kim (1987: 3000; 1988: 988); Green & Palma (1991: 5, 33); Butler & O'Connor (1994: 450); Forrester *et al.* (1995: 29); Palma (1996b: 119); Martín-Mateo (2002: 59, figs 18D,F, 19C, 20); Price *et al.* (2003: 93); Palma & Jensen (2005: 52, 65); González-Acuña *et al.* (2011: 300); Palma & Peck (2013: 16).

Remarks: This species is the only *Austromenopon* known from all species of gulls (Price *et al.* 2003: 93, 290).

**Austromenopon species 1**

"*Austromenopon popellus*" Pilgrim & Palma, 1982: 9 (not *Menopon popellus* Piaget, 1890a).

*Austromenopon* sp.: Pilgrim & Palma 1982: 9.

*Austromenopon* sp.: Murray *et al.* 1990: 1370.

"*Austromenopon popellus*" Murray *et al.*, 1990: 1370 (not *Menopon popellus* Piaget, 1890a).

*Austromenopon* sp.: Palma 1999: 377.

*Austromenopon* spp.: Palma 2010: 407.

New Zealand hosts: *Pterodroma nigripennis* (Rothschild, 1893); *Pterodroma cookii* (G.R. Gray, 1843); *Pterodroma longirostris* (Stejneger, 1888); *Pterodroma pycrofti* Falla, 1933; *Pterodroma leucoptera caledonica* Imber & Jenkins, 1981.

Other host: *Pterodroma brevipes* (Peale, 1848).

New Zealand localities: ND, CL, WO, BP, GB, KE.

Geographic distribution: New Zealand; New Caledonia; Cook Islands.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999); Palma (2010).

Other significant references: None.

Remarks: These records of *Austromenopon* from small species of *Pterodroma* represent an undescribed, unnamed species, although male lice are needed from *Pt. longirostris* and *Pt. pycrofti* to confirm their taxonomic status. Records of *A. popellus* from *Pterodroma pycrofti* in Pilgrim & Palma (1982: 9) and in Murray *et al.* (1990: 1370) are erroneous.

**Austromenopon species 2**

*Austromenopon* sp.: Horning *et al.* 1980: 4, 9.

*Austromenopon* sp.: Pilgrim & Palma 1982: 8.

*Austrmenopon* [sic] sp.: Murray *et al.* 1990: 1370.

*Austromenopon* spp.: Palma 2010: 407.

New Zealand host: *Pterodroma inexpectata* (J.R. Forster, 1844).

Other hosts: None.

New Zealand localities: AK, WN, SI, SN.

Geographic distribution: New Zealand.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: None.

Remarks: This undescribed, unnamed species of *Austromenopon* is exclusively parasitic on the mottled petrel, a host species which only breeds in New Zealand but ranges widely over the Pacific Ocean (Checklist Committee 2010: 93).

### ***Austromenopon* species 3**

*Austromenopon* sp.: Pilgrim & Palma 1982: 21.

*Austromenopon* species 2: Marris 2000: 188.

*Austromenopon* sp.: Murray *et al.* 2006a: 1964.

*Austromenopon* spp.: Palma 2010: 407.

New Zealand hosts: *Coenocorypha aucklandica meinertzhagenae* Rothschild, 1927; *Coenocorypha aucklandica aucklandica* (G.R. Gray, 1845).

Other hosts: None.

New Zealand localities: AN, AU.

Geographic distribution: Subantarctic Islands of New Zealand.

New Zealand references: Pilgrim & Palma (1982); Marris (2000); Murray *et al.* (2006a); Palma (2010).

Other significant references: None.

Remarks: These records of *Austromenopon* from two New Zealand Subantarctic snipes comprise three samples totalling 11 females only (voucher specimens in MONZ), and may represent an undescribed endemic taxon (Theresa Clay pers. comm. 1971). Males are needed to confirm the taxonomic status of these lice.

### **Genus *Bonomiella* Conci, 1942**

*Bonomiella* Conci, 1942a. *Studi Trentini Scienze Naturali* 23(2): 124. Type species: *Bonomiella insolitunguicolata* Conci, 1942a (by original designation).

### ***Bonomiella columbae* Emerson, 1957**

Figs 20–21

*Bonomiella columbae* Emerson, 1957: 63, figs 1–3.

*Bonomiella*; Pilgrim 1970: 76.

*Bonomiella columbae* Emerson, 1957; Pilgrim 1976: 162, figs 4–5.

*Bonomiella columbae* Emerson, 1957; Wise 1977: 56.

*Bonomiella columbae* Emerson, 1957; Pilgrim & Palma 1982: 23.

*Bonomiella columbae* Emerson, 1957; Price *et al.* 2003: 93.

*Bonomiella columbae* Emerson, 1957; Murray *et al.* 2006a: 1965.

*Bonomiella columbae* Emerson, 1957; Palma 2010: 407.

Holotype ♂ in USNM.

Type host: “domestic pigeon” = *Columba livia domestica* J.F. Gmelin, 1789.

New Zealand host: *Columba livia* J.F. Gmelin, 1789.

Other hosts: *Zenaida macroura* (Linnaeus, 1758).

New Zealand localities: WN, MC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim (1970); Pilgrim (1976); Wise (1977); Hill & Tuff (1978: 316, 324); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Murray *et al.* (2006a); Palma (2010).

Other significant references: Złotorzycka & Lucińska (1967: 341, figs 1–5); Selim *et al.* (1968: 79, fig. 7); Ribbeck (1972: 129, figs 1–5); Conti & Forrester (1981: 531); Selva *et al.* (1987: 246, 249); Price *et al.* (2003); Cicchino & González-Acuña (2012: 49, figs 1, 3, 5).

Remarks: *Bonomiella columbae* was introduced to New Zealand and other countries with rock pigeons by human agency (Checklist Committee 2010: 245).

### Genus *Ciconiphilus* Bedford, 1939

*Ciconiphilus* Bedford, 1939. *Onderstepoort Jour. Vet. Sci.* 12(1): 141. Type species: *Colpocephalum quadripustulatus* Nitzsch [sic] = *Ciconiphilus quadripustulatus* (Burmeister, 1838) (by original designation).

*Anseriphilus* Eichler, 1944a. *Deutsch. Entomol. Zeit.* 1943: 57. Type species: *Colpocephalum pectiniventre* Harrison, 1916 = *Ciconiphilus pectiniventris* (Harrison, 1916) (by original designation).

### *Ciconiphilus decimfasciatus* (Boisduval & Lacordaire, 1835)

Figs 22–23

*Liotheum (Colpocephalum) 10-fasciatum* Boisduval & Lacordaire, 1835: 123.

*Colpocephalum decimfasciatum* Boisduval & Lacordaire, 1835 [sic]; Harrison 1916: 48.

*Pseudocolpocephalum doriabagla* Ansari, 1951: 154, fig. 11.

*Ciconiphilus decimfasciatus* (Boisduval & Lacordaire, 1835); Hopkins & Clay 1952: 71.

*Ciconiphilus doriabagla* (Ansari, 1951); Hopkins & Clay 1953: 436.

*Ciconiphilus decimfasciatus* (Boisduval & Lacordaire, 1835); Price & Beer 1965a: 661, figs 10–12.

*Ciconiphilus decimfasciatus* (Boisduval & Lacordaire, 1835); Pilgrim & Palma 1982: 15.

*Ciconiphilus decimfasciatus* (Boisduval & Lacordaire, 1835); Murray *et al.* 1990: 1373.

*Ciconiphilus decimfasciatus* (Boisduval & Lacordaire, 1835); Palma 2010: 407.

Status, sex and repository of types unknown, probably lost (Palma 1996b: 119).

Type host: *Ardea cinerea cinerea* Linnaeus, 1758.

New Zealand hosts: *Ardea modesta* J.E. Gray, 1831; *Ardea ibis coromanda* (Boddaert, 1783); *Egretta garzetta immaculata* (Gould, 1846); *Egretta novaehollandiae novaehollandiae* (Latham, 1790); *Egretta sacra sacra* (J.F. Gmelin, 1789); *Botaurus poiciloptilus* (Wagler, 1827).

Other hosts: Over 25 species in the genera *Agamia*, *Ardea*, *Ardeola*, *Botaurus*, *Butorides*, *Egretta*, *Ixobrychus*, *Nycticorax*, *Pilherodius*, and *Tigrisoma* (see Price *et al.* 2003: 95).

New Zealand localities: ND, BP, HB, WA, SD, MB, NN, NC, MC, SC, WD, CO, DN, KE, AU.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 221); Palma (1999: 379); Palma (2010).

Other significant references: Ansari (1951); Price & Beer (1965a); Price & Emerson (1966: 432); Price & Emerson (1967: 249); Moreby (1976: 92); Clay (1976b: 537); Butler & O'Connor (1994: 450); Forrester *et al.* (1995: 9); Palma (1996b: 119); Martín-Mateo (2002: 84); Price *et al.* (2003: 94); Palma & Jensen (2005: 52, 61); Adam (2007: 168).

Remarks: *Ciconiphilus decimfasciatus* is a widespread species on many members of the family Ardeidae, usually found in large numbers per host.

### *Ciconiphilus pectiniventris* (Harrison, 1916)

*Menopon pectinatum* Neumann, 1912b: 368, fig. 15.

*Colpocephalum pectinatum* Neumann, 1912 [sic]; Harrison 1916: 53. Preoccupied by *Colpocephalum pectinatum* Osborn, 1902: 201.

*Colpocephalum pectiniventre* Harrison, 1916: 53. *Nomen novum* for *Colpocephalum pectinatum* (Neumann, 1912).

*Ciconiphilus pectiniventris* (Harrison, 1916); Hopkins & Clay 1952: 72.

*Ciconiphilus pectiniventris* (Harrison, 1916); Price & Beer 1965a: 664.

*Ciconiphilus pectiniventris* (Harrison, 1916); Pilgrim & Palma 1982: 16.

*Ciconiphilus pectiniventris* (Harrison, 1916); Murray *et al.* 1990: 1374.

*Ciconiphilus pectiniventris* (Harrison, 1916); Palma 1999: 379, 383, note 4.

*Ciconiphilus pectiniventris* (Harrison, 1916); Palma 2010: 407.

Syntypes ♂♀, repository not confirmed, probably in the Ecole Vétérinaire de Toulouse, France.

Type host: *Anser anser domesticus* (Linnaeus, 1758).

New Zealand hosts: *Anser anser* (Linnaeus, 1758); *Branta canadensis maxima* Delacour, 1951; *Cygnus atratus* (Latham, 1790).

Other hosts: Seven other species of *Anser* and two other species of *Branta* (see Price *et al.* 2003: 95).

New Zealand localities: BP, WO, WA, SD, MB, MC.

Geographic distribution: North America; Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 219); Palma (1999); Palma (2010).

Other significant references: Price & Beer (1965a); Palma (1996b: 119); Price *et al.* (2003: 95); Palma & Jensen (2005: 52, 61).

Remarks: *Ciconiphilus pectiniventris* was introduced to New Zealand with geese by human agency (Checklist Committee 2010: 35). The association of *C. pectiniventris* with *Cygnus atratus* is likely to be the result of a natural host-switch from one or both goose species living in New Zealand.

### Genus *Colpocephalum* Nitzsch, 1818

*Colpocephalum* Nitzsch, 1818. *Germer's Mag. Entomol.* 3: 298. Type species: *Colpocephalum zebra* Burmeister, 1838a (by subsequent designation).

*Ferrisia* Uchida, 1926. *Jour. Coll. Agric. Tokyo* 9: 43. Type species: *Colpocephalum turbinatum* Denny, 1842 (by original designation). Preoccupied by *Ferrisia* Fullaway, 1923.

*Neocolpocephalum* Ewing, 1933. *Jour. Parasit.* 20: 65. *Nomen novum* for *Ferrisia* Uchida, 1926.

*Corvocolpocephalum* Conci, 1942b. *Boll. Soc. Entomol. Italiana* 74: 30. Type species: *Colpocephalum subaequale* "Nitzsch in Burm. 1838" = *Colpocephalum fregili* Denny, 1842 (by original designation).

*Liothella* Eichler, 1947. *Arkiv Zool.* 39A(2): 15. Type species: *Colpocephalum leptopygos* Nitzsch [in Giebel], 1874 (by original designation).

*Pelecanigogus* Eichler, 1949a. *Boll. Soc. Entomol. Italiana* 79: 12. Type species: *Colpocephalum eucarenum* "Ntz i. Brm." (by original designation).

*Galliferrisia* Ansari, 1951. *Proc. Nat. Inst. Sci. India* 17: 150. Type species: *Galliferrisia tausi* Ansari, 1951 = *Colpocephalum tausi* (Ansari, 1951) (by original designation).

### *Colpocephalum eucarenum* Burmeister, 1838

*Colpocephalum eucarenum* Burmeister, 1838a: 439.

*Colpocephalum eucarenum* Burmeister, 1838; Hopkins & Clay 1952: 78.

*Colpocephalum eucarenum* Burmeister, 1838; Price 1967a: 274, figs 1–7.

*Colpocephalum eucarenum* Burmeister, 1838; Pilgrim & Palma 1982: 14.

*Colpocephalum eucarenum* Burmeister, 1838; Murray *et al.* 1990: 1372.

*Colpocephalum eucarenum* Burmeister, 1838; Price *et al.* 2003: 98.

*Colpocephalum eucarenum* Burmeister, 1838; Palma 2010: 407.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Pelecanus onocrotalus* Linnaeus, 1758.

New Zealand host: *Pelecanus conspicillatus* Temminck, 1824.

Other hosts: *Pelecanus rufescens* J.F. Gmelin, 1789; *Pelecanus philippensis* J.F. Gmelin, 1789.

New Zealand locality: SL.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Price (1967a); Moreby (1976: 91); Green & Palma (1991: 5, 28); Palma (1996b: 121); Price *et al.* (2003).

Remarks: *Pelecanus conspicillatus* breeds in Australia and is a straggler to New Zealand (Checklist Committee (2010: 138), with only one record of *Colpocephalum eucarenum* from this country.



**Colpocephalum fregili Denny, 1842**

“*Colpocephalum subaequale*” Burmeister, 1838a: 438 (not *Liotheum (Colpocephalum) subaequale* Haan, 1829).

*Liotheum (Colpocephalum) fregili* Denny, 1842: 198, 208, pl. 20: fig. 4.

*Corvocolpocephalum subaequale* (Nitzsch [in Burmeister], 1838); Conci 1942b: 30.

*Colpocephalum fregili* Denny, 1842; Hopkins & Clay 1952: 79.

*Colpocephalum fregili* Denny, 1842; Price & Beer 1965b: 7, figs 1–4.

*Colpocephalum fregili* Denny, 1842; Pilgrim & Palma 1982: 28.

*Colpocephalum fregili* Denny, 1842; Murray *et al.* 2006b: 1957.

*Colpocephalum fregili* Denny, 1842; Palma 2010: 407.

Syntypes in NHML (Palma 1996b: 121).

Type host: *Pyrrhocorax pyrrhocorax pyrrhocorax* (Linnaeus, 1758).

New Zealand host: *Corvus frugilegus* Linnaeus, 1758.

Other hosts: *Pyrrhocorax graculus* (Linnaeus, 1766); *Cyanopica cyanus* (Pallas, 1776); 18 other species of *Corvus* (see Price *et al.* 2003: 98).

New Zealand localities: HB, WA, NC, MC, SC, CH.

Geographic distribution: Eurasia; Africa; North America; Australasia.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999: 221); Murray *et al.* (2006b); Palma (2010).

Other significant references: Price & Beer (1965b); Clay (1976b: 537); Green & Palma (1991: 5, 41); Butler & O'Connor (1994: 450); Palma (1996b: 121); Price *et al.* (2003: 98); Palma & Jensen (2005: 52, 68).

Remarks: *Colpocephalum fregili* was introduced to New Zealand with rooks by human agency (Checklist Committee 2010: 300). It is remarkable that, from a complement of six louse species known to parasitise *Corvus frugilegus* in its home range (Price *et al.* 2003: 338), only *C. fregili* became established on the New Zealand rook populations.

**Colpocephalum leptopygos Nitzsch [in Giebel], 1874**

*Colpocephalum leptopygos* Nitzsch [in Giebel], 1874: 273.

*Liothella leptopygos* (Nitzsch [in Giebel], 1874); Eichler 1947: 15.

*Colpocephalum leptopygos* Nitzsch [in Giebel], 1874; Hopkins & Clay 1952: 80.

*Colpocephalum leptopygos* Nitzsch [in Giebel], 1874; Price & Beer 1965c: 122, figs 58–60.

*Colpocephalum leptopygos* Nitzsch, 1874 [sic]; Pilgrim & Palma 1982: 16.

*Colpocephalum leptopygos* Nitzsch, 1866 [sic]; Murray *et al.* 1990: 1373.

*Colpocephalum leptopygos* Nitzsch [in Giebel], 1874; Price *et al.* 2003: 100.

*Colpocephalum leptopygos* Nitzsch [in Giebel], 1874; Palma 2010: 407.

Syntypes NN presumed lost. See Clay (1949a: 1), Palma & Pilgrim (1984: 150) and Palma (1996b: 122).

Type host: *Plegadis falcinellus* (Linnaeus, 1766).

New Zealand host: *Plegadis falcinellus* (Linnaeus, 1766).

Other hosts: *Plegadis chihi* (Vieillot, 1817); *Plegadis ridgwayi* (Allen, 1876).

New Zealand localities: WI, SL.

Geographic distribution: Eurasia; Africa; Americas; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Price & Beer (1965c); Forrester *et al.* (1995: 12); Palma (1996b: 122); Price *et al.* (2003).

Remarks: Although the glossy ibis is a regular vagrant to New Zealand (Checklist Committee 2010: 166), there are only two records of *Colpocephalum leptopygos* from this country.

**Colpocephalum pilgrimi Price, 1967**

Figs 24–25

“*Colpocephalum setosum*” Kellogg, 1907: 122 (not *Colpocephalum setosum* Piaget, 1880).

*Colpocephalum pilgrimi* Price, 1967b: 11, figs 1–3.

*Colpocephalum pilgrimi* Price, 1967; Wise 1977: 56.

*Colpocephalum pilgrimi* Price, 1967; Pilgrim & Palma 1982: 24.

*Colpocephalum pilgrimi* Price, 1967; Murray *et al.* 1999: 1241.

*Colpocephalum pilgrimi* Price, 1967; Price *et al.* 2003: 101.

*Colpocephalum pilgrimi* Price, 1967; Palma 2010: 407.

Holotype ♀ in CMNZ (Nicholls *et al.* 1998: 30).

Type host: *Nestor notabilis* Gould, 1856.

New Zealand host: *Nestor notabilis* Gould, 1856.

Other hosts: None.

New Zealand localities: NN, NC, MC, WD.

Geographic distribution: South Island, New Zealand.

New Zealand references: Kellogg (1907); Price (1967b); Pilgrim (1970: 75); Wise (1977); Pilgrim & Palma (1982); Nicholls *et al.* (1998: 30); Murray *et al.* (1999); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003).

Remarks: *Colpocephalum pilgrimi* is an endemic and “at risk” species (Buckley *et al.* 2012), exclusively parasitic on the endemic kea.

### ***Colpocephalum subzerafae* Tendeiro, 1988**

“*Colpocephalum zerafae* Ansari, ?1955 *sen. lat.*” Price & Beer 1963a: 758, fig. 55. In part.

“*Colpocephalum zerafae*” Pilgrim & Palma, 1982: 17 (not *Colpocephalum zerafae* Ansari, 1955).

*Colpocephalum subzerafae subzerafae* Tendeiro, 1988: 88, figs 4–6, 16–19.

*Colpocephalum subzerafae* Tendeiro, 1988; Murray *et al.* 1993: 960.

*Colpocephalum subzerafae* Tendeiro, 1988; Palma 1999: 384, note D.

*Colpocephalum subzerafae* Tendeiro, 1988; Price *et al.* 2003: 102.

*Colpocephalum subzerafae* Tendeiro, 1988; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Falco naumanni naumanni* Fleischer, 1818.

New Zealand host: *Falco cenchroides cenchroides* Vigors & Horsfield, 1827.

Other hosts: Nine species of *Falco* (see Price *et al.* 2003: 102).

New Zealand locality: WN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (1999); Palma (2010).

Other significant references: Price & Beer (1963a); Palma (1996b: 122); Price *et al.* (2003).

Remarks: *Falco cenchroides cenchroides* breeds in Australia and is an infrequent straggler to New Zealand (Checklist Committee 2010: 174), with only one sample of *Colpocephalum subzerafae* collected from a nankeen kestrel in this country (voucher specimens in MONZ).

### ***Colpocephalum tausi* (Ansari, 1951)**

*Galliferrisia tausi* Ansari, 1951: 151, fig. 10.

*Colpocephalum tausi* (Ansari, 1951); Hopkins & Clay 1953: 436.

*Colpocephalum tausi* (Ansari, 1951); Price & Beer 1964: 394, figs 12–14.

*Colpocephalum tausi* (Ansari, 1951); Pilgrim & Palma 1982: 18.

*Colpocephalum tausi* (Ansari, 1951); Murray *et al.* 1993: 960.

*Colpocephalum tausi* (Ansari, 1951); Palma 2010: 407.

Holotype ♀ in NHML (Vincent S. Smith pers. comm. December 2014).

Type host: *Pavo cristatus* Linnaeus, 1758.

New Zealand host: *Pavo cristatus* Linnaeus, 1758.

Other host: *Meleagris gallopavo* Linnaeus, 1758.

New Zealand locality: WA.

Geographic distribution: Eurasia; Americas; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Price & Beer (1964); Price *et al.* (2003: 102).

Remarks: *Colpocephalum tausi* was introduced to New Zealand and other countries with its primary host (the peafowl) by human agency (Checklist Committee 2010: 28).

***Colpocephalum turbinatum* Denny, 1842**

- Liotheum (Colpocephalum) turbinatum* Denny, 1842: 198, 209, pl. 21: fig. 1.  
*Colpocephalum turbinatum* Denny, 1842; Hopkins & Clay 1952: 85.  
*Colpocephalum turbinatum* Denny, 1842, *sens. lat.*; Price & Beer 1963a: 754, figs 49, 53, 57.  
*Colpocephalum turbinatum* Denny, 1842; Nelson & Murray 1971: 23, 25, figs 2, 5.  
*Colpocephalum turbinatum* Denny, 1842; Pilgrim 1976: 160, figs 6–7.  
*Colpocephalum turbinatum* Denny, 1842; Pilgrim & Palma 1982: 17, 23.  
*Colpocephalum turbinatum* Denny, 1842; Murray *et al.* 1993: 960.  
*Colpocephalum turbinatum* Denny, 1842; Palma 2010: 407.

Syntypes ♀♀ in NHML (Price & Beer 1963a: 756).

Type host: *Columba livia domestica* J.F. Gmelin, 1789.

New Zealand hosts: *Columba livia* J.F. Gmelin, 1789; *Circus approximans* Peale, 1848.

Other hosts: Over 50 species in the orders Columbiformes, Falconiformes and Strigiformes (see Price *et al.* 2003: 102).

New Zealand localities: WO, HB, WN, SD, MB, NN, NC, MC, SC, NN, WD, CH.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Price & Beer (1963a: 757); Pilgrim (1976); Wise (1977: 56); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 220); Galloway (2005: 17); Murray *et al.* (2006a: 1965); Palma (2010).

Other significant references: Emerson (1957: 64); Nelson & Murray (1971); Moreby (1976: 92); Clay (1976b: 537); Palma (1996b: 123); Price *et al.* (2003: 102); Palma & Peck (2013: 18).

Remarks: *Colpocephalum turbinatum* is exceptional among lice in regard to its wide range of host associations, including many species of pigeons, eagles, hawks, harriers, vultures, osprey and owls (see Price *et al.* 2003: 102).

**Genus *Cuculiphilus* Uchida, 1926****Subgenus *Cuculiphilus* Uchida, 1926**

- Cuculiphilus* Uchida, 1926. *Jour. Coll. Agric. Tokyo* 9: 47. Type species: *Pediculus fasciatus* Scopoli, 1763 = *Cuculiphilus (Cuculiphilus) fasciatus* (Scopoli, 1763) (by original designation).

***Cuculiphilus (Cuculiphilus) fasciiventris* Carriker, 1955**

Figs 26–27

- Cuculiphilus fasciiventris* Carriker, 1955: 11.  
*Cuculiphilus (Cuculiphilus) fasciiventris* Carriker, 1955; Scharf & Price 1965: 551, figs 19, 36.  
*Cuculiphilus (Cuculiphilus) fasciiventris* Carriker, 1955; Palma 1999: 381.  
*Cuculiphilus (Cuculiphilus) fasciiventris* Carriker, 1955; Price *et al.* 2003: 104.  
*Cuculiphilus (Cuculiphilus) fasciiventris* Carriker, 1955; Palma 2010: 408 (incorrectly listed under “Philopteridae”).

Holotype ♂ in Pablo Anduze’s private collection, Caracas, Venezuela (see Emerson 1967: 76)

Type host: *Piaya cayana mehleri* Bonaparte, 1850.

New Zealand host: *Eudynamys taitensis* (Sparrman, 1787).

Other hosts: *Eudynamys scolopacea* (Linnaeus, 1758); *Rhopodytes sumatranus* (Raffles, 1822).

New Zealand locality: WN.

Geographic distribution: Central and South America; Asia; Indonesia; Australasia.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Scharf & Price (1965); Price *et al.* (2003).

Remarks: *Cuculiphilus (Cuculiphilus) fasciiventris* is an infrequently collected species (see Scharf & Price 1965: 551).

Only one small sample of this louse species has been collected from over 50 long-tailed cuckoos searched for lice in New Zealand (voucher specimens in MONZ).

***Cuculiphilus (Cuculiphilus) platygaster* (Giebel, 1874)**

- Menopon platygaster* Giebel, 1874: 290.  
*Cuculiphilus platygaster* (Giebel, 1874); Hopkins & Clay 1952: 99.  
*Cuculiphilus (Cuculiphilus) platygaster* (Giebel, 1874); Scharf & Price 1965: 547, figs 7, 14.  
*Cuculiphilus cuculiphilus* [sic] *platygaster*; Tennyson & Brackenbury 1998: 225.

*Cuculiphilus (Cuculiphilus) platygaster* (Giebel, 1874); Palma 1999: 381.

*Cuculiphilus (Cuculiphilus) platygaster* (Giebel, 1874); Price *et al.* 2003: 105.

*Cuculiphilus (C.) platygaster* (Giebel, 1874); Palma 2010: 408 (incorrectly listed under “Phloptera”).

Holotype ♀, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Scythrops novaehollandiae* Latham, 1790.

New Zealand host: *Scythrops novaehollandiae* Latham, 1790.

Other hosts: None.

New Zealand locality: WO.

Geographic distribution: Indonesia; Australasia.

New Zealand references: Tennyson & Brackenbury (1998); Palma (1999); Palma (2010).

Other significant references: Scharf & Price (1965); Palma (1996b: 124); Price *et al.* (2003).

Remarks: *Scythrops novaehollandiae* is an infrequent straggler to New Zealand, with only six birds recorded from 1924 to 2002 (Checklist Committee 2010: 263). Only one small sample of *Cuculiphilus (Cuculiphilus) platygaster* has been collected from a channel-billed cuckoo in New Zealand (voucher specimens in MONZ).

### Genus *Dennyus* Neumann, 1906

*Nitzschia* Denny, 1842. *Mon. Anopl. Brit.*: 230. Type species: *Nitzschia burmeisteri* Denny, 1842 = *Dennyus (Dennyus) hirundinis* (Linnaeus, 1761) (by monotypy). Preoccupied by *Nitzschia* Baer, 1827. As a subgenus of *Liotheum* Nitzsch.

*Dennyus* Neumann, 1906. *Bull. Soc. Zool. France* 31: 60. *Nomen novum* for *Nitzschia* Denny, 1842.

#### Subgenus *Takamatsuia* Uchida, 1926

*Takamatsuia* Uchida, 1926. *Jour. Coll. Agric. Tokyo* 9: 32. Type species: *Takamatsuia major* Uchida, 1926 = *Dennyus (Takamatsuia) major* (Uchida, 1926) (by original designation).

### *Dennyus (Takamatsuia) species*

### New Record

Fig. 28

New Zealand host: *Hirundapus caudacutus caudacutus* (Latham, 1802).

New Zealand locality: WN.

Geographic distribution: Asia; Australasia; Oceania.

New Zealand reference: This paper.

Other significant references: Uchida (1926: 32, figs 10–11); Nakagawa (1959b: 23); Emerson & Price (1968: 87, fig. 5); Price *et al.* (2003: 105).

Material examined and repository: 1 pharate, 1 nymph (1 sample, MONZ).

Remarks: *Hirundapus caudacutus caudacutus* is a straggler to New Zealand (Checklist Committee 2010: 270), but birds are seldom available for collecting lice. The subgenus *Takamatsuia* includes two species (Price *et al.* 2003: 106) of which, *Dennyus (Takamatsuia) major* (Uchida, 1926) has been recorded in Japan by Uchida (1926) and Nakagawa (1959b). It is likely that the single New Zealand sample belongs to this species but, in the absence of an adult specimen, this record should remain at the subgeneric level only.

### Genus *Eidmanniella* Kéler, 1938

*Eidmanniella* Kéler, 1938b. *Ann. Mus. Zool. Polon.* 13: 81. Type species: *Menopon brevivalpe* Piaget, 1880 = *Eidmanniella pellucida* (Rudow, 1869a) (by original designation).

#### *Eidmanniella albescens* (Piaget, 1880)

*Menopon albescens* Piaget, 1880: 491, pl. 41: fig. 4.

*Menopon singularis* Kellogg & Kuwana, 1902: 485, pl. 31: fig. 1.

*Eidmanniella albescens* (Piaget, 1880); Hopkins & Clay 1952: 129.

*Eidmanniella sula* Tendeiro 1958: 443, figs 1–2, photos 1–4.

*Eidmanniella albescens* (Piaget, 1880); Ryan & Price 1969: 822, figs 5, 13, 26, 29.

*Eidmanniella albescens* (Piaget, 1880); Pilgrim & Palma 1982: 14.

*Eidmanniella albescens* (Piaget, 1880); Murray *et al.* 1990: 1372.

*Eidmanniella albescens* (Piaget, 1880); Palma 2010: 407.

Lectotype ♂ in NHML (Clay 1949b: 816).

Type host: “*Morus serratus*”, in error (see Ryan & Price 1969: 822).

New Zealand hosts: *Sula leucogaster plotus* (J.R. Forster, 1844); *Sula dactylatra tasmani* van Tets, *et al.* 1988.

Other hosts: *Sula sula* (Linnaeus, 1766); *Sula variegata* (Tschudi, 1843); *Sula granti* Rothschild, 1902; *Sula neboxii excisa* Todd, 1948; *Papasula abbotti* (Ridgway, 1893).

New Zealand localities: ND, AK, WO, Westland, KE.

Geographic distribution: Tropical and subtropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999: 378); Palma (2010).

Other significant references: Tendeiro (1958); Ryan & Price (1969: 823); Amerson & Emerson (1971: 10, 29); Moreby (1976: 91); Forrester *et al.* (1995: 6); Palma (1996b: 124); Price *et al.* (2003: 107); Palma & Peck (2013: 19); Rivera-Parra *et al.* (2014: 571); Silva *et al.* (2014: 942); Rivera-Parra *et al.* (2015: 3267).

Remarks: *Eidmanniella albescens* is a frequently collected louse restricted to several species of boobies.

### ***Eidmanniella eurygaster* (Nitzsch [in Giebel], 1866)**

Figs 29–30

*Menopon eurygaster* Nitzsch [in Giebel], 1866: 393.

*Menopon subrotundum* Piaget, 1880: 453, pl. 35: fig. 2.

*Eidmanniella eurygaster* (Nitzsch, 1866) [sic]; Guimarães 1943: 424.

*Eidmanniella eurygaster* (Nitzsch, 1866) [sic]; Hopkins & Clay 1952: 129.

*Eidmanniella subrotunda* (Piaget, 1880); Hopkins & Clay 1952: 130.

*Eidmanniella subrotunda* (Piaget, 1880); Ryan & Price 1969: 820, figs 18–19, 22, 27–28.

*Eidmanniella eurygaster* (Nitzsch, 1866) [sic]; Ryan & Price 1969: 820.

*Eidmanniella subrotunda* (Piaget, 1880); Pilgrim & Palma 1982: 14.

*Eidmanniella subrotunda* (Piaget, 1880); Murray *et al.* 1990: 1372–1373.

*Eidmanniella subrotunda* (Piaget, 1880); Green & Palma 1991: 6, 28.

*Eidmanniella subrotunda* (Piaget, 1880); Palma 1996b: 125.

*Eidmanniella eurygaster* (Nitzsch, 1866) [sic]; Castro & Cicchino 1996: 139.

*Eidmanniella eurygaster* (Nitzsch [in Giebel], 1866); Price *et al.* 2003: 107.

*Eidmanniella subrotunda* (Piaget, 1880); Palma 2010: 407.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Lectotype ♂ of *Menopon subrotundum* in NHML (Palma 1996b: 125).

Type host: *Phalacrocorax brasilianus* (J.F. Gmelin, 1789).

New Zealand hosts: *Phalacrocorax melanoleucos brevirostris* Gould, 1837; *Phalacrocorax sulcirostris* (J.F. Brandt, 1837).

Other hosts: *Phalacrocorax africanus* (J.F. Gmelin, 1789); *Phalacrocorax fuscicollis* Stephens, 1826; *Phalacrocorax niger* (Vieillot, 1817).

New Zealand localities: BP, WO, WA, NC, WD.

Geographic distribution: Americas; Asia; Africa; Australasia; Pacific Islands.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Guimarães (1943); Ryan & Price (1969); Green & Palma (1991); Castro & Cicchino (1996); Palma (1996b); Price *et al.* (2003).

Remarks: This is the first publication where the species name *Eidmanniella eurygaster* is applied to the New Zealand populations of this louse. *Eidmanniella subrotunda* was synonymised under *E. eurygaster* by Castro & Cicchino (1996: 139), but Palma (2010: 407) overlooked that synonymy.

### ***Eidmanniella pellucida* (Rudow, 1869)**

*Menopon pellucidum* Rudow, 1869a: 400.

*Menopon brevialpis* Piaget, 1880: 498, pl. 40: fig. 5.

*Menopon sigmoidalis* Picaglia, 1885: 87.

*Menopon kuwani* Kellogg & Chapman, 1902: 26, pl. 3: fig. 4.

- Eidmanniella brevipalpis* (Piaget, 1880); Hopkins & Clay 1952: 129.  
*Eidmanniella kuwani* (Kellogg & Chapman, 1902); Hopkins & Clay 1952: 129.  
*Eidmanniella pellucida* (Rudow, 1869); Hopkins & Clay 1952: 129.  
*Eidmanniella pellucida* (Rudow, 1869); Ryan & Price 1969: 819, figs 1–2, 4, 6, 14–15, 21, 24, 31, 33.  
*Eidmanniella pellucida* (Rudow, 1869); Wise 1977: 57.  
*Eidmanniella pellucida* (Rudow, 1869); Pilgrim & Palma 1982: 14.  
*Eidmanniella pellucida* (Rudow, 1869); Murray *et al.* 1990: 1372.  
*Eidmanniella pellucida* (Rudow, 1869); Palma 2010: 407.

Neotype ♀ in NHML (Ryan & Price 1969: 820).

Type host: *Phalacrocorax capensis* (Sparrman, 1788).

New Zealand hosts: *Phalacrocorax varius varius* (J.F. Gmelin, 1789); *Phalacrocorax carbo novaehollandiae* Stephens, 1826; *Leucocarbo carunculatus* (J.F. Gmelin, 1789); *Leucocarbo chalconotus* (G.R. Gray, 1845); *Stictocarbo punctatus punctatus* (Sparrman, 1786); *Stictocarbo punctatus oliveri* Mathews, 1930; *Stictocarbo featherstoni* (Buller, 1873).

Other hosts: *Leucocarbo albiventer* (Lesson, 1831); *Leucocarbo bougainvillii* (Lesson, 1837); *Phalacrocorax aristotelis* (Linnaeus, 1761); *Phalacrocorax auritus* (Lesson, 1831); *Phalacrocorax fuscescens* (Vieillot, 1817); *Phalacrocorax gaimardi* (Lesson & Garnot, 1828); *Phalacrocorax magellanicus* (J.F. Gmelin, 1789); *Phalacrocorax pelagicus* Pallas, 1811; *Phalacrocorax penicillatus* (J.F. Brandt, 1837).

New Zealand localities: AK, HB, WN, SD, MB, KA, NC, MC, SC, CO, DN, CH, SI.

Geographic distribution: Europe; Africa; North & South America; Australasia.

New Zealand references: Ryan & Price (1969: 820); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999: 378); Paterson *et al.* (1999: 222); Galloway (2005: 17); Palma (2010).

Other significant references: Clay (1976b: 537); Green & Palma (1991: 6, 28); Forrester *et al.* (1995: 8); Palma (1996b: 125); Price *et al.* (2003: 107); Palma & Jensen (2005: 52, 61).

Remarks: *Eidmanniella pellucida* is a widespread species and highly prevalent on its New Zealand hosts.

#### ***Eidmanniella pustulosa* (Nitzsch [*in* Giebel], 1866)**

- Menopon pustulosum* Nitzsch [*in* Giebel], 1866: 393.  
*Eidmanniella pustulosa* (Nitzsch, 1866) [sic]; Hopkins & Clay 1952: 129.  
*Eidmanniella pustulosa* (Nitzsch, 1866) [sic]; Ryan & Price 1969: 821, figs 7, 16–17, 20, 23, 32.  
*Eidmanniella pustulosa* (Nitzsch, 1866) [sic]; Wise 1977: 57.  
*Eidmanniella pustulosa* (Nitzsch, 1866) [sic]; Pilgrim & Palma 1982: 14.  
*Eidmanniella pustulosa* (Nitzsch [*in* Giebel], 1866); Murray *et al.* 1990: 1372.  
*Eidmanniella pustulosa* (Nitzsch, 1866) [sic]; Palma 2010: 407.

Neotype ♂ in NHML (Ryan & Price 1969: 822).

Type host: *Morus bassanus* (Linnaeus, 1758).

New Zealand host: *Morus serrator* (G.R. Gray, 1843).

Other host: *Morus capensis* (Lichtenstein, 1823).

New Zealand localities: AK, BP, HB, TK, WN, SD, MB, NC, MC, SC, CO, DN.

Geographic distribution: Europe; Africa; Australasia.

New Zealand references: Ryan & Price (1969: 822); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Tendeiro (1958: 446, pl. 3: figs 5–6); Green & Palma (1991: 6, 28); Martín-Mateo (1992a: 40, figs 4–9); Forrester *et al.* (1995: 7); Palma (1996b: 125); Martín-Mateo (2002: 51, figs 14–15); Price *et al.* (2003: 108); Palma & Jensen (2005: 52, 61).

Remarks: *Eidmanniella pustulosa* is highly prevalent on the Australasian gannet, and exclusively parasitic on all species of gannets.

#### **Genus *Eucolpocephalum* Bedford, 1930**

- Eucolpocephalum* Bedford, 1930. *16th Report Director Vet. Services Animal Ind. Union of South Africa*: 161. Type species: *Eucolpocephalum robustum* Bedford, 1930 = *Eucolpocephalum femorale* (Piaget, 1880) (by original designation).

***Eucolpocephalum femorale* (Piaget, 1880)**

Figs 31–32

*Menopon femorale* Piaget, 1880: 484, pl. 39: fig. 8.*Eucolpocephalum robustum* Bedford, 1930: 161: fig. 6.*Eucolpocephalum femorale* (Piaget, 1880); Hopkins & Clay 1952: 133.*Eucolpocephalum femorale* (Piaget, 1880); Pilgrim & Palma 1982: 16.*Eucolpocephalum femorale* (Piaget, 1880); Murray *et al.* 1990: 1373.*Eucolpocephalum femorale* (Piaget, 1880); Martín-Mateo 1994: 110, figs 1c,d, 2c.*Eucolpocephalum femorale* (Piaget, 1880); Price *et al.* 2003: 108.*Eucolpocephalum femorale* (Piaget, 1880); Palma 2010: 408.

Lectotype ♂ in NHML (Clay 1949b: 828).

Type host: *Platalea leucorodia* Linnaeus, 1758.New Zealand host: *Platalea regia* Gould, 1838.Other hosts: *Platalea alba* Scopoli, 1786; *Ajaia ajaja* (Linnaeus, 1758); *Phimosus infuscatus berlepschi* Hellmayr, 1903.

New Zealand locality: Unknown.

Geographic distribution: Eurasia; Africa; Americas; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).Other significant references: Tuff (1966: 379, figs 1–3); Palma (1996b: 126); Martín-Mateo (1994; 2002: 48, figs 12–13); Price *et al.* (2003).

Remarks: *Eucolpocephalum femorale* is a monotypic louse species parasitic on spoonbills and ibises. Since the first record of the royal spoonbill in New Zealand in 1861, this host has become well established with many breeding localities in this country (Checklist Committee 2010: 167). However, there is only one sample of *E. femorale* from this country, without collection data other than the host name.

**Genus *Franciscoloa* Conci, 1942****Subgenus *Franciscoloa* Conci, 1942**

*Franciscoloa* Conci, 1942c. *Boll. Soc. Entomol. Italiana* 74: 35. Type species: *Franciscoloa cacatuae* Conci, 1942c = *Franciscoloa pallida* (Piaget, 1880) (by original designation).

***Franciscoloa (Franciscoloa) pallida* (Piaget, 1880)**

Figs 33–34

*Colpocephalum pallidum* Piaget, 1880: 526, pl. 43: fig. 9.*Franciscoloa cacatuae* Conci, 1942c: 35, figs 5–12.*Psittacomenopon pallidum* (Piaget, 1880); Hopkins & Clay 1952: 305.*Franciscoloa pallida* (Piaget, 1880); Price & Beer 1966: 636, figs 5–8, 10–12, 19.*Franciscoloa (Franciscoloa) pallida* (Piaget, 1880); Price 1967c: 511.*Franciscoloa (Franciscoloa) pallida* (Piaget, 1880); Palma 1999: 381.*Franciscoloa (Franciscoloa) pallida* (Piaget, 1880); Price *et al.* 2003: 109.*Franciscoloa (Franciscoloa) pallida* (Piaget, 1880); Palma 2010: 408.

Lectotype ♂ in NHML (Palma 1996b: 127).

Type host: *Cacatua moluccensis* (J.F. Gmelin, 1788).New Zealand host: *Cacatua galerita* (Latham, 1790).Other host: *Cacatua ducorpsii* Pucheran, 1853.

New Zealand locality: WI.

Geographic distribution: Australasia.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Price & Beer (1966); Price (1967c); Green & Palma (1991: 6, 34); Palma (1996b: 127); Price *et al.* (2003).

Remarks: It is not possible to assert if *Franciscoloa (Franciscoloa) pallida* was introduced to New Zealand with sulphur-crested cockatoos by human agency, or if it was self-introduced with its host (Checklist Committee 2010: 252).

**Genus *Heteromenopon* Carriker, 1954**

*Heteromenopon* Carriker, 1954. *Rev. Bras. Entomol.* 2: 170. Type species: *Heteromenopon sincipitalis* Carriker, 1954 (by original designation).

**Subgenus *Keamenopon* Price & Beer, 1967**

*Keamenopon* Price & Beer, 1967. *Ann. Entomol. Soc. America* 60: 335. Type species: *Heteromenopon kea* (Kellogg, 1907) (by original designation).

***Heteromenopon (Keamenopon) kea* (Kellogg, 1907)**

Figs 35–36

*Menopon fulvofasciatum* var. *kea* Kellogg, 1907: 122.

*Menopon fulvofasciatum kea* Kellogg, 1907; Harrison 1916: 39.

*Menopon kea* Kellogg, 1907; Conci 1942c: 36.

*Psittacomenopon kea* (Kellogg, 1907); Hopkins & Clay 1952: 305.

*Heteromenopon (Keamenopon) kea* (Kellogg, 1907); Price & Beer 1967: 336, figs 12, 25, 32.

*Psittacomenopon kea*; Miller 1971: 132.

*Heteromenopon kea* (Kellogg, 1907); Wise 1977: 57.

*Heteromenopon (Keamenopon) kea* (Kellogg, 1907); Pilgrim & Palma 1982: 24.

*Heteromenopon (Keamenopon) kea* (Kellogg, 1907); Murray *et al.* 1999: 1241.

*Heteromenopon (Keamenopon) kea* (Kellogg, 1907); Palma 2010: 408.

Lectotype ♂ in EMEC (Price & Beer, 1967: 337).

Type host: *Nestor notabilis* Gould, 1856.

New Zealand hosts: *Strigops habroptilus* G.R. Gray, 1845; *Nestor meridionalis septentrionalis* Lorenz, 1896; *Nestor meridionalis meridionalis* (J.F. Gmelin, 1788); *Nestor notabilis* Gould, 1856.

Other hosts: None.

New Zealand localities: ND, CL, HB, WN, NN, NC, MC, WD, SI.

Geographic distribution: New Zealand.

New Zealand references: Kellogg (1907); Conci (1942c); Price & Beer (1967: 337); Pilgrim (1970: 75); Miller (1971: 132); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1999); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Price *et al.* (2003: 110).

Remarks: *Heteromenopon (Keamenopon) kea* is an endemic and “at risk” species (Buckley *et al.* 2012), exclusively parasitic on members of the parrot family Strigopidae (*sensu* Checklist Committee 2010: 249).

***Heteromenopon (Keamenopon) species***

*Heteromenopon (Keamenopon) sp.*; Pilgrim & Palma 1982: 24.

*Heteromenopon (Keamenopon) sp.*; Murray *et al.* 1999: 1241.

*Heteromenopon (Keamenopon) species*; Marris 2000: 188.

New Zealand host: *Cyanoramphus unicolor* (Lear, 1831).

Other hosts: None.

New Zealand locality: AN.

Geographic distribution: Antipodes Islands, New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1999); Marris (2000).

Other significant references: Price & Beer (1967: 335); Price (1969).

Remarks: The single record of this louse (voucher specimens in MONZ) could not be identified beyond the subgenus due to the poor condition of the only two available specimens which does not allow an identification of the species.

**Genus *Hohorstiella* Eichler, 1940**

*Hohorstiella* Eichler, 1940. *Zentralbl. Bakter. Parasitenkd. Infekt.* 145: 362. Type species: *Menopon latum* Piaget, 1880 = *Hohorstiella lata* (Piaget, 1880) (by original designation).



**Hohorstiella lata (Piaget, 1880)**

*Menopon latum* Piaget, 1880: 457, pl. 37: fig. 1.

*Menopon latum* Piaget, 1880; Harrison 1916: 39 (as junior synonym of *Menopon giganteum* Denny, 1842).

*Hohorstiella lata* (Piaget, 1880); Hopkins & Clay 1952: 173.

*Hohorstiella lata* (Piaget, 1880); Pilgrim 1976: 160, fig. 3.

*Hohorstiella lata* (Piaget, 1880); Wise 1977: 57.

*Hohorstiella lata* (Piaget, 1880); Pilgrim & Palma 1982: 23.

*Hohorstiella lata* (Piaget, 1880); Murray *et al.* 2006a: 1965.

*Hohorstiella lata* (Piaget, 1880); Palma 2010: 408.

Lectotype ♂ in NHML (Clay 1949b: 835).

Type host: *Columba livia domestica* Linnaeus, 1758.

New Zealand host: *Columba livia* J.F. Gmelin, 1789.

Other hosts: None.

New Zealand locality: MC, SC.

Geographic distribution: Eurasia; north Africa; Americas; Australasia.

New Zealand references: Pilgrim (1970: 76); Pilgrim (1976); Wise (1977); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Martín-Mateo (2002: 105); Murray *et al.* (2006a); Palma (2010).

Other significant references: Emerson (1972b: 64); Nelson & Murray (1971: 22, 25 figs 1, 6); Hill & Tuff (1978: 310); Palma (1996b: 129); Price *et al.* (2003: 111); Galloway & Lamb (2014: 445; 2015: 715).

Remarks: *Hohorstiella lata* was introduced to New Zealand and other countries with rock pigeons by human agency (Checklist Committee 2010: 245). Notwithstanding the abundance of rock pigeons in New Zealand, and unlike other lice from these hosts, records of *H. lata* are few and only from one region.

**Hohorstiella timorensis Tendeiro, 1980****New Record**

Figs 37–38

*Hohorstiella timorensis* Tendeiro, 1980: 9, figs 3–4, 7–8, photos 2–3.

*Hohorstiella* sp.; Pilgrim & Palma 1982: 25.

*Hohorstiella* sp.; Paterson *et al.* 1999: 223.

*Hohorstiella* sp.; Murray *et al.* 2006a: 1966.

*Hohorstiella* sp.; Palma 2010: 408.

Holotype ♂ in CZLP.

Type host: *Ducula cineracea* (Temminck, 1835).

New Zealand host: *Hemiphaga novaeseelandiae* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: ND, AK, HB, TK, WN, MB.

Geographic distribution: Timor & Wetar Islands (Indonesia); New Zealand.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999: 223); Murray *et al.* (2006a); Palma (2010).

Other significant reference: Price *et al.* (2003: 111).

Material examined and repository: 69♂, 72♀ (13 samples, MONZ).

Remarks: This is the first record of *Hohorstiella timorensis* for New Zealand because the New Zealand references cited above reported this louse as “*Hohorstiella* sp.” only. The type host of *H. timorensis* is endangered (Trainor 2002: 10). However, the New Zealand pigeon is common and widespread (Checklist Committee 2010: 247).

**Genus Holomenopon Eichler, 1941**

*Holomenopon* Eichler, 1941c. *Stettin. Entomol. Zeit.* 102: 125. Type species: *Menopon albofasciatum* Piaget, 1880 = *Holomenopon leucoxanthum* (Burmeister, 1838) (by original designation).

**Holomenopon leucoxanthum (Burmeister, 1838)**

*Menopon leucoxanthum* Burmeister, 1838a: 440.

*Menopon leucoxanthum* Nitzsch [in Burmeister], 1838; Séguy 1944: 92, figs 109–111.

*Holomenopon leucoxanthum* (Burmeister, 1838); Hopkins & Clay 1952: 174.

*Holomenopon leucoxanthum* (Burmeister, 1838); Price 1971: 635, figs 2–13.

*Holomenopon leucoxanthum* (Burmeister, 1838); Pilgrim & Palma 1982: 16–17.

*Holomenopon leucoxanthum* (Burmeister, 1838); Palma 1999: 379.

*Holomenopon leucoxanthum* (Burmeister, 1838); Palma 2010: 408.

Neotype ♀ in USNM (Price 1971: 635).

Type host: *Anas crecca crecca* Linnaeus, 1758.

New Zealand hosts: *Dendrocygna eytoni* (Eyton, 1838); *Cygnus atratus* (Latham, 1790); *Branta canadensis maxima* Delacour, 1951; *Anas gracilis* Buller, 1869; *Anas aucklandica* (G.R. Gray, 1849); *Anas platyrhynchos platyrhynchos* Linnaeus, 1758.

Other hosts: Nine other species of *Anas*; seven species of *Aythya*; four other species of *Dendrocygna*; two species of *Netta*; two species of *Tadorna*; *Anser anser* (Linnaeus, 1758); *Bucephala albeola* (Linnaeus, 1758); *Cairina moschata* (Linnaeus, 1758); *Clangula hyemalis* (Linnaeus, 1758); *Melanitta nigra* (Linnaeus, 1758); *Nettapus cormandelianus* (J.F. Gmelin, 1789); *Oxyura jamaicensis* (J.F. Gmelin, 1789); *Sarkidiornis melanotos* (Pennant, 1769) (see Price *et al.* 2003: 112).

New Zealand localities: ND, AK, BP, WO, HB, TK, WA, CH, AU.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990: 1374); Paterson *et al.* (1999: 219); Palma (1999); Palma (2010).

Other significant references: Price (1971); Humphreys (1975: 96); Moreby (1976: 92); Clay (1976b: 537); Green & Palma (1991: 6, 29); Forrester *et al.* (1995: 14); Palma (1996b: 130); Martín-Mateo (2002: 54, figs 16B,D, 17); Price *et al.* (2003: 112); Palma & Jensen (2005: 52, 62); Ahmad *et al.* (2015: 568).

Remarks: *Dendrocygna eytoni* is a straggler to New Zealand, recorded occasionally since 1871 (Checklist Committee 2010: 31), and is a new host record for *Holomenopon leucoxanthum* in New Zealand (voucher specimens in MONZ).

### ***Holomenopon tadornae* (Gervais, 1844)**

Figs 39–40

*Philopterus tadornae* Gervais, 1844: 323, pl. 49: fig. 6.

*Menopon tadornae* Gervais, 1844 [sic]; Harrison 1916: 45.

*Holomenopon tadornae* (Gervais, 1844); Hopkins & Clay 1952: 175.

*Holomenopon tadornae* (Gervais, 1844) sensu lato; Price 1971: 640, figs 17–18.

*Holomenopon tadornae* (Gervais, 1844); Pilgrim & Palma 1982: 16.

*Holomenopon tadornae* (Gervais, 1844); Palma 1999: 379.

*Holomenopon tadornae* (Gervais, 1844); Palma 2010: 408.

Status, sex and repository of types unknown.

Type host: *Tadorna tadorna* (Linnaeus, 1758).

New Zealand hosts: *Tadorna variegata* (J.F. Gmelin, 1789); *Tadorna tadornoides* (Jardine & Selby, 1828).

Other hosts: *Branta bernicla* (Linnaeus, 1758); *Alopochen aegyptiaca* (Linnaeus; 1766); *Chloephaga melanoptera* (Eyton, 1838); *Tadorna ferruginea* (Pallas).

New Zealand localities: HB, NN, NC, MC, SC, FD.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990: 1374); Palma (1999); Palma (2010).

Other significant references: Price (1971); Price *et al.* (2003: 113).

Remarks: *Holomenopon tadornae* appears to be restricted to species of the tribe Tadornini.

### ***Holomenopon* species 1**

“*Holomenopon clypeilargum*” Price, 1971: 643 (not *Holomenopon clypeilargum* Eichler, 1943).

“*Holomenopon clypeilargum*” Wise, 1977: 57 (not *Holomenopon clypeilargum* Eichler, 1943).

*Holomenopon* sp.; Pilgrim & Palma 1982: 17, 30, note 19.

*Holomenopon* sp.; Murray *et al.* 1990: 1374.

*Holomenopon clypeilargum* Eichler, 1943; Price *et al.* 2003: 112, 279. In part.

*Holomenopon* sp.; Palma 2010: 408.

New Zealand host: *Aythya novaeseelandiae* (J.F. Gmelin, 1789).

New Zealand locality: BP.

Geographic distribution: New Zealand.

New Zealand references: Price (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Price *et al.* (2003: 112, 279); Palma (2010).

Other significant references: None.

Remarks: This entry represents an undescribed and unnamed endemic species, but more samples are needed before a proper description of the species can be prepared and published. Price (1971: 643) misidentified a female from *Aythya novaeseelandiae* as “*Holomenopon clypeilargum*”, establishing an incorrect host-louse association which was adopted by Wise (1977: 57) and Price *et al.* (2003: 279). Therefore, the host listed as “*A. novaeseelandiae* (Gmelin)” under *Holomenopon clypeilargum* in Price *et al.* (2003: 112) is incorrect. See also “Species & subspecies deleted from the New Zealand louse fauna” below.

### **Holomenopon species 2**

*Holomenopon* sp.; Pilgrim & Palma 1982: 17.

*Holomenopon* sp.; Mourik & Norman 1985: 2.

*Holomenopon* sp.; Murray *et al.* 1990: 1374.

*Holomenopon* sp.; Palma 2010: 408.

New Zealand host: *Anas rhynchotis* Latham, 1802.

New Zealand locality: MC.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant reference: Mourik & Norman (1985).

Remarks: This entry represents an undescribed and unnamed species, but the material available is insufficient to publish a formal description.

### **Genus *Kurodaia* Uchida, 1926**

*Kurodaia* Uchida, 1926. *Jour. Coll. Agric. Tokyo* 9: 50. Type species: *Colpocephalum haliaeeti* Denny, 1842 = *Kurodaia (Kurodaia) haliaeeti* (Denny, 1842) (by original designation).

#### **Subgenus *Conciella* Eichler, 1949**

*Conciella* Eichler, Eichler, 1949a. *Boll. Soc. Entomol. Italiana* 79: 11. Type species: *Colpocephalum painei* McGregor, 1912 = *Kurodaia (Conciella) painei* (McGregor, 1912) (by original designation).

### ***Kurodaia (Conciella) cryptostigmatia* (Nitzsch [*in* Giebel], 1861)**

Figs 41–42

*Menopon cryptostigmatia* Nitzsch [*in* Giebel], 1861a: 529.

*Conciella cryptostigmatia*; Eichler 1949a: 11.

*Kurodaia cryptostigmatia* (Nitzsch, 1861) [sic]; Hopkins & Clay 1952: 181.

*Kurodaia cryptostigmatia* (Nitzsch, 1861) [sic]; Price & Beer 1963b: 850, 1–4, 10B.

*Kurodaia cryptostigmatia* (Nitzsch, 1861) [sic]; Pilgrim & Palma 1982: 25.

*Kurodaia cryptostigmatia* (Nitzsch, 1861) [sic]; Murray *et al.* 1999: 1241.

*Kurodaia (Conciella) cryptostigmatia* (Nitzsch [*in* Giebel], 1861); Price *et al.* 2003: 114.

*Kurodaia cryptostigmatia* (Nitzsch, 1861) [sic]; Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Athene noctua* (Scopoli, 1769).

New Zealand host: *Ninox novaeseelandiae novaeseelandiae* (J.F. Gmelin, 1788).

Other hosts: *Otus senegalensis* (Swainson, 1837); *Otus scops* (Linnaeus, 1758); *Otus rutilus* (Pucheran, 1849); *Otus cooperi* (Ridgway, 1878); *Strix aluco* Linnaeus, 1758; *Glaucidium capense* (A. Smith, 1834); *Glaucidium passerinum* (Linnaeus, 1758); *Aegolius funereus* (Linnaeus, 1758); *Asio madagascariensis* (A. Smith, 1834).

New Zealand localities: AK, BP, BP, TO, TK, WN, SD, MB, WD.

Geographic distribution: Eurasia; Africa; Central America; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1999); Palma (2010).

Other significant references: Price & Beer (1963b); Green & Palma (1991: 6, 36); Palma (1996b: 131); Martín-Mateo (2002: 83, fig. 27C); Price *et al.* (2003).

Remarks: *Kurodaia (C.) cryptostigmatia* has not yet been found on the New Zealand population of *Athene noctua*, which was originally introduced by humans in 1906–1911 (Checklist Committee 2010: 267).

### Genus *Longimenopon* Thompson, 1948

“New genus F” Clay, 1947. *Proc. Zool. Soc. London* 117: 466, 473.

*Longimenopon* Thompson, 1948d. *Occ. Pap. Bishop Mus. Honolulu* 19: 197. Type species: *Longimenopon puffinus* Thompson, 1948 (by original designation).

### *Longimenopon galeatum* Timmermann, 1957

*Longimenopon galeatum* Timmermann, 1957b: 9, figs 2c, 7–8.

*Longimenopon galeatum* Timmermann, 1957; Timmermann 1965: 182, fig. 117c, pl. 12: fig. 5.

*Longimenopon galeatum* Timmermann, 1957; Clay & Moreby 1967: 159, 168, fig. 54.

*Longimenopon galeatum* Timmermann, 1957; Watson 1967: 72.

*Longimenopon galeatum* Timmermann, 1957; Clay & Moreby 1970: 218.

*Longimenopon galeatum* Timmermann, 1957; Wise 1977: 57.

*Longimenopon galeatum* Timmermann, 1957; Palma & Horning 2002: 6, 16.

*Longimenopon galeatum* Timmermann, 1957; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Pelagodroma marina* (Latham, 1790).

New Zealand host: *Pachyptila desolata* (J.F. Gmelin, 1789).

Other host: *Lugensa brevirostris* (Lesson, 1833).

New Zealand localities: AU, Macquarie Island.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Timmermann (1957b: 12); Watson (1967); Clay & Moreby (1970); Gressitt (1970: 326); Wise (1977); Palma (1996b: 132); Palma & Horning (2002); Palma (2010).

Other significant references: Timmermann (1965); Clay & Moreby (1967); Price *et al.* (2003: 115); Hänel & Palma (2007: 112, 122, 130).

Remarks: I have included *Longimenopon galeatum* following records from New Zealand by Timmermann (1957b), Watson (1967) and others (see above). However, until a complete revision of the genus *Longimenopon* becomes available, I am not able to specifically name with confidence any sample of this genus from any host (see below).

### *Longimenopon* species

Figs 43–44

*Longimenopon* sp.; Nelson 1969: 199.

*Longimenopon* sp.; Watt 1971: 233, 242.

*Longimenopon* sp.; Palma & Pilgrim 1977: 290.

*Longimenopon* sp.; Horning *et al.* 1980: 4, 9.

*Longimenopon* sp.; Pilgrim & Palma 1982: 8–11, 13.

*Longimenopon* sp.; Murray *et al.* 1990: 1369–1370, 1372.

*Longimenopon* sp.; Paterson *et al.* 1999: 222.

*Longimenopon* sp.; Palma 1999: 376.

New Zealand hosts: *Lugensa brevirostris* (Lesson, 1833); *Pterodroma magentae* (Giglioli & Salvadori, 1869); *Pterodroma neglecta neglecta* (Schlegel, 1863); *Pterodroma inexpectata* (J.R. Forster, 1844); *Pterodroma nigripennis* (Rothschild, 1893); *Pterodroma cookii* (G.R. Gray, 1843); *Pterodroma longirostris* (Stejneger, 1888); *Pterodroma pycrofti* Falla, 1933; *Halobaena caerulea* (J.F. Gmelin, 1789); *Pachyptila vittata* (G. Forster, 1777); *Pachyptila salvini salvini* (Mathews, 1912); *Pachyptila desolata* (J.F. Gmelin, 1789); *Pachyptila belcheri*

(Mathews, 1912); *Pachyptila turtur* (Kuhl, 1820); *Pachyptila crassirostris pyramidalis* Fleming, 1939; *Pelagodroma marina maoriana* Mathews, 1912.

Other host: *Pterodroma mollis* (Gould, 1844).

New Zealand localities: ND, CL, AK, BP, GB, WI, WN, NN, NC, MC, SC, WD, SL, KE, Norfolk Island, CH, SI, SN, Macquarie Island.

Geographic distribution: Atlantic, Indian and Pacific Oceans.

New Zealand references: Nelson (1969); Watt (1971); Palma & Pilgrim (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Palma & Pilgrim (1983: 148); Murray *et al.* (1990); Paterson *et al.* (1999); Palma (1999); Palma & Imber (2000: 230); Buckley *et al.* (2012: App. 2).

Other significant references: Séguy (1953: 590, fig. 47); Timmermann (1965: 179, figs 116–117, pl. 12: figs 1–4); Green & Palma (1991: 7, 27); Zonfrillo (1993: 327); Furness & Palma (1992: 35, 41); Jensen & Palma (2005: 228).

Remarks: A complete systematic revision of the six species of *Longimenopon* currently accepted as valid (see Price *et al.* 2003: 115), including type material and samples available from many other hosts, needs to be made before any specimen can be confidently named (Pilgrim & Palma 1982: 2). A preliminary examination of many specimens from all the hosts listed above from New Zealand has shown a remarkably uniform morphology, especially regarding chaetotaxy, male genitalia and head shape. Timmermann (1957b: 8) expressed difficulty in deciding where to draw the limits between species, and proceeded to separate the five species he recognised as valid by the shape of the head and overall size. However, both these characters are liable to change significantly during the slide-mounting process, especially because of the very soft exoskeleton of these lice. I have examined specimens from the *same* sample that show a range of variation in head shape—due to the different levels of pressure exerted by the cover-slip—equivalent to that depicted by Timmermann (1957b: 10, fig. 2) for three different species!

### Genus *Menacanthus* Neumann 1912

*Menopon (Menacanthus)* Neumann, 1912a. *Arch. Parasitol., Paris* 15(3): 354. Type species: *Menopon robustum* Kellogg, 1896b = *Menacanthus robustus* (Kellogg, 1896) (by original designation).

*Eomenacanthus* Uchida, 1926. *Jour. Coll. Agric. Tokyo* 9: 30. Type species: “*Eomenacanthus biseriatum* (Piaget)” = *Menacanthus stramineus* (Nitzsch, 1818) (by original designation).

### *Menacanthus eurysternus* (Burmeister, 1838)

*Menopon eurysternum* Burmeister, 1838a: 439.

*Menopon* sp.; Johnston & Harrison 1912: 364.

*Menacanthus mutabilis* Blagoveshtchensky, 1940: 31, 78, fig. 5.

*Menacanthus eurysternum* [sic] (Burmeister, 1838); Hopkins & Clay 1952: 210.

*Menacanthus mutabilis* Blagoveshtchensky, 1940; Watt 1971: 233, 244, fig. 3.

*Menacanthus* sp.; Watt 1971: 233, 244.

*Menacanthus eurysternus* (Burmeister, 1838); Price 1975: 617, figs 1–9.

*Menacanthus eurysternus* (Burmeister, 1838); Wise 1977: 57.

*Menacanthus eurysternus* (Burmeister, 1838); Pilgrim & Palma 1982: 26.

*Menacanthus eurysternus* (Burmeister, 1838); Murray *et al.* 2001: 1263.

*Menacanthus eurysternus* (Burmeister, 1838); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1), Palma & Pilgrim (1984: 150) and Palma (1996b: 133).

Type host: *Pica pica pica* (Linnaeus, 1758).

New Zealand hosts: *Anthornis melanura melanura* (Sparrman, 1786); *Prosthemadera novaeseelandiae novaeseelandiae* (Gmelin, 1788); *Petroica (Miro) australis australis* (Sparrman, 1788); *Zosterops lateralis lateralis* (Latham, 1802); *Turdus merula merula* Linnaeus, 1758; *Turdus philomelos* Brehm, 1831; *Sturnus vulgaris vulgaris* Linnaeus, 1758; *Acridotheres tristis* (Linnaeus, 1766); *Passer domesticus domesticus* (Linnaeus, 1758); *Anthus novaeseelandiae novaeseelandiae* (Gmelin, 1789); *Carduelis chloris* (Linnaeus, 1758); *Serinus canaria* (Linnaeus, 1758) captive.

Other hosts: Over 150 species of the order Passeriformes and seven species of the order Piciformes (see Price *et al.* 2003: 119, 364).

New Zealand localities: ND, CL, AK, HB, TK, WI, WN, KA, NC, MC, SC, WD, CO, DN, KE, CH, SI, SN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Johnston & Harrison (1912); Watt (1971); Price (1975: 621); Wise (1977); Horning *et al.* (1980: 4, 12); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Palma (1999: 381); Murray *et al.* (2001); Murray *et al.* (2002: 1216); Galloway (2005: 14, 16); Murray *et al.* (2006b: 1958, 1960); Palma (2010).

Other significant references: Price (1977: 219); Castro & Cicchino (1978: 78); Kettle (1983: 403); Green & Palma (1991: 7, 38); Palma (1996b: 133); Palma *et al.* (1998: 311, 317); Martín-Mateo (2002: 108); Price *et al.* (2003: 119); Palma & Jensen (2005: 52, 68); Adam (2007: 161, figs 5d, 6a); Martín *et al.* (2015: 64).

Remarks: *Menacanthus eurysternus* is the louse species with the largest number of hosts among all Phthiraptera, with over 170 host species, belonging to more than 20 families in two orders (Price 1975: 620; Price *et al.* 2003: 119). *Carduelis chloris* is a new host record for this louse species in New Zealand (voucher specimens in MONZ). Also, *Serinus canaria* is only a cage bird in New Zealand.

### ***Menacanthus pallidulus* (Neumann, 1912)**

Figs 45–46

*Menopon (Menacanthus) pallidulum* Neumann, 1912a: 361, figs 7–9.

*Menacanthus pallidulus* (Neumann, 1912); Hopkins & Clay 1952: 213.

*Menacanthus* sp.; Pilgrim & Palma 1982: 18.

*Menacanthus* sp.; Murray *et al.* 1993: 960.

*Menacanthus* sp.; Paterson *et al.* 1999: 219.

*Menacanthus pallidulus* Neumann, 1912 [sic]; Palma 1999: 380, 383, note 5.

*Menacanthus pallidulus* (Neumann, 1912); Price *et al.* 2003: 124.

*Menacanthus pallidulus* Neumann, 1912 [sic]; Palma 2010: 408.

Syntypes ♂♀, repository not confirmed, probably in the Ecole Vétérinaire de Toulouse, France.

Type host: *Gallus gallus* (Linnaeus, 1758).

New Zealand hosts: *Gallus gallus gallus* (Linnaeus, 1758); *Phasianus colchicus* Linnaeus, 1758.

Other hosts: *Bambusicola thoracicus* (Temminck, 1815); *Gallus sonneratii* Temminck, 1813.

New Zealand locality: WO, WN, NC, MC, SC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999); Palma (1999); Palma (2010).

Other significant references: Emerson (1956a: 77, pl. 8); Emerson & Ward (1958: 52); Marconcini & Macchioni (1975: 108, figs 9–10); Forrester *et al.* (1995: 20); Martín-Mateo (2002: 110); Price *et al.* (2003).

Remarks: *Menacanthus pallidulus* was introduced to New Zealand and other countries with chickens by human agency (Checklist Committee 2010: 27). Records of this louse from *Phasianus colchicus* may be due to the human practice of using chickens to incubate pheasant eggs in captivity.

*Gallus gallus gallus* has been listed for the first time in the latest edition of the New Zealand Checklist of Birds (Checklist Committee 2010: 27; see also Palma 1999: 383, note 5).

### ***Menacanthus rhipidurae* Palma & Price, 2005**

*Menacanthus* sp.; Pilgrim & Palma 1982: 26.

*Menacanthus* sp.; Paterson *et al.* 1999: 221.

*Menacanthus* sp.; Galloway 2005: 17.

*Menacanthus rhipidurae* Palma & Price, 2005: 112, figs 1–5.

*Menacanthus rhipidurae* Palma & Price, 2005; Murray *et al.* 2006b: 1956.

*Menacanthus rhipidurae* Palma & Price, 2005; Palma 2010: 408.

Holotype ♂ in MONZ.

Type host: *Rhipidura fuliginosa fuliginosa* (Sparman, 1787).

New Zealand host: *Rhipidura fuliginosa fuliginosa* (Sparman, 1787).

Other hosts: None.

New Zealand localities: NN, NC, WD.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999); Galloway (2005); Palma & Price (2005); Murray *et al.* (2006b); Palma (2010).

Other significant references: None.

Remarks: *Menacanthus rhipidurae* is an endemic species, although it is likely that some Australian species of *Rhipidura* closely related to *Rhipidura fuliginosa* (see Checklist Committee 2010: 298) may harbour this species of *Menacanthus* as well.

### ***Menacanthus stramineus* (Nitzsch, 1818)**

*Pediculus meleagridis* Panzer, 1793: 51, fig. 20. Preoccupied by *Pediculus meleagridis* Linnaeus, 1758: 613.

*Liotheum (Menopon) stramineum* Nitzsch, 1818: 300. *Nomen novum* for *Pediculus meleagridis* Panzer, 1793.

*Menopon biseriatum* Piaget, 1880: 469, pl. 37: fig. 2.

*Menopon stramineum* Nitzsch [*in* Giebel], 1874 [*sic*]; Harrison 1916: 45.

*Menopon biseriatim* [*sic*] Nitzsch [*sic*]; Thomson 1922: 270.

*Menacanthus stramineus* (Nitzsch, 1818); Hopkins & Clay 1952: 215.

*Eomenacanthus stramineus* (Nitzsch); Neuffer 1954: 452, figs 5, 8, 19, 28, 33, 42, 44.

*Eomenacanthus* [*sic*] *stramineus*; Helson 1956: 13.

*Menacanthus stramineus* (Nitzsch); Whitten 1971: 383.

*Menacanthus stramineus* (Nitzsch, 1818); Wise 1977: 57.

*Menacanthus stramineus* (Nitzsch, 1818); Pilgrim & Palma 1982: 18.

*Menacanthus stramineus* (Nitzsch, 1818); Murray *et al.* 1993: 960.

*Menacanthus stramineus* (Nitzsch, 1818); Palma 1999: 384, note G.

*Menacanthus stramineus* (Nitzsch, 1818); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1960: 48).

Type host: *Meleagris gallopavo* Linnaeus, 1758.

New Zealand hosts: *Meleagris gallopavo* Linnaeus, 1758; *Gallus gallus gallus* (Linnaeus, 1758).

Other hosts: *Numida meleagris* (Linnaeus, 1758); *Tragopan satyra* (Linnaeus, 1758); *Lophura leucomelanos* (Latham, 1790); *Phasianus colchicus* Linnaeus, 1758; *Pavo cristatus* Linnaeus, 1758.

New Zealand localities: BP, WO, HB, WN, NC, MC, SC, CO, DN, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Thomson (1922); Helson (1956); Whitten (1971); Watt (1971: 233); Pilgrim (1974: 1035, fig. 6); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 220); Palma (1999); Palma (2010).

Other significant references: Webb (1946: 51); Neuffer (1954); Emerson (1956a: 77, pl. 9); Clay & Hopkins (1960: 47, figs 73–74); Emerson (1962: 196, figs 1–3); Forrester *et al.* (1995: 21); Palma (1996b: 135); Martín-Mateo (2002: 112, fig. 36); Price *et al.* (2003: 125); Palma & Jensen (2005: 53, 62); Adam (2007: 163, figs 6d, 7a).

Remarks: *Menacanthus stramineus* was introduced to New Zealand and other countries with chickens or turkeys by human agency. This louse has spread onto a number of adventive hosts due to the human practice of mixing various species of game birds in captivity. *Menacanthus stramineus* is haematophagous (Wilson 1933); consequently, it is a serious pest of captive birds by causing anaemia and transmitting microorganisms that cause disease (Derylo 1970, 1977).

*Gallus gallus gallus* has been listed for the first time in the latest edition of the New Zealand Checklist of Birds (Checklist Committee 2010: 27; see also Palma 1999: 383, note 5).

### ***Menacanthus* species**

*Menacanthus* sp.; Pilgrim & Palma 1982: 25.

*Menacanthus* sp.; Murray *et al.* 2001: 1262.

New Zealand host: *Acanthisitta chloris chloris* (Sparrman, 1787).

Other hosts: None.

New Zealand localities: NN, FD.

Geographic distribution: New Zealand: South Island.

New Zealand references: Pilgrim & Palma (1982); Murray et al. (2001).

Other significant reference: Price (1977).

Remarks: The two available records of *Menacanthus* from the South Island rifleman could not be identified to species because the samples contain females only (voucher specimens in MONZ).

### Genus *Menopon* Nitzsch, 1818

*Menopon* Nitzsch, 1818. *German's Mag. Entomol.* 3: 299. Type species: *Menopon gallinae* (Linnaeus, 1758) (by subsequent designation).

#### *Menopon gallinae* (Linnaeus, 1758)

Figs 47–48

*Pediculus gallinae* Linnaeus, 1758: 613.

*Ricinus gallinae* (Linnaeus, 1758); Latreille 1804: 109.

*Liotheum (Menopon) pallidum* Nitzsch, 1818: 299.

*Menopon pallidum* Nitzsch; Thomson 1922: 269.

*Menopon gallinae* (Linnaeus, 1758); Hopkins & Clay 1952: 219.

*Menopon gallinae* (L.); Helson 1956: 13, 17.

*Menopon gallinae* (Linnaeus, 1758); Watt 1971: 233, 243.

*Menopon gallinae* (Linnaeus, 1758); Wise 1977: 57.

*Menopon gallinae* (Linnaeus, 1758); Palma 1999: 384, note G.

*Menopon gallinae* (Linnaeus, 1758); Price et al. 2003: 126.

*Menopon gallinae* (Linnaeus, 1758); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1950: 262).

Type host: *Gallus gallus* (Linnaeus, 1758).

New Zealand hosts: *Gallus gallus gallus* (Linnaeus, 1758); *Numida meleagris* (Linnaeus, 1758) captive.

Other hosts: *Meleagris gallopavo* Linnaeus, 1758; *Caloperdix ocellus* (Temminck, 1815); *Tragopan satyra* (Linnaeus, 1758); *Gallus sonneratii* Temminck, 1813; *Gallus lafayettei* Lesson, 1831; *Syrmaticus mikado* (Ogilvie-Grant, 1906) and seven species of *Lophura* (see Price et al. 2003: 126).

New Zealand localities: AK, TO, WI, WN, NC, MC, SC, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Thomson (1922); Tillyard (1926: 134); Helson (1956); Miller (1971: 131); Whitten (1971: 383); Watt (1971); Wise (1977); Palma (1999); Palma (2010).

Other significant references: Ferris (1924: 57, fig. 1); Clay & Hopkins (1950: 262, fig. 56); Emerson (1954: 225, fig. 8); Emerson (1956a: 77, pl. 10); Emerson & Ward (1958: 54); Rudolph (1983: 16); Price (1987: 218, figs 22.2, 22.8, 22.11–22.13); Palma (1996b: 136); Martín-Mateo (2002: 103, fig. 34); Adam (2007: 158, figs 4a,b); Price et al. (2003: 126); Palma & Jensen (2005: 53, 62); Palma & Peck (2013: 21).

Remarks: *Menopon gallinae* was introduced to New Zealand and other countries with chickens by human agency. *Gallus gallus gallus* has been listed for the first time in the latest edition of the New Zealand Checklist of Birds (Checklist Committee 2010: 27; see also Palma 1999: 383, note 5).

### Genus *Myrsidea* Waterston, 1915

*Myrsidea* Waterston, 1915. *Entomol. Month. Mag.* 51: 12. Type species: *Myrsidea victrix* Waterston, 1915 (by original designation).

#### *Myrsidea hihī* Sychra, Kolencik & Palma, 2016

*Myrsidea* sp.; Pilgrim & Palma 1982: 27.

*Myrsidea* sp.; Murray et al. 2001: 1263.

*Myrsidea* sp.; Palma 2010: 408.

*Myrsidea hihī* Sychra, Kolencik & Palma, 2016: 405, figs 12–13, 22, 29–30.

Holotype ♀ in MONZ.



Type host: *Notiomystis cincta* (du Bus de Gisignies, 1839).

New Zealand host: *Notiomystis cincta* (du Bus de Gisignies, 1839).

Other hosts: None.

New Zealand localities: CL, WA (captive).

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2001); Palma (2010); Sychra *et al.* (2016).

Other significant references: None.

Remarks: Both *Myrsidea hihi* and its single host, the stitchbird, are endemic to New Zealand and regarded as vulnerable species (Robertson *et al.* 2013: 11).

#### ***Myrsidea ivanliteraki* Sychra, Kolencik & Palma, 2016**

*Myrsidea* sp.; Pilgrim & Palma 1982: 28.

*Myrsidea* sp.; Murray *et al.* 2006b: 1957.

*Myrsidea* sp.; Palma 2010: 408.

*Myrsidea ivanliteraki* Sychra, Kolencik & Palma, 2016: 399, figs 1–6, 14–16, 25–26.

Holotype ♀ in MONZ.

Type host: *Gymnorhina tibicen* (Latham, 1802).

New Zealand host: *Gymnorhina tibicen* (Latham, 1802).

Other hosts: None.

New Zealand localities: ND, CL, BP, GB, TK, WN.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999: 221, 223); Murray *et al.* (2006b); Palma (2010); Sychra *et al.* (2016).

Other significant references: None

Remarks: *Myrsidea ivanliteraki* is native to Australia, and was introduced to New Zealand with Australian magpies by human agency (Checklist Committee 2010: 297).

#### ***Myrsidea novaeseelandiae* Sychra, Kolencik & Palma, 2016**

*Myrsidea* sp.; Pilgrim & Palma 1982: 27.

*Myrsidea* sp.; Murray *et al.* 2001: 1263.

*Myrsidea* sp.; Palma 2010: 408.

*Myrsidea novaeseelandiae* Sychra, Kolencik & Palma, 2016: 401, figs 9–11, 19–21, 27–28.

Holotype ♀ in MONZ.

Type host: *Anthornis melanura oneho* Bartle & Sagar, 1987.

New Zealand hosts: *Anthornis melanura obscura* Falla, 1948; *Anthornis melanura melanura* (Sparrman, 1786); *Prosthemadera novaeseelandiae novaeseelandiae* (J.F. Gmelin, 1788).

Other hosts: None.

New Zealand localities: AK, ND, CL, BP, WN, NN, NC, MC, SC, WD, CO, DN, TH, AU.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2001); Palma (2010); Sychra *et al.* (2016).

Other significant references: None.

Remarks: Together with its hosts, *Myrsidea novaeseelandiae* is endemic to New Zealand. Notwithstanding the large number of species placed in the bird family Meliphagidae (Dickinson 2003: 431), and some records of “*Myrsidea* sp.” from Australian meliphagids in Murray *et al.* (2001: 1262), *Myrsidea novaeseelandiae* is so far the only described and named species of *Myrsidea* from that host family.

#### ***Myrsidea serini* (Séguy, 1944)**

*Menopon serini* Séguy 1944: 80, fig. 84a,b,c.

*Myrsidea serini* (Séguy, 1944); Hopkins & Clay 1952: 233.

*Myrsidea* sp.; Pilgrim & Palma 1982: 27.

*Myrsidea serini* (Séguy, 1944); Klockenhoff 1984: 18, figs 1–4.

*Myrsidea serini* (Séguy, 1944); Palma 1999: 382.

*Myrsidea serini* (Séguy, 1944); Murray *et al.* 2006b: 1959.

*Myrsidea serini* (Séguy, 1944); Palma 2010: 407.

Lectotype ♂ in MNHN (Klockenhoff 1984: 20).

Type host: *Serinus serinus* (Linnaeus, 1766).

New Zealand hosts: *Carduelis chloris* (Linnaeus, 1758); *Carduelis carduelis britannica* (Hartert, 1903); *Emberiza citrinella* Linnaeus, 1758; *Serinus canaria* (Linnaeus, 1758) captive.

Other hosts: *Carduelis barbata* (Molina, 1782); *Chrysomus thilius petersii* (Laubmann, 1934); *Agelaioides badius badius* (Vieillot, 1819).

New Zealand localities: WN, MC, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Klockenhoff (1984); Palma (1999); Murray *et al.* (2006b); Price & Dalgleish (2007: 13, fig. 39); Palma (2010); Cicchino & Valim (2015: 240, figs 1–33); Sychra *et al.* (2016: 397, 409).

Other significant references: Price *et al.* (2003: 131).

Remarks: *Myrsidea serini* was introduced to New Zealand with its hosts by human agency (Checklist Committee 2010: 320, 322). In New Zealand, *Serinus canaria* is only a cage bird, and releases to establish it in the wild were unsuccessful (Checklist Committee 2010: 346).

### ***Myrsidea thoracica* (Giebel, 1874)**

Figs 49–50

*Menopon thoracicum* Giebel, 1874: 287.

*Myrsidea thoracica* (Giebel, 1874); Hopkins & Clay 1952: 234.

*Myrsidea thoracica* (Giebel, 1874); Clay 1966c: 342, figs 1, 5–8, 25–27, 64, pl. 1: figs 1, 5, 6.

*Myrsidea thoracica* (Giebel, 1874); Pilgrim & Palma 1982: 26.

*Myrsidea thoracica* (Giebel, 1874); Price *et al.* 2003: 132.

*Myrsidea thoracica* (Giebel, 1874); Murray *et al.* 2006b: 1959.

*Myrsidea thoracica* (Giebel, 1874); Palma 2010: 408.

Syntypes ♂♀, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Turdus viscivorus* Linnaeus, 1758.

New Zealand host: *Turdus merula merula* Linnaeus, 1758.

Other hosts: *Zoothera sibirica* (Pallas, 1776); *Turdus boulboul* (Latham, 1790); *Turdus obscurus* J.F. Gmelin, 1789; *Turdus pallidus* J.F. Gmelin, 1789; *Turdus chrysolaus* Temminck, 1832; *Turdus ruficollis* Pallas, 1776.

New Zealand localities: AK, WN, CH.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Murray *et al.* (2006b); Palma (2010); Sychra *et al.* (2016: 397, 409).

Other significant references: Clay (1966c); Baum (1968: 143, fig. 4); Martín-Mateo (2002: 120); Price *et al.* (2003).

Remarks: *Myrsidea thoracica* was introduced to New Zealand with Eurasian blackbirds by human agency (Checklist Committee 2010: 313).

### ***Myrsidea* species**

*Myrsidea* sp.; Pilgrim & Palma 1982: 26.

*Myrsidea* sp.; Palma 1999: 381.

*Myrsidea* sp.; Murray *et al.* 2002: 1215.

*Myrsidea* sp.; Murray *et al.* 2006b: 1959.

*Myrsidea* sp.; Palma 2010: 408.

New Zealand hosts: *Gerygone igata* (Quoy & Gaymard, 1830); *Mohoua albicilla* (Lesson, 1830); *Anthus novaeseelandiae novaeseelandiae* (J.F. Gmelin, 1789).

New Zealand localities: BP, GB, WN.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Palma (1999); Murray *et al.* (2002); Palma (2010); Sychra *et al.* (2016: 409).

Other significant references: None.

Remarks: The available samples of *Myrsidea* from the three hosts listed above cannot be identified to species because they contain nymphs only (voucher specimens in MONZ).

### Genus *Nosopon* Hopkins, 1950

*Nosopon* Hopkins, 1950. *Ann. Mag. Nat. Hist.* (Ser. 12) 3: 239. Type species: *Menopon* “*fulvofasciatum* var.” *minor* Piaget, 1880 = *Nosopon lucidum* (Rudow, 1869) (by original designation).

### *Nosopon lucidum* (Rudow, 1869)

Figs 51–52

*Menopon lucidum* Rudow, 1869a: 34, 402.

*Menopon fulvofasciatum* var. *minor* Piaget, 1880: 418.

*Nosopon lucidum* (Rudow, 1869); Hopkins & Clay 1952: 249.

*Nosopon minus* (Piaget, 1880); Hopkins & Clay 1952: 249.

*Nosopon lucidum lucidum* (Rudow, 1869); Tendeiro 1959: 194, figs 1–5.

*Nosopon lucidum* (Rudow, 1869); Pilgrim & Palma 1982: 17.

*Nosopon* sp.; Pilgrim & Palma 1982: 17.

*Nosopon lucidum* (Rudow, 1869); Murray *et al.* 1993: 960.

*Nosopon* sp.; Murray *et al.* 1993: 960.

*Nosopon lucidum* (Rudow, 1869); Palma 1999: 379.

*Nosopon lucidum* (Rudow, 1869); Palma 2010: 408.

Lectotype ♂ in ZMHG (Clay & Hopkins 1955: 51; Weidner 1966: 259).

Type host: *Falco vespertinus* Linnaeus, 1766.

New Zealand hosts: *Circus approximans* Peale, 1848; *Falco novaeseelandiae* J.F. Gmelin, 1788.

Other hosts: *Accipiter gentilis* (Linnaeus, 1758); *Accipiter nisus* (Linnaeus, 1758); *Elanus caeruleus* (Desfontaines, 1789); *Lophætus occipitalis* (Daudin, 1800), three other species of *Circus* and five other species of *Falco* (see Price *et al.* 2003: 133).

New Zealand localities: TK, WN, SD, MB, NC, MC, SC, WD.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (1999); Paterson *et al.* (1999: 221); Palma (2010).

Other significant references: Tendeiro (1959); Green & Palma (1991: 8, 30); Palma (1996b: 140); Martín-Mateo (2002: 100, fig. 33A); Price *et al.* (2003: 133); Palma & Jensen (2005: 53, 62).

Remarks: *Nosopon lucidum* is widespread on many host species, but not frequently collected.

### Genus *Plegadiphilus* Bedford, 1939

*Plegadiphilus* Bedford, 1939. *Onderstepoort Jour. Vet. Sci. Animal Ind.* 12(1): 138. Type species: *Plegadiphilus threskiornis* Bedford, 1939 (by original designation).

### *Plegadiphilus plegadis* (Dubinin, 1938)

Figs 53–54

*Menopon plegadis* Dubinin, 1938: 178, fig. 12.

*Plegadiphilus plegadis* (Dubinin, 1938); Hopkins & Clay 1952: 295.

*Plegadiphilus plegadis* (Dubinin, 1938); Pilgrim & Palma 1982: 16.

*Plegadiphilus plegadis* (Dubinin, 1938); Murray *et al.* 1990: 1373.

*Plegadiphilus plegadis* (Dubinin, 1938); Price *et al.* 2003: 135.

*Plegadiphilus plegadis* (Dubinin, 1938); Palma 2010: 408.

Syntypes ♂♀, repository unknown.

Type host: *Plegadis falcinellus* (Linnaeus, 1766).

New Zealand host: *Plegadis falcinellus* (Linnaeus, 1766).

Other hosts: *Plegadis chihi* (Vieillot, 1817).

New Zealand localities: WI, SL.

Geographic distribution: Eurasia; Africa; Americas; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Forrester *et al.* (1995: 12); Palma (1996b: 140); Price *et al.* (2003).

Remarks: Although the glossy ibis is a regular vagrant to New Zealand (Checklist Committee 2010: 166), there are only two records of *Plegadiphilus plegadis* from this country.

### ***Plegadiphilus threskiornis* Bedford, 1939**

*Plegadiphilus threskiornis* Bedford, 1939: 139, figs 7–8.

*Plegadiphilus threskiornis* Bedford, 1939; Hopkins & Clay 1952: 295.

*Plegadiphilus threskiornis* Bedford, 1939; Pilgrim & Palma 1982: 16.

*Plegadiphilus threskiornis* Bedford, 1939; Murray *et al.* 1990: 1373.

*Plegadiphilus threskiornis* Bedford, 1939; Price *et al.* 2003: 135.

*Plegadiphilus threskiornis* Bedford, 1939; Palma 2010: 408.

Holotype ♀, probably in SAIM.

Type host: *Threskiornis aethiopicus* (Latham, 1790).

New Zealand host: *Threskiornis molucca strictipennis* (Gould, 1838).

Other hosts: *Threskiornis melanocephalus* (Latham, 1790); *Threskiornis molucca pygmaeus* Mayr, 1931; *Threskiornis spinicollis* (Jameson, 1835); *Bostrychia hagedash* (Latham, 1790).

New Zealand locality: SL.

Geographic distribution: Asia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Clay (1958c: 144); Moreby (1976: 92); Palma (1996b: 141); Price *et al.* (2003).

Remarks: The Australian white ibis is an occasional visitor to New Zealand (Checklist Committee 2010: 166), with only a single record of *Plegadiphilus threskiornis* from this country.

### **Genus *Pseudomenopon* Mjöberg, 1910**

*Pseudomenopon* Mjöberg, 1910a. *Arkiv Zool.* 6(13): 50. Type species *Menopon tridens* “N.” = *Pseudomenopon pilosum* (Scopoli, 1763) (by original designation).

### ***Pseudomenopon concretum* (Piaget, 1880)**

*Menopon concretum* Piaget, 1880: 481, pl. 38: fig. 9.

*Pseudomenopon concretum* (Piaget, 1880); Hopkins & Clay 1952: 302.

*Pseudomenopon concretum* (Piaget, 1880); Price 1974: 78, fig. 32.

*Pseudomenopon concretum* (Piaget, 1880); Pilgrim & Palma 1982: 19.

*Pseudomenopon concretum* (Piaget, 1880); Murray *et al.* 1993: 961.

*Pseudomenopon concretum* (Piaget, 1880); Price *et al.* 2003: 135.

*Pseudomenopon concretum* (Piaget, 1880); Palma 2010: 408.

Lectotype ♀ in NHML (Clay 1949b: 824).

Type host: *Porphyrio melanotus melanopterus* Bonaparte, 1856.

New Zealand host: *Porphyrio melanotus melanotus* Temminck, 1820.

Other hosts: *Porphyrio porphyrio* (Linnaeus, 1758); *Porphyrio poliocephalus* (Latham, 1802); *Porphyrio madagascariensis* (Latham, 1802); *Megacrex inepta* D'Albertis & Salvadori, 1879.

New Zealand localities: AK, BP, SD, MB, NC, MC, SC, WD.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Price (1974); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 221, 223); Palma (2010).

Other significant references: Clay (1949b); Tendeiro (1965d: 29, photos 10–11, 26, 36, 49, 56); Lakshminarayana (1977: 54); Green & Palma (1991: 8, 32); Palma (1996b: 141); Price *et al.* (2003: 135).

Remarks: *Pseudomenopon concretum* has been frequently collected from pukekos in New Zealand.

### ***Pseudomenopon pilgrimi* Price, 1974**

Figs 55–56

*Pseudomenopon pilgrimi* Price, 1974: 81, fig. 29.

*Pseudomenopon pilgrimi* Price, 1974; Pilgrim & Palma 1982: 18–19.

*Pseudomenopon pilgrimi* Price, 1974; Murray *et al.* 1993: 961.

*Pseudomenopon pilgrimi* Price, 1974; Price *et al.* 2003: 135.

*Pseudomenopon pilgrimi* Price, 1974; Palma 2010: 408.

Holotype ♂ in CMNZ (Nicholls *et al.* 1998: 30).

Type host: *Gallirallus australis australis* (Sparrman, 1786).

New Zealand hosts: *Gallirallus australis greyi* (Buller, 1888); *Gallirallus australis hectori* (Hutton, 1873).

Other hosts: None.

New Zealand localities: HB, SD, MB, NN, WD, CH.

Geographic distribution: New Zealand.

New Zealand references: Price (1974); Pilgrim & Palma (1982); Murray *et al.* (1993); Nicholls *et al.* (1998: 30); Paterson *et al.* (1999: 223); Palma (2010).

Other significant references: Lakshminarayana (1977: 54); Price *et al.* (2003: 131).

Remarks: *Pseudomenopon pilgrimi* is an endemic species, exclusively parasitic on wekas but not frequently collected.

### ***Pseudomenopon pilosum* (Scopoli, 1763)**

*Pediculus pilosus* Scopoli, 1763: 384.

*Menopon tridens* Burmeister, 1838a: 440.

*Pseudomenopon tridens* (Nitzsch) [sic]; Ferris 1924: 63, fig. 4.

*Pseudomenopon pilosum* (Scopoli, 1763); Hopkins & Clay 1952: 302.

*Pseudomenopon pilosum* (Scopoli, 1763); Price 1974: 73, figs 1–6, 11, 18, 21, 22.

*Pseudomenopon pilosum* (Scopoli, 1763); Pilgrim & Palma 1982: 19.

*Pseudomenopon pilosum* (Scopoli, 1763); Murray *et al.* 1993: 961.

*Pseudomenopon pilosum* (Scopoli, 1763); Price *et al.* 2003: 135.

*Pseudomenopon pilosum* (Scopoli, 1763); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1951: 19).

Type host: *Fulica atra* Linnaeus, 1758.

New Zealand host: *Fulica atra australis* Gould, 1845.

Other hosts: *Hydrophasianus chirurgus* (Scopoli, 1786); *Podica senegalensis* (Vieillot, 1817); five species of *Gallinula*; and six species of *Fulica* (see Price *et al.* 2003: 136).

New Zealand localities: NC, MC, SC, CO, DN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Ferris (1924); Clay & Hopkins (1951: 19); Tendeiro (1965d: 19, photos 2–3, 22, 32, 43, 52); Price (1974); Lakshminarayana (1977: 56); Benoit (1976: 235); Green & Palma (1991: 8, 32); Butler & O'Connor (1994: 451); Forrester *et al.* (1995: 23); Palma (1996b: 142); Martín-Mateo (2002: 42, figs 9–10); Price *et al.* (2003); Palma & Jensen (2005: 53, 63); Adam (2007: 166, fig. 9).

Remarks: *Pseudomenopon pilosum* is widespread on many host species and frequently collected.

### ***Pseudomenopon scopulacorne* (Denny, 1842)**

*Liotheum (Menopon) scopulacorne* Denny, 1842: 200, 221, pl. 18: fig. 9.

*Pseudomenopon scopulacorne* Denny, 1842 [sic]; Harrison 1916: 63 (as junior synonym of *Pseudomenopon tridens* Nitzsch [in Burmeister]).

*Pseudomenopon scopulacorne* (Denny, 1842); Hopkins & Clay 1952: 303.

*Pseudomenopon scopulacorne* (Denny, 1842); Price 1974: 75, figs 10, 12, 20, 25.

*Pseudomenopon scopulacorne* (Denny, 1842); Pilgrim & Palma 1982: 18.

*Pseudomenopon scopulacorne* (Denny, 1842); Murray *et al.* 1993: 961.

*Pseudomenopon scopulacorne* (Denny, 1842); Price *et al.* 2003: 136.

*Pseudomenopon scopulacorne* (Denny, 1842); Palma 2010: 408.

Syntypes ♀♀ in NHML (Thompson 1937a: 76; Palma 1996b: 142).

Type host: *Rallus aquaticus aquaticus* Linnaeus, 1758.

New Zealand host: *Gallirallus philippensis assimilis* (G.R. Gray, 1843).

Other hosts: *Atlantisia rogersi* P.R. Lowe, 1923; *Rallina eurizonoides* (Lafresnaye, 1845); four species of *Laterallus*; three species of *Porzana*; and five species of *Rallus* (see Price *et al.* 2003: 136).

New Zealand locality: BP.

Geographic distribution: Eurasia; Americas; Australasia; Oceania.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Thompson (1937a: 76); Price (1974); Lakshminarayana (1977: 56); Forrester *et al.* (1995: 23); Green & Palma (1991: 8, 31); Palma (1996b: 142); Price *et al.* (2003); Palma & Jensen (2005: 53, 63); Palma & Peck (2013: 25).

Remarks: Although *Pseudomenopon scopulacorne* is widespread on many host species, it has only been found twice on banded rails in New Zealand.

### Genus *Trinoton* Nitzsch, 1818

*Trinoton* Nitzsch, 1818. *German's Mag. Entomol.* 3: 300. Type species: *Liotheum (Trinoton) conspurcatum* Nitzsch, 1818 = *Trinoton anserinum* (J.C. Fabricius, 1805) (by monotypy).

*Trinotum* Burmeister, 1838a. *Handb. Entomol.* 2(1): 440. Invalid emendation.

### *Trinoton nigrum* Le Souëf, 1902

Figs 57–58

*Trinoton* [sic] *niger* [sic] Le Souëf, 1902b: 90.

*Trinoton nigrum* Le Souëf, 1902; Harrison 1916: 62. Emendation.

*Trinoton nigrum* Le Souëf, 1902; Hopkins & Clay 1952: 358.

*Trinoton nigrum* Le Souëf, 1902; Pilgrim & Palma 1982: 16.

*Trinoton nigrum* Le Souëf, 1902; Murray *et al.* 1990: 1374.

*Trinoton nigrum* Le Souëf, 1902; Palma 2010: 408.

Holotype ♀ in SAMA (Palma 1996b: 143).

Type host: *Cygnus atratus* (Latham, 1790).

New Zealand host: *Cygnus atratus* (Latham, 1790).

Other hosts: None.

New Zealand localities: AK, WO, BP, WA, WN, NC, MC, SC, CH.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Eichler & Vasjukova (1981: 35); Green & Palma (1991: 8, 29); Palma (1996b: 143); Price *et al.* (2003: 139).

Remarks: The population of black swans living in New Zealand is probably the result of a mixture of self-introduced birds and others introduced by human agency (Checklist Committee 2010: 33). Therefore, it is not possible to determine if *Trinoton nigrum* is a native or an introduced species.

### *Trinoton querquedulae* (Linnaeus, 1758)

*Pediculus querquedulae* Linnaeus, 1758: 612.

*Trinotum lituratum* Burmeister, 1838a: 441.

*Trinotum luridum* Burmeister, 1838a: 441.

*Trinoton querquedulae* (Linnaeus, 1758); Hopkins & Clay 1952: 358.

*Trinoton querquedulae* (Linnaeus, 1758); Watt 1971: 233, 243.

*Trinoton querquedulae* (Linnaeus, 1758); Wise 1977: 57.

*Trinoton querquedulae querquedulae* (Linnaeus, 1758); Eichler & Vasjukova 1981: 37, figs 12, 22, 25, 33, 40, 41, pl. 2: fig. 1, pl. 4: fig. 2, pl. 7: fig. 2, pl. 8: fig. 2.

*Trinoton querquedulae* (Linnaeus, 1758) s. l.; Pilgrim & Palma 1982: 16–17.

*Trinoton querquedulae* (Linnaeus, 1758); Murray *et al.* 1990: 1374.

*Trinoton querquedulae* (Linnaeus, 1758); Palma 2010: 408.

Neotype ♀ in NHML (Clay & Hopkins 1950: 244, pl. 2: fig. 2).

Type host: *Anas crecca crecca* Linnaeus, 1758.

New Zealand hosts: *Branta canadensis maxima* Delacour, 1951; *Tadorna variegata* (J.F. Gmelin, 1789); *Anas gracilis* Buller, 1869; *Anas chlorotis* G.R. Gray, 1845; *Anas platyrhynchos platyrhynchos* Linnaeus, 1758; *Anas superciliosa* J.F. Gmelin, 1789; *Anas rhynchotis* Latham, 1802; *Aythya novaeseelandiae* (J.F. Gmelin, 1789).

Other hosts: Twenty-two other species of *Anas*, nine other species of *Aythya*, two species of *Aix*, three species of *Bucephala*, three species of *Melanitta*, four species of *Mergus*, two species of *Netta*, three species of *Oxyura*, three species of *Somateria*, two species of *Tadorna* and over 10 species of other genera (see Price *et al.* 2003: 139).

New Zealand localities: ND, AK, BP, WO, BP, HB, TK, WA, WN, NN, NC, MC, SC, CO, DN, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 219); Galloway (2005: 16); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Clay & Hopkins (1950: 243, figs 26–28, pl. 2: fig. 2); Kéler (1957c: fig. 34a); Eichler & Vasjukova (1981); Rudolph (1983: 16); Price (1987: 218, fig. 22.14); Green & Palma (1991: 8, 29); Butler & O'Connor (1994: 451); Forrester *et al.* (1995: 15); Palma (1996b: 143); Martín-Mateo (2002: 39, figs 7f,g,h,i, 8); Price *et al.* (2003: 139); Palma & Jensen (2005: 53, 62); Adam (2007: 167); Palma & Peck (2013: 25).

Remarks: *Trinoton querquedulae* is an extremely widespread species with some morphological variability. Pilgrim & Palma (1982: 16–17) regarded the populations of *T. querquedulae* from seven host species somewhat different from that of the type host, and qualified them as *sensu lato*; however, my examination of more samples, including one from the type host, shows that making such difference is not warranted. *Anas chlorotis* is a new host record for *Trinoton querquedulae* in New Zealand (voucher specimens in MONZ).

## Family RICINIDAE Neumann, 1890

Ricinidae Neumann, 1890. *Bull. Soc. d'Hist. Nat., Toulouse* 24: 55. Type genus: *Ricinus* De Geer, 1778b.

### Genus *Ricinus* De Geer, 1778

*Ricinus* De Geer, 1778b. *Mém. Hist. Ins.* 7: 69. Type species: *Ricinus fringillae* De Geer, 1778 (by subsequent designation).

*Physostomum* Nitzsch, 1818. *Germar's Mag. Entomol.* 3: 302. Type species: *Ricinus nitidissimus* Nitzsch, 1818 = *Ricinus fringillae* De Geer, 1778 (by subsequent designation).

### *Ricinus* species

Fig. 59

*Ricinus* sp.; Gill 1980: 246

*Ricinus* sp.; Pilgrim & Palma 1982: 26.

*Ricinus* sp.; Murray *et al.* 2002: 1215.

New Zealand host: *Gerygone igata* (Quoy & Gaymard, 1830).

New Zealand localities: KA, SI.

Geographic distribution: New Zealand.

New Zealand references: Gill (1980); Pilgrim & Palma (1982); Murray *et al.* (2002).

Other significant references: Rheinwald (1968); Nelson (1972); Price *et al.* (2003: 246).

Remarks: The only two available records of *Ricinus* from New Zealand could not be identified to species because the samples contain females only (voucher specimens in MONZ).

**Suborder ISCHNOCERA Kellogg, 1896**

Ischnocera Kellogg, 1896a. *Proc. Calif. Acad. Sci.* 6: 63.

**Family PHILOPTERIDAE Burmeister, 1838**

Philopteridae Burmeister, 1838a. *Handb. Entomol.* 2(1): 422.

**Genus *Acidoproctus* Piaget, 1878**

*Acidoproctus* Piaget, 1878. *Tijdschrift Entomologie* 21: 178. Type species: *Acidoproctus marginatus* Piaget, 1878 = *Acidoproctus rostratus* (Rudow, 1866b) (by subsequent designation).

***Acidoproctus emersoni* Timmermann, 1962****New Record**

*Acidoproctus emersoni* Timmermann, 1962b: 145, figs 11, 14.

*Acidoproctus emersoni* Timmermann, 1962; Price *et al.* 2003: 140.

*Acidoproctus emersoni* Timmermann, 1962; Arnold 2006: 278, 281, figs 13, 21.

Holotype ♂ in NHML (Palma 1996b: 151).

Type host: *Dendrocygna javanica* (Horsfield, 1821).

New Zealand host: *Dendrocygna eytoni* (Eyton, 1838).

Other hosts: *Dendrocygna arcuata* (Horsfield, 1824); *Dendrocygna guttata* Schlegel, 1866.

New Zealand locality: ND.

Geographic distribution: Asia; Australasia.

New Zealand reference: This paper.

Other significant references: Palma (1996b: 151); Price *et al.* (2003); Arnold (2006).

Material examined and repository: 7♂, 5♀ (1 sample, MONZ).

Remarks: *Acidoproctus emersoni* is a new louse species for New Zealand. *Dendrocygna eytoni* is a straggler to New Zealand, recorded occasionally since 1871 (Checklist Committee 2010: 31).

***Acidoproctus gottwaldhirschi* Eichler, 1958**

Figs 60–61

*Acidoproctus gottwaldhirschi* Eichler, 1958: 60, figs 1–2.

*Acidoproctus gottwaldhirschi* Eichler, 1958; Pilgrim & Palma 1982: 17.

*Acidoproctus gottwaldhirschi* Eichler, 1958; Murray *et al.* 1990: 1374.

*Acidoproctus gottwaldhirschi* Eichler, 1958; Arnold 2006: 280, figs 4, 17, 25.

*Acidoproctus gottwaldhirschi* Eichler, 1958; Palma 2010: 408.

Holotype ♂, presumed lost. Eichler collected it from a mounted bird in the Zoologischen Museum der Universität Göttingen, but the holotype of *A. gottwaldhirschi* has not be located in that museum (Gert Tröster pers. comm. October 2013).

Type host: *Hymenolaimus malacorhynchos* (J.F. Gmelin, 1789).

New Zealand host: *Hymenolaimus malacorhynchos* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: BP, GB, HB, WN, NC.

Geographic distribution: New Zealand.

New Zealand references: Eichler (1958); Pilgrim & Palma (1982); Murray *et al.* (1990); Arnold (2006); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003: 140).

Remarks: *Acidoproctus gottwaldhirschi* is an endemic and vulnerable species (Buckley *et al.* 2012), exclusively parasitic on blue ducks.

**Genus *Alcedoecus* Clay & Meinertzhagen, 1939**

*Alcedoecus* Clay & Meinertzhagen, 1939a. *Entomologist* 72: 162. Type species: *Philopterus capistratus* Neumann, 1912b = *Alcedoecus capistratus* (Neumann, 1912) (by original designation).



*Halcyonicola* Uchida 1948. *Japan. Med. Jour.* 1: 312. Type species: *Docophorus alatoclypeatus* Piaget, 1885 = *Alcedoecus alatoclypeatus* (Piaget, 1885) (by original designation).

***Alcedoecus alatoclypeatus* (Piaget, 1885)**

Figs 62–63

- Docophorus alatoclypeatus* Piaget, 1885: 10, pl. 1: fig. 11.  
*Philopterus alatoclypeatus* Piaget, 1885 [sic]; Harrison 1916: 87.  
*Philopterus alatoclypeatus* (Piaget, 1885); Séguy 1944: 235, fig. 347.  
*Halcyonicola alatoclypeatus* (Piaget, 1885); Uchida 1948: 312.  
*Alcedoecus alatoclypeatus* (Piaget, 1885); Hopkins & Clay 1952: 25.  
*Alcedoecus alatoclypeatus* (Piaget, 1885); Tendeiro 1965c: 44, photos 23–25.  
*Alcedoecus* sp.; Nelson 1969: 199.  
*Alcedoecus* sp.; Watt 1971: 235, 244.  
*Alcedoecus alatoclypeatus* (Piaget, 1885) *s. l.*; Pilgrim & Palma 1982: 25.  
*Alcedoecus alatoclypeatus* (Piaget, 1885); Murray *et al.* 1999: 1242.  
*Alcedoecus alatoclypeatus* (Piaget, 1885); Palma 2010: 408.

Lectotype ♂ in NHML (Palma 1996b: 152).

Type host: *Halcyon malimbica* (Shaw, 1811).

New Zealand hosts: *Todiramphus sanctus vagans* (Lesson, 1828); *Todiramphus sanctus norfolkiensis* (Tristram, 1885).

Other hosts: *Todiramphus sanctus sanctus* Vigors & Horsfield, 1827.

New Zealand localities: BP, HB, TK, WI, WA, WN, NN, NC, MC, SC, WD, KE, Norfolk Island.

Geographic distribution: Australasia.

New Zealand references: Nelson (1969); Watt (1971); Pilgrim & Palma (1982); Murray *et al.* (1999); Paterson *et al.* (1999: 221); Palma (2010).

Other significant references: Séguy (1944); Uchida (1948); Tendeiro (1965c); Palma (1996b: 152); Price *et al.* (2003: 141).

Remarks: Pilgrim & Palma (1982: 25) regarded the population of *Alcedoecus alatoclypeatus* from *Todiramphus sanctus vagans* (as *Halcyon sancta vagans*) as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of samples from other hosts shows that making such difference is not warranted.

***Alcedoecus delphax* (Nitzsch [*in* Giebel], 1866)**

**New Record**

- Docophorus delphax* Nitzsch [*in* Giebel], 1866: 360  
*Philopterus delphax* Nitzsch [*in* Giebel], 1866 [sic]; Harrison 1916: 92.  
*Halcyonicola delphax* (Nitzsch) [sic]; Uchida 1948: 312.  
*Alcedoecus delphax* (Nitzsch [*in* Giebel], 1866); Hopkins & Clay 1952: 26.  
*Alcedoecus delphax* (Nitzsch [*in* Giebel], 1866); Palma 1996b: 152.  
*Alcedoecus delphax* (Nitzsch [*in* Giebel], 1866); Price *et al.* 2003: 141.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1), Palma & Pilgrim (1984: 150) and Palma (1996b: 152).

Type host: *Dacelo novaeguineae* (Hermann, 1783).

New Zealand host: *Dacelo novaeguineae novaeguineae* (Hermann, 1783).

Other hosts: None.

New Zealand locality: ND.

Geographic distribution: Australasia.

New Zealand reference: This paper.

Other significant references: Uchida (1948); Green & Palma (1991: 10, 36); Palma (1996b); Price *et al.* (2003).

Material examined and repository: 4♂, 4♀ (1 sample, MONZ).

Remarks: *Alcedoecus delphax* is a new louse species for New Zealand, introduced to this country from Australia with laughing kookaburras by human agency (Checklist Committee 2010: 272).

**Genus *Anaticola* Clay, 1936**

*Anaticola* Clay, 1936. *Proc. Zool. Soc. London* [no volume number]: 617. Type species: *Esthiopterum crassicornis* (Scopoli, 1763) = *Anaticola crassicornis* (Scopoli, 1763) (by original designation).

***Anaticola anseris* (Linnaeus, 1758)**

“Pollini del oca reale” Redi, 1668: pl. 11.

*Pediculus anseris* Linnaeus, 1758: 612. *Nomen novum* for “Pollini del oca reale” Redi, 1668.

*Ricinus anseris* (Linnaeus, 1758); Latreille 1804: 106.

*Esthiopterum anseris* Linnaeus, 1758 [sic]; Harrison 1916: 130.

*Anaticola anseris* (Linnaeus, 1758); Clay & Hopkins 1950: 239, figs 18–21, pl. 2: fig. 1.

*Anaticola anseris* (Linnaeus, 1758); Hopkins & Clay 1952: 31.

*Anaticola anseris anseris* (Linnaeus, 1758); Eichler & Vasjukova 1980: 341, pl. 17: figs 5–8.

*Anaticola anseris* (Linnaeus, 1758); Palma 1999: 379, 383, note 4.

*Anaticola anseris* (Linnaeus, 1758); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1950: 239).

Type host: *Anser anser* (Linnaeus, 1758).

New Zealand hosts: *Anser anser* (Linnaeus, 1758); *Branta canadensis maxima* Delacour, 1951.

Other hosts: Six other species of *Anser*; *Branta leucopsis* (Bechstein, 1803); *Branta sandvicensis* (Vigors, 1833).

New Zealand localities: SD, MB, MC.

Geographic distribution: Eurasia; North America; Australasia.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Wundrig (1936: 84, figs 49–51); Clay & Hopkins (1950); Hackman & Nyholm (1968: 80); Butler & O’Connor (1994: 452); Palma (1996b: 152); Price *et al.* (2003: 142); Palma & Jensen (2005: 53, 61); Martín-Mateo (2009: 227, fig. 46); Escalante *et al.* (2016: 203).

Remarks: *Anaticola anseris* was introduced to New Zealand with greylag geese and Canada geese by human agency (Checklist Committee 2010: 35).

***Anaticola crassicornis* (Scopoli, 1763)**

*Pediculus crassicornis* Scopoli, 1763: 383.

*Esthiopterum crassicornis* Scopoli, 1763 [sic]; Harrison 1916: 132.

*Anaticola crassicornis* [sic] (Scopoli, 1763); Séguy 1944: 361, figs 536–538.

*Anaticola crassicornis* (Scopoli, 1763); Hopkins & Clay 1952: 32.

*Anaticola crassicornis* (Scopoli, 1763); Watt 1971: 235, 243.

*Anaticola crassicornis* (Scopoli, 1763) *s. l.*; Horning *et al.* 1980: 4, 11.

*Anaticola crassicornis crassicornis* (Scopoli, 1763); Eichler & Vasjukova 1980: 345, figs 2b, 37–42, pl. 21: figs 43–46, pl. 22: figs 57–58.

*Anaticola crassicornis* (Scopoli, 1763); Pilgrim & Palma 1982: 16.

*Anaticola crassicornis* (Scopoli, 1763) *s. l.*; Pilgrim & Palma 1982: 17.

*Anaticola crassicornis* (Scopoli, 1763); Murray *et al.* 1990: 1374

*Anaticola crassicornis* (Scopoli, 1763); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1951: 19).

Type host: *Anas platyrhynchos platyrhynchos* Linnaeus, 1758.

New Zealand hosts: *Anas gracilis* Buller, 1869; *Anas platyrhynchos platyrhynchos* Linnaeus, 1758; *Anas superciliosa* J.F. Gmelin, 1789; *Anas rhynchotis* Latham, 1802; *Aythya novaeseelandiae* (J.F. Gmelin, 1789).

Other hosts: Twenty other species of *Anas* (see Price *et al.* 2003: 143); *Cygnus cygnus* (Linnaeus, 1758); *Oxyura jamaicensis* (J.F. Gmelin, 1789); *Somateria spectabilis* (Linnaeus, 1758).

New Zealand localities: ND, WO, BP, HB, TK, WI, WA, WN, SD, MB, NN, NC, MC, SC, CO, DN, KE, SN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Watt (1971); Wise (1977: 58); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Clay & Hopkins (1951: 17, figs 23–24); Eichler & Vasjukova (1980); Rudolph (1983: 16); Mey (1986: 37, fig. 6); Green & Palma (1991: 10, 29); Butler & O’Connor (1994: 452); Forrester *et al.* (1995: 14);

Palma (1996b: 153); Price *et al.* (2003: 143); Palma & Jensen (2005: 54, 61); Martín-Mateo (2009: 225, figs 44–45); Palma & Peck (2013: 27); Ahmad *et al.* (2015: 568); Escalante *et al.* (2016: 203).

Remarks: *Anaticola crassicornis* is a very widespread species with some morphological variability. Pilgrim & Palma (1982: 17) regarded the populations of *A. crassicornis* from four host species as somewhat different from that of the type host, and qualified them as *sensu lato*; however, my examination of more samples from those and other hosts shows that making such difference is not warranted. Price *et al.* (2003: 143, 279) incorrectly identified the population from *Aythya novaeseelandiae* as “*Anaticola mergiserrati* (De Geer, 1778)” (see Escalante *et al.* 2016: 205, fig. 1).

### ***Anaticola magnificus* Ansari, 1955**

Figs 64–65

*Anaticola magnificus* Ansari, 1955b: 51.

*Anaticola magnificus* Ansari, 1955; Pilgrim & Palma 1982: 16.

*Anaticola magnificus* Ansari, 1955; Murray *et al.* 1990: 1374.

*Anaticola magnificus* Ansari, 1955; Palma 1999: 379.

*Anaticola magnificus* Ansari, 1955; Palma 2010: 408.

Holotype ♂, repository unknown (see Remarks).

Type host: *Tadorna ferruginea* (Pallas, 1764).

New Zealand hosts: *Tadorna variegata* (J.F. Gmelin, 1789); *Tadorna tadornoides* (Jardine & Selby, 1828).

Other hosts: None.

New Zealand localities: HB, WI, WA, SD, MB, NN, NC, MC, SC, WD, FD.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999); Palma (2010).

Other significant references: Price *et al.* (2003: 143); Escalante *et al.* (2016: fig. 1).

Remarks: The holotype ♂ of *Anaticola magnificus* is not in the collection of the NHML (Vincent S. Smith pers. comm. December 2014).

### ***Anaticola* species**

*Anaticola* sp.; Pilgrim & Palma 1982: 17.

*Anaticola* sp.; Murray *et al.* 1990: 1374.

*Anaticola* sp.; Palma 1999: 379.

New Zealand hosts: *Chenonetta jubata* (Latham, 1802); *Anas chlorotis* G.R. Gray, 1845.

Other hosts: None.

New Zealand localities: ND, CL, MB.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999); Buckley *et al.* (2012: App. 2).

Other significant references: None.

Remarks: These records possibly represent undescribed, unnamed species, especially those from *Chenonetta jubata* (see Escalante *et al.* 2016: 203, fig. 1), but available samples are not adequate for proper descriptions of the taxa involved. The population from *Anas chlorotis* is regarded “at risk” (Buckley *et al.* 2012).

### **Genus *Anatoecus* Cummings, 1916**

*Anatoecus* Cummings, 1916b. *Proc. Zool. Soc. London 1916*: 653. Type species: *Anatoecus icterodes* (Nitzsch, 1818) (by original designation).

*Benatoecus* Zlotorzyczka, 1970. *Polskie Pismo Entomol.* 40(1): 10. Type species: *Anatoecus dentatus* (Scopoli, 1763) (by original designation).

### ***Anatoecus dentatus* (Scopoli, 1763) *sensu lato***

*Pediculus dentatus* Scopoli, 1763: 383.

*Philopterus dentatus* Scopoli, 1763 [sic]; Harrison 1916: 93.

*Anatoecus dentatus* (Scopoli, 1763); Hopkins & Clay 1952: 35.

*Anatoecus dentatus dentatus* (Scopoli, 1763); Kéler 1960a: 305, figs 6, 10(A), 15, 22, 34.

*Anatoecus (Benatoecus) dentatus dentatus* (Scopoli, 1763); Złotorzycka 1970: 13, figs 3–4.

*Anatoecus dentatus* (Scopoli, 1763); Watt 1971: 235, 243.

*Anatoecus dentatus* (Scopoli, 1763); Wise 1977: 58.

*Anatoecus dentatus* (Scopoli, 1763) *s. l.*; Pilgrim & Palma 1982: 16.

*Anatoecus dentatus* (Scopoli, 1763); Pilgrim & Palma 1982: 16.

*Anatoecus dentatus dentatus* (Scopoli, 1763); Palma 1999: 379.

*Anatoecus dentatus dentatus* (Scopoli, 1763); Palma & Horning 2002: 6, 17.

*Anatoecus dentatus* (Scopoli, 1763); Price *et al.* 2003: 144.

*Anatoecus dentatus dentatus* (Scopoli, 1763); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1951: 17).

Type host: *Anas platyrhynchos platyrhynchos* Linnaeus, 1758.

New Zealand hosts: *Tadorna variegata* (J.F. Gmelin, 1789); *Chenonetta jubata* (Latham, 1802); *Anas gracilis* Buller, 1869; *Anas chlorotis* G.R. Gray, 1845; *Anas platyrhynchos platyrhynchos* Linnaeus, 1758; *Anas superciliosa* J.F. Gmelin, 1789; *Aythya novaeseelandiae* (J.F. Gmelin, 1789).

Other hosts: Two species of *Aix*; *Alopochen aegyptiaca* (Linnaeus, 1766); 14 other species of *Anas*; six species of *Anser*; nine other species of *Aythya*; three species of *Branta*; three species of *Bucephala*; *Clangula hyemalis* (Linnaeus, 1758); *Coscoroba coscoroba* (Molina, 1782); two species of *Dendrocygna*; three species of *Melanitta*; four species of *Mergus*; *Netta rufina* (Pallas, 1776); two species of *Oxyura*; *Polysticta stelleri* (Pallas, 1769); *Sarkidiornis melanotos* (Pennant, 1769); two species of *Somateria*; *Tadorna tadorna* (Linnaeus, 1758); *Thalassornis leuconotus* Eyton, 1838 (see Price *et al.* 2003: 144).

New Zealand localities: AK, BP, TK, WA, WN, SD, MB, NN, NC, MC, SC, KE, Macquarie Island.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Gressitt (1970: 326); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1374); Palma (1999); Paterson *et al.* (1999: 219); Palma & Horning (2002); Galloway (2005: 16); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Clay & Hopkins (1951: 15, figs 21–22); Kéler (1960a); Złotorzycka (1970); Moreby (1976: 92); Rudolph (1983: 16); Green & Palma (1991: 10, 29); Forrester *et al.* (1995: 14); Palma (1996b: 154); Price *et al.* (2003); Palma & Jensen (2005: 54, 61); Martín-Mateo (2009: 211, fig. 41); Palma & Peck (2013: 28); Grossi *et al.* (2014); Ahmad *et al.* (2015: 568).

Remarks: Many subspecies of *Anatoecus dentatus* have been described (see Price *et al.* 2003: 145) but, with a few exceptions, their type hosts are not New Zealand hosts. Therefore, until a detailed revision of the New Zealand populations of *A. dentatus* is available, I regard them as “*sensu lato*”, except for one subspecies (see below). *Anas chlorotis* is a new host record for *Anatoecus dentatus sensu lato* in New Zealand (voucher specimens in MONZ).

### ***Anatoecus dentatus magnicornutus* Złotorzycka, 1970**

*Anatoecus (Benatoecus) dentatus magnicornutus* Złotorzycka, 1970: 52, figs 53–54, photo 11.

*Anatoecus dentatus magnicornutus* Złotorzycka, 1970; Pilgrim & Palma 1982: 16.

*Anatoecus dentatus magnicornutus* Złotorzycka, 1970; Murray *et al.* 1990: 1374.

*Anatoecus dentatus magnicornutus* Złotorzycka, 1970; Palma 2010: 408.

Holotype ♂ in MNHW (Palma 1996b: 155; Jałoszyński *et al.* 2014: 654).

Type host: *Cygnus olor* (J.F. Gmelin, 1789).

New Zealand host: *Cygnus atratus* (Latham, 1790).

Other hosts: None.

New Zealand locality: WO.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010); Palma (2015a: 146).

Other significant references: Palma (1996b: 155).

Remarks: Contrary to Price *et al.* (2003: 145), I recognise *Anatoecus dentatus magnicornutus* as a valid taxon. It is not possible to determine if this louse was introduced to New Zealand with mute swans by human agency (Checklist Committee 2010: 32) or with black swans, because the New Zealand population of black swans is probably the

result of a mixture of self-introduced birds and others introduced by humans (Checklist Committee 2010: 33). Therefore, it is uncertain if *Anatoecus d. magnicornutus* is a native or an introduced louse.

### ***Anatoecus icterodes* (Nitzsch, 1818) sensu lato**

*Philoaterus* (*Docophorus*) *icterodes* Nitzsch, 1818: 290.

*Philoaterus icterodes* Nitzsch, 1818; Harrison 1916: 96 (as junior synonym of *Philoaterus dentatus* (Scopoli, 1763)).

*Anatoecus icterodes* Nitzsch, 1818 [sic]; Cummings 1916b: 655, figs 7B, 10, 12.

*Anatoecus icterodes* (Nitzsch, 1818); Hopkins & Clay 1952: 35.

*Anatoecus icterodes knechteli* Kéler, 1960a: 247.

*Anatoecus icterodes icterodes* (Nitzsch, 1818); Kéler 1960a: 299, figs 1, 2, 12(1), 30, 42, 48.

*A. icteroides* [sic] (Nitzsch, 1818); Gressitt 1970: 327.

*Anatoecus* (*Anatoecus*) *icterodes icterodes* (Nitzsch, 1818); Złotorzycka 1970: 58, figs 56–59.

*Anatoecus icterodes* (Nitzsch, 1818); Wise 1977: 58.

*Anatoecus icterodes* (Nitzsch, 1818) *s. l.*; Pilgrim & Palma 1982: 16–17.

*Anatoecus icterodes* (Nitzsch, 1818); Palma 1999: 379.

*Anatoecus icterodes icterodes* (Nitzsch, 1818); Palma & Horning 2002: 6.

*Anatoecus icterodes* (Nitzsch, 1818); Price *et al.* 2003: 145.

*Anatoecus icterodes icterodes* (Nitzsch, 1818); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1960: 39, pl. 2: fig. 3).

Type host: *Mergus serrator* Linnaeus, 1758.

New Zealand hosts: *Tadorna variegata* (J.F. Gmelin, 1789); *Tadorna tadornoides* (Jardine & Selby, 1828); *Hymenolaimus malacorhynchos* (J.F. Gmelin, 1789); *Anas gracilis* Buller, 1869; *Anas chlorotis* G.R. Gray, 1845; *Anas platyrhynchos platyrhynchos* Linnaeus, 1758; *Anas superciliosa* J.F. Gmelin, 1789; *Anas rhynchotis* Latham, 1802; *Aythya novaeseelandiae* (J.F. Gmelin, 1789).

Other hosts: Two species of *Aix*; *Alopothen aegyptiaca* (Linnaeus, 1766); 13 other species of *Anas*; five species of *Anser*; 10 other species of *Aythya*; *Biziura lobata* (Shaw, 1796); four species of *Branta*; two species of *Bucephala*; *Clangula hyemalis* (Linnaeus, 1758); two species of *Dendrocygna*; three species of *Melanitta*; three species of *Mergus*; two species of *Netta*; three species of *Oxyura*; *Plectropterus gambensis* (Linnaeus, 1766); *Sarkidiornis melanotos* (Pennant, 1769); two species of *Somateria*; *Tadorna tadorna* (Linnaeus, 1758); *Thalassornis leucotis* Eyton, 1838 (see Price *et al.* 2003: 145).

New Zealand localities: ND, CL, AK, WO, BP, TK, WA, WN, NN, NC, MC, SC, FD, Macquarie Island.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Kéler (1960a: 247); Gressitt (1970); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1374); Paterson *et al.* (1999: 219); Palma (1999); Palma & Horning (2002); Galloway (2005: 16); Palma (2010).

Other significant references: Cummings (1916b); Clay & Hopkins (1960: 37, figs 59–61); Kéler (1960a); Złotorzycka (1970); Moreby (1976: 92); Green & Palma (1991: 10, 29); Palma (1996b: 155); Price *et al.* (2003); Palma & Jensen (2005: 54, 61); Adam (2007: 176); Martín-Mateo (2009: 213); Palma & Peck (2013: 28); Grossi *et al.* (2014).

Remarks: Many subspecies of *Anatoecus icterodes* have been described (see Price *et al.* 2003: 146) but, with a few exceptions, their type hosts are not New Zealand hosts. Therefore, until a detailed revision of the New Zealand populations of *A. icterodes* is available, I regard them as “*sensu lato*”, except for one subspecies (see below).

### ***Anatoecus icterodes oloris* Złotorzycka, 1970**

Figs 66–67

*Anatoecus icterodes* ssp. ?; Kéler 1960a: 234.

*Anatoecus* (*Anatoecus*) *icterodes oloris* Złotorzycka, 1970: 50, fig. 48, photo 10.

*Anatoecus icterodes oloris* Złotorzycka, 1970; Pilgrim & Palma 1982: 16.

*Anatoecus icterodes oloris* Złotorzycka, 1970; Murray *et al.* 1990: 1374.

*Anatoecus i. oloris* Złotorzycka, 1970; Palma 2010: 408.

Holotype ♂ in MNHW (Palma 1996b: 155; Jałoszyński *et al.* 2014: 655).

Type host: *Cygnus olor* (J.F. Gmelin, 1789).

New Zealand hosts: *Cygnus olor* (J.F. Gmelin, 1789); *Cygnus atratus* (Latham, 1790).

Other hosts: None.

New Zealand localities: WO, WA, MC.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 219); Palma (2010); Palma (2015a: 146).

Other significant references: Kéler (1960a); Palma (1996b: 155).

Remarks: Contrary to Price *et al.* (2003: 146), I recognise *Anatoecus icterodes olor* as a valid taxon. This louse was introduced to New Zealand with mute swans by human agency (Checklist Committee 2010: 32). However, the population of black swans living in New Zealand is probably the result of a mixture of self-introduced birds and others introduced by human agency (Checklist Committee 2010: 33).

### ***Anatoecus singhi* Palma, 2015**

*Anatoecus* sp.; Pilgrim & Palma 1982: 16.

*Anatoecus* sp.; Murray *et al.* 1990: 1374.

*Anatoecus singhi* Palma, 2015a: 143, figs 1–4.

Holotype ♂ in MONZ.

Type host: *Cygnus atratus* (Latham, 1790).

New Zealand host: *Cygnus atratus* (Latham, 1790).

Other hosts: None.

New Zealand localities: WO, BP, WA, MB, NC, MC.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2015a).

Other significant references: None.

Remarks: The population of black swans living in New Zealand is probably the result of a mixture of self-introduced birds and others introduced by human agency (Checklist Committee 2010: 33). Therefore, it is not possible to determine if *Anatoecus singhi* is a native or an introduced species.

### **Genus *Aquanirmus* Clay & Meinertzhagen, 1939**

*Aquanirmus* Clay & Meinertzhagen, 1939a. *Entomologist* 72: 163. Type species: *Degeeriella runcinata* (Nitzsch [*in* Giebel], 1866) = *Aquanirmus runcinatus* (Nitzsch [*in* Giebel], 1866) (by original designation).

### ***Aquanirmus australis* Kettle, 1974**

Figs 68–69

*Aquanirmus australis* Kettle, 1974b: 337, figs 1–2.

*Aquanirmus australis* Kettle, 1974; Wise 1977: 58.

*Aquanirmus australis* Kettle, 1974; Pilgrim & Palma 1982: 5.

*Aquanirmus australis* Kettle, 1974; Murray *et al.* 1990: 1367.

*Aquanirmus australis* Kettle, 1974; Palma 2010: 408.

Holotype ♂ in NZAC.

Type host: *Poliocephalus rufopectus* (G.R. Gray, 1843).

New Zealand host: *Poliocephalus rufopectus* (G.R. Gray, 1843).

Other hosts: None.

New Zealand localities: AK, HB, TK, WA.

Geographic distribution: New Zealand.

New Zealand references: Kettle (1974b); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1367); Storer (2000: 37); Palma (2010).

Other significant references: Price *et al.* (2003: 147).

Remarks: *Aquanirmus australis* is an endemic species, exclusively parasitic on the New Zealand dabchick.

**Aquanirmus podiceps (Denny, 1842)****New Record**

- Philopterus (Nirmus) podiceps* Denny, 1842: 53, 142, pl. 10: fig. 9.  
*Degeeriella podicipis* [sic] Denny, 1842 [sic]; Harrison 1916: 121.  
*Aquanirmus podicipis* [sic] (Denny, 1842); Hopkins & Clay 1952: 37.  
*Aquanirmus podicipis* [sic] (Denny, 1842); Edwards 1965: 929, figs 2, 4, 6.  
*Aquanirmus* sp.; Pilgrim & Palma 1982: 5.  
*Aquanirmus* sp.; Murray *et al.* 1990: 1367.  
*Aquanirmus podiceps* (Denny, 1842); Price *et al.* 2003: 147.  
*Aquanirmus* sp.; Palma 2010: 408.

Lectotype ♀ in NHML (Edwards 1965: 930).

Type host: *Podiceps cristatus cristatus* (Linnaeus, 1758)

New Zealand host: *Podiceps cristatus australis* Gould, 1844.

Other host: *Podiceps cristatus infuscatus* Salvadori, 1844.

New Zealand localities: MC, OL, FD.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Edwards (1965); Clay (1976b: 543); Storer (2000: 37); Price *et al.* (2003).

Material examined and repository: 39♂, 33♀, 6N (4 samples, MONZ).

Remarks: This is the first record of *Aquanirmus podiceps* for New Zealand, because the New Zealand references cited above reported this louse as “*Aquanirmus* sp.” only. Also, *Podiceps cristatus australis* is a new host record for *A. podiceps* (voucher specimens in MONZ).

**Genus Ardeicola Clay, 1936**

- Ardeicola* Clay, 1936. *Proc. Zool. Soc. London* [no volume number]: 615. Type species: *Esthiopterum ardeae* (Linnaeus, 1758) = *Ardeicola ardeae* (Linnaeus, 1758) (by original designation).

**Ardeicola expallidus Blagoveshtchensky, 1940**

- Ardeicola expallida* [sic] Blagoveshtchensky, 1940: 69, 89, fig. 21.  
*Ardeicola gaibagla* Ansari, 1947: 256, fig. 1.  
*Ardeicola albulus* Eichler, 1948: 107, figs 2–3.  
*Ardeicola albulus* Eichler, 1948; Hopkins & Clay 1952: 38.  
*Ardeicola expallidus* Blagoveshtchensky, 1940; Hopkins & Clay 1952: 39. Emendation.  
*Ardeicola gaibagla* Ansari, 1947; Hopkins & Clay 1952: 39.  
*Ardeicola gaibagla* Ansari, 1947; Tuff 1967: 251, figs 17–19.  
*Ardeicola expallida* Blagoveshtchensky, 1940; Tuff 1970: 484.  
*Ardeicola expallidus* Blagoveshtchensky, 1940; Pilgrim & Palma 1982: 15.  
*Ardeicola expallidus* Blagoveshtchensky, 1940; Murray *et al.* 1990: 1373.  
*Ardeicola albulus* Eichler, 1948; Mey 1994: 33, figs 15–16.  
*Ardeicola expallidus* Blagoveshtchensky, 1940; Forrester *et al.* 1995: 9.  
*Ardeicola expallidus* Blagoveshtchensky, 1940; Palma 1999: 379.  
*Ardeicola expallidus* Blagoveshtchensky, 1940; Palma 2010: 408.

Syntypes ♂♀, repository unknown (Palma 1996b: 156), but probably in ZMAS.

Type host: *Egretta garzetta garzetta* (Linnaeus, 1766).

New Zealand hosts: *Ardea modesta* J.E. Gray, 1831; *Ardea ibis coromanda* (Boddaert, 1783); *Egretta garzetta immaculata* (Gould, 1846).

Other hosts: *Ardea alba* Linnaeus, 1758; *Ardea ibis ibis* Linnaeus, 1758; *Egretta garzetta garzetta* (Linnaeus, 1766); *Egretta thula* (Molina, 1782).

New Zealand localities: HB, NN, NC, MC, SC, WD, CO, DN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 221); Palma (1999); Palma (2010).

Other significant references: Tuff (1967); Tuff (1970); Green & Palma (1991: 11, 29); Mey (1994); Forrester *et al.* (1995); Palma (1996b: 156); Price *et al.* (2003: 148); Palma & Jensen (2005: 54, 61); Ahmad *et al.* (2013).

Remarks: *Ardeicola expallidus* is widespread on a number of heron species, it but has been collected infrequently in New Zealand.

***Ardeicola neopallidus* Price, Hellenthal & Palma, 2003**

Figs 70–71

*Lipeurus signatus* var. *pallida* Piaget, 1880: 312. Preoccupied by *Lipeurus pallidus* Giebel, 1874: 219.

*Esthiopterum pallidum* Piaget, 1880 [sic]; Harrison 1916: 139.

*Ardeicola pallidus* (Piaget, 1880) “*nec L. pallidus* Giebel, 1866” [sic]; Hopkins & Clay 1952: 40.

*Ardeicola pallidus* (Piaget, 1880); Palma 1999: 379.

*Ardeicola neopallidus* Price, Hellenthal & Palma, 2003: 149. *Nomen novum* for *Lipeurus pallidus* Piaget, 1880: 312.

*Ardeicola neopallidus* Price, Hellenthal & Palma, 2003; Palma 2010: 408.

Syntypes ♂♀ in NHML (Thompson 1937–1939: 342).

Type host: *Egretta sacra* (J.F. Gmelin, 1789).

New Zealand host: *Egretta sacra sacra* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: AK, WN.

Geographic distribution: Asia, Australasia and tropical Pacific Islands.

New Zealand references: Palma (1999); Palma (2010).

Other significant reference: Price *et al.* (2003: 149).

Remarks: The reef heron has been searched for lice in very few occasions in New Zealand, resulting in only two records of *Ardeicola neopallidus*.

***Ardeicola pilgrimi* Tandan, 1972**

*Ardeicola pilgrimi* Tandan, 1972b: 52, figs 1–9.

*Ardeicola pilgrimi* Tandan, 1972; Wise 1977: 58.

*Ardeicola pilgrimi* Tandan, 1972; Horning *et al.* 1980: 5, 11.

*Ardeicola pilgrimi* Tandan, 1972; Pilgrim & Palma 1982: 15.

*Ardeicola pilgrimi* Tandan, 1972; Murray *et al.* 1990: 1373.

*Ardeicola pilgrimi* Tandan, 1972; Palma 2010: 408.

Holotype ♂ in CMNZ (Nicholls *et al.* 1998: 30).

Type host: *Egretta novaehollandiae novaehollandiae* (Latham, 1790).

New Zealand host: *Egretta novaehollandiae novaehollandiae* (Latham, 1790).

Other hosts: None.

New Zealand localities: WA, WN, NN, KA, NC, MC, SC, KE, SN.

Geographic distribution: Australasia.

New Zealand references: Tandan (1972); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Nicholls *et al.* (1998: 30); Paterson *et al.* (1999: 221); Palma (2010).

Other significant references: Green & Palma (1991: 11, 28); Palma (1996b: 157); Price *et al.* (2003: 149).

Remarks: Although the white-faced heron has become established in New Zealand from Australia in relatively recent times (Checklist Committee 2010: 161), *Ardeicola pilgrimi* was first described from New Zealand material.

***Ardeicola plataleae* (Linnaeus, 1758)**

*Pediculus plataleae* Linnaeus, 1758: 613.

*Esthiopterum plataleae* Linnaeus, 1758 [sic]; Harrison 1916: 139.

*Ardeicola plataleae* (Linnaeus, 1758); Clay & Hopkins 1950: 245, figs 29–33, pl. 1: fig. 5.

*Ardeicola plataleae* (Linnaeus, 1758); Pilgrim & Palma 1982: 16.

*Ardeicola plataleae* (Linnaeus, 1758); Murray *et al.* 1990: 1373.

*Ardeicola plataleae* (Linnaeus, 1758); Price *et al.* 2003: 149.

*Ardeicola plataleae* (Linnaeus, 1758); Palma 2010: 408.



Neotype ♀ in NHML (Clay & Hopkins 1950: 246, pl. 1: fig. 5).

Type host: *Platalea leucorodia* Linnaeus, 1758.

New Zealand host: *Platalea regia* Gould, 1838.

Other hosts: None.

New Zealand localities: HB, WD.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Séguy (1944: 356, figs 529–530); Clay & Hopkins (1950); Moreby (1976: 92); Martín-Mateo (1994: 110, figs 1e, 3a,b); Palma (1996b: 157); Price *et al.* (2003); Martín-Mateo (2009: 231, fig. 47).

Remarks: Since the first record of the royal spoonbill in New Zealand in 1861, this host has become well established with many breeding localities in this country (Checklist Committee 2010: 167). However, *Ardeicola plataleae* has been collected only twice in this country.

### ***Ardeicola raphidius* (Nitzsch [*in* Giebel], 1866)**

*Lipeurus raphidius* Nitzsch [*in* Giebel], 1866; 384.

*Esthiopterum raphidium* Nitzsch [*in* Giebel], 1866 [sic]; Harrison 1916: 141.

*Ardeicola raphidius* (Nitzsch [*in* Giebel], 1866); Hopkins & Clay 1952: 41.

*Ardeicola raphidius* (Nitzsch, 1866) [sic]; Pilgrim & Palma 1982: 16.

*Ardeicola raphidius* (Nitzsch, 1866) [sic]; Murray *et al.* 1990: 1373.

*Ardeicola raphidius* (Nitzsch [*in* Giebel], 1866); Price *et al.* 2003: 149.

*Ardeicola raphidius* (Nitzsch [*in* Giebel], 1866); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1), Palma & Pilgrim (1984: 150) and Palma (1996b: 157).

Type host: *Plegadis falcinellus* (Linnaeus, 1766).

New Zealand host: *Plegadis falcinellus* (Linnaeus, 1766).

Other host: *Plegadis chihi* (Vieillot, 1817).

New Zealand localities: WI.

Geographic distribution: Eurasia; Africa; Americas; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Dubinin (1938: fig. 15); Séguy (1944: 356, figs 531–532); Butler & O'Connor (1994: 453); Forrester *et al.* (1995: 12); Palma (1996b: 157); Price *et al.* (2003).

Remarks: Although the glossy ibis is a regular vagrant to New Zealand (Checklist Committee 2010: 166), there is only one record of *Ardeicola raphidius* from this country.

### ***Ardeicola stellaris* (Denny, 1842)**

*Philopterus* (*Lipeurus*) *stellaris* Denny, 1842: 59, 178, pl. 15: fig. 3.

*Esthiopterum stellare* Denny, 1842; [sic]; Harrison 1916: 142.

*Ardeicola stellaris* (Denny, 1842); Hopkins & Clay 1952: 41.

*Ardeicola stellaris* (Denny, 1842) *s. l.*; Pilgrim & Palma 1982: 15.

*Ardeicola stellaris* (Denny, 1842); Murray *et al.* 1990: 1373.

*Ardeicola stellaris* (Denny, 1842); Price *et al.* 2003: 149.

*Ardeicola stellaris* (Denny, 1842); Palma 2010: 408.

Syntypes ♂♀ in NHML (Palma 1996b: 157).

Type host: *Botaurus stellaris* (Linnaeus, 1758).

New Zealand host: *Botaurus poiciloptilus* (Wagler, 1827).

Other hosts: None.

New Zealand localities: ND, BP, HB, NN.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990: 1373); Paterson *et al.* (1999: 221); Palma (2010).

Other significant references: Séguy (1944: 357, figs 533–534); Green & Palma (1991: 11, 29); Palma (1996b: 157); Price *et al.* (2003: 149).

Remarks: Pilgrim & Palma (1982: 15) regarded the population of *Ardeicola stellaris* from the Australasian bittern as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of more samples, including one from the type host, shows that making such difference is not warranted.

### Genus *Austrogoniodes* Harrison, 1915

*Austrogoniodes* Harrison, 1915a. *Parasitology* 7: 398. Type species: *Goniocotes waterstoni* Cummings, 1914 = *Austrogoniodes waterstoni* (Cummings, 1914) (by original designation).

*Cesareus* Kéler, 1952. *Jour. Entomol. Soc. Southern Africa* 15: 221. Type species: *Cesareus concii* Kéler, 1952 = *Austrogoniodes concii* (Kéler, 1952) (by original designation).

### *Austrogoniodes antarcticus* Harrison, 1937

*Austrogoniodes antarcticus* Harrison, 1937: 20, pl. 1: figs 6–7.

*Austrogoniodes antarcticus* Harrison, 1937; Clay 1967: 149, figs 1–2, 6, 8–9, 16, 18–20, 28.

*Austrogoniodes antarcticus* Harrison, 1937; Clay & Moreby 1967: 161, 167.

*Austrogoniodes antarcticus* Harrison, 1937; Pilgrim & Palma 1982: 4.

*Austrogoniodes antarcticus* Harrison, 1937; Murray *et al.* 1990: 1368.

*Austrogoniodes antarcticus* Harrison, 1937; Banks & Paterson 2004: 94, figs 3a, 5, 7d, 9–11.

*Austrogoniodes antarcticus* Harrison, 1937; Palma 2010: 408.

Holotype ♂ in AMSA (Palma 1996b: 158).

Type host: *Pygoscelis adeliae* (Hombron & Jacquinet, 1841).

New Zealand host: *Pygoscelis adeliae* (Hombron & Jacquinet, 1841).

Other hosts: None.

New Zealand localities: RO.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Clay (1967: 150); Clay & Moreby (1967); Pilgrim & Palma (1982); Murray *et al.* (1990); Banks & Paterson (2004; 2005: 744); Banks *et al.* (2006: 158); Palma (2010).

Other significant references: Kéler (1952: 221); Palma (1996b: 158); Price *et al.* (2003: 151).

Remarks: Collections of *Austrogoniodes antarcticus* from the Ross Dependency (Antarctica) show that this louse species has both low abundance and low prevalence of infestation.

### *Austrogoniodes concii* (Kéler, 1952)

*Cesareus concii* Kéler, 1952: 223, figs 14–19.

*Austrogoniodes concii* (Kéler, 1952); Hopkins & Clay 1953: 435.

*Austrogoniodes concii* [sic] (Kéler, 1952); Clay 1967: 151, 154, figs 29, 36.

*Austrogoniodes concii* (Kéler, 1952); Wise 1977: 59.

*Austrogoniodes concii* (Kéler, 1952) *s. l.*; Horning *et al.* 1980: 5, 8.

*Austrogoniodes concii* (Kéler, 1952) *s. l.*; Pilgrim & Palma 1982: 4.

*Austrogoniodes concii* (Kéler, 1952); Murray *et al.* 1990: 1368.

*Austrogoniodes concii* (Kéler, 1952); Banks & Paterson 2004: 93, figs 4a, 9–11.

*Austrogoniodes concii* (Kéler, 1952); Palma 2010: 409.

Holotype ♂ in SAIM (Palma 1996b: 159).

Type host: *Eudyptes moseleyi* Mathews & Iredale, 1921.

New Zealand hosts: *Eudyptes pachyrhynchus* G.R. Gray, 1845; *Eudyptes robustus* Oliver, 1953; *Eudyptes sclateri* Buller, 1888; *Megadyptes antipodes* (Hombron & Jacquinet, 1841).

Other hosts: None.

New Zealand localities: HB, WN, KA, NC, MC, SC, WD, FD, SN, AN, CA.

Geographic distribution: Australasia; Antarctica; Subantarctic Islands; Southern Oceans.

New Zealand references: Gressitt (1964: 538); Clay (1967); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Pérez (1985: 160); Murray *et al.* (1990); Marris (2000: 187); Banks & Paterson (2004; 2005: 744); Banks *et al.* (2006: 158); Palma (2010).

Other significant references: Séguy (1953: 558, figs 1–2); Green & Palma (1991: 11, 24); Palma (1996b: 159); Price *et al.* (2003: 151); Hänel & Palma (2007: 112, 122, 131).

Remarks: Pilgrim & Palma (1982: 4–5) regarded the populations of *Austrogoniodes concii* from four penguin species as somewhat different from that of the type host, and qualified them as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted.

The type locality of *Austrogoniodes concii* is Tristan da Cunha, and its type host has been referred to as *Eudyptes chrysocome* (J.R. Forster, 1781) or as its junior synonym *Eudyptes crestatus* Miller, 1784 in most publications. However, the species *E. chrysocome* is now restricted to the population of rockhopper penguins breeding in southern South America (Checklist Committee 2010: 56). Therefore, the type host of *A. concii* is the population of rockhopper penguins from Tristan da Cunha, i.e. *Eudyptes moseleyi* (Checklist Committee 2010: 57).

The record of *Austrogoniodes concii* on *Eudyptes crestatus* (= *chrysocome*) from “New Zealand–Christchurch” in Clay (1967: 154) is erroneous (see Pilgrim & Palma 1982: 4, 29, Note 3).

### ***Austrogoniodes cristati* Kéler, 1952**

“*Austrogoniodes struthus*” Harrison, 1937: 15 (not *Austrogoniodes struthus* Harrison, 1915a).

“*Austrogoniodes waterstoni*” Harrison, 1937: 15 (not *Goniocotes waterstoni* Cummings, 1914).

*Austrogoniodes cristati* Kéler, 1952: 230, figs 20–22.

*Austrogoniodes cristati* Kéler, 1952; Hopkins & Clay 1953: 435.

*Austrogoniodes cristati* Kéler, 1952; Clay 1967: 152, 154, figs 42, 48.

*Austrogoniodes cristati* Kéler, 1952; Wise 1977: 59.

*Austrogoniodes cristati* Kéler, 1952; Horning *et al.* (1980: 5, 8).

*Austrogoniodes cristati* Kéler, 1952; Pilgrim & Palma 1982: 4, 29, note 6.

*Austrogoniodes cristati* Kéler, 1952; Murray *et al.* 1990: 1368.

*Austrogoniodes cristati* Kéler, 1952; Banks & Paterson 2004: 92, figs 9–11.

*Austrogoniodes cristati* Kéler, 1952; Palma 2010: 408.

Holotype ♂ in SAIM (Palma 1996b: 159).

Type host: *Eudyptes moseleyi* Mathews & Iredale, 1921.

New Zealand hosts: *Eudyptes filholi* Hutton, 1879; *Eudyptes pachyrhynchus* G.R. Gray, 1845; *Eudyptes robustus* Oliver, 1953; *Eudyptes sclateri* Buller, 1888; *Eudyptes chrysolophus* (Brandt, 1837); *Eudyptes schlegeli* Finsch, 1876.

Other hosts: None.

New Zealand localities: HB, KA, NC, MC, SC, WD, CO, DN, FD, SN, AN, CA, Macquarie Island.

Geographic distribution: Australasia; Antarctica; Subantarctic Islands; Southern Oceans.

New Zealand references: Kéler (1954: 58); Clay (1964a: 230); Gressitt (1964: 538); Clay (1967); Watson (1967: 71); Clay & Moreby (1970: 216); Gressitt (1970: 327); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Pérez (1985: 160); Murray *et al.* (1990); Palma (1996b: 159); Palma (1999: 384, note E); Marris (2000: 187); Palma & Horning (2002: 6, 17); Banks & Paterson (2004; 2005: 744); Banks *et al.* (2006: 158); Palma (2010).

Other significant references: Clay & Moreby (1967: 161, 167); Green & Palma (1991: 11, 24); Price *et al.* (2003: 151); Hänel & Palma (2007: 112, 122, 131).

Remarks: The two misidentifications listed above from Harrison (1937: 15) have been clarified by Palma & Horning (2002: 6, 20, notes 1–2). The type locality of *Austrogoniodes cristati* is Tristan da Cunha, and its type host has been referred to as *Eudyptes chrysocome* (J.R. Forster, 1781) or as its junior synonym *Eudyptes crestatus* Miller, 1784 in most publications. However, the species *E. chrysocome* is now restricted to the population of rockhopper penguins breeding in southern South America (Checklist Committee 2010: 56). Therefore, the type host of *A. cristati* is the population of rockhopper penguins from Tristan da Cunha, i.e. *Eudyptes moseleyi* (Checklist Committee 2010: 57).

### ***Austrogoniodes hamiltoni* Harrison, 1937**

Figs 72–73

*Austrogoniodes hamiltoni* Harrison, 1937: 18, pl. 1: figs 8–9, pl. 2: figs 1–2.

*Cesareus hamiltoni* (Harrison, 1937); Kéler 1952: 237.

*Cesareus hamiltoni* (Harrison, 1937); Kéler 1954: 54, figs 4–7.

- ? *Austrogoniodes hamiltoni* Harrison, 1937; Clay 1940a: 297.  
*Austrogoniodes hamiltoni* Harrison, 1937; Clay 1967: 152, 154, figs 15, 23, 30, 35, 44.  
*Austrogoniodes hamiltoni* Harrison, 1937; Wise 1977: 59.  
*Austrogoniodes hamiltoni* Harrison, 1937; Pilgrim & Palma 1982: 4, 29, notes 5, 7.  
*Austrogoniodes hamiltoni* Harrison, 1937; Murray *et al.* 1990: 1368.  
*Austrogoniodes hamiltoni* Harrison, 1937; Banks & Paterson 2004: 93, figs 9–11.  
*Austrogoniodes hamiltoni* Harrison, 1937; Palma 2010: 408.

Holotype ♂ in AMSA (Palma 1996b: 159)

Type host: *Eudyptes filholi* Hutton, 1879.

New Zealand hosts: *Eudyptes filholi* Hutton, 1879; *Eudyptes schlegeli* Finsch, 1876.

Other hosts: None.

New Zealand localities: CO, DN, AN, CA, Macquarie Island.

Geographic distribution: Australasia; Antarctica; Subantarctic Islands; Southern Oceans.

New Zealand references: Harrison (1937); Kéler (1954); Clay (1967: 154); Watson (1967: 71); Clay & Moreby (1970: 216); Gressitt (1970: 327); Wise (1977); Pilgrim & Palma (1982); Pérez (1985: 161); Murray *et al.* (1990); Palma (1996b: 159); Marris (2000: 187); Palma & Horning (2002: 7, 17); Banks & Paterson (2004; 2005: 745); Banks *et al.* (2006: 158); Palma (2010).

Other significant references: Kéler (1952); Clay (1940a); Clay & Moreby (1967: 180, fig. 79); Green & Palma (1991: 11, 24); Price *et al.* (2003: 151).

Remarks: The type locality of *Austrogoniodes hamiltoni* is Macquarie Island, and its type host has been referred to as *Eudyptes chrysocome* (J.R. Forster, 1781) or as its junior synonym *Eudyptes crestatus* Miller, 1784 in most publications. However, the species *E. chrysocome* is now restricted to the population of rockhopper penguins breeding in southern South America (Checklist Committee 2010: 56). Therefore, the type host of *A. hamiltoni* is the population of rockhopper penguins from Macquarie Island, i.e. *Eudyptes filholi* Hutton, 1879 (Checklist Committee 2010: 56).

Records of *Austrogoniodes hamiltoni* on *Eudyptes pachyrhynchus* from “Macquarie I.” and on *Eudyptes sclateri* from “Antipodes Is” in Clay (1967: 154) are the result of straggling or contaminations (see Pilgrim & Palma 1982: 4–5, 29, notes 4–5, 7; Palma 1999: 383, note A).

### ***Austrogoniodes macquariensis* Harrison, 1937**

- Austrogoniodes macquariensis* Harrison, 1937: 17, pl. 1: figs 4–5.  
*Austrogoniodes macquariensis* Harrison, 1937; Clay 1967: 154.  
*Austrogoniodes macquariensis* Harrison, 1937; Watson 1967: 71.  
*Austrogoniodes macquariensis* Harrison, 1937; Wise 1977: 59.  
*Austrogoniodes macquariensis* Harrison, 1937; Pilgrim & Palma 1982: 4, 29, note 5.  
*Austrogoniodes macquariensis* Harrison, 1937; Murray *et al.* 1990: 1368.  
*Austrogoniodes macquariensis* Harrison, 1937; Palma & Horning 2002: 7, 17.  
*Austrogoniodes macquariensis* Harrison, 1937; Banks & Paterson 2004: 94, figs 3c, 9–11. In part.  
*Austrogoniodes macquariensis* Harrison, 1937; Palma 2010: 408.

Holotype ♂ in AMSA (Palma 1996b: 159).

Type host: *Eudyptes filholi* Hutton, 1879.

New Zealand hosts: *Eudyptes filholi* Hutton, 1879; *Eudyptes schlegeli* Finsch, 1876.

Other hosts: None.

New Zealand localities: WA, CO, DN, SN, AN, CA, Macquarie Island.

Geographic distribution: Australasia; Antarctica; Subantarctic Islands; Southern Oceans.

New Zealand references: Harrison (1937); Kéler (1952: 221); Clay (1967); Watson (1967); Clay & Moreby (1970: 216); Gressitt (1970: 326); Wise (1977); Lowry *et al.* (1978: 138); Pilgrim & Palma (1982); Pérez (1985: 163); Murray *et al.* (1990); Palma (1996b: 159); Marris (2000: 187); Palma & Horning (2002: 7, 17); Banks & Paterson (2004); Banks *et al.* (2006: 158); Palma (2010).

Other significant references: Price *et al.* (2003: 151).

Remarks: The type locality of *Austrogoniodes macquariensis* is Macquarie Island, and its type host has been referred to as *Eudyptes chrysocome* (J.R. Forster, 1781) or as its junior synonym *Eudyptes crestatus* Miller, 1784 in most publications. However, the species name *E. chrysocome* is now restricted to the population of rockhopper penguins breeding in southern South America (Checklist Committee 2010: 56). Therefore, the type host of *A. macquariensis* is the population of rockhopper penguins from Macquarie Island, i.e. *Eudyptes filholi* Hutton, 1879 (Checklist Committee 2010: 56). Also, see “Remarks” below, under *Austrogoniodes macquariensis* Harrison, 1937 *sensu lato*.

***Austrogoniodes macquariensis* Harrison, 1937 *sensu lato***

*Cesareus macquariensis* (Harrison, 1937); Kéler 1954: 50, figs 2–3.

*Austrogoniodes macquariensis* Harrison, 1937; Clay 1967: 152, 154, figs 31, 34, 47.

*Austrogoniodes macquariensis* Harrison, 1937; Horning *et al.* 1980: 5, 8.

*Austrogoniodes* sp.; Horning *et al.* 1980: 5, 8.

*Austrogoniodes macquariensis* Harrison, 1937 *s. l.*; Pilgrim & Palma 1982: 4.

*Austrogoniodes* sp.; Pilgrim & Palma 1982: 4.

*Austrogoniodes* sp.; Murray *et al.* 1990: 1368.

*Austrogoniodes macquariensis* Harrison, 1937; Banks & Paterson 2004: 94, figs 3c, 9–11. In part.

New Zealand host: *Eudyptes chrysolophus* (J.F. Brandt, 1837).

Other host: *Eudyptes chrysocome* (J.R. Forster, 1781).

New Zealand locality: SN.

Geographic distribution: Australasia; Antarctica; Subantarctic Islands; Southern Oceans.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Banks & Paterson (2004).

Other significant references: Kéler (1954); Clay (1967); Clay & Moreby (1967: 161, 167).

Remarks: I regard the populations of *Austrogoniodes macquariensis* from *Eudyptes chrysocome* and *E. chrysolophus* as “*sensu lato*” because the males show subtle but consistent differences from those parasitising *Eudyptes filholi* and *E. schlegeli*.

Records of *Austrogoniodes macquariensis* from other penguin species listed in Clay (1967: 154) and from two species of *Pygoscelis* in Price *et al.* (2003: 151) are cases of either straggling or contaminations (see Pilgrim & Palma 1982: 4–5, 29, notes 4–5; Banks & Paterson 2004: 100).

***Austrogoniodes mawsoni* Harrison, 1937**

*Austrogoniodes mawsoni* Harrison, 1937: 15, pl. 1: figs 2–3.

*Cesareus mawsoni* (Harrison, 1937); Kéler 1952: 237.

*Austrogoniodes mawsoni* Harrison, 1937; Clay 1967: 151, 154, figs 13, 27, 40–41, 45.

*Austrogoniodes mawsoni* Harrison, 1937; Pilgrim & Palma 1982: 4.

*Austrogoniodes mawsoni* Harrison, 1937; Murray *et al.* 1990: 1367.

*Austrogoniodes mawsoni* Harrison, 1937; Banks & Paterson 2004: 93, figs 3d, 9–11.

*Austrogoniodes mawsoni* Harrison, 1937; Palma 2010: 408.

Holotype ♂ in AMSA (Palma 1996b: 160).

Type host: *Aptenodytes forsteri* G.R. Gray, 1844.

New Zealand host: *Aptenodytes forsteri* G.R. Gray, 1844.

Other hosts: None.

New Zealand locality: RO.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Clay (1967: 154); Clay & Moreby (1967: 161, 167); Pilgrim & Palma (1982); Murray *et al.* (1990); Banks & Paterson (2004; 2005: 744); Banks *et al.* (2006: 158); Palma (2010).

Other significant references: Kéler (1952: 221); Palma (1996b: 160); Price *et al.* (2003: 151).

Remarks: Collections of *Austrogoniodes mawsoni* from the Ross Dependency (Antarctica) show that this louse species has both low abundance and low prevalence of infestation.

***Austrogoniodes strutheus* Harrison, 1915 nomen dubium**

*Austrogoniodes strutheus* Harrison, 1915a: 399, fig. 2, pl. 27: fig. 15.

*Austrogoniodes strutheus* Harrison, 1915; Guimarães 1938: 43 (as junior synonym of *A. bifasciatus* (Piaget, 1885))

*Austrogoniodes strutheus* Harrison, 1915; Kéler 1952: 220, 231, 236.

*Austrogoniodes strutheus* Harrison, 1915; Clay 1967: 153.

*Austrogoniodes ?strutheus* Harrison, 1915; Clay & Moreby 1970: 216.

*Austrogoniodes ?strutheus* Harrison, 1915; Gressitt 1970: 327.

*Austrogoniodes strutheus* Harrison, 1915 *nom. dub.*; Banks & Paterson 2004: 90.

*Austrogoniodes strutheus* Harrison, 1915; Pilgrim & Palma 1982: 5, 29, note 7.

*Austrogoniodes strutheus* Harrison, 1915; Murray *et al.* 1990: 1368.

*Austrogoniodes strutheus* Harrison, 1915; Palma 2010: 408.

Syntypes ♂♀, presumed lost. See Clay (1967: 153).

Type host: *Eudyptes sclateri* Buller, 1888.

New Zealand host: *Eudyptes sclateri* Buller, 1888.

Other hosts: None.

New Zealand locality: "Probably from one of the southern islands of New Zealand" (Harrison, 1915a: 402)

Geographic distribution: New Zealand.

New Zealand references: Harrison (1915a); Harrison (1937); Clay (1967); Clay & Moreby (1970); Gressitt (1970); Wise (1977: 59); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma & Horning (2002: 20, note 1); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Guimarães (1938); Kéler (1952); Price *et al.* (2003: 151); Banks & Paterson (2004).

Remarks: In agreement with Banks & Paterson (2004: 90), I regard this species as a *nomen dubium* following the discussion on its uncertain taxonomic status in Clay (1967: 153).

***Austrogoniodes vanalphenae* Banks & Palma, 2003**

*Austrogoniodes* sp.; Pilgrim & Palma 1982: 4.

*Austrogoniodes* sp.; Murray *et al.* 1990: 1368.

*Austrogoniodes vanalphenae* Banks & Palma, 2003: 69, figs 1, 3–7A, 8–9.

*Austrogoniodes vanalphenae* Banks & Palma, 2003; Banks & Paterson 2004: 92 figs 1, 3b, 4b, 6, 7c.

*Austrogoniodes vanalphenae* Banks & Palma, 2003; Palma 2010: 408.

Holotype ♀ in MONZ.

Type host: *Megadyptes antipodes* (Hombron & Jacquinet, 1841).

New Zealand host: *Megadyptes antipodes* (Hombron & Jacquinet, 1841).

Other hosts: None.

New Zealand localities: WN, KA, MC, SC, DN, AU.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Banks & Palma (2003); Banks & Paterson (2004; 2005: 744); Banks *et al.* (2006: 158); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: None.

Remarks: *Austrogoniodes vanalphenae* is an endemic and vulnerable species (Buckley *et al.* 2012), exclusively parasitic on yellow-eyed penguins.

***Austrogoniodes waterstoni* (Cummings, 1914)**

*Goniocotes waterstoni* Cummings, 1914: 173, fig. 8.

*Austrogoniodes waterstoni* (Cummings, 1914); Harrison 1915a: 399.

*Cesareus waterstoni* (Cummings, 1914); Kéler 1952: 237.

*Austrogoniodes waterstoni* (Cummings, 1914); Clay 1967: 152, 155, figs 14, 39.

*Austrogoniodes waterstoni* (Cummings, 1914); Wise 1977: 59.

*Austrogoniodes waterstoni* (Cummings, 1914); Horning *et al.* 1980: 5, 8.

*Austrogoniodes waterstoni* (Cummings, 1914); Pilgrim & Palma 1982: 4.

*Austrogoniodes waterstoni* (Cummings, 1914); Murray *et al.* 1990: 1368.

*Austrogoniodes waterstoni* (Cummings, 1914); Banks & Paterson 2004: 94, figs 9–11.

*Austrogoniodes waterstoni* (Cummings, 1914); Palma 2010: 408.

Syntypes ♂♀ in NHML (Palma 1996b: 160).

Type host: *Eudiptula minor* (J.R. Forster, 1781).

New Zealand host: *Eudiptula minor* (J.R. Forster, 1781).

Other hosts: None.

New Zealand localities: AK, BP, TK, WN, SD, MB, NN, KA, NC, MC, SC, WD, CH, SI, SN.

Geographic distribution: Australasia.

New Zealand references: Kéler (1952: 220); Clay (1967); Gressitt (1970: 327); Pilgrim (1974: 1035, fig. 5); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Banks & Paterson (2004); Banks *et al.* (2006: 158); Palma (2010).

Other significant references: Harrison (1915a); Thompson (1940b: 498, pl. 10: figs 3–4); Green & Palma (1991: 11, 24); Palma (1996b: 160); Price *et al.* (2003: 151).

Remarks: Checklist Committee (2010: 61) do not recognise any subspecies within *Eudiptula minor*. However, the type host of *Austrogoniodes waterstoni* is from the Australian population, sometimes referred to as *Eudiptula minor novaehollandiae* (Stephens, 1826).

### Genus *Bedfordiella* Thompson, 1937

*Bedfordiella* Thompson, 1937b. *Ann. Mag. Nat. Hist.* (Ser. 10) 20: 434. Type species: *Bedfordiella unica* Thompson, 1937b (by original designation).

#### *Bedfordiella unica* Thompson, 1937

Figs 74–75

*Bedfordiella unica* Thompson, 1937b: 434, figs 1–2.

*Bedfordiella unica* Thompson, 1937; Hopkins & Clay 1952: 50.

*Bedfordiella unica* Thompson, 1937; Séguy 1953: 562, figs 6–8.

*Bedfordiella unica* Thompson, 1937; Timmermann 1961c: 38, figs 7, 9–10.

*Bedfordiella simsi* Timmermann, 1961c: 39, figs 8, 11.

*Bedfordiella unica* Thompson, 1937; Timmermann 1965: 126, figs 64–65, pl. 8: fig. 4.

*Bedfordiella simsi* Timmermann, 1961; Timmermann 1965: 126, fig. 66, pl. 9: fig. 1.

*Bedfordiella simsi* Timmermann, 1961; Clay & Moreby 1967: 160, figs 134, 142.

*Bedfordiella unica* Thompson, 1937; Pilgrim & Palma 1982: 8.

*Bedfordiella unica* Thompson, 1937; Palma & Pilgrim 1983: 145, figs 1a,b.

*Bedfordiella unica* Thompson, 1937; Murray *et al.* 1990: 1369.

*Bedfordiella unica* Thompson, 1937; Palma 2010: 408.

Holotype ♀ in NHML. Holotype ♂ of *Bedfordiella simsi* in NHML (Palma & Pilgrim 1983: 146).

Type host: *Lugensa brevisrostris* (Lesson, 1833).

New Zealand host: *Lugensa brevisrostris* (Lesson, 1833).

Other hosts: None.

New Zealand localities: ND, AK, BP, WI, WN, WD, SL, SI.

Geographic distribution: Southern Oceans; Atlantic, Indian and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Palma & Pilgrim (1983); Murray *et al.* (1990: 1369); Page *et al.* (2004: 639, 650); Palma (2010).

Other significant references: Séguy (1953); Timmermann (1961c; 1965); Clay & Moreby (1967); Ledger (1980: 114, fig. 135); Green & Palma (1991: 11, 26); Palma (1996b: 160); Furness & Palma (1992: 35, 39); Price *et al.* (2003: 152); Hänel & Palma (2007: 112, 123, 130).

Remarks: The two sexes of this louse were described as two different species, and later synonymised by Palma & Pilgrim (1983: 145). Timmermann (1961c) described the male as *Bedfordiella simsi*, unaware that Séguy (1953: 563, figs 7–8) had already recorded, described and figured the main differences between the male and the female of *Bedfordiella unica*.

**Genus *Brueelia* Kéler, 1936**

*Brüelia* Kéler, 1936. *Arb. Morph. tax. Entomol. Berlin-Dahlem* 3: 257. Type species: *Brüelia rossittensis* Kéler, 1936 = *Brueelia brachythorax* (Giebel, 1874) (by original designation).

*Allobrüelia* Eichler, 1951a. *Bedeutung der Vogelwelt in Forschung und Praxis*, Berlin: 36. Type species: *Allobrüelia amsel* Eichler, 1951a = *Brueelia amsel* (Eichler, 1951) (by original designation).

*Turdinirmus* Eichler, 1951a. *Bedeutung der Vogelwelt in Forschung und Praxis*, Berlin: 41. Type species: *Philoapterus (Nirmus) merulensis* Denny, 1842 = *Brueelia merulensis* (Denny, 1842) (by original designation).

*Brueelia* Kéler, 1936; Ansari 1956: 102. Invalid emendation (see Palma 1996b: 162).

*Brueelia* Kéler, 1936; Złotorzycka 1964a: 252. Emendation.

***Brueelia amsel* (Eichler, 1951)**

*Allobrüelia amsel* Eichler, 1951a: 37, fig. 8.

*Allobrüelia amsel* Eichler, 1952b: 74, fig. 1.

*Brüelia amsel* (Eichler, 1951); Hopkins & Clay 1953: 435.

*Brueelia amsel* (Eichler, 1951); Ansari 1956: 135, figs 86–92.

*Allobrueelia amsel* Eichler, 1952; Złotorzycka 1964a: 265, fig. 7b.

*Brueelia* sp.; Pilgrim & Palma 1982: 26.

*Brueelia amsel* (Eichler, 1951); Palma 1999: 382.

*Brueelia amsel* (Eichler, 1951); Price *et al.* 2003: 153.

*Brueelia amsel* (Eichler, 1951); Murray *et al.* 2006b: 1959.

*Brueelia amsel* (Eichler, 1951); Palma 2010: 408.

Holotype ♂ in ZMHU (Jürgen Deckert pers. comm. November 2013).

Type host: *Turdus merula merula* Linnaeus, 1758.

New Zealand host: *Turdus merula merula* Linnaeus, 1758.

Other hosts: None

New Zealand locality: KE.

Geographic distribution: Eurasia; New Zealand.

New Zealand references: Pilgrim & Palma (1982); Palma (1999); Paterson *et al.* (1999: 220); Murray *et al.* (2006b); Palma (2010).

Other significant references: Ansari (1956); Złotorzycka (1964a); Baum (1968: 143, fig. 8); Price *et al.* (2003); Martín-Mateo (2009: 198).

Remarks: *Brueelia amsel* was introduced to New Zealand with blackbirds by human agency (Checklist Committee 2010: 313). Eichler (1951a; 1952b) described “*Allobrüelia amsel*” as a new species twice.

***Brueelia breueri* Balát, 1955****New Record**

*Brüelia breueri* Balát, 1955: 505, pl. 1: fig. 2, pl. 2: fig. 2.

*Brueelia breueri* Balát, 1955; Złotorzycka 1964a: 255, fig. 4c.

*Brueelia* sp.; Pilgrim & Palma 1982: 27.

*Brueelia breueri* Balát, 1955; Price *et al.* 2003: 153.

*Brueelia* sp.; Murray *et al.* 2006b: 1958.

*Brueelia* sp.; Palma 2010: 408.

Syntypes ♂♀ in Moravian Museum, Brno, Czech Republic (Oldřich Sychra pers. comm. December 2014).

Type host: *Carduelis chloris chloris* (Linnaeus, 1758).

New Zealand host: *Carduelis chloris* (Linnaeus, 1758).

Other hosts: None.

New Zealand localities: WN, MC, SL.

Geographic distribution: Eurasia; New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006b); Palma (2010).

Other significant references: Złotorzycka (1964a); Price *et al.* (2003).

Material examined and repository: 22♂, 34♀ (7 samples, MONZ).



Remarks: This is the first record of *Brueelia breueri* for New Zealand because the New Zealand references cited above reported this louse as “*Brueelia* sp.” only. *Brueelia breueri* was introduced to New Zealand with European greenfinches by human agency (Checklist Committee 2010: 320).

***Brueelia callaeincola* Valim & Palma, 2015**

*Brueelia* sp.; Pilgrim & Palma 1982: 28.

*Brueelia* sp.; Murray *et al.* 2006b: 1958.

*Brueelia* sp.; Palma 2010: 408.

*Brueelia* sp.; Buckley *et al.* 2012: App. 2.

*Brueelia callaeincola* Valim & Palma, 2015: 490, figs 4, 5, 6d, 7c,d.

Holotype ♂ in MONZ.

Type host: *Philesturnus carunculatus* (J.F. Gmelin, 1788).

New Zealand hosts: *Callaeas wilsoni* (Bonaparte, 1851); *Callaeas cinerea* (J.F. Gmelin, 1788); *Philesturnus rufusater* (Lesson, 1828); *Philesturnus carunculatus* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: ND, CL, AK, FD.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006b); Palma (2010); Buckley *et al.* (2012); Valim & Palma (2015).

Other significant references: none.

Remarks: *Brueelia callaeincola* is an endemic and vulnerable species (Buckley *et al.* 2012), exclusively parasitic on members of the endemic bird family Callaeidae (Checklist Committee 2010: 280).

***Brueelia cyclothorax* (Burmeister, 1838)**

*Nirmus cyclothorax* Burmeister, 1838a: 429.

*Degeeriella cyclothorax* Nitzsch in Burmeister, 1838 [sic]; Harrison 1916: 111.

*Bruëlia cyclothorax* (Burmeister, 1838); Hopkins & Clay 1952: 55.

*Brueelia cyclothorax* (Burmeister, 1838); Złotorzycka 1964a: 255, fig. 4d.

*Brueelia cyclothorax* (Burmeister, 1838); Horning *et al.* 1980: 5, 12.

*Brueelia cyclothorax* (Burmeister, 1838); Pilgrim & Palma 1982: 27.

*Brueelia cyclothorax* (Burmeister, 1838); Murray *et al.* 2006b: 1958.

*Brueelia cyclothorax* (Burmeister, 1838); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Passer domesticus domesticus* (Linnaeus, 1758).

New Zealand host: *Passer domesticus domesticus* (Linnaeus, 1758).

Other host: *Passer montanus* (Linnaeus, 1758).

New Zealand localities: HB, WN, NC, MC, SC, WD, SN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Galloway (2005: 16); Murray *et al.* (2006b); Palma (2010).

Other significant references: Złotorzycka (1964a); Price *et al.* (2003: 154); Palma & Jensen (2005: 54, 68); Adam (2007: 182).

Remarks: *Brueelia cyclothorax* was introduced to New Zealand with house sparrows by human agency (Checklist Committee 2010: 316).

***Brueelia delicata* (Nitzsch [*in* Giebel], 1866)**

*Nirmus delicatus* Nitzsch [*in* Giebel], 1866: 368.

*Degeeriella delicata* Nitzsch [*in* Giebel], 1866 [sic]; Harrison 1916: 111.

*Bruëlia delicata* (Nitzsch [*in* Giebel], 1866); Hopkins & Clay 1952: 55.

*Brueelia delicata* (Nitzsch, 1866) [sic]; Złotorzycka 1964a: 256.

*Brueelia delicata* (Nitzsch, 1866) [sic]; Pilgrim & Palma 1982: 27.

*Brueelia delicata* (Nitzsch [*in* Giebel], 1866); Price *et al.* 2003: 154.

*Brueelia delicata* (Nitzsch [*in* Giebel], 1866); Murray *et al.* 2006b: 1959.

*Brueelia delicata* (Nitzsch [*in* Giebel], 1866); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Emberiza citrinella* Linnaeus, 1758.

New Zealand host: *Emberiza citrinella* Linnaeus, 1758.

Other hosts: *Emberiza leucocephalos* S.G. Gmelin, 1771; *Emberiza rustica* Pallas, 1776.

New Zealand localities: WN, MB, MC, KE.

Geographic distribution: Eurasia; New Zealand.

New Zealand references: Pilgrim & Palma (1982); Galloway (2005: 17); Murray *et al.* (2006b); Palma (2010).

Other significant references: Złotorzycka (1964a); Mey (1982: 171, fig. 18); Soler-Cruz *et al.* (1982: 256, figs 1, 5b,e,g); Price *et al.* (2003); Martín-Mateo (2009: 192).

Remarks: *Brueelia delicata* was introduced to New Zealand with yellowhammers by human agency (Checklist Committee 2010: 322).

### ***Brueelia densilimba* (Nitzsch [*in* Giebel], 1866)**

### **New Record**

*Nirmus densilimbus* Nitzsch [*in* Giebel], 1866: 368.

*Degeeriella densilimba* Nitzsch [*in* Giebel], 1866 [sic]; Harrison 1916: 111

*Bruëlia densilimba* (Nitzsch [*in* Giebel], 1866); Hopkins & Clay 1952: 55.

*Brueelia* sp.; Pilgrim & Palma 1982: 27.

*Brueelia densilimba* (Nitzsch [*in* Giebel], 1866); Rékási & Kiss 1984: 107, 111.

*Brueelia densilimba* (Nitzsch [*in* Giebel], 1866); Price *et al.* 2003: 154.

*Brueelia* sp.; Murray *et al.* 2006b: 1959.

*Brueelia* sp.; Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Carduelis carduelis* (Linnaeus, 1758).

New Zealand host: *Carduelis carduelis britannica* (Hartert, 1903).

Other hosts: None.

New Zealand locality: WD.

Geographic distribution: Eurasia; New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006b); Palma (2010).

Other significant references: Soler-Cruz & Guevara-Benítez (1981: 275, figs 1–2); Price *et al.* (2003); Martín-Mateo (2009: 191).

Material examined and repository: 2♂, 10♀ (1 sample, MONZ).

Remarks: This is the first record of *Brueelia densilimba* for New Zealand because the New Zealand references cited above reported this louse as “*Brueelia* sp.” only. *Brueelia densilimba* was introduced to New Zealand with European goldfinches by human agency (Checklist Committee 2010: 320).

### ***Brueelia merulensis* (Denny, 1842)**

*Philopterus (Nirmus) merulensis* Denny, 1842: 51, 128, pl. 7: fig. 1.

*Degeeriella merulensis* Denny, 1842 [sic]; Harrison 1916: 118.

*Turdinirmus merulensis* Denny [sic]; Eichler 1952b: 78, fig. 3.

*Bruëlia merulensis* (Denny, 1842); Hopkins & Clay 1952: 58.

*Brueelia merulensis* (Denny, 1842); Ansari 1956: 119, figs 35–42.

*Turdinirmus merulensis* (Denny, 1842); Złotorzycka 1964a: 267, figs 8a,b.

*Brueelia merulensis* (Denny, 1842); Tendeiro 1965c: 73, figs 48–51.

*Brueelia merulensis* (Denny, 1842); Watt 1971: 235, 244, fig. 8.

*Brueelia merulensis* (Denny, 1842); Horning *et al.* 1980: 5, 12.

*Brueelia merulensis* (Denny, 1842); Pilgrim & Palma 1982: 26.

*Brueelia merulensis* (Denny, 1842); Murray *et al.* 2006b: 1959.

*Brueelia merulensis* (Denny, 1842); Palma 2010: 408.

Syntypes ♂♀ in NHML (Vincent S. Smith pers. comm. December 2014).

Type host: *Turdus merula* Linnaeus, 1758.

New Zealand host: *Turdus merula merula* Linnaeus, 1758.

Other hosts: *Turdus merula syriacus* Hemprich & Ehrenberg, 1833; *Turdus philomelos* Brehm, 1831.

New Zealand localities: WN, KE, CH, SN.

Geographic distribution: Eurasia; New Zealand.

New Zealand references: Watt (1971); Wise (1977: 59); Horning *et al.* (1980); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Murray *et al.* (2006b); Palma (2010).

Other significant references: Clay (1951b: 188, figs 22–23, 25); Ansari (1956); Złotorzycka (1964a); Tendeiro (1965c); Baum (1968: 143, fig. 9); Rékási & Kiss (1984: 107, 111); Soler-Cruz *et al.* (1985: 108, fig. 1); Price *et al.* (2003: 156); Palma & Jensen (2005: 54, 67); Adam (2007: 182); Martín-Mateo (2009: 200); Bartlow *et al.* (2016: 222).

Remarks: *Brueelia merulensis* was introduced to New Zealand with blackbirds by human agency (Checklist Committee 2010: 313).

### ***Brueelia nebulosa* (Burmeister, 1838)**

*Nirmus nebulosus* Burmeister, 1838a: 429.

*Degeeriella nebulosa* Burmeister [sic]; Johnston & Harrison 1912: 368.

*Degeeriella nebulosa* (Burmeister, 1838); Thompson 1939: 120.

*Bruëlia nebulosa* (Burmeister, 1838); Hopkins & Clay 1952: 59.

*Brueelia nebulosa* (Burmeister, 1838); Watt 1971: 235, 244, fig. 7.

*Brueelia nebulosa* (Burmeister, 1838); Pilgrim & Palma 1982: 27.

*Brueelia nebulosa* (Burmeister, 1838); Murray *et al.* 2006b: 1960.

*Brueelia nebulosa* (Burmeister, 1838); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1), Palma & Pilgrim (1984: 150) and Palma (1996b: 162).

Type host: *Sturnus vulgaris* Linnaeus, 1758.

New Zealand host: *Sturnus vulgaris vulgaris* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: HB, WI, WN, KA, NC, MC, SC, WD, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Johnston & Harrison (1912); Watt (1971); Wise (1977: 59); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Murray *et al.* (2006b); Palma (2010).

Other significant references: Thompson (1939); Kettle (1983: 403); Green & Palma (1991: 11, 41); Butler & O'Connor (1994: 453); Palma (1996b: 162); Price *et al.* (2003: 157); Palma & Jensen (2005: 54, 68); Adam (2007: 183); Martín-Mateo (2009: 189).

Remarks: *Brueelia nebulosa* was introduced to New Zealand with starlings by human agency (Checklist Committee 2010: 315).

### ***Brueelia parviguttata* (Blagoveshtchensky, 1940)**

**New record**

*Degeeriella parviguttata* Blagoveshtchensky, 1940: 62, 86, fig. 17.

*Bruëlia parviguttata* (Blagoveshtchensky, 1940); Hopkins & Clay 1952: 60.

*Brueelia* sp.; Pilgrim & Palma 1982: 25.

*Brueelia parviguttata* (Blagoveshtchensky, 1940); Price *et al.* 2003: 157.

*Brueelia* sp.; Murray *et al.* 2006b: 1958.

Syntypes ♂♀, repository unknown.

Type host: *Alauda arvensis cantarella* Bonaparte, 1850.

New Zealand host: *Alauda arvensis* Linnaeus, 1758.

Other host: *Galerida cristata* (Linnaeus, 1758).

New Zealand locality: MC.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006b).

Other significant references: Balát (1955: 502); Soler-Cruz *et al.* (1979: 165); Price *et al.* (2003).

Material examined and repository: 2♂, 11♀ (2 samples, MONZ).

Remarks: This is the first record of *Brueelia parviguttata* for New Zealand, because the New Zealand references cited above reported this louse as “*Brueelia* sp.” only. *Brueelia parviguttata* was introduced to New Zealand with Eurasian skylarks by human agency (Checklist Committee 2010: 306).

### ***Brueelia semiannulata* (Piaget, 1883)**

Figs 76–77

*Nirmus semiannulatus* Piaget, 1883: 156, pl. 9: fig. 3.

*Degeeriella semiannulata* Piaget, 1883 [sic]; Harrison 1916: 123.

*Bruëlia semiannulata* (Piaget, 1883); Hopkins & Clay 1952: 61.

*Brueelia elegans* Ansari, 1957: 122, figs 1–25.

*Brueelia semiannulata* (Piaget, 1883); Pilgrim & Palma 1982: 28.

*Brueelia semiannulata* (Piaget, 1883); Murray *et al.* 2006b: 1957.

*Brueelia semiannulata* (Piaget, 1883); Palma 2010: 408.

Syntypes ♀♀ in NHML (Palma 1996b: 162). Holotype ♂ of *Brueelia elegans* in NHML (Palma 1996b: 162).

Type host: *Gymnorhina tibicen leuconota* Gould, 1844.

New Zealand host: *Gymnorhina tibicen* (Latham, 1802).

Other host: *Gymnorhina tibicen hypoleuca* (Gould, 1837).

New Zealand localities: ND, CL, HB, WN, NC, MC, SC.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999: 221); Galloway (2005: 17); Murray *et al.* (2006b); Palma (2010).

Other significant references: Ansari (1957); Hughes (1984a: 459); Hughes (1984b: 467); Palma (1996b: 162); Price *et al.* (2003: 158); Toon & Hughes (2008: 128).

Remarks: *Brueelia semiannulata* was introduced to New Zealand with Australian magpies by human agency (Checklist Committee 2010: 296).

### ***Brueelia turdinulae* Ansari, 1956**

*Brueelia turdinulae* Ansari, 1956: 126, figs 59–65.

*Brueelia turdinulae eternitatus* Ansari, 1956: 129, figs 66–71.

*Brueelia* sp.; Watt 1971: 235, 244.

*Brueelia turdinulae eternitatus* Ansari, 1956; Jiménez-González *et al.* 1980: 212, figs 15–20.

*Brueelia* sp.; Pilgrim & Palma 1982: 26.

*Brueelia turdinulae* Ansari, 1956; Palma 1999: 382.

*Brueelia turdinulae* Ansari, 1956; Murray *et al.* 2006b: 1959.

*Brueelia turdinulae* Ansari, 1956; Palma 2010: 408.

Holotype ♂ in NHML. Holotype ♂ of *Brueelia turdinulae eternitatus* in NHML.

Type host: *Turdus philomelos philomelos* Brehm, 1831.

New Zealand host: *Turdus philomelos* Brehm, 1831.

Other host: *Turdus philomelos clarkei* Hartert, 1909.

New Zealand localities: WD, KE.

Geographic distribution: Eurasia; Africa; New Zealand.

New Zealand references: Watt (1971); Pilgrim & Palma (1982); Palma (1999); Paterson *et al.* (1999: 220); Murray *et al.* (2006b); Palma (2010).

Other significant references: Jiménez-González *et al.* (1980); Price *et al.* (2003: 159); Palma & Jensen (2005: 54, 67); Adam (2007: 183); Martín-Mateo (2009: 197); Bartlow *et al.* (2016: 222).

Remarks: *Brueelia turdinulae* was introduced to New Zealand with song thrushes by human agency (Checklist Committee 2010: 313).

**Brueelia species**

*Brueelia* sp.; Tennyson 1997: 268.

*Brueelia* sp.; Palma 1999: 382, 383 note 6.

*Brueelia* sp.; Murray *et al.* 2002: 1216.

*Brueelia* sp.; Murray *et al.* 2006b: 1956, 1958.

*Brueelia* sp.; Palma 2010: 408.

New Zealand hosts: *Turnagra capensis capensis* (Sparrman, 1787); *Mohoua albicilla* (Lesson, 1830); *Monarcha melanopsis* (Vieillot, 1818).

New Zealand localities: ND, CL, TK, South Island.

Geographic distribution: Australasia.

New Zealand references: Tennyson (1997); Paterson *et al.* (1999: 223); Palma (1999); Murray *et al.* (2002; 2006b); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: None.

Remarks: The samples of *Brueelia* from the three hosts listed above may represent different species which are not identifiable at present because the available material contain females only (voucher specimens in MONZ).

**Genus *Campanulotes* Kéler, 1939**

*Campanulotes* Kéler, 1939. *Nova Acta Leop.-Carol. (N.F.)* 8: 157. Type species: *Goniocotes compar* "(Nitzsch)" = *Campanulotes bidentatus compar* (Burmeister, 1838) (by original designation).

***Campanulotes bidentatus compar* (Burmeister, 1838)**

Figs 78–79

*Goniocotes compar* Burmeister, 1838a: 431.

*Philopterus (Goniocotes) compar* Burmeister, 1838 [sic]; Denny 1842: 152, pl. 13: fig. 2.

*Goniocotes compar* Nitzsch [in Giebel], 1874 [sic]; Kellogg 1908: 32, fig. 4.

*Campanulotes compar* (Nitzsch, 1818) [sic]; Kéler 1939: 157, figs 89–91.

*Campanulotes compar* (Burmeister, 1838); Hopkins & Clay 1952: 64.

"*Campanulotes bidentatus*" Pilgrim 1970: 76 (not *Pediculus bidentatus* Scopoli, 1763).

"*Goniocotes gallinae*" Heath *et al.*, 1971: 91 (not *Ricinus gallinae* De Geer, 1778b).

*Campanulotes bidentatus compar* (Burmeister, 1838); Pilgrim 1976: 162, fig. 2.

*Campanulotes bidentatus compar* (Burmeister, 1838); Wise 1977: 59.

*Campanulotes bidentatus compar* (Burmeister, 1838); Pilgrim & Palma 1982: 23.

*Campanulotes compar* (Burmeister, 1838); Price *et al.* 2003: 160.

*Campanulotes bidentatus compar* (Burmeister, 1838); Murray *et al.* 2006a: 1965.

*Campanulotes bidentatus compar* (Burmeister, 1838); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1), Palma & Pilgrim (1984: 150) and Palma (1996b: 163).

Type host: *Columba livia* J.F. Gmelin, 1789.

New Zealand host: *Columba livia* J.F. Gmelin, 1789.

Other hosts: None.

New Zealand localities: WN, SD, MB, NC, MC, SC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim (1970); Heath *et al.* (1971); Pilgrim (1976); Wise (1977); Tendeiro (1978: 117); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Murray *et al.* (2006a); Palma (2010).

Other significant references: Emerson (1957: 64); Kéler (1939); Nelson & Murray (1971: 22, 25); Rudolph (1983: 16); Forrester *et al.* (1995: 32); Palma (1996b: 163); Price *et al.* (2003); Palma & Jensen (2005: 55, 66); Martín-Mateo (2009: 30, fig. 6); Galloway & Lamb (2014: 445; 2015: 715).

Remarks: *Campanulotes bidentatus compar* was introduced to New Zealand and other countries with rock pigeons by human agency (Checklist Committee 2010: 245). In agreement with Pilgrim & Palma (1982: 23) but contrary to Price *et al.* (2003: 160), I regard this louse taxon as a subspecies of *Campanulotes bidentatus* (Scopoli, 1763).

**Genus *Capraiella* Conci, 1941**

*Capraiella* Conci, 1941. *Boll. Soc. Entomol. Italiana* 73: 126. Type species: “*Degeeriella subcuspidata* Nitzsch in Burm. 1838” = *Capraiella subcuspidata* (Burmeister, 1838a) (by original designation).

***Capraiella* species**

Fig. 80

*Capraiella* sp.; Pilgrim & Palma 1982: 25.

*Capraiella* sp.; Murray *et al.* 1999: 1242.

*Capraiella* sp.; Catanach & Johnson 2015: 839.

New Zealand host: *Eurystomus orientalis pacificus* (Latham, 1802).

New Zealand locality: NN.

Geographic distribution: Indonesia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1999).

Other significant references: Hopkins & Clay (1952: 65); Ansari (1955a: 46); Tandan (1955: 425, figs 13–16); Ledger (1980: 170, fig. 178); Price *et al.* (2003: 160); Catanach & Johnson (2015: 839).

Remarks: *Eurystomus orientalis pacificus* is a frequent straggler to New Zealand (Checklist Committee 2010: 274). The single available New Zealand record of *Capraiella* could not be identified to species because the sample contains females only (voucher specimens in MONZ). Furthermore, the Australian record in Catanach & Johnson (2015: 839) also refers to females only (Therese A. Catanach pers. comm. September 2015).

**Genus *Carduiceps* Clay & Meinertzhagen, 1939**

*Carduiceps* Clay & Meinertzhagen, 1939b. *Ann. Mag. Nat. Hist. (Ser. 11) 4*: 451. Type species: *Degeeriella complexiva* (Kellogg & Chapman, 1899) = *Carduiceps complexivus* (Kellogg & Chapman, 1899) (by original designation).

***Carduiceps cingulatus cingulatus* (Denny, 1842)**

Figs 81–82

*Philopterus (Nirmus) cingulatus* Denny, 1842: 54, 146, pl. 11: fig. 3.

*Degeeriella cingulata* Nitzsch in Denny, 1842 [sic]; Harrison 1916: 110.

*Carduiceps cingulatus* (Denny, 1842); Hopkins & Clay 1952: 65.

*Carduiceps cingulatus cingulatus* (Denny, 1842); Emerson 1972a: 37.

*Carduiceps cingulatus* (Denny, 1842); Wise 1977: 59.

*Carduiceps cingulatus* (Denny, 1842); Pilgrim & Palma 1982: 20.

*Carduiceps cingulatus* (Denny, 1842); Murray *et al.* 2006a: 1964. In part.

*Carduiceps cingulatus* (Denny, 1842); Palma 2010: 408. In part.

Neotype ♂ in NHML (Timmermann 1955: 529).

Type host: *Limosa limosa limosa* (Linnaeus, 1758).

New Zealand host: *Limosa limosa melanuroides* Gould, 1846.

Other hosts: *Limnodromus griseus* (J.F. Gmelin, 1789); *Limnodromus scolopaceus* (Say, 1823).

New Zealand locality: AU.

Geographic distribution: Eurasia; Americas; Australasia.

New Zealand references: Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1954a: 40, figs 2–3); Timmermann (1955: 527); Emerson (1972a); Hunter & Colwell (1994: 402); Price *et al.* (2003: 161).

Remarks: In agreement with Emerson (1972a: 37) and contrary to Price *et al.* (2003: 161), I regard this louse taxon as a subspecies. The Asiatic black-tailed godwit is an uncommon annual visitor to New Zealand (Checklist Committee 2010: 203).

***Carduiceps cingulatus lapponicus* Emerson, 1953**

*Carduiceps lapponicus* Emerson, 1953: 209, fig. 2.

*Carduiceps cingulatus* (Denny, 1842); Timmermann 1955: 529. In part.

- Carduiceps cingulatus lapponicus* Emerson, 1953; Watt 1971: 236, 243.  
*Carduiceps cingulatus lapponicus* Emerson, 1953; Emerson 1972a: 37.  
*Carduiceps cingulatus lapponicus* Emerson, 1953; Wise 1977: 59.  
*Carduiceps cingulatus* (Denny, 1842) *s. l.*; Pilgrim & Palma 1982: 21.  
*Carduiceps lapponicus* Emerson, 1953; Price *et al.* 2003: 161.  
*Carduiceps cingulatus* (Denny, 1842); Murray *et al.* 2006a: 1964. In part.  
*Carduiceps cingulatus* (Denny, 1842); Palma 2010: 408. In part.

Holotype ♂ in USNM.

Type host: *Limosa lapponica lapponica* (Linnaeus, 1758).

New Zealand host: *Limosa lapponica baueri* Naumann, 1836.

Other hosts: None.

New Zealand localities: SD, MB, NC, MC, SC, WD, SL, KE, SN, CA.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Emerson (1972a); Price *et al.* (2003).

Remarks: In agreement with Emerson (1972a: 37) and Wise (1977: 59), and contrary to Price *et al.* (2003: 161), I regard this louse taxon as a subspecies of *Carduiceps cingulatus*. The eastern bar-tailed godwit is the most numerous wader that visits New Zealand every year (Checklist Committee 2010: 203).

### ***Carduiceps zonarius* (Nitzsch [*in* Giebel], 1866)**

- Nirmus zonarius* Nitzsch [*in* Giebel], 1866: 374.  
*Degeeriella zonaria* Nitzsch [*in* Giebel], 1866 [sic]; Harrison 1916: 126.  
*Carduiceps zonarius* (Nitzsch, 1866) [sic]; Hopkins & Clay 1952: 66.  
*Carduiceps zonarius* (Nitzsch [*in* Giebel], 1866); Timmermann 1954a: 44, fig. 6a.  
*Carduiceps zonarius* (Nitzsch, 1866) [sic] *s. l.*; Pilgrim & Palma 1982: 21.  
*Carduiceps zonarius* (Nitzsch [*in* Giebel], 1866); Palma 1999: 380.  
*Carduiceps zonarius* (Nitzsch [*in* Giebel], 1866); Murray *et al.* 2006a: 1964.  
*Carduiceps zonarius* (Nitzsch [*in* Giebel], 1866); Palma 2010: 408.

Neotype ♂ in NHML (Timmermann 1954a: 45; Palma 1996b: 164).

Type host: *Calidris minuta* (Leisler, 1812) (see Timmermann 1954a: 44).

New Zealand hosts: *Calidris canutus rogersi* Mathews, 1913; *Calidris acuminata* (Horsfield, 1821).

Other hosts: *Eurynorhynchus pygmaeus* (Linnaeus, 1758); *Tryngites subruficollis* (Vieillot, 1819) and 12 other species of *Calidris* (see Price *et al.* 2003: 161).

New Zealand localities: WN, NC, MC, SC, KE, CA.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Palma (1999); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1954a); Hackman & Nyholm (1968: 79); Amerson & Emerson (1971: 15, 22); Green & Palma (1991: 12, 33); Hunter & Colwell (1994: 402); Forrester *et al.* (1995: 27); Palma (1996b: 164); Price *et al.* (2003: 161); Palma & Jensen (2005: 55, 63).

Remarks: Pilgrim & Palma (1982: 21) regarded the population of *Carduiceps zonarius* from *Calidris canutus* as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted.

### **Genus *Chelopistes* Kéler, 1939**

- Chelopistes* Kéler, 1939. *Nova Acta Leop.-Carol. (N.F.)* 8: 180. Type species: *Rhopaloceras styliifer* (Nitzsch, 1818) = *Chelopistes meleagridis* (Linnaeus, 1758) (by original designation).  
*Virgula* Clay, 1941. *Parasitology* 33: 119. Type species: *Goniodes meleagridis* (Linnaeus, 1758) = *Chelopistes meleagridis* (Linnaeus, 1758) (by original designation).

***Chelopistes meleagridis* (Linnaeus, 1758)**

Figs 83–84

*Pediculus meleagridis* Linnaeus, 1758: 613.*Ricinus melagris* (Linnaeus, 1758); Latreille 1804: 108.*Philopterus (Goniodes) stylifer* Nitzsch, 1818: 294. *Nomen novum* for *Pediculus meleagridis* Schrank, 1781.*Rhopaloceras styliferum* Nitzsch [sic]; Taschenberg 1882: 47. Emendation.*Rhopaloceras stylifer* (Nitzsch, 1818); Kéler 1939: 180.*Chelopistes meleagridis* (Linnaeus, 1758); Kéler 1939: 181, figs 103–104.*Virgula meleagridis* (Linnaeus, 1758); Clay 1941: 120, figs 1–4.*Goniodes meleagridis* (Linnaeus, 1758); Séguy 1944: 170, fig. 243.*Chelopistes meleagridis* (Linnaeus, 1758); Hopkins & Clay 1952: 69.*Chelopistes meleagridis* (Linnaeus, 1758); Pilgrim & Palma 1982: 18.*Chelopistes meleagridis* (Linnaeus, 1758); Murray *et al.* 1993: 960.*Chelopistes meleagridis* (Linnaeus, 1758); Palma 2010: 408.

Neotype ♂ in NHML (Clay 1941: 124).

Type host: *Meleagris gallopavo* Linnaeus, 1758.New Zealand host: *Meleagris gallopavo gallopavo* Linnaeus, 1758.Other host: *Meleagris ocellata* Cuvier, 1820.

New Zealand localities: WO, HB, WA, WN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 220); Palma (2010).Other significant references: Kéler (1939); Clay (1941); Séguy (1944); Clay & Hopkins (1950: 261); Emerson (1962: 198, figs 4–6); Price (1987: 220, fig. 22.25); Butler & O'Connor (1994: 453); Forrester *et al.* (1995: 21); Palma (1996b: 164); Price *et al.* (2003: 162); Martín-Mateo (2009: 40, figs 5d, 10); Maturano & Daemon (2014).Remarks: *Chelopistes meleagridis* was introduced to New Zealand and other countries with turkeys by human agency (Checklist Committee 2010: 29).**Genus *Coloceras* Taschenberg, 1882***Coloceras* Taschenberg, 1882. *Nova Acta Leop.-Carol.* 44: 42. Type species: *Goniodes damicornis* Nitzsch, 1866 = *Coloceras damicorne* (Nitzsch, 1866) (by subsequent designation).*Nitzschiella* Kéler, 1939. *Nova Acta Leop.-Carol. (N.F.)* 8: 67. Type species: *Goniocotes menadensis* Piaget, 1880 = *Coloceras menadense* (Piaget, 1880) (by original designation).*Patellinirmus* Tendeiro, 1972b. *Rev. Ciênc. Vet. (Lourenço Marques)* (Series A) 5: 63. Type species: *Patellinirmus novaeseelandiae* Tendeiro, 1972 = *Coloceras novaeseelandiae* (Tendeiro, 1972) (by original designation).***Coloceras chinense* (Kellogg & Chapman, 1902)****New Record***Goniocotes chinensis* Kellogg & Chapman, 1902: 160, pl. 13: fig. 5.*Coloceras chinense* (Kellogg & Chapman, 1902); Hopkins & Clay 1952: 74.*Coloceras chinense* (Kellogg & Chapman, 1902); Palma 1996b: 166.*Coloceras chinense* (Kellogg & Chapman, 1902); Price *et al.* 2003: 164.

Holotype ♀ in EMEC (Palma 1996b: 166).

Type host: *Streptopelia chinensis chinensis* (Scopoli, 1786).New Zealand host: *Streptopelia chinensis tigrina* (Temminck, 1810).Other hosts: *Gallicolumba luzonica* (Scopoli, 1786); *Leptotila rufaxilla* (Richard & Bernard, 1792); *Macropygia unchall* (Wagler, 1827); 14 other species of *Streptopelia*, and five species of *Turtur* (see Price *et al.* 2003: 164).

New Zealand locality: AK.

Geographic distribution: Eurasia; Africa; South America; Pacific Ocean.

New Zealand reference: This paper.

Other significant references: Tendeiro (1973: 353, figs 34–36, photos 91–101, 188–189, 221–222); Green & Palma (1991: 12, 34); Palma (1996b); Price *et al.* (2003).

Material examined and repository: 1♂, 2♀, 8 nymphs (1 sample, MONZ).



Remarks: *Coloceras chinense* is a new louse species for New Zealand, introduced to this country with spotted doves by human agency (Checklist Committee 2010: 245).

#### ***Coloceras harrisoni* (Tendeiro, 1972)**

*Patellinirmus harrisoni* Tendeiro, 1972b: 74, fig. 3, photos 10–17, 24–25.

*Patellinirmus harrisoni* Tendeiro, 1972; Pilgrim & Palma 1982: 23.

*Coloceras harrisoni* (Tendeiro, 1972); Price *et al.* 2003: 164.

*Coloceras harrisoni* (Tendeiro, 1972); Murray *et al.* 2006a: 1966.

*Coloceras harrisoni* (Tendeiro, 1972); Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Hemiphaga chathamensis* (Rothschild, 1891).

New Zealand host: *Hemiphaga chathamensis* (Rothschild, 1891).

Other hosts: None.

New Zealand localities: CH.

Geographic distribution: Chatham Islands, New Zealand.

New Zealand references: Tendeiro (1972b); Pilgrim & Palma (1982); Paterson *et al.* (1999: 223); Murray *et al.* (2006a); Palma (2010); Buckley *et al.* (2012: 137, App. 2).

Other significant references: Price *et al.* (2003).

Remarks: *Coloceras harrisoni* is an endemic and critically threatened species (Buckley *et al.* 2012), exclusively parasitic on Chatham Island pigeons.

#### ***Coloceras hemiphagae* (Tendeiro, 1972)**

*Nitzschiella hemiphagae* Tendeiro, 1972a: 2, figs 1–3.

*Coloceras hemiphagae* (Tendeiro, 1972); Palma 1996b: 166.

*Coloceras hemiphagae* (Tendeiro, 1972); Price *et al.* 2003: 164.

*Nitzschiella hemiphagae* Tendeiro, 1972; Mey 2005: 213.

*Coloceras hemiphagae* (Tendeiro, 1972); Murray *et al.* 2006a: 1966.

Holotype ♀ in NHML.

Type host: *Hemiphaga spadicea* (Latham, 1802).

New Zealand host: *Hemiphaga spadicea* (Latham, 1802).

Other hosts: None

New Zealand localities: Norfolk Island.

Geographic distribution: Norfolk Island.

New Zealand references: Tendeiro (1972a); Palma (1996b); Mey (2005); Murray *et al.* (2006a); Rózsa & Vas (2015b: 108).

Other significant references: Price *et al.* (2003).

Remarks: *Coloceras hemiphagae* is known from the holotype only. Its host, *Hemiphaga spadicea*, was endemic to Norfolk Island, but it is now extinct (Checklist Committee 2010: 248). Considering that there is no other known host for *Coloceras hemiphagae*, this species is also regarded as extinct (Rózsa & Vas 2015b: 108).

#### ***Coloceras novaeseelandiae* (Tendeiro, 1972)**

Figs 85–86

*Patellinirmus novaeseelandiae* Tendeiro, 1972b: 63, figs 1–2, photos 1–9, 23.

*Patellinirmus novaeseelandiae* Tendeiro, 1972; Pilgrim & Palma 1982: 23.

*Coloceras novaeseelandiae* (Tendeiro, 1972); Price *et al.* 2003: 165.

*Coloceras novaeseelandiae* (Tendeiro, 1972); Murray *et al.* 2006a: 1966.

*Coloceras novaeseelandiae* (Tendeiro, 1972); Palma 2010: 408.

Holotype ♂ in CMNZ (Nicholls *et al.* 1998: 30).

Type host: *Hemiphaga novaeseelandiae* (J.F. Gmelin, 1789).

New Zealand host: *Hemiphaga novaeseelandiae* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: ND, CL, AK, HB, TK, WI, WA, WN, SD, MB, NN, NC, MC, SC, WD, CO, DN.

Geographic distribution: New Zealand.

New Zealand references: Tendeiro (1972b); Pilgrim & Palma (1982); Nicholls *et al.* (1998: 30); Paterson *et al.* (1999: 223); Murray *et al.* (2006a); Palma (2010).

Other significant references: Price *et al.* (2003: 165).

Remarks: *Coloceras novaeseelandiae* is an endemic species exclusively parasitic on the New Zealand pigeon, with a relatively high prevalence of infestation.

### ***Coloceras restinctum* (Tendeiro, 1972)**

*Patellinirmus restinctus* Tendeiro, 1972b: 81, fig. 4, photos 18–21, 26.

*Coloceras restinctum* (Tendeiro, 1972); Palma 1996b: 167.

*Coloceras restinctus* [sic] (Tendeiro, 1972); Price *et al.* 2003: 165.

*Patellinirmus restinctus* Tendeiro, 1972; Mey 2005: 213.

*Coloceras restinctum* (Tendeiro, 1972); Murray *et al.* 2006a: 1966.

Holotype ♀ in NHML.

Type host: *Hemiphaga spadicea* (Latham, 1802).

New Zealand host: *Hemiphaga spadicea* (Latham, 1802).

Other hosts: None

New Zealand localities: Norfolk Island.

Geographic distribution: Norfolk Island.

New Zealand references: Tendeiro (1972b); Palma (1996b); Mey (2005); Murray *et al.* (2006a); Rózsa & Vas (2015b: 108).

Other significant references: Price *et al.* (2003).

Remarks: *Coloceras restinctum* is known from the holotype only. Its host, *Hemiphaga spadicea*, was endemic to Norfolk Island, but it is now extinct (Checklist Committee 2010: 248). Considering that there is no other known host for *Coloceras restinctum*, this species is also regarded as extinct (Rózsa & Vas 2015b: 108).

### **Genus *Columbicola* Ewing, 1929**

*Columbicola* Ewing, 1929. *Manual External Parasites*: 112, 190. Type species: *Esthiopterum columbae* (Linnaeus, 1758) = *Columbicola columbae* (Linnaeus, 1758) (by original designation).

### ***Columbicola columbae columbae* (Linnaeus, 1758)**

Figs 87–88

“Pollino del piccion grosso” Redi, 1668: pl. 2.

*Pediculus columbae* Linnaeus, 1758: 614. *Nomen novum* for “Pollino del piccion grosso” Redi, 1668.

*Ricinus columbae* (Linnaeus, 1758); Latreille 1804: 110.

*Philopterus (Lipeurus) baculus* Nitzsch, 1818: 293.

*Philopterus (Lipeurus) baculus* (Nitzsch, 1818); Denny 1842: 172, pl. 14: fig. 3.

*Lipeurus columbae* (Linnaeus, 1758); Séguy 1924: 40.

*Esthiopterum columbae* Linnaeus, 1758 [sic]; Harrison 1916: 132.

*Columbicola columbae* (Linnaeus, 1758); Hopkins & Clay 1952: 86.

*Columbicola columbae columbae* (Linnaeus, 1758); Tendeiro 1965a: 77, figs 1–12, photos 1–6, 189–190.

*Columbicola columbae* (Linnaeus, 1758); Pilgrim 1976: 162, fig. 1.

*Columbicola columbae columbae* (Linnaeus, 1758); Wise 1977: 59.

*Columbicola columbae columbae* (Linnaeus, 1758); Pilgrim & Palma 1982: 23.

*Columbicola columbae* (Linnaeus, 1758); Price *et al.* 2003: 166.

*Columbicola columbae* (Linnaeus, 1758); Murray *et al.* 2006a: 1965.

*Columbicola columbae columbae* (Linnaeus, 1758); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1950: 265).

Type host: *Columba livia* J.F. Gmelin, 1789.

New Zealand hosts: *Columba livia* J.F. Gmelin, 1789; *Streptopelia chinensis tigrina* (Temminck, 1810).

Other hosts: *Columba eversmanni* Bonaparte, 1856; *Columba guinea* Linnaeus, 1758; *Columba oenas* Linnaeus, 1758.

New Zealand localities: AK, WN, NN, NC, MC, SC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim (1970: 76); Heath *et al.* (1971: 91); Pilgrim (1976); Wise (1977); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Murray *et al.* (2006a); Palma (2010).

Other significant references: Wigglesworth (1932: 365); Martin (1934: 6, figs 1–6); Clay & Hopkins (1950: 264, figs 57–59); Conci (1956: 47); Stenram (1956: 170, figs 1–7); Emerson (1957: 64); Tendeiro (1965a); Nelson & Murray (1971: 22, 25, fig. 4); Rudolph (1983: 16); Green & Palma (1991: 12, 34); Butler & O'Connor (1994: 453); Forrester *et al.* (1995: 32); Palma (1996b: 167); Price *et al.* (2003); Palma & Jensen (2005: 55, 66); Adams *et al.* (2005: 3548, figs 1, 7–10); Adam (2007: 178); Martín-Mateo (2009: 216, fig. 42); Galloway & Lamb (2014: 445; 2015: 715); Bartlow *et al.* (2016: 222).

Remarks: *Columbicola columbae columbae* was introduced to New Zealand with its primary host (*Columba livia*) by human agency (Checklist Committee 2010: 245). Contrary to Price *et al.* (2003: 166), I regard this louse taxon as a subspecies. The New Zealand record of *Columbicola columbae* from *Streptopelia chinensis tigrina* is based on a large sample (37 specimens) from two individual hosts, which is unlikely to be the result of contamination or accidental straggling from *Columba livia*. Instead, it is likely that *Columbicola columbae* host-switched and has become established on *Streptopelia chinensis tigrina* in New Zealand.

### Genus *Cuclotogaster* Carriker, 1936

*Cuclotogaster* Carriker, 1936. *Proc. Acad. Nat. Sci. Philad.*: 88: 67. Type species: *Cuclotogaster laticorpus* Carriker, 1936 = *Cuclotogaster heterographus* (Nitzsch [*in* Giebel], 1866) (by original designation).

*Gallipeurus* Clay, 1938. *Proc. Zool. Soc. London* 108: 135. Type species: *Lipeurus heterographus* Giebel, 1866 [sic] = *Cuclotogaster heterographus* (Nitzsch [*in* Giebel], 1866) (by original designation).

### *Cuclotogaster heterographus* (Nitzsch [*in* Giebel], 1866)

*Lipeurus heterographus* Nitzsch [*in* Giebel], 1866: 381.

*Lipeurus heterographus* Nitzsch [*in* Giebel], 1866; Harrison 1916: 84.

*Gallipeurus heterographus heterographus* Giebel, 1866 [sic]; Clay 1938: 136, 15–17, 18a.

*Lipeurus (Gallipeurus) heterographus* Nitzsch [*in* Giebel], 1866; Séguy 1944: 185, figs 268–269.

*Lipeurus heterographus* Nitz. [sic]; Helson 1956: 13, 17.

*Cuclotogaster heterographus* (Nitzsch, 1866) [sic]; Hopkins & Clay 1952: 95.

*Cuclotogaster heterographus heterographus* (Nitzsch, 1866) [sic]; Emerson 1956a: 64, pl. 1.

*Cuclotogaster heterographus*; Whitten 1971: 383.

*Cuclotogaster heterographus* (Nitzsch, 1866) [sic]; Wise 1977: 60.

*Cuclotogaster heterographus* (Nitzsch, 1866) [sic]; Pilgrim & Palma 1982: 18.

*Cuclotogaster heterographus* (Nitzsch, 1866) [sic]; Murray *et al.* 1993: 960.

*Cuclotogaster heterographus* (Nitzsch [*in* Giebel], 1866); Palma 1999: 384, note G.

*Cuclotogaster heterographus* (Nitzsch [*in* Giebel], 1866); Palma 2010: 408.

Syntypes probably lost (Palma 1996b: 170).

Type host: *Gallus gallus* (Linnaeus, 1758).

New Zealand hosts: *Alectoris chukar* (J.E. Gray, 1830); *Gallus gallus gallus* (Linnaeus, 1758).

Other host: *Phasianus colchicus* Linnaeus, 1758.

New Zealand localities: NC, MC, SC, CO, DN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Helson (1956); Whitten (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (1999); Palma (2010).

Other significant references: Webb (1946: 54); Conci (1952: 17); Emerson (1956a); Price (1987: 220, fig. 22.23); Martín-Mateo (1990: 180, figs 4–6); Palma (1996b: 170); Price *et al.* (2003: 171); Martín-Mateo (2009: 70, fig. 16); Dik *et al.* (2015: 796, figs 3–5).

Remarks: *Cuclotogaster heterographus* was introduced to New Zealand and other countries with its hosts by human agency. *Gallus gallus gallus* has been listed for the first time in the latest edition of the New Zealand Checklist of Birds (Checklist Committee 2010: 27; see also Palma 1999: 383, note 5).

***Cuclotogaster synoicus* (Clay, 1938)**

Figs 89–90

*Gallipeurus synoicus* Clay, 1938: 150, figs 23c, 24a, pl. 7: fig. 3.*Cuclotogaster synoicus* (Clay, 1938); Hopkins & Clay 1952: 96.*Cuclotogaster synoicus* (Clay, 1938); Pilgrim & Palma 1982: 18.*Cuclotogaster synoicus* (Clay, 1938); Murray *et al.* 1993: 961.*Cuclotogaster synoicus* (Clay, 1938); Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Coturnix ypsilophora australis* (Latham, 1802).New Zealand host: *Coturnix ypsilophora australis* (Latham, 1802).

Other hosts: None.

New Zealand localities: BP, HB.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).Other significant references: Green & Palma (1991: 13, 31); Palma (1996b: 170); Price *et al.* (2003: 171).Remarks: *Cuclotogaster synoicus* and its host are native to Australia, and were introduced to New Zealand by human agency (Checklist Committee 2010: 27).**Genus *Cuculicola* Clay & Meinertzhagen, 1939.***Cuculicola* Clay & Meinertzhagen, 1939a. *Entomologist* 72: 165. Type species: *Degeeriella latirostris* (Burmeister, 1838a) = *Cuculicola latirostris* (Burmeister, 1838a) (by original designation).***Cuculicola kui* Kettle, 1980**

Figs 91–92

*Cuculicola kui* Kettle, 1980: 89, figs 1–3a.*Cuculicola kui* Kettle, 1980; Pilgrim & Palma 1982: 25.*Cuculicola kui* Kettle, 1980; Murray *et al.* 1999: 1241.*Cuculicola kui* Kettle, 1980; Palma 2010: 408.Holotype ♂ in MONZ (Palma *et al.* 1989: 45).Type host: *Chrysococcyx lucidus lucidus* (J.F. Gmelin, 1788).New Zealand host: *Chrysococcyx lucidus lucidus* (J.F. Gmelin, 1788).

Other hosts: None.

New Zealand localities: BP, BP, WA, WN, NN, NC, MC, SC, WD, CO, DN.

Geographic distribution: Australasia.

New Zealand references: Kettle (1980); Pilgrim & Palma (1982); Palma *et al.* (1989: 45); Murray *et al.* (1999); Palma (2010).Other significant references: Palma (1996b: 171); Price *et al.* (2003: 172).Remarks: *Cuculicola kui* is currently known from New Zealand records only, but its host has a much wider geographical distribution (Checklist Committee 2010: 261). Therefore, although at present this louse is regarded as endemic to New Zealand, it may prove to be native if collected elsewhere in the future.***Cuculicola latirostris* (Burmeister, 1838)***Nirmus latirostris* Burmeister, 1838a: 429.*Degeeriella latirostris* Burmeister, 1838 [sic]; Harrison 1916: 116.*Cuculicola latirostris* (Burmeister, 1838); Hopkins & Clay 1952: 97.*Cuculicola latirostris* (Burmeister, 1838); Palma 1999: 381.*Cuculicola latirostris* (Burmeister, 1838); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma &amp; Pilgrim (1984: 150).

Type host: *Cuculus canorus* Linnaeus, 1758.New Zealand host: *Cuculus optatus* Gould, 1845.Other host: *Cuculus saturatus* Blyth, 1843.

New Zealand localities: ND, WD.

Geographic distribution: Eurasia, Australasia.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Clay & Meinertzhagen (1939a: 165, fig. 5); Séguy (1944: 306, fig. 466); Tendeiro (1961: 300, pl. 8: photo 17); Kettle (1980: 89, fig. 3b); Rékasi (1986: 123, figs 16–17); Brooke & Nakamura (1998: 168); Price *et al.* (2003: 172).

Remarks: The taxonomic nomenclature of the hosts of *Cuculicola latirostris* is unstable (Checklist Committee 2010: 259).

### Genus *Degeeriella* Neumann, 1906

*Nirmus* Nitzsch, 1818. *Germer's Mag. Entomol.* 3: 291. Type species: "*Degeeriella discocephalus* N." = *Degeeriella discocephalus* (Burmeister, 1838a) (by subsequent designation). Preoccupied by *Nirmus* Hermann, 1804.

*Degeeriella* Neumann, 1906. *Bull. Soc. Zool. France* 31: 59. *Nomen novum* for *Nirmus* Nitzsch, 1818.

### *Degeeriella fusca* (Denny, 1842)

Figs 93–94

*Philopterus* (*Nirmus*) *fuscus* ? [sic] Denny, 1842: 48, 118, pl. 9: fig. 8.

*Nirmus fuscus* N. [sic]; Piaget 1880: 130, pl. 10: fig. 9.

*Degeeriella fusca* Nitzsch [in Denny], 1842 [sic]; Harrison 1916: 113.

*Degeeriella fusca* (Denny, 1842); Hopkins & Clay 1952: 112.

*Degeeriella fusca* (Denny, 1842); Clay 1958: 162, figs 3, 30, 48, pl. 4: fig. 3, pl. 8: fig. 6.

*Degeeriella fusca* (Denny, 1842); Pilgrim & Palma 1982: 17.

*Degeeriella fusca* (Denny, 1842); Murray *et al.* 1993: 960.

*Degeeriella fuscus* [sic] (Denny, 1842); Galloway 2005: 17.

*Degeeriella fusca* (Denny, 1842); Palma 2010: 408.

Lectotype ♀ in NHML (Clay 1958: 164).

Type host: *Circus aeruginosus aeruginosus* (Linnaeus, 1758).

New Zealand host: *Circus approximans* Peale, 1848.

Other hosts: *Accipiter fasciatus* (Vigors & Horsfield, 1827) and five other species of *Circus* (see Price *et al.* 2003: 174)

New Zealand localities: AK, WO, HB, SD, MB, NN, NC, MC, SC, WD, CO, DN, CH.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Galloway (2005); Palma (2010).

Other significant references: Séguy (1944: 271, figs 411–413); Clay (1958); Green & Palma (1991: 13, 30); Palma (1996b: 173); Price *et al.* (2003: 174); Martín-Mateo (2009: 92, fig. 20I); Catanach & Johnson (2015: 839).

Remarks: *Degeeriella fusca* is frequently collected from New Zealand swamp harriers.

### *Degeeriella rufa rufa* (Burmeister, 1838)

*Nirmus rufus* Burmeister, 1838a: 430.

*Degeeriella rufa* Nitzsch [in Burmeister], 1838 [sic]; Harrison 1916: 122.

*Degeeriella rufa* (Burmeister, 1838); Hopkins & Clay 1952: 113.

*Degeeriella rufa rufa* (Burmeister, 1838); Clay 1958: 180, figs 5, 10, 18–22, 38, 55, 75, 95, 104, 107, 111, 120, 129, 139–164, pl. 6: fig. 3, pl. 8: fig. 7.

*Degeeriella rufa rufa* (Burmeister, 1838); Pilgrim & Palma 1982: 17.

*Degeeriella rufa rufa* (Burmeister, 1838); Murray *et al.* 1993: 960.

*Degeeriella rufa* (Burmeister, 1838); Price *et al.* 2003: 175.

*Degeeriella rufa rufa* (Burmeister, 1838); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1), Palma & Pilgrim (1984: 150) and Palma (1996b: 174).

Type host: *Falco tinnunculus* Linnaeus, 1758.

New Zealand host: *Falco novaeseelandiae* J.F. Gmelin, 1788.

Other hosts: *Polihierax insignis* Walden, 1872 and over twenty other species of *Falco* (Price *et al.* 2003: 175).

New Zealand localities: TK, WA, WN, SD, MB, NC, MC, SC, WD.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 221); Palma (2010).

Other significant references: Clay (1958); Rékasi (1986: 124, figs 22–23); Green & Palma (1991: 13, 31); Butler & O'Connor (1994: 454); Palma (1996b: 174); Price *et al.* (2003); Palma & Jensen (2005: 55, 62); Martín-Mateo (2009: 87, fig. 20C); Catanach & Johnson (2015: 839).

Remarks: *Degeeriella rufa* is a morphologically variable species (Clay 1958: 181). Contrary to Price *et al.* (2003: 175), I regard some subspecies of *Degeeriella rufa* as valid taxa.

### Genus *Docophoroides* Giglioli, 1864

*Docophoroides* “Denny MSS” Giglioli, 1864. *Quart. Jour. Sci.* 4: 21. Type species: *Philopterus brevis* Dufour, 1835 = *Docophoroides brevis* (Dufour, 1835) (by monotypy).

*Eurymetopus* Taschenberg, 1882. *Nova Acat Leop.-Carol.* 44: 182. Type species: *Lipeurus taurus* Nitzsch [in Giebel], 1866 = *Docophoroides brevis* (Dufour, 1835) (by subsequent designation). Preoccupied by *Eurymetopus* Schönherr, 1840.

### *Docophoroides brevis* (Dufour, 1835)

*Philopterus brevis* Dufour, 1835: 674, pl. 21: fig. 3.

*Docophoroides brevis*; Giglioli 1864: 21, pl. 1: figs 3–4.

*Lipeurus taurus* Nitzsch [in Giebel], 1866: 385.

*Docophoroides brevis* Dufour, 1835 [sic]; Harrison 1937: 40, pl. 3: fig. 6a.

*Docophoroides* sp. n.; Eichler 1941a: 354, fig. 16.

*Docophoroides brevis* (Dufour, 1834) [sic]; Séguy 1944: 382, figs 571–572.

*Docophoroides brevis* (Dufour, 1835); Hopkins & Clay 1952: 117.

*Docophoroides brevis* (Dufour, 1835); Timmermann 1959c: 59, figs 2a, 3, 7c.

*Docophoroides brevis* (Dufour, 1835); Clay 1964a: 232.

*Docophoroides brevis* (Dufour, 1835); Timmermann 1965: 83, figs 19, 24c, 26a.

*Docophoroides brevis* (Dufour, 1835); Wise 1977: 60.

*Docophoroides brevis* (Dufour, 1835); Pilgrim & Palma 1982: 5.

*Docophoroides brevis* (Dufour, 1835); Palma 2010: 408.

Syntypes ♂♀, repository unknown (Palma 1996b: 174), probably lost.

Type host: *Diomedea exulans* Linnaeus, 1758.

New Zealand hosts: *Diomedea exulans* Linnaeus, 1758; *Diomedea antipodensis antipodensis* Robertson & Warham, 1992; *Diomedea antipodensis gibsoni* Robertson & Warham, 1992; *Diomedea epomophora* Lesson, 1825; *Diomedea sanfordi* Murphy, 1917.

Other hosts: *Diomedea amsterdamensis* Roux *et al.* 1983; *Diomedea dabbenena* Mathews, 1929.

New Zealand localities: ND, AK, BP, HB, TK, WA, WN, KA, NC, MC, SC, WD, CO, DN, CH, AU, AN, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Harrison (1937); Clay (1940a: 298); Clay (1964a); Gressitt (1964: 538); Watson (1967: 71); Clay & Moreby (1970: 217); Gressitt (1970: 327); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1368); Palma (1996b: 174); Marris (2000: 187); Palma & Horning (2002: 7, 15); Page *et al.* (2004: 637, 650); Palma (2010).

Other significant references: Séguy (1944); Séguy (1953: 585, fig. 43); Timmermann (1959c; 1965); Clay & Moreby (1967: 161, 168, figs 108, 111–112); Green & Palma (1991: 13, 25); Furness & Palma (1992: 35, 38); Green & Turner (2003b: 77); Price *et al.* (2003: 176); Hänel & Palma (2007: 112, 123, 129).

Remarks: *Docophoroides brevis* is a highly prevalent species, mostly found on the head and neck of large albatrosses.

### *Docophoroides harrisoni* Waterston, 1917

*Docophoroides harrisoni* Waterston, 1917: 99, fig.

*Docophoroides harrisoni* Waterston, 1917; Harrison 1937: 41, pl. 3: figs 1, 6f.

*Docophoroides harrisoni* Waterston, 1917; Hopkins & Clay 1952: 118.

*Docophoroides harrisoni* Waterston, 1917; Timmermann 1959c: 62, figs 1, 6.

*Docophoroides harrisoni* Waterston, 1917; Timmermann 1965: 85, figs 20, pl. 3: fig. 1.

*Docophoroides harrisoni* Waterston, 1917; Horning *et al.* 1980: 5, 9.

*Docophoroides harrisoni* Waterston, 1917; Pilgrim & Palma 1982: 6.

*Docophoroides harrisoni* Waterston, 1917; Murray *et al.* 1990: 1368–1369.

*Docophoroides harrisoni* Waterston, 1917; Palma 2010: 408.

Holotype ♂ in SAMS (Palma 1996b: 174).

Type host: *Thalassarche melanophris* (Temminck, 1828).

New Zealand hosts: *Thalassarche melanophris* (Temminck, 1828); *Thalassarche bulleri bulleri* (Rothschild, 1893); *Thalassarche bulleri platei* (Reichenow, 1898); *Thalassarche cauta steadi* Falla, 1933; *Thalassarche eremita* Murphy, 1930; *Thalassarche salvini* (Rothschild, 1893).

Other host: *Thalassarche cauta cauta* (Gould, 1841).

New Zealand localities: ND, AK, BP, HB, WI, WA, WN, NN, KA, NC, MC, SC, WD, CO, DN, SL, CH, SI, BO, SN, AU.

Geographic distribution: Southern Hemisphere.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Page *et al.* (2004: 637, 650); Palma (2010).

Other significant references: Harrison (1937); Timmermann (1959c; 1965); Clay & Moreby (1967: 161, 168, fig. 106); Green & Palma (1991: 13, 25); Palma (1996b: 174); Price *et al.* (2003: 176).

Remarks: *Docophoroides harrisoni* is a highly prevalent species, mostly found on the head and neck of small albatrosses, also known as mollymawks.

#### ***Docophoroides murphyi* (Kellogg, 1914)**

*Eurymetopus murphyi* Kellogg, 1914: 87, pl. 16: figs 4–5.

*Docophoroides murphyi* Kellogg, 1914 [sic]; Harrison 1916: 144.

*Docophoroides hunteri* Harrison, 1937: 42, pl. 3: figs 3–6c.

*Docophoroides hunteri* Harrison, 1937; Hopkins & Clay 1952: 118.

*Docophoroides murphyi* (Kellogg, 1914); Hopkins & Clay 1952: 118.

*Docophoroides hunteri* Harrison, 1937; Timmermann 1959c: 67.

*Docophoroides murphyi* (Kellogg, 1914); Timmermann 1959c: 68.

*Docophoroides murphyi* (Kellogg, 1914); Clay 1964a: 232.

*Docophoroides hunteri* Harrison, 1937; Timmermann 1965: 88.

*Docophoroides murphyi* (Kellogg, 1914); Clay & Moreby 1967: 161, 168, figs 107, 109, 113.

*Docophoroides murphyi* (Kellogg, 1914); Wise 1977: 60.

*Docophoroides murphyi* (Kellogg, 1914); Pilgrim & Palma 1982: 7.

*Docophoroides murphyi* (Kellogg, 1914); Palma 2010: 408.

Lectotype ♂ in USNM (Emerson 1961b: 250). Holotype ♂ of *Docophoroides hunteri* in AMSA (Palma 1996b: 175).

Type host: *Macronectes giganteus* (J.F. Gmelin, 1789) (see Emerson 1961b: 250).

New Zealand hosts: *Macronectes giganteus* (J.F. Gmelin, 1789); *Macronectes halli* Mathews, 1912.

Other hosts: None.

New Zealand localities: ND, AK, BP, WO, TK, WA, WN, KA, NC, MC, SC, WD, SN, AN, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Harrison (1937); Clay (1964a); Gressitt (1964: 539); Watson (1967: 71); Clay & Moreby (1970: 217); Gressitt (1970: 327); Wise (1977); Pilgrim & Palma (1982); Horning *et al.* (1980: 5, 9); Murray *et al.* (1990: 1369); Palma (1996b: 174); Paterson *et al.* (1999: 222); Marris (2000: 187); Palma (2001: 65, fig. 2); Palma & Horning (2002: 7, 16); Palma (2010).

Other significant references: Timmermann (1959c); Emerson (1961b: 250); Timmermann (1965); Clay & Moreby (1967); Green & Palma (1991: 13, 25); Price *et al.* (2003: 176); Hänel & Palma (2007: 112, 123, 129).

Remarks: *Docophoroides murphyi* is a highly prevalent species, mostly found on the head and neck of giant petrels.

***Docophoroides simplex* (Waterston, 1914)**

Figs 95–96

*Eurymetopus simplex* Waterston, 1914: 302, fig. 2.*Docophoroides simplex* Waterston, 1914 [sic]; Harrison 1937: 41, pl. 3: figs 2, 6b.*Docophoroides simplex* (Waterston, 1914); Hopkins & Clay 1952: 118.*Docophoroides simplex* (Waterston, 1914); Timmermann 1959c: 66, figs 2c, 5, 7b.*Docophoroides simplex* (Waterston, 1914); Timmermann 1965: 87, figs 23, 24b, 26c, pl. 3: fig. 2.*Docophoroides simplex* (Waterston, 1914); Clay & Moreby 1967: 161, 168, figs 110, 114.*Docophoroides simplex* (Waterston, 1914); Pilgrim & Palma 1982: 6.*Docophoroides simplex* (Waterston, 1914) s. l.; Pilgrim & Palma 1982: 6.*Docophoroides simplex* (Waterston, 1914); Murray *et al.* 1990: 1368–1369.*Docophoroides simplex* (Waterston, 1914); Palma 2010: 408.

Syntypes ♂♀ in SAMS (Palma 1996b: 175).

Type host: *Thalassarche melanophris* (Temminck, 1828).New Zealand hosts: *Thalassarche chrysostoma* (J.R. Forster, 1785); *Thalassarche melanophris* (Temminck, 1828); *Thalassarche impavida* Mathews, 1912; *Phoebetria palpebrata* (J.R. Forster, 1785).Other host: *Thalassarche chlororhynchos* (J.F. Gmelin, 1789).

New Zealand localities: ND, AK, WI, WN, NN, NC, MC, SC, SL, CH, CA, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b: 175); Palma & Horning (2002: 8, 15); Palma (2010).Other significant references: Timmermann (1959c; 1965); Clay & Moreby (1967); Green & Palma (1991: 13, 25); Price *et al.* (2003: 176); Page *et al.* (2004: 647); Hänel & Palma (2007: 112, 123, 129).Remarks: Pilgrim & Palma (1982: 6) regarded the populations of *Docophoroides simplex* from *Thalassarche chrysostoma* and *Phoebetria palpebrata* as somewhat different from that of the type host, and qualified them as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted. *Docophoroides simplex* is mostly found on the head and neck of small albatrosses.**Genus *Emersoniella* Tendeiro, 1965***Emersoniella* Tendeiro, 1965c. *Revista Estudos Gerais Universitários Moçambique* 2 (Série 4): 69. Type species: *Emersoniella halcyonis* Tendeiro, 1965c (by original designation).***Emersoniella bracteata* (Nitzsch [*in* Giebel], 1866)****New Record**

Figs 97–98

*Nirmus bracteatus* Nitzsch [*in* Giebel], 1866: 369.*Nirmus bracteatus* Nitzsch, 1866 [sic]; Hopkins & Clay 1952: 241.*Emersoniella bracteata* (Nitzsch, 1866) [sic]; Clay 1971: 45.*Brueelia* [sic] *bracteata* (Nitzsch, 1866) [sic]; Clay 1971: figs 1–2.*Emersoniella bracteata* (Nitzsch [*in* Giebel], 1866); Palma 1996b: 176.*Emersoniella bracteata* (Nitzsch [*in* Giebel], 1866); Price *et al.* 2003: 177.

Syntypes ♂♀, presumed lost. See Clay (1949a: 1), Palma &amp; Pilgrim (1984: 150) and Palma (1996b: 176).

Type host: *Dacelo novaeguineae* (Hermann, 1783).New Zealand host: *Dacelo novaeguineae novaeguineae* (Hermann, 1783).

Other hosts: None.

New Zealand localities: ND, AK.

Geographic distribution: Australasia.

New Zealand reference: This paper.

Other significant references: Clay (1971); Palma (1996b); Price *et al.* (2003); Gustafsson & Bush (2014: 528, 543).

Material examined and repository: 4♂, 7♀, 5 nymphs (2 samples, MONZ).



Remarks: *Emersoniella bracteata* is a new louse species for New Zealand, introduced to this country with kookaburras from Australia by human agency (Checklist Committee 2010: 272). Also, *Emersoniella* is a new genus record for New Zealand.

### Genus *Episbates* Thompson, 1935

*Episbates* Thompson, 1935c. *Ann. Mag. Nat. Hist.* (Ser. 10) 16: 485. Type species: *Philopterus pederiformis* Dufour, 1835 = *Episbates pederiformis* (Dufour, 1835) (by original designation).

#### *Episbates pederiformis* (Dufour, 1835)

Figs 99–100

*Philopterus pederiformis* Dufour, 1835: 676, pl. 21: fig. 4.

*Nirmus angulicollis* Giebel, 1876: 388.

*Lipeurus macilhennyi* Kellogg & Kuwana, 1901: 155, pl. 7: fig. 3.

*Esthiopterum pederiforme* Dufour, 1835 [sic]; Harrison 1916: 139.

*Episbates pederiformis* (Dufour, 1835); Thompson 1935c: 486, fig. 1.

*Episbates pederiformis* (Dufour, 1835); Thompson 1948c: 662, figs 1–8, pl. 19: figs, 1–3.

*Episbates macilhennyi* (Kellogg & Kuwana, 1901); Hopkins & Clay 1952: 131.

*Episbates pederiformis* (Dufour, 1835); Hopkins & Clay 1952: 131.

*Episbates pederiformis* (Dufour, 1835); Timmermann 1961c: 44, figs 12–13.

*Episbates pederiformis* (Dufour, 1835); Timmermann 1965: 105, fig. 41, pl. 10: fig. 4.

*Episbates pederiformis* (Dufour, 1835); Pilgrim & Palma 1982: 5.

*Episbates pederiformis* (Dufour, 1835); Palma 2010: 408.

Status, sex and repository of types unknown (Palma 1996b: 176), probably lost.

Type host: *Diomedea exulans* Linnaeus, 1758.

New Zealand hosts: *Diomedea exulans* Linnaeus, 1758; *Diomedea antipodensis antipodensis* Robertson & Warham, 1992; *Diomedea antipodensis gibsoni* Robertson & Warham, 1992; *Diomedea epomophora* Lesson, 1825; *Diomedea sanfordi* Murphy, 1917.

Other hosts: *Phoebastria albatrus* (Pallas, 1769); *Phoebastria nigripes* (Audubon, 1839); *Phoebastria irrorata* (Salvin, 1883); *Phoebastria immutabilis* (Rothschild, 1893).

New Zealand localities: AK, WN, KA, WD, CO, DN, CH, AU, AN, Macquarie Island.

Geographic distribution: Southern Hemisphere and north Pacific Ocean.

New Zealand references: Harrison (1937: 27, fig. 3); Pilgrim & Palma (1982); Murray *et al.* (1990: 1368); Marris (2000: 187); Palma (2001: 66, fig. 4); Page *et al.* (2004: 638, 648); Palma (2010).

Other significant references: Thompson (1935c); Clay (1940a: 298); Séguy (1944: 369, figs 548–549); Thompson (1948c); Hopkins & Clay (1952); Séguy (1953: 595, figs 51–52); Clay (1958a: 251); Timmermann (1961c; 1965); Clay & Moreby (1967: 159, fig. 81); Amerson & Emerson (1971: 2, 23); Palma (1996b: 176); Price *et al.* (2003: 178); Palma & Peck (2013: 36).

Remarks: *Episbates pederiformis* is a monotypic louse species parasitic on a wide range of albatross species. Thompson's (1948c: 667) designation of "... a male and a female of the British Museum specimens as neotypes ..." is invalid because the neotype has to be a single specimen, which needs to be clearly identifiable (see Palma 1996b: 176).

### Genus *Forficuloecus* Conci, 1941

*Forficuloecus* Conci, 1941. *Boll. Soc. Entomol. Italiana* 73: 126. Type species: *Philopterus forficula* Piaget, 1871 = *Forficuloecus forficula* (Piaget, 1871) (by original designation).

#### *Forficuloecus meinertzhageni* Guimarães, 1974

*Forficuloecus meinertzhageni* Guimarães, 1974b: 177, figs 25–29.

*Forficuloecus meinertzhageni* Guimarães, 1974; Wise 1977: 60.

*Forficuloecus meinertzhageni* Guimarães, 1974; Pilgrim & Palma 1982: 24.

*Forficuloecus meinertzhageni* Guimarães, 1974 s. l.; Pilgrim & Palma 1982: 24.

*Forficuloecus meinertzhageni* Guimarães, 1974; Guimarães 1985: 41, figs 1, 9.

*Forficuloecus meinertzhageni* Guimarães, 1974; Murray *et al.* 1999: 1241.

*Forficuloecus meinertzhageni* Guimarães, 1974; Price *et al.* 2008: 53, fig. 6.

*Forficuloecus meinertzhageni* Guimarães, 1974; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Nestor meridionalis* (J.F. Gmelin, 1788).

New Zealand hosts: *Nestor meridionalis septentrionalis* Lorenz, 1896; *Nestor meridionalis meridionalis* (J.F. Gmelin, 1788); *Nestor notabilis* Gould, 1856.

Other hosts: None.

New Zealand localities: HB, WN, NC, MC, WD.

Geographic distribution: New Zealand.

New Zealand references: Guimarães (1974b); Wise (1977); Pilgrim & Palma (1982); Guimarães (1985); Murray *et al.* (1999); Price *et al.* (2008); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Price *et al.* (2003: 176).

Remarks: *Forficuloecus meinertzhageni* is an endemic and “at risk” species (Buckley *et al.* 2012), exclusively parasitic on the kea and on both kaka subspecies. Pilgrim & Palma (1982: 24) regarded the population of *Forficuloecus meinertzhageni* from the kea as somewhat different from that of the type host, the kaka, and qualified it as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted

### ***Forficuloecus pilgrimi* Guimarães, 1985**

Figs 101–102

*Forficuloecus* sp.; Pilgrim & Palma 1982: 24.

*Forficuloecus pilgrimi* Guimarães, 1985: 43, figs 3, 7, 11, 15.

*Forficuloecus pilgrimi* Guimarães, 1985; Murray *et al.* 1999: 1241.

*Forficuloecus pilgrimi* Guimarães, 1985; Palma 1999: 381.

*Forficuloecus pilgrimi* Guimarães, 1985; Price *et al.* 2008: 53, figs 7–8.

*Forficuloecus pilgrimi* Guimarães, 1985; Palma 2010: 408.

Holotype ♂ in MONZ (Palma *et al.* 1989: 45).

Type host: *Cyanoramphus novaezelandiae chathamensis* Oliver, 1930.

New Zealand hosts: *Cyanoramphus novaezelandiae novaezelandiae* (Sparrman, 1787); *Cyanoramphus novaezelandiae chathamensis* Oliver, 1930; *Cyanoramphus auriceps* (Kuhl, 1820); *Cyanoramphus malherbi* Souancé, 1857; *Cyanoramphus forbesi* Rothschild, 1893; *Cyanoramphus unicolor* (Lear, 1831); *Cyanoramphus hochstetteri* (Reischek, 1889).

Other hosts: None.

New Zealand localities: CL, BP, WN, NN, NC, MC, SC, FD, CH, AN.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Guimarães (1985); Palma *et al.* (1989: 45); Murray *et al.* (1999); Palma (1999); Marris (2000: 187); Price *et al.* (2008); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Price *et al.* (2003: 180).

Remarks: *Forficuloecus pilgrimi* is an endemic and “at risk” species (Buckley *et al.* 2012), exclusively parasitic on several species and subspecies of New Zealand parakeets.

### ***Forficuloecus* species**

### **New Record**

New Zealand host: *Platycercus eximius* (Shaw, 1792).

New Zealand locality: WN.

New Zealand reference: This paper.

Material examined and repository: 3 nymphs (1 sample, MONZ).

Remarks: The natural and regular species of *Forficuloecus* parasitising *Platycercus eximius* in Australia is *F. forficula* (Piaget, 1871) (see Price *et al.* 2003: 179). However, the single and first New Zealand record of this louse could not be identified to species because the sample contains nymphs only.

### Genus *Fulicoffula* Clay & Meinertzhagen, 1938

*Fulicoffula* Clay & Meinertzhagen, 1938b. *Entomologist* 71: 279. Type species: *Esthiopterum luridum* (Denny) [sic] = *Fulicoffula lurida* (Nitzsch, 1818) (by original designation).

#### *Fulicoffula lurida* (Nitzsch, 1818)

“Pollino della folaga” Redi, 1668: pl. 4: fig. 2.

*Philopterus* (*Lipeurus*) *luridus* Nitzsch, 1818: 292. *Nomen novum* for “Pollino della folaga” Redi, 1668: pl. 4: fig. 2.

*Lipeurus luridus* Nitzsch, 1818; Giebel 1874: 230, pl. 16: fig. 4.

*Esthiopterum luridum* Nitzsch in Denny, 1842 [sic]; Harrison 1916: 137.

*Esthiopterum luridum* (Denny) [sic]; Clay & Meinertzhagen, 1938b: 279.

*Fulicoffula lurida* (Nitzsch, 1818); Hopkins 1940: 425.

*Fulicoffula lurida* (Nitzsch, 1818); Hopkins & Clay 1952: 141.

*Fulicoffula lurida* (Nitzsch, 1818); Clay & Hopkins 1960: 40, figs 63–71, pl. 6: figs 1–2.

*Fulicoffula lurida* (Nitzsch, 1818); Tendeiro 1963: 86, photos 69–70, 74, 76.

*Fulicoffula lurida* (Nitzsch, 1818); Pilgrim & Palma 1982: 19.

*Fulicoffula lurida* (Nitzsch, 1818); Murray *et al.* 1993: 961.

*Fulicoffula lurida* (Nitzsch, 1818); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1960: 43, pl. 6: fig. 1).

Type host: *Fulica atra* Linnaeus, 1758.

New Zealand host: *Fulica atra australis* Gould, 1845.

Other hosts: *Fulica cristata* J.F. Gmelin, 1789; *Fulica leucoptera* Vieillot, 1817.

New Zealand localities: WO, SD, MB, NC, MC, SC, CO, DN.

Geographic distribution: Africa; Eurasia; Australasia; South America.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Hopkins (1940); Clay & Hopkins (1960); Tendeiro (1963); Palma (1996b: 178); Green & Palma (1991: 14, 32); Butler & O’Connor (1994: 455); Price *et al.* (2003: 181); Palma & Jensen (2005: 55, 63); Adam (2007: 175); Martín-Mateo (2009: 234).

Remarks: Hopkins (1940: 421) gives a complete account of the early taxonomic confusion among several species of lice described from *Fulica atra*, including a clarification of the synonymy of *Fulicoffula lurida*.

#### *Fulicoffula stammeri* Eichler, 1958

#### New Record

Figs 103–104

*Fulicoffula stammeri* Eichler, 1958: 61, fig. 3.

*Fulicoffula* sp.; Pilgrim & Palma 1982: 19.

*Fulicoffula* sp.; Murray *et al.* 1993: 961.

*Fulicoffula stammeri* Eichler, 1958; Price *et al.* 2003: 181.

Holotype ♂, probably in the University of Erlangen-Nuremberg, Germany (Jürgen Schmidl pers. comm. July 2012).

Type host: *Porzana porzana* (Linnaeus, 1766).

New Zealand hosts: *Porzana pusilla affinis* (J.E. Gray, 1845); *Porzana tabuensis tabuensis* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: ND, AK, WO, TO, HB, RI, SD, MB, NC, MC, SC, WD, SL, FD, AN.

Geographic distribution: Eurasia; Africa; New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993).

Other significant references: Price *et al.* (2003).

Material examined and repository: 36♂, 54♀, 5 nymphs (22 samples, MONZ).

Remarks: This is the first record of *Fulicoffula stammeri* for New Zealand because the New Zealand references cited above reported this louse as “*Fulicoffula* sp.” only.

**Genus *Goniocotes* Burmeister, 1838**

*Goniocotes* Burmeister, 1838a. *Handb. Entomol.* 2(1): 431. Type species: *Goniocotes gallinae* (De Geer, 1778b) (by subsequent designation).

***Goniocotes chrysocephalus* Giebel, 1874**

*Goniocotes chrysocephalus* Giebel, 1874: 189.  
*Goniocotes chrysocephalus* Giebel, 1874; Kéler 1939: 152, fig. 85.  
*Goniocotes chrysocephalus* Giebel, 1874; Hopkins & Clay 1952: 147.  
*Goniocotes chrysocephalus* Giebel, 1874; Pilgrim & Palma 1982: 18.  
*Goniocotes chrysocephalus* Giebel, 1874; Murray *et al.* 1993: 960.  
*Goniocotes chrysocephalus* Giebel, 1874; Palma 2010: 408.

Type material: one nymph of doubtful status in MLUH (Karla Schneider pers. comm. May 2014).

Type host: *Phasianus colchicus* Linnaeus, 1758.

New Zealand host: *Phasianus colchicus* Linnaeus, 1758.

Other host: *Bonasa umbellus* (Linnaeus, 1766).

New Zealand localities: RI, WI.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Kéler (1939); Marconcini & Macchioni (1975: 105, figs 3–4); Modrzejewska & Złotorzycka (1987: 661, fig. 3); Kopociński *et al.* (1998: 81); Price *et al.* (2003: 181); Adam (2007: 171); Martín-Mateo (2009: 36).

Remarks: *Goniocotes chrysocephalus* was introduced to New Zealand and other countries with common pheasants by human agency (Checklist Committee 2010: 28).

***Goniocotes gallinae* (De Geer, 1778)**

*Ricinus gallinae* De Geer, 1778b: 79, pl. 4: fig. 15.  
*Philopterus (Goniodes) hologaster* Nitzsch, 1818: 294. Unnecessary *nomen novum* for *Ricinus gallinae* De Geer, 1778.  
*Goniocotes gallinae* De Geer, 1778 [sic]; Harrison 1916: 81 (as junior synonym of *Goniocotes hologaster* Nitzsch).  
*Goniocotes hologaster* (Nitzsch, 1818); Kéler 1939: 135, figs 72–73.  
*Goniocotes gallinae* (De Geer, 1778); Hopkins & Clay 1952: 147.  
*Goniocotes gallinae*; Whitten 1971: 383.  
*Goniocotes gallinae* (De Geer, 1778); Murray *et al.* 1993: 960.  
*Goniocotes gallinae* (De Geer, 1778); Palma 1999: 384, note G.  
*Goniocotes gallinae* (De Geer, 1778); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1954: 242).

Type host: *Gallus gallus* (Linnaeus, 1758).

New Zealand host: *Gallus gallus gallus* (Linnaeus, 1758).

Other hosts: *Caloperdix oculeus* (Temminck, 1815); *Meleagris gallopavo* Linnaeus, 1758.

New Zealand localities: WI, WN, NC, MC, SC, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Whitten (1971: 383); Murray *et al.* (1993); Palma (1999); Palma (2010).

Other significant references: Kéler (1939); Clay & Hopkins (1954: 242, figs 29–31); Emerson (1956a: 67, pl. 2); Emerson & Ward (1958: 56); Price (1987: 220, fig. 22.24); Palma (1996b: 179); Price *et al.* (2003: 181); Palma & Jensen (2005: 55, 62); Martín-Mateo (2009: 32, figs 5b, 7).

Remarks: *Goniocotes gallinae* was introduced to New Zealand and other countries with chickens by human agency. *Gallus gallus gallus* has been listed for the first time in the latest edition of the New Zealand Checklist of Birds (Checklist Committee 2010: 27; see also Palma 1999: 383, note 5).

***Goniocotes pusillus* (Nitzsch [in Giebel], 1866)**

Figs 105–106

*Goniodes pusillus* Nitzsch [in Giebel], 1866: 387.

- Goniocotes pusillus* Nitzsch [*in* Giebel], 1874 [sic]; Harrison 1916: 82.  
*Goniocotes pusillus* Nitzsch in Burmeister, 1838a [sic]; Kéler 1939: 149, fig. 83.  
*Goniocotes pusillus* (Nitzsch, 1866) [sic]; Hopkins & Clay 1952: 149.  
*Goniocotes pusillus* (Nitzsch, 1866) [sic]; Pilgrim & Palma 1982: 18.  
*Goniocotes pusillus* (Nitzsch, 1866) [sic]; Murray *et al.* 1993: 961.  
*Goniocotes pusillus* (Nitzsch [*in* Giebel], 1866); Price *et al.* 2003: 182.  
*Goniocotes pusillus* (Nitzsch [*in* Giebel], 1866); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Alectoris barbara* (Bonnaterre, 1792).

New Zealand host: *Alectoris chukar* (J.E. Gray, 1830).

Other hosts: None.

New Zealand locality: CO, DN.

Geographic distribution: Eurasia; Africa, New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Kéler (1939); Price *et al.* (2003).

Remarks: *Goniocotes pusillus* was introduced to New Zealand with chukors by human agency (Checklist Committee 2010: 26).

### Genus *Goniodes* Nitzsch, 1818

- Goniodes* Nitzsch, 1818. *German's Mag. Entomol.* 3: 293. Type species: *Goniodes pavonis* (Linnaeus, 1758) (by subsequent designation).  
*Gonocephalus* "Nitzsch" in Kéler, 1937. *Arb. Morph. tax. Entomol. Berlin-Dahlem* 4: 130. Type species: *Goniodes chelicornis* "Nitzsch" = *Goniodes bituberculatus* Rudow, 1869b (by original designation).  
*Oulocrepis* Kéler, 1939. *Nova Acta Leop.-Carol. (N.F.)* 8: 97. Type species: *Goniodes dissimilis* Nitzsch [sic] = *Goniodes dissimilis* (Denny, 1842) (by original designation).  
*Solenodes* Kéler, 1939. *Nova Acta Leop.-Carol. (N.F.)* 8: 101. Type species: *Goniodes dispar* Nitzsch [sic] = *Goniodes dispar* Burmeister, 1838a (by original designation).

### *Goniodes colchici* (Denny, 1842)

- Philoferus* (*Goniodes*) *colchici* Denny, 1842: 56, 158, pl. 12: fig. 4.  
*Gonocephalus colchici* (Denny, 1842); Kéler 1939: 94, figs 49–50.  
*Goniodes colchici* Denny, 1842 [sic]; Clay 1940b: 50, figs 32, 34b.  
*Goniodes colchici* Denny, 1842 [sic]; Hopkins & Clay 1952: 152.  
*Goniodes colchici* Denny, 1842 [sic]; Pilgrim & Palma 1982: 18.  
*Goniodes colchici* Denny, 1842 [sic]; Murray *et al.* 1993: 960.  
*Goniodes colchici* Denny, 1842 [sic]; Palma 2010: 408.

Lectotype ♂ in NHML (Clay 1940b: 50).

Type host: *Phasianus colchicus* Linnaeus, 1758.

New Zealand host: *Phasianus colchicus* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: WI, WN, NC, MC, SC.

Geographic distribution: Americas; Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Kéler (1939); Clay (1940b); Williams (1970a,b; 1971); Marconcini & Macchioni (1975: 104, figs 1–2); Rudolph (1983: 16); Modrzejewska & Złotorzycka (1987: 662, fig. 4); Butler & O'Connor (1994: 455); Palma (1996b: 180); Kopociński *et al.* (1998: 81); Price *et al.* (2003: 183); Martín-Mateo (2009: 49, fig. 11C,F).

Remarks: *Goniodes colchici* was introduced to New Zealand and other countries with common pheasants by human agency (Checklist Committee 2010: 28). Denny (1842: 155) clearly regarded *Goniodes* as a subgenus of *Philoferus*. However, subsequent authors ignored that fact and did not add parentheses around author and date when *Goniodes* is used at generic level.

***Goniodes dispar* Burmeister, 1838**

- Goniodes dispar* Burmeister, 1838a: 432.  
*Solenodes dispar* (Nitzsch) [sic]; Kéler 1939: 102, fig. 53.  
*Goniodes dispar* Burmeister, 1838; Clay 1940b: 87, fig. 60a.  
*Goniodes dispar* Burmeister, 1838; Hopkins & Clay 1952: 153.  
*Goniodes dispar* Burmeister, 1838; Pilgrim & Palma 1982: 18.  
*Goniodes dispar* Burmeister, 1838; Murray *et al.* 1993: 961.  
*Goniodes dispar* Burmeister, 1838; Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Perdix perdix perdix* (Linnaeus, 1758).

New Zealand host: *Alectoris chukar* (J.E. Gray, 1830).

Other hosts: *Alectoris graeca* (Mesiner, 1804); *Alectoris rufa* (Linnaeus, 1758).

New Zealand locality: CO, DN.

Geographic distribution: Eurasia; New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Clay (1940b); Price *et al.* (2003: 184); Millán *et al.* (2004: 79); Martín-Mateo (2009: 48, fig. 11B).

Remarks: *Goniodes dispar* was introduced to New Zealand with chukors by human agency (Checklist Committee 2010: 26).

***Goniodes dissimilis* (Denny, 1842)**

- Philopterus (Goniodes) dissimilis* Denny, 1842: 57, 162, pl. 12: fig. 6.  
*Goniodes dissimilis* Nitzsch [sic]; Thomson 1922: 269.  
*Oulocrepis dissimilis* (Nitzsch) [sic]; Kéler 1939: 98, figs 51–52.  
*Goniodes dissimilis* Denny, 1842 [sic]; Clay 1940b: 62, figs 41–43.  
*Goniodes dissimilis* Denny, 1842 [sic]; Hopkins & Clay 1952: 153.  
*Goniodes dissimilis*; Whitten 1971: 383.  
*Goniodes dissimilis* Denny, 1842 [sic]; Palma 1999: 384, note G.  
*Goniodes dissimilis* Denny, 1842 [sic]; Murray *et al.* 1993: 960.  
*Goniodes dissimilis* Denny, 1842 [sic]; Palma 2010: 408.

Neotype ♀ in NHML (Clay 1940b: 65).

Type host: *Gallus gallus* (Linnaeus, 1758).

New Zealand host: *Gallus gallus gallus* (Linnaeus, 1758).

Other hosts: *Gallus sonneratii* Temminck, 1813; *Gallus lafayettii* Lesson, 1831.

New Zealand localities: AK, WN, CO, DN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Thomson (1922); Kéler (1939); Whitten (1971); Murray *et al.* (1993); Palma (1999); Palma (2010).

Other significant references: Clay (1940b); Emerson (1956a: 69, pl. 3); Emerson & Ward (1958: 56); Palma (1996b: 181); Price *et al.* (2003: 184); Martín-Mateo (2009: 49, fig. 11E); Palma & Peck (2013: 37).

Remarks: *Goniodes dissimilis* was introduced to New Zealand and other countries with chickens by human agency. *Gallus gallus gallus* has been listed for the first time in the latest edition of the New Zealand Checklist of Birds (Checklist Committee 2010: 27; see also Palma 1999: 383, note 5).

Denny (1842: 155) clearly regarded *Goniodes* as a subgenus of *Philopterus*. However, subsequent authors ignored that fact and did not add parentheses around author and date when *Goniodes* is used at generic level.

***Goniodes ortygis* (Denny, 1842)**

- Philopterus (Goniodes) ortygis* Denny, 1842: 56, 158, pl. 13: fig. 6.  
*Goniodes ortygis* Denny, 1842 [sic]; Clay 1940b: 91, fig. 62.  
*Goniodes ortygis* Denny, 1842 [sic]; Hopkins & Clay 1952: 157.  
*Goniodes* sp.; Pilgrim & Palma 1982: 18.  
*Goniodes ortygis* Denny, 1842 [sic]; Palma 1999: 379.

*Goniodes ortygis* Denny, 1842 [sic]; Palma 2010: 408.

Lectotype ♂ in NHML (Clay 1940: 93).

Type host: *Colinus virginianus* (Linnaeus, 1758).

New Zealand host: *Colinus virginianus taylori* Lincoln, 1915.

Other hosts: *Colinus virginianus virginianus* (Linnaeus, 1758); *Colinus virginianus floridanus* (Coues, 1872); *Colinus virginianus texanus* (Lawrence, 1853).

New Zealand locality: WN.

Geographic distribution: North and Central America.

New Zealand references: Pilgrim & Palma (1982); Palma (1999); Palma (2010).

Other significant references: Clay (1940b); Forrester *et al.* (1995: 22); Price *et al.* (2003: 185).

Remarks: *Goniodes ortygis* was introduced to New Zealand with bobwhite quails by human agency, but these quails appear to have died out in New Zealand (Checklist Committee 2010: 349). Denny (1842: 155) clearly regarded *Goniodes* as a subgenus of *Philopterus*. However, subsequent authors ignored that fact and did not add parentheses around author and date when *Goniodes* is used at generic level.

### ***Goniodes pavonis* (Linnaeus, 1758)**

*Pediculus pavonis* Linnaeus, 1758: 613.

*Ricinus pavonis* (Linnaeus, 1758); Latreille 1804: 103.

*Goniodes pavonis* Linnaeus, 1758 [sic]; Harrison 1916: 78.

*Goniodes pavonis* (Linnaeus, 1758); Kéler 1939: 39, figs 11, 13–15.

*Goniodes pavonis* (Linnaeus, 1758); Clay 1940b: 5, figs 1a, 3, 4a, 5, 9a.

*Goniodes pavonis* (Linnaeus, 1758); Hopkins & Clay 1952: 157.

*Goniodes pavonis* (Linnaeus, 1758); Pigrim & Palma 1982: 18.

*Goniodes pavonis* (Linnaeus, 1758); Murray *et al.* 1993: 960.

*Goniodes pavonis* (Linnaeus, 1758); Palma 2010: 408.

Neotype ♂ in NHML (Clay 1940b: 7).

Type host: *Pavo cristatus* Linnaeus, 1758.

New Zealand host: *Pavo cristatus* Linnaeus, 1758.

Other host: *Pavo muticus* Linnaeus, 1766.

New Zealand localities: ND, WA, WN, NC, MC, SC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pigrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 220); Palma (2010).

Other significant references: Kéler (1939); Clay (1940b); Clay & Hopkins (1950: 261); Green & Palma (1991: 14, 31); Butler & O'Connor (1994: 455); Palma (1996b: 182); Price *et al.* (2003: 185); Adam (2007: 171); Martín-Mateo (2009: 46, figs 5e, 12).

Remarks: *Goniodes pavonis* is native to Asia, and was introduced to New Zealand and other countries with peafowl by human agency (Checklist Committee 2010: 28).

### ***Goniodes retractus* Le Souëf, 1902**

Figs 107–108

*Gonoides* [sic] *retractus* Le Souëf, 1902b: 90.

*Goniodes retractus* Le Souëf, 1902; Clay 1940b: 98, fig. 67.

*Goniodes retractus* Le Souëf, 1902; Hopkins & Clay 1952: 158.

*Goniodes retractus* Le Souëf, 1902; Pigrim & Palma 1982: 18.

*Goniodes retractus* Le Souëf, 1902; Murray *et al.* 1993: 961.

*Goniodes retractus* Le Souëf, 1902; Palma 2010: 408.

Holotype ♀, repository unknown (Palma 1996b: 183).

Type host: *Coturnix ypsilophora australis* (Latham, 1802).

New Zealand host: *Coturnix ypsilophora australis* (Latham, 1802).

Other hosts: None.

New Zealand localities: BP, HB.

Geographic distribution: Australasia.

New Zealand references: Pigrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Clay (1940b); Green & Palma (1991: 14, 31); Palma (1996b: 182); Price *et al.* (2003: 186).

Remarks: *Goniodes retractus* and its host are native to Australia, and were introduced to New Zealand by human agency (Checklist Committee 2010: 27).

### ***Goniodes stefani* Clay & Hopkins, 1955**

“*Goniodes mamillatus*” Taschenberg, 1882: 25, pl. 1: figs 1a,b (not *Goniodes mamillatus* Rudow, 1870: 483).

“*Goniodes mammillatus* [sic]” Kellogg, 1896: 509, pl. 69: fig. 2 (not *Goniodes mamillatus* Rudow, 1870: 483).

“*Gonocephalus mamillatus*” Kéler, 1939: 88, figs 45–46 (not *Goniodes mamillatus* Rudow, 1870: 483).

*Goniodes stefani* Clay & Hopkins 1955: 57. *Nomen novum* for “*Goniodes*” *mamillatus* Kéler, 1939.

*Goniodes stefani* Clay & Hopkins 1955; Pilgrim & Palma 1982: 18.

*Goniodes stefani* Clay & Hopkins 1955; Murray *et al.* 1993: 960.

*Goniodes stefani* Clay & Hopkins 1955; Palma 2010: 408.

Syntypes ♂♀ in ZMHG (Clay & Hopkins 1955: 57).

Type host: *Callipepla californica* (Shaw, 1798).

New Zealand host: *Callipepla californica brunnescens* (Ridgway, 1884).

Other hosts: None.

New Zealand localities: AK, BP, TK, SD, MB, NN, NC, MC, SC, CO, DN.

Geographic distribution: North & South America; Europe; Australasia, Hawaiian Islands.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Clay & Hopkins (1955); Price *et al.* (2003: 186).

Remarks: *Goniodes stefani* was introduced to New Zealand and other countries with California quails by human agency (Checklist Committee 2010: 25).

### **Genus *Haffneria* Timmermann, 1966**

*Haffneria* Timmermann, 1966. *Mitt. Hamburg. Zool. Mus. Inst.* 63: 87. Type species: *Perineus piratae* Timmermann, 1955 (by original designation).

#### ***Haffneria grandis* (Piaget, 1880)**

Figs 109–110

*Lipeurus grandis* Piaget, 1880: 323, pl. 26: fig. 7.

*Lipeurus laculatus* Kellogg & Chapman, 1899: 93, pl. 7: fig. 1.

*Esthiopterum grande* Piaget, 1880 [sic]; Harrison 1916: 135 (as junior synonym of *Esthiopterum modestum* Giebel, 1874).

“*Perineus modestus*” Séguy, 1944: 365, figs 544–545 (not *Lipeurus modestus* Giebel, 1874).

*Perineus grandis* (Piaget, 1880); Hopkins & Clay 1952: 277.

*Perineus piratae* Timmermann, 1955: 532, fig. 16.

*Perineus laculatus* (Kellogg & Chapman, 1899); Timmermann 1955: 532.

*Diomedicola grandis* (Piaget, 1880); Kéler 1957b: 509, figs 1d, 2a,b, 5, 11–12.

*Perineus antarcticus* Carriker, 1958: 186, fig. 7.

*Harrisoniella grandis* (Piaget, 1880); Clay 1964a: 231.

*H. grandis* (Piaget, 1880); Gressitt 1964: 538.

*Haffneria grandis* (Piaget, 1880); Timmermann 1966: 86, figs 1a, 3.

*Haffneria piratae* (Timmermann, 1955); Timmermann 1966: 87, fig. 2a.

*Haffneria grandis* (Piaget, 1880); Clay & Moreby 1967: 161, 169, figs 82–83.

*Haffneria grandis* (Piaget, 1880); Clay & Moreby 1970: 220.

“*H.*” *grandis* (Piaget, 1880); Gressitt 1970: 327.

*Haffneria grandis* (Piaget, 1880); Wise 1977: 60.

*Haffneria grandis* (Piaget, 1880) *s. l.*; Pilgrim & Palma 1982: 22.

*Haffneria grandis* (Piaget, 1880); Murray *et al.* 2006a: 1964.

*Haffneria grandis* (Piaget, 1880); Palma 2010: 408.

Lectotype ♂ in NHML (Timmermann, 1955: 534). Holotype nymph of *Perineus antarcticus* in FMLA.



Type host: “*Hydrobates pelagicus*”, in error (see Hopkins & Clay 1952: 277).

New Zealand hosts: *Catharacta antarctica lonnbergi* Mathews, 1912; *Catharacta maccormicki* (Saunders, 1893); *Stercorarius parasiticus* (Linnaeus, 1758).

Other hosts: *Catharacta antarctica hamiltoni* Hagen, 1952; *Catharacta chilensis* (Bonaparte, 1857); *Catharacta skua* Brünnich, 1764; *Coprotheres pomarinus* (Temminck, 1815); *Stercorarius longicaudus* Vieillot, 1819.

New Zealand localities: ND, WI, CH, SI, SN, AU, CA, Macquarie Island, RO.

Geographic distribution: Cosmopolitan.

New Zealand references: Clay (1964a); Gressitt (1964); Watson (1967: 72); Clay & Moreby (1967; 1970); Gressitt (1970); Spellerberg (1971: 19); Schaefer & Strandtmann (1971: 16); Wise (1977); Horning *et al.* (1980: 5, 11); Pilgrim & Palma (1982); Palma (1996b: 183); Palma & Horning (2002: 8, 17); Murray *et al.* (2006a); Palma (2010).

Other significant references: Hopkins (1942b: 100); Séguy (1944); Hopkins & Clay (1952); Séguy (1953: 568, figs 18–20); Kéler (1957b); Timmermann (1966); Furness & Palma (1992: 35, 42); Cohen *et al.* (1997: 186); Ramli *et al.* (2000: 71); Price *et al.* (2003: 186); Page *et al.* (2004: 648, 650); Palma & Jensen (2005: 55, 64); Hänel & Palma (2007: 112, 124, 131).

Remarks: Pilgrim & Palma (1982: 22) regarded the populations of *Haffneria grandis* from the three New Zealand hosts listed above as somewhat different from the lectotype, and qualified them as *sensu lato*. However, my examination of more samples, including some from non-New Zealand hosts, shows that making such difference is not warranted.

### Genus *Halipeurus* Thompson, 1936

*Halipeurus* Thompson, 1936. *Ann. Mag. Nat. Hist.* (Ser. 10) 18: 40. Type species: *Lipeurus angusticeps* Piaget, 1880 (by original designation).

*Synnautes* Thompson, 1936. *Ann. Mag. Nat. Hist.* (Ser. 10) 18: 43. Type species: *Lipeurus pelagicus* Denny, 1842 (by original designation).

*Anamias* Timmermann, 1965. *Abhandl. Verhandl. Naturwiss. Vereins Hamburg, N.F.* 8 (Supplement): 155. Type species: *Halipeurus raphanus* Timmermann, 1961a (by original designation).

*Halipeurus* Thompson, 1936; Palma 2011b: 16. Synonymy.

### *Halipeurus angusticeps* (Piaget, 1880)

*Lipeurus angusticeps* Piaget, 1880: 306, pl. 25: fig. 4.

*Esthiopteris angusticeps* Piaget, 1880 [sic]; Harrison 1916: 130.

*Halipeurus angusticeps* (Piaget, 1880); Hopkins & Clay 1952: 163.

*Halipeurus sawadai* Nakagawa, 1959a: 384, figs 1a,b,c, 2b,d.

*Halipeurus (Halipeurus) angusticeps angusticeps* (Piaget, 1880); Edwards 1961: 135, figs 3a–7a.

*Halipeurus (Halipeurus) angusticeps fosteri* Edwards, 1961: 137, figs 3b–7b.

*Halipeurus (Halipeurus) angusticeps* (Piaget, 1880); Timmermann 1965: 139.

*Halipeurus (Halipeurus) sawadai* Nakagawa, 1959; Price *et al.* 2003: 188.

*Halipeurus angusticeps* (Piaget, 1880); Scofield *et al.* 2011: 214.

*Halipeurus angusticeps* (Piaget, 1880); Palma 2011b: 19, figs 38–40, 62.

Lectotype in NHML (Palma 2011b: 19).

Type host: “*Procellaria cinerea*”, in error (see Palma 2011b: 21).

New Zealand host: *Calonectris leucomelas* (Temminck, 1836).

Other hosts: None.

New Zealand locality: WO.

Geographic distribution: Japan; Korea; eastern China; Pacific Ocean.

New Zealand references: Scofield *et al.* (2011); Palma (2011b).

Other significant references: Timmermann (1961a: 402); Edwards (1961); Timmermann (1965); Tsurumi (1989: 281); Price *et al.* (2003: 187).

Remarks: *Halipeurus angusticeps* is a “wing” louse exclusively parasitic on the streaked shearwater, *Calonectris leucomelas*, a host recorded only once in New Zealand (Checklist Committee 2010: 111; Scofield *et al.* 2011). Wise (1977: 60) listed “*Halipeurus (Halipeurus) angusticeps*” from Macquarie Island, but Pilgrim & Palma (1982: 30, note 16) showed that it was a misidentification of *H. procellariae* (see below).

#### ***Halipeurus bulweriae* Timmermann, 1960**

*Halipeurus* sp.; Clay 1940a: 309.

*Naubates* sp.; Clay 1940a: pl. 1: fig. 3.

*Halipeurus bulweriae* Timmermann, 1960: 328, figs 11, 15a.

*Halipeurus (Halipeurus) bulweriae* Edwards, 1961: 145, figs 3h–7h.

*Halipeurus (Halipeurus) bulweriae* Timmermann, 1960; Timmermann 1965: 149, fig. 90.

*Halipeurus (Halipeurus) bulweriae* Timmermann, 1960; Palma 1999: 376, 383, note 3.

*Halipeurus (Halipeurus) bulweriae* Timmermann, 1960; Palma 2010: 408.

*Halipeurus bulweriae* Timmermann, 1960; Palma 2011b: 17.

Holotype ♂ in NHML.

Type host: *Bulweria bulwerii* (Jardine & Selby, 1828).

New Zealand host: *Bulweria bulwerii* (Jardine & Selby, 1828).

Other hosts: None.

New Zealand localities: WN, MC.

Geographic distribution: Temperate and tropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Palma (1999); Palma (2010); Palma (2011b).

Other significant references: Clay (1940a); Edwards (1961); Timmermann (1965); Amerson & Emerson (1971: 5, 23); Price *et al.* (2003: 187).

Remarks: *Halipeurus bulweriae* is a “wing” louse exclusively parasitic on Bulwer’s petrel, a host which, with a single record until 2013, is regarded as a very rare straggler to New Zealand (Checklist Committee 2010: 107). A second record occurred in North Brighton, Canterbury, in January 2014. Both specimens were parasitised by *Halipeurus bulweriae*.

#### ***Halipeurus confusus* Palma, 2011**

Figs 111–112

*Halipeurus (Halipeurus) accentor* Edwards, 1961: 151, figs 3r–7r. In part.

“*Halipeurus accentor*” Nelson, 1969: 199 (not *Halipeurus accentor* Edwards, 1961).

“*Halipeurus leucophryna*” Watt, 1971: 236, 242 (not *Halipeurus leucophryna* Timmermann, 1960).

“*Halipeurus (Halipeurus) leucophryna*” Wise, 1977: 61 (not *Halipeurus leucophryna* Timmermann, 1960).

*Halipeurus (Halipeurus)* sp.; Pilgrim & Palma 1982: 9, 30.

*Halipeurus (Halipeurus)* sp.; Murray *et al.* 1990: 1370.

*Halipeurus* sp.; Palma 2010: 408.

*Halipeurus confusus* Palma, 2011b: 3, figs 1–3, 28, 43, 57.

Holotype ♂ in MONZ.

Type host: *Pterodroma nigripennis* (Rothschild, 1893).

New Zealand host: *Pterodroma nigripennis* (Rothschild, 1893).

Other hosts: None.

New Zealand localities: ND, AK, BP, GB, NC, MC, SC, KE, Norfolk Island, CH.

Geographic distribution: Pacific Ocean.

New Zealand references: Nelson (1969); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010); Palma (2011b).

Other significant reference: Edwards (1961).

Remarks: *Halipeurus confusus* is a “wing” louse exclusively parasitic on black-winged petrels. This *Halipeurus* species was initially misidentified by Edwards (1961) and confused with other species by several authors for many years (see Palma 2011b: 3, 6).

***Halipeurus consimilis* Timmermann, 1960**

*Halipeurus consimilis* Timmermann, 1960: 326, fig. 9.

*Halipeurus (Halipeurus) consimilis* Timmermann, 1960; Timmermann 1965: 147, fig. 88.

*Halipeurus (Halipeurus) consimilis* Timmermann, 1960; Palma & Pilgrim 1977: 290, figs 1–6.

*Halipeurus consimilis* Timmermann, 1960; Horning *et al.* 1980: 5, 9.

*Halipeurus (Halipeurus) consimilis* Timmermann, 1960; Pilgrim & Palma 1982: 8.

*Halipeurus (H.) consimilis* Timmermann, 1960; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Pterodroma inexpectata* (J.R. Forster, 1844).

New Zealand host: *Pterodroma inexpectata* (J.R. Forster, 1844).

Other hosts: None.

New Zealand localities: ND, AK, WN, NC, MC, SC, WD, SL, SI, SN.

Geographic distribution: Pacific Ocean.

New Zealand references: Timmermann (1960); Horning *et al.* (1980); Palma & Pilgrim (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1370); Page *et al.* (2004: 643, 648); Palma (2010).

Other significant references: Timmermann (1965); Palma (1996b: 184); Price *et al.* (2003: 187); Hammer *et al.* (2010: 1113).

Remarks: *Halipeurus consimilis* is a “wing” louse exclusively parasitic on the mottled petrel, a host which only breeds in New Zealand but which ranges widely over the Pacific Ocean (Checklist Committee 2010: 93).

***Halipeurus diversus* (Kellogg, 1896)**

*Lipeurus diversus* Kellogg, 1896a: 123, pl. 8: figs 3–4.

*Lipeurus limitatus* Kellogg, 1896a: 124, pl. 8: fig. 5, 6.

*Esthiopterum diversum* Kellogg, 1896 [sic]; Harrison 1916: 133.

*Esthiopterum constrictiventre* Pessôa & Guimarães, 1935a: 313, fig. 6, 7.

*Halipeurus diversus* (Kellogg, 1896); Hopkins & Clay 1952: 163.

*Halipeurus hanáki* [sic] Balát, 1958: 415.

*Halipeurus diversus* (Kellogg, 1896); Timmermann 1961a: 408, fig. 6.

*Halipeurus (Halipeurus) diversus* (Kellogg, 1896); Edwards, 1961: 142, figs 3f–7f.

*Halipeurus* sp.; Clay 1964a: 231.

*Halipeurus* sp.; Gressitt 1964: 538.

*Halipeurus (Halipeurus) diversus* Timmermann, 1965: 142, fig. 83.

*Halipeurus diversus hanáki* [sic] Balát, 1958; Timmermann 1965: 142.

“*Halipeurus turtur*” Watson, 1967: 72 (not *Halipeurus turtur* Edwards, 1961).

“*Halipeurus turtur*” Gressitt, 1970: 327 (not *Halipeurus turtur* Edwards, 1961).

*Halipeurus (Halipeurus) diversus* (Kellogg, 1896); Wise 1977: 60.

*Halipeurus diversus* (Kellogg, 1896); Horning *et al.* 1980: 5, 10.

*Halipeurus (Halipeurus) diversus* (Kellogg, 1896); Pilgrim & Palma 1982: 12.

*Halipeurus (Halipeurus) diversus* (Kellogg, 1896); Palma & Horning 2002: 8, 15, 20.

*Halipeurus (H.) diversus* (Kellogg, 1896); Palma 2010: 408.

*Halipeurus diversus* (Kellogg, 1896); Palma 2011b: 32.

Syntypes ♂♀ in USNM and EMEC (Emerson 1961b: 251; Palma 2011b: 32).

Type host: “*Puffinus opisthomelas*”, in error (see Hopkins & Clay 1952: 163).

New Zealand hosts: *Puffinus griseus* (J.F. Gmelin, 1789); *Puffinus tenuirostris* (Temminck, 1835); *Puffinus puffinus* (Brünnich, 1764).

Other hosts: *Puffinus yelkouan* (Acerbi, 1827); *Puffinus mauretanicus* Lowe, 1921; *Puffinus assimilis baroli* (Bonaparte, 1857); *Puffinus assimilis boydi* Mathews, 1912.

New Zealand localities: AK, BP, WI, WA, WN, NC, MC, SC, WD, CO, DN, CH, SI, SN, AN, Macquarie Island.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Edwards (1961); Clay (1964a); Gressitt (1964); Watson (1967); Clay & Moreby (1970: 218); Gressitt (1970); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Tennyson (1986: 60); Murray *et al.*

(1990: 1371); Palma (1996b: 184); Palma (1999: 384); Palma & Horning (2002); Page *et al.* (2004: 643, 648); Palma (2010); Hammer *et al.* (2010: 1113); Palma (2011b).

Other significant references: Timmermann (1961a; 1965); Fowler & Miller (1984: 24, 27, fig. 1); Fowler & Shaw (1990: 15); Green & Palma (1991: 14, 27); Price *et al.* (2003: 187); Palma & Jensen (2005: 56, 60); Palma & Peck (2013: 38).

Remarks: *Halipeurus diversus* is a “wing” louse with a range of variable dimensions, a feature congruent with having a large number of host species. However, size differences only do not justify subdividing it into more than one taxon. The record of “*Halipeurus turtur* Edwards, 1961” in Watson (1967: 72), repeated in subsequent publications, is a misidentification of *H. diversus* (see Palma 1999: 384, note F).

#### ***Halipeurus falsus pacificus* Edwards, 1961**

*Halipeurus (Halipeurus) falsus pacificus* Edwards, 1961: 147, figs 3J2–7J2.

*Halipeurus (Halipeurus) falsus pacificus* Edwards, 1961: Timmermann 1965: 144, fig. 85.

*Halipeurus falsus pacificus* Edwards, 1961; Pilgrim 1970: 75.

*Halipeurus falsus pacificus* Edwards, 1961; Horning *et al.* 1980: 6, 11.

*Halipeurus (Halipeurus) falsus pacificus* Edwards, 1961; Pilgrim & Palma 1982: 13.

*Halipeurus (Halipeurus) falsus pacificus* Edwards, 1961; Murray *et al.* 1990: 1372.

*Halipeurus (H.) falsus pacificus* Edwards, 1961; Palma 2010: 408.

Holotype ♂ in AMNH (Palma 1996b: 185).

Type host: *Pelecanoides urinatrix* (J.F. Gmelin, 1789).

New Zealand hosts: *Pelecanoides urinatrix urinatrix* (J.F. Gmelin, 1789); *Pelecanoides urinatrix chathamensis* Murphy & Harper, 1916; *Pelecanoides urinatrix exsul* Salvin, 1896.

Other hosts: *Pelecanoides urinatrix dacunhae* Nicholl, 1906; *Pelecanoides magellani* (Mathews, 1912).

New Zealand localities: ND, CL, BP, GB, TO, WI, WA, WN, NC, MC, SC, WD, CO, DN, FD, CH, SN, AN.

Geographic distribution: South Atlantic and Pacific Oceans.

New Zealand references: Edwards (1961: 147); Pilgrim (1970); Wise (1977: 60); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Marris (2000: 187); Palma (2010).

Other significant references: Timmermann (1965); Green & Palma (1991: 14, 28); Furness & Palma (1992: 35, 41); Palma (1996b: 185); Price *et al.* (2003: 187); Page *et al.* (2004: 638, 648); Hänel & Palma (2007: 112, 124, 131).

Remarks: Both subspecies of *Halipeurus falsus* are “wing” lice and the smallest members of the genus, exclusively parasitic on diving petrels (Family Pelecanoididae).

#### ***Halipeurus gravis priapulus* Timmermann, 1961**

*Halipeurus priapulus* Timmermann, 1961a: 406, fig. 3.

*Halipeurus (Halipeurus) micariproctus* Edwards, 1961: 148, figs 3I–7I.

? *Halipeurus priapulus* Timmermann, 1961; Timmermann 1965: 141 (as a questionable junior synonym of *Halipeurus (H.) gravis* Timmermann, 1961a).

“*Halipeurus (Halipeurus) gravis*” Wise, 1977: 60 (not *Halipeurus gravis gravis* Timmermann, 1961a).

*Halipeurus (Halipeurus) gravis priapulus* Timmermann, 1961; Pilgrim & Palma 1982: 11.

*Halipeurus (Halipeurus) gravis priapulus* Timmermann, 1961; Murray *et al.* 1990: 1371.

*Halipeurus (Halipeurus) priapulus* Timmermann, 1961; Price *et al.* 2003: 188.

*Halipeurus (H.) gravis priapulus* Timmermann, 1961; Palma 2010: 408.

Holotype ♂ in NHML. Holotype ♂ of *Halipeurus (Halipeurus) micariproctus* in AMNH (Palma 1996b: 185).

Type host: *Puffinus carneipes* Gould, 1844.

New Zealand host: *Puffinus carneipes* Gould, 1844.

Other host: *Puffinus creatopus* Coues, 1864.

New Zealand localities: ND, AK, CL, BP, WN.

Geographic distribution: Pacific Ocean.

New Zealand references: Edwards (1961); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Page *et al.* (2004: 648, 650); Palma (2010).

Other significant references: Timmermann (1965); Palma (1996b: 185); Price *et al.* (2003); Hammer *et al.* (2010: 1113).

Remarks: Contrary to Price *et al.* (2003: 188), I regard this louse taxon as a subspecies. The two subspecies of *Halipeurus gravis* are “wing” lice and morphologically extremely similar: females are identical, and males only differ in the shape of the mesosomal sclerite of the genitalia (“Hakensklerit” in Timmermann 1961a: figs 2–3).

### ***Halipeurus kermadecensis* (Johnston & Harrison, 1912)**

*Lipeurus kermadecensis* Johnston & Harrison, 1912: 365, fig. 1.

*Lipeurus diversus* var. *excavatus* Johnston & Harrison, 1912: 366, fig. 2.

*Esthiopterum excavatum* Johnston & Harrison, 1912 [sic]; Harrison 1916: 133 (as junior synonym of *Esthiopterum kermadecense* Johnston & Harrison, 1912 [sic]).

*Esthiopterum kermadecense* Johnston & Harrison, 1912 [sic]; Harrison 1916: 136.

*Halipeurus kermadecense* [sic] (Johnston & Harrison, 1912); Thompson 1938d: 488.

*Halipeurus kermadecensis* (Johnston & Harrison, 1912); Hopkins & Clay 1952: 164.

*Halipeurus kermadecensis* (Johnston & Harrison, 1912); Timmermann 1960: 327, fig. 8.

*Halipeurus (Halipeurus) kermadecense* [sic] (Johnston & Harrison, 1912); Edwards, 1961: 150, figs 3q–7q.

*Halipeurus (Halipeurus) kermadecensis* (Johnston & Harrison, 1912); Timmermann 1965: 137, fig. 87.

*Halipeurus kermadecensis* (Johnston & Harrison, 1912); Watt 1971: 236, 242, fig. 5.

*Halipeurus (Halipeurus) kermadecensis* (Johnston & Harrison, 1912) *s. l.*; Pilgrim & Palma 1982: 8.

*Halipeurus (Halipeurus) kermadecensis* (Johnston & Harrison, 1912); Pilgrim & Palma 1982: 9.

*Halipeurus (H.) kermadecensis* (Johnston & Harrison, 1912); Palma 2010: 408.

*Halipeurus kermadecensis* (Johnston & Harrison, 1912); Palma 2011b: 29.

Holotype nymph in MONZ (Palma *et al.* 1989: 45). Syntypes ♂♀ of *Lipeurus diversus excavatus* in MONZ (Palma *et al.* 1989: 45).

Type host: *Pterodroma neglecta neglecta* (Schlegel, 1863).

New Zealand hosts: *Pterodroma externa* (Salvin, 1875); *Pterodroma neglecta neglecta* (Schlegel, 1863).

Other hosts: *Pterodroma neglecta juana* Mathews, 1935; *Pterodroma arminjoniana arminjoniana* (Giglioli & Salvadori, 1869); *Pterodroma sandwichensis* (Ridgway, 1884).

New Zealand localities: ND, AK, CL, KE.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Johnston & Harrison (1912); Thompson (1938d); Thompson (1939: 123); Pilgrim (1970: 75); Watt (1971); Wise (1977: 60); Pilgrim & Palma (1982); Palma *et al.* (1989: 45); Murray *et al.* (1990: 1369); Palma (2001: 67, fig. 8); Palma (2010); Palma (2011b).

Other significant references: Thompson (1938d); Timmermann (1960; 1965); Edwards (1961); Timmermann (1965); Price *et al.* (2003: 187); Hammer *et al.* (2010: 1113).

Remarks: *Halipeurus kermadecensis* is a “wing” louse parasitic on several gadfly petrel species. The main New Zealand population of *H. kermadecensis* is in the Kermadec Islands, where the type host breeds. Records of *H. kermadecensis* from *Pterodroma externa* were qualified as *sensu lato* by Pilgrim & Palma (1982: 8); however, my examination of more samples from that and three other host species (see above) showed that such qualification is not warranted.

### ***Halipeurus leucophryna* Timmermann, 1960**

*Halipeurus leucophryna* Timmermann, 1960: 327, fig. 10.

*Halipeurus (Halipeurus) accentor* Edwards, 1961: 151.

*Halipeurus (Halipeurus) leucophryna* Timmermann, 1960; Timmermann 1965: 148, fig. 89.

*Halipeurus (Halipeurus) leucophryna* Timmermann, 1960; Pilgrim & Palma 1982: 9.

*Halipeurus (Halipeurus) leucophryna* Timmermann, 1960; Murray *et al.* 1990: 1370.

*Halipeurus (H.) leucophryna* Timmermann, 1960; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Pterodroma longirostris* (Stejneger, 1888).

New Zealand hosts: *Pterodroma longirostris* (Stejneger, 1888); *Pterodroma pycrofti* Falla, 1933.

Other host: *Pterodroma defilippiana* (Giglioli & Salvadori, 1869).

New Zealand localities: ND, CL, AK, BP.

Geographic distribution: Pacific Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Timmermann (1965); Ward & Downey (1973: 394); Price *et al.* (2003: 187); Palma (2011b: 6).

Remarks: *Halipeurus leucophryna* is a “wing” louse parasitic on several gadfly petrel species. Records of “*Halipeurus leucophryna*” in Watt (1971: 236) and “*Halipeurus (Halipeurus) leucophryna*” in Wise (1977: 61) are misidentifications. See above under *Halipeurus confusus* Palma, 2011.

### ***Halipeurus marquesanus* (Ferris, 1932)**

*Esthiopterum marquesanum* Ferris, 1932a: 62, figs 14a,b, 15a,b,c,d,e,f.

*Halipeurus marquesanus* (Ferris, 1932); Thompson 1936: 41.

*Halipeurus marquesanus* (Ferris, 1932); Hopkins & Clay 1952: 164.

*Halipeurus marquesanus* (Ferris, 1932); Timmermann 1960: 329, fig. 6a.

*Halipeurus (Halipeurus) marquesanus* (Ferris, 1932); Edwards 1961: 152, figs 3t–7t.

*Halipeurus (Halipeurus) marquesanus* (Ferris, 1932); Timmermann 1965: 151, fig. 77b.

*Halipeurus (Halipeurus) marquesanus* (Ferris, 1932); Palma 1999: 376.

*Halipeurus (H.) marquesanus* (Ferris, 1932); Palma 2010: 408.

Holotype ♀ in EMEC (Peter T. Oboyski pers. comm. October 2014).

Type host: *Pseudobulweria rostrata* (Peale, 1848).

New Zealand host: *Pseudobulweria rostrata* (Peale, 1848).

Other hosts: None.

New Zealand locality: ND.

Geographic distribution: South Pacific Ocean.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Thompson (1936); Timmermann (1960; 1965); Edwards (1961); Timmermann (1965); Price *et al.* (2003: 187).

Remarks: *Halipeurus marquesanus* is a “wing” louse exclusively parasitic on the Tahiti petrel. *Pseudobulweria rostrata* is a rare vagrant to New Zealand (Checklist Committee 2010: 110), with a single record of *H. marquesanus* from this country.

### ***Halipeurus mirabilis* Thompson, 1940**

*Halipeurus mirabilis* Thompson, 1940b: 499, text figs 1–5, pl. 10: figs 1–2.

*Halipeurus (Halipeurus) mirabilis* Thompson, 1940; Timmermann 1961a: 407, fig. 4.

*Halipeurus (Halipeurus) mirabilis* Thompson, 1940; Edwards 1961: 139, figs 3d–7d.

*Halipeurus (Halipeurus) mirabilis* Thompson, 1940; Timmermann 1965: 141, fig. 82.

*Halipeurus mirabilis* Thompson, 1940; Watt 1971: 236, 243.

*Halipeurus (Halipeurus) mirabilis* Thompson, 1940; Wise 1977: 61.

*Halipeurus (Halipeurus) mirabilis* Thompson, 1940; Pilgrim & Palma 1982: 11.

*Halipeurus (Halipeurus) mirabilis* Thompson, 1940; Murray *et al.* 1990: 1371.

*Halipeurus (H.) mirabilis* Thompson, 1940; Palma 2010: 408.

Syntypes ♂♀ in NHML (Vincent S. Smith pers. comm. July 2014).

Type host: *Puffinus pacificus chlororhynchus* Lesson, 1831.

New Zealand hosts: *Puffinus pacificus pacificus* (J.F. Gmelin, 1789); *Puffinus pacificus chlororhynchus* Lesson, 1831.

Other hosts: None.

New Zealand localities: ND, WI, WN, KE.

Geographic distribution: Indian and Pacific Oceans.

New Zealand references: Edwards (1961); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 222); Palma (2010).

Other significant references: Timmermann (1961a; 1965); Amerson & Emerson (1971: 6, 23); Ward & Downey (1973: 394); Palma (1996b: 185); Price *et al.* (2003: 187).

Remarks: *Halipeurus mirabilis* is a “wing” louse parasitic on all subspecies of the wedge-tailed shearwater. Palma (1996b: 185) assumed incorrectly that there was a holotype designated by Thompson (1940b: 499), but the latter author failed to make such a designation. In his original description of *H. mirabilis*, Thompson (1940b: 499) only stated “♀ and ♂ types off the type host ...” and then proceeded to designate 18 “paratypes”, but without mentioning either a “holotype” or a single “type”.

#### ***Halipeurus munda* Edwards, 1961**

- Halipeurus (Halipeurus) munda* Edwards, 1961: 149, figs 3n–7n.  
*Halipeurus (Halipeurus) munda* Edwards, 1961; Timmermann 1965: 146.  
*Halipeurus (Halipeurus) munda* Edwards, 1961; Pilgrim & Palma 1982: 12.  
*Halipeurus (Halipeurus) munda* Edwards, 1961; Murray *et al.* 1990: 1371.  
*Halipeurus (Halipeurus) munda* Edwards, 1961; Marris 2000: 187.  
*Halipeurus (H.) munda* Edwards, 1961; Palma 2010: 408.

Holotype ♂ in AMNH.

Type host: *Puffinus elegans* Giglioli & Salvadori, 1869.

New Zealand hosts: *Puffinus assimilis haurakiensis* Fleming & Serventy, 1943; *Puffinus elegans* Giglioli & Salvadori, 1869.

Other hosts: None.

New Zealand localities: ND, AK, CL, BP, WD, CH, AN.

Geographic distribution: Southern Pacific, Atlantic and Indian Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Marris (2000); Palma (2010).

Other significant references: Timmermann (1965); Furness & Palma (1992: 35, 39); Price *et al.* (2003: 187); Hänel & Palma (2007: 112, 124, 129).

Remarks: *Halipeurus munda* is a “wing” louse parasitic on some of the smaller shearwater species. Both *H. munda* and *H. placodus* (see below) are useful “diagnostic” species to confirm the identity of hosts, especially juveniles and beach-wrecked remains of specimens belonging to the *Puffinus assimilis* species complex (see Holdaway *et al.* 2001: 127).

#### ***Halipeurus noctivagus* Timmermann, 1960**

- “*Lipeurus diversus* var. *major*” Kellogg & Kuwana, 1902: 477 (not *Lipeurus major* Piaget, 1880).  
*Halipeurus noctivagus* Timmermann, 1960: 331, fig. 13, 16b.  
*Halipeurus (Halipeurus) intermedius* Edwards, 1961: 151, figs 3s–7s.  
*Halipeurus (Halipeurus) noctivagus* Timmermann, 1960; Timmermann 1965: 151, fig. 92.  
*Halipeurus (Halipeurus) noctivagus* Timmermann, 1960; Wise 1977: 61.  
*Halipeurus (Halipeurus) noctivagus* Timmermann, 1960; Pilgrim & Palma 1982: 8.  
*Halipeurus (Halipeurus) noctivagus* Timmermann, 1960; Price *et al.* 2003: 188.  
*Halipeurus (H.) noctivagus* Timmermann, 1960; Palma 2010: 408.  
*Halipeurus noctivagus* Timmermann, 1960; Palma 2011b: 41, fig. 53.

Holotype ♂ in NHML.

Type host: *Pterodroma phaeopygia* (Salvin, 1876).

New Zealand host: *Pterodroma cervicalis* (Salvin, 1891).

Other host: *Pterodroma occulta* Imber & Tennyson, 2001.

New Zealand localities: AK, KE.

Geographic distribution: Pacific Ocean.

New Zealand references: Edwards (1961: 152); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1370); Palma (2010: 408); Palma (2011b: 42).

Other significant references: Timmermann (1965); Imber & Tennyson (2001: 125); Price *et al.* (2003); Palma & Peck (2013: 38).

Remarks: *Halipeurus noctivagus* is a “wing” louse parasitic on several gadfly petrel species. There is now sufficient evidence, both genetic (Browne *et al.* 1997) and parasitological (Palma 2011b: 30), to treat *Pterodroma phaeopygia*—the type host of *H. noctivagus*—as a full species.

***Halipeurus pelagicus* (Denny, 1842)**

- Philopterus (Lipeurus) pelagicus* Denny, 1842: 58, 173, pl. 14: fig. 2.  
*Lipeurus subangusticeps* Piaget, 1880: 308, pl. 25: fig. 5.  
*Lipeurus languidus* Kellogg & Kuwana, 1902: 475, pl. 29: fig. 8.  
*Lipeurus exiguus* Kellogg & Kuwana, 1902: 479, pl. 30: fig. 2.  
*Esthiopterum pelagicum* Denny, 1842 [sic]; Harrison 1916: 139.  
*Synnautes pelagicus* (Denny, 1842); Thompson 1939: 209.  
*Halipeurus languidus* (Kellogg & Kuwana, 1902); Hopkins & Clay 1952: 164.  
*Halipeurus pelagicus* (Denny, 1842); Hopkins & Clay 1952: 164.  
*Halipeurus subangusticeps* (Piaget, 1880); Hopkins & Clay 1952: 164.  
*Halipeurus pelagicus* (Denny, 1842); Timmermann 1961a: 413, figs 9–10. In part.  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842); Edwards, 1961: 155, figs 2b, 3v–7v. In part.  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842); Timmermann 1965: 153, fig. 94, pl. 7: figs 3–4.  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842) s. l.; Pilgrim & Palma 1982: 13. In part.  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842); Murray *et al.* 1990: 1372. In part.  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842); Marris 2000: 187.  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842); Palma 2010: 408. In part.  
*Halipeurus pelagicus* (Denny, 1842); Palma 2011b: 27, figs 7, 23, 25, 33, 48.

Lectotype ♀ in NHML (Palma 2011b: 29).

Type host: *Hydrobates pelagicus* (Linnaeus, 1758).

New Zealand hosts: *Fregetta grallaria* (Vieillot, 1817); *Fregetta tropica* (Gould, 1844); *Oceanodroma leucorhoa* (Vieillot, 1817).

Other hosts: *Oceanites oceanicus oceanicus* (Kuhl, 1820); *Oceanites gracilis gracilis* (Elliot, 1859); eight other species of *Oceanodroma* (see Edwards 1961: 156; Palma 2011b: 28).

New Zealand localities: ND, CL, KE, AN.

Geographic distribution: Cosmopolitan.

New Zealand references: Thompson (1939: 210); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999: 378); Marris (2000); Palma (2010); Palma (2011b: 28).

Other significant references: Séguy (1953: 576, figs 36–37); Edwards (1961); Timmermann (1961a; 1965); Clay & Moreby (1967: 160, 169, fig. 87); Fowler *et al.* (1984: 126, figs 1–2); Benoit (1976: 234); Fowler & Miller (1984: 24, figs 1–2a); Fowler & Price (1987: 44); Fowler & Hodson (1988: 48); Forrester *et al.* (1995: 6); Palma (1996b: 186); Furness & Palma (1992: 35, 41); Price *et al.* (2003: 188); Page *et al.* (2004: 648, 650); Palma & Jensen (2005: 56, 60); Hänel & Palma (2007: 112, 124, 130); Hammer *et al.* (2010: 1113); Palma & Peck (2013: 39).

Remarks: *Halipeurus pelagicus* is a “wing” louse parasitic on many storm petrel species. *Halipeurus pelagicus* was placed in the subgenus *Synnautes* Thompson, 1936 in many publications, but Palma (2011b: 16) formally synonymised this subgenus under the genus *Halipeurus*.

***Halipeurus pelagodromae* Palma, 2011**

- “*Lipeurus languidus*” Johnston & Harrison, 1912: 367 (not *Lipeurus languidus* Kellogg & Kuwana, 1902).  
 “*Lipeurus exiguus*” Johnston & Harrison, 1912: 367 (not *Lipeurus exiguus* Kellogg & Kuwana, 1902).  
*Halipeurus pelagicus* (Denny, 1842); Timmermann 1961a: 413. In part.  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842); Edwards 1961: 155. In part.  
 “*Halipeurus pelagicus*” Watt, 1971: 236, 243 (not *Philopterus (Lipeurus) pelagicus* Denny, 1842).  
 “*Halipeurus (Synnautes) pelagicus*” Wise, 1977: 61 (not *Philopterus (Lipeurus) pelagicus* Denny, 1842).  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842) s. l.; Pilgrim & Palma 1982: 13. In part.  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842); Murray *et al.* 1990: 1372. In part.  
 “*H. (Synnautes) pelagicus*” Green & Palma, 1991: 14 (not *Philopterus (Lipeurus) pelagicus* Denny, 1842).  
 “*Halipeurus pelagicus*” Paterson *et al.*, 1999: 222 (not *Philopterus (Lipeurus) pelagicus* Denny, 1842).  
*Halipeurus (Synnautes) pelagicus* (Denny, 1842); Palma 2010: 408. In part.  
*Halipeurus pelagodromae* Palma, 2011b: 11, figs 6, 22, 24, 26–27, 32, 47, 63.

Holotype in MONZ.

Type host: *Pelagodroma marina maoriana* Mathews, 1912.



New Zealand hosts: *Pelagodroma marina maoriana* Mathews, 1912; *Pelagodroma marina dulciae* Mathews, 1912; *Pelagodroma albiclunis* Murphy & Irving, 1951.

Other hosts: *Pelagodroma marina marina* (Latham, 1790); *Pelagodroma marina hypoleuca* (Moquin-Tandon, 1841); *Pelagodroma marina eadesi* Bourne, 1953.

New Zealand localities: ND, CL, WA, KE, CH, AU.

Geographic distribution: Pacific and Atlantic Oceans.

New Zealand references: Johnston & Harrison (1912); Edwards (1961); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999); Palma (2010); Palma (2011b: 15).

Other significant references: Green & Palma (1991); Furness & Palma (1992: 35, 42); Palma (1996b: 186); Page *et al.* (2004: 648, 650); Hänel & Palma (2007: 112, 124, 130); Hammer *et al.* (2010: 1113); Palma & Peck (2013: 40).

Remarks: *Halipeurus pelagodromae* is a “wing” louse parasitic on all the subspecies of the white-faced storm petrel. This louse was recorded as “*Halipeurus (Synnautes) pelagicus*”, or “*Halipeurus (Synnautes) pelagicus s. l.*”, or “*Halipeurus pelagicus*” in papers published until 2010.

### ***Halipeurus placodus* Edwards, 1961**

*Halipeurus (Halipeurus) placodus* Edwards, 1961: 141, figs 3e–7e.

*Halipeurus (Halipeurus) placodus* Edwards, 1961; Timmermann 1965: 144.

*Halipeurus placodus* Edwards, 1961; Watt, 1971: 236, 243.

*Halipeurus (Halipeurus) placodus* Edwards, 1961; Wise, 1977: 61

*Halipeurus (Halipeurus) placodus* Edwards, 1961; Pilgrim & Palma 1982: 12.

*Halipeurus (Halipeurus) placodus* Edwards, 1961; Palma 1999: 376, 383, note 2.

*Halipeurus (H.) placodus* Edwards, 1961; Palma 2010: 408

Holotype ♂ in MCZC (Palma 1996b: 186).

Type host: *Puffinus assimilis tunneyi* Mathews, 1912.

New Zealand hosts: *Puffinus newelli* Henshaw, 1900; *Puffinus assimilis kermadecensis* Murphy, 1927.

Other hosts: *Puffinus auricularis* Townsend, 1890; *Puffinus lherminieri gunax* Mathews, 1930; *Puffinus assimilis assimilis* Gould, 1838.

New Zealand localities: ND, AK, WI, WA, WN, KE.

Geographic distribution: Pacific Ocean.

New Zealand references: Edwards (1961); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1371); Palma (1999); Palma (2010).

Other significant references: Timmermann (1965); Palma (1996b: 186); Price *et al.* (2003: 188).

Remarks: *Halipeurus placodus* is a “wing” louse parasitic on several species of small shearwaters. Both *H. placodus* and *H. munda* (see above) are useful species to confirm the identity of the hosts, especially juveniles and beach-wrecked remains of specimens belonging to the *Puffinus assimilis* species complex (see Holdaway *et al.* 2001: 127).

### ***Halipeurus pricei* Palma, 2011**

*Halipeurus (Halipeurus)* sp.; Pilgrim & Palma 1982: 9.

*Halipeurus (Halipeurus)* sp.; Murray *et al.* 1990: 1370.

*Halipeurus* sp.; Palma 2010: 408.

*Halipeurus pricei* Palma, 2011b: 7, figs 4, 13–15, 29, 44, 65.

Holotype ♂ in MONZ.

Type host: *Pterodroma brevipes brevipes* (Peale, 1848).

New Zealand host: *Pterodroma leucoptera caledonica* Imber & Jenkins, 1981.

Other hosts: *Pterodroma brevipes magnificens* Bretagnolle & Shirihai, 2010; *Pterodroma leucoptera leucoptera* (Gould, 1844).

New Zealand localities: WI, WN.

Geographic distribution: Pacific Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010); Palma (2011b: 9).

Other significant references: None.

Remarks: *Halipeurus pricei* is a “wing” louse parasitic on two gadfly petrel species. Pilgrim & Palma (1982: 9) listed “*Halipeurus (Halipeurus) sp.*” (now = *Halipeurus pricei*) under the host “*Pterodroma leucoptera* subspecies”, a petrel now known as *Pterodroma leucoptera caledonica* (see Palma & Tennyson 2005).

### ***Halipeurus procellariae* (J.C. Fabricius, 1775)**

*Pediculus procellariae* J.C. Fabricius, 1775: 808.

*Ricinus procellariae* (J.C. Fabricius, 1775); Latreille 1804: 107.

*Esthiopeterum procellariae* J.C. Fabricius, 1775 [sic]; Harrison 1916: 140 (as “Not recognisable”).

“*Halipeurus angusticeps*” Harrison, 1937: 31 (not *Lipeurus angusticeps* Piaget, 1880).

*Halipeurus procellariae* (J.C. Fabricius, 1775); Hopkins & Clay 1952: 164.

*Halipeurus procellariae* (J.C. Fabricius, 1775); Timmermann 1960: 325, figs 1–2, 4a, 5, 7.

*Halipeurus (Halipeurus) procellariae* (J.C. Fabricius, 1775); Edwards 1961: 149, figs 3p–7p.

*Halipeurus (Halipeurus) procellariae* (J.C. Fabricius, 1775); Timmermann 1965: 147, figs 75, 77a, 78a, 86, pl. 7: figs 1–2.

*Halipeurus procellariae* (J.C. Fabricius, 1775); Clay & Moreby 1970: 218.

“*Halipeurus (Halipeurus) angusticeps*” Wise, 1977: 60 (not *Lipeurus angusticeps* Piaget, 1880).

*Halipeurus (Halipeurus) procellariae* (J.C. Fabricius, 1775); Wise 1977: 61.

*Halipeurus sp.*; Lowry *et al.* 1978: 138.

*Halipeurus (Halipeurus) procellariae* (J.C. Fabricius, 1775); Pilgrim & Palma 1982: 8, 30.

*Halipeurus (Halipeurus) procellariae* (J.C. Fabricius, 1775) *s. l.*; Pilgrim & Palma 1982: 8.

*Halipeurus (Halipeurus) procellariae* (J.C. Fabricius, 1775); Marris 2000: 187.

*Halipeurus (H.) procellariae* (J.C. Fabricius, 1775); Palma 2010: 408.

Type/s lost. A neotype has not been designated (see Clay & Hopkins 1951: 34 and Edwards, 1961: 150).

Type host: “*Brasiliae procellaris*” = *Pterodroma macroptera macroptera* (Smith, 1840) (see Clay & Hopkins 1951: 34; Edwards, 1961: 150).

New Zealand hosts: *Pterodroma macroptera gouldi* (Hutton, 1869); *Pterodroma lessonii* (Garnot, 1826); *Pterodroma magentae* (Giglioli & Salvadori, 1869); *Pterodroma mollis* (Gould, 1844).

Other host: *Pterodroma incerta* (Schlegel, 1863).

New Zealand localities: ND, AK, CL, BP, WO, TK, WI, WN, NC, MC, SC, WD, CH, AN, Macquarie Island.

Geographic distribution: Atlantic, Indian and Pacific Oceans.

New Zealand references: Harrison (1937); Edwards (1961: 150); Watson (1967: 72); Clay & Moreby (1970); Gressitt (1970: 327); Pilgrim (1974: 1034, fig. 3); Wise (1977); Lowry *et al.* (1978); Pilgrim & Palma (1982); Murray *et al.* (1990: 1369); Marris (2000); Palma & Imber (2000: 229); Palma & Horning (2002: 8, 16); Page *et al.* (2004: 643, 648); Palma (2010).

Other significant references: Clay & Hopkins (1951: 34); Timmermann (1960; 1965); Green & Palma (1991: 14, 26); Palma (1996b: 186); Furness & Palma (1992: 35, 40); Price *et al.* (2003: 188); Hänel & Palma (2007: 113, 124, 130); Hammer *et al.* (2010: 1113).

Remarks: *Halipeurus procellariae* is a “wing” louse parasitic on several species of large gadfly petrels. Pilgrim & Palma (1982: 8) regarded the population of *Halipeurus procellariae* from *Pterodroma mollis* as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of more samples of this species shows that making such difference is not warranted. Harrison’s (1937: 31) record of “*Halipeurus angusticeps*” from Macquarie Island —repeated by Wise (1977: 60)—is a misidentification of *H. procellariae* (see Pilgrim & Palma 1982: 30, note 16).

### ***Halipeurus spadix spadix* Timmermann, 1961**

*Halipeurus spadix* Timmermann, 1961a: 409, fig. 7.

*Halipeurus intestatus* Timmermann, 1961a: 410.

*Halipeurus (Halipeurus) taxosetus* Edwards, 1961: 145, figs 2a, 3i–7i.

*Halipeurus (Halipeurus) spadix* Timmermann, 1961; Timmermann 1965: 142, fig. 84.

*Halipeurus spadex* [sic] Timmermann, 1961; Pilgrim 1970: 74.

*Halipeurus (Halipeurus) spadix* Timmermann, 1961; Wise 1977: 61.

*Halipeurus (Halipeurus) spadix* Timmermann, 1961; Pilgrim & Palma 1982: 12. In part.

*Halipeurus (Halipeurus) spadix* Timmermann, 1961; Murray *et al.* 1990: 1371. In part.

*Halipeurus (H.) spadix* Timmermann, 1961; Palma 2010: 408. In part.

*Halipeurus spadix spadix* Timmermann, 1961; Palma 2011b: 22.

Holotype ♂ in NHML.

Type host: *Puffinus opisthomelas* Coues, 1864.

New Zealand host: *Puffinus huttoni* Mathews, 1912.

Other hosts: *Puffinus nativitatis* Streets, 1877; *Puffinus lherminieri lherminieri* Lesson, 1839; *Puffinus lherminieri bannermani* Mathews & Iredale, 1915.

New Zealand localities: AK, WN, KA, NC, MC, SC.

Geographic distribution: Pacific and Atlantic Oceans.

New Zealand references: Pilgrim (1970: 75); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Page *et al.* (2004: 643, 648); Galloway (2005: 16); Palma (2010); Palma (2011b).

Other significant references: Edwards (1961); Timmermann (1965); Amerson & Emerson (1971: 8, 24); Ward & Downey (1973: 394); Palma (1996b: 186); Price *et al.* (2003: 188); Hammer *et al.* (2010: 1113).

Remarks: *Halipeurus spadix spadix* is a “wing” louse parasitic on several shearwater species. All New Zealand references, except for Palma (2011b), refer to this taxon as “*Halipeurus (Halipeurus) spadix* Timmermann, 1961”.

#### ***Halipeurus spadix subclavus* Timmermann, 1961**

*Halipeurus subclavus* Timmermann, 1961a: 411.

*Halipeurus (Halipeurus) subclavus* Timmermann, 1961; Timmermann 1965: 144.

*Halipeurus (Halipeurus) spadix* Timmermann, 1961; Pilgrim & Palma 1982: 12. In part.

*Halipeurus (Halipeurus) spadix* Timmermann, 1961; Murray *et al.* 1990: 1371. In part.

“*Halipeurus (Halipeurus) spadix*” Green & Palma 1991: 14, 27 (not *Halipeurus spadix* Timmermann, 1961).

*Halipeurus (Halipeurus) subclavus* Timmermann, 1961; Price *et al.* 2003: 188.

*Halipeurus (H.) spadix* Timmermann, 1961; Palma 2010: 408. In part.

*Halipeurus spadix subclavus* Timmermann, 1961; Palma 2011b: 23, figs 9, 30, 45, 64, 66–67.

Holotype ♂ in NMHL.

Type host: *Puffinus lherminieri (?persicus)* Hume, 1873).

New Zealand host: *Puffinus gavia* (J.R. Forster, 1844).

Other hosts: *Puffinus lherminieri bailloni* (Bonaparte, 1857); *Puffinus lherminieri nicolae* Jouanin, 1971.

New Zealand localities: CL, BP, GB, WN, SD, MB, NN.

Geographic distribution: Indian Ocean and Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010); Palma (2011b).

Other significant references: Green & Palma (1991); Timmermann (1965); Price *et al.* (2003).

Remarks: *Halipeurus spadix subclavus* is a “wing” louse parasitic on several shearwater species. All New Zealand references, except for Palma (2011b), refer to this taxon as “*Halipeurus (Halipeurus) spadix* Timmermann, 1961”.

#### ***Halipeurus theresae* Timmermann, 1969**

*Halipeurus theresae* Timmermann, 1969b: 249, figs 4–5.

“*Halipeurus leucophryna*” Ward & Downey, 1973: 394 (not *Halipeurus leucophryna* Timmermann, 1960).

*Halipeurus (Halipeurus) theresae* Timmermann, 1969; Pilgrim & Palma 1982: 9.

*Halipeurus (Halipeurus) theresae* Timmermann, 1969; Palma 1999: 378.

*Halipeurus (H.) theresae* Timmermann, 1969; Palma 2010: 408.

Holotype ♂ in BPBM (Tenorio 1979: 13).

Type host: *Pterodroma hypoleuca* (Salvin, 1888).

New Zealand hosts: *Pterodroma magentae* (Giglioli & Salvadori, 1869); *Pterodroma axillaris* (Salvin, 1893).

Other hosts: *Pterodroma hasitata hasitata* (Kuhl, 1820); *Pterodroma hasitata caribbaea* Carte, 1866; *Pterodroma feae* (Salvadori, 1899); *Pterodroma deserta* Mathews, 1934.

New Zealand locality: CH.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990: 1370); Palma (1999); Palma & Imber (2000: 230); Palma (2010).

Other significant references: Tenorio (1979: 13); Zonfrillo (1993: 327); Monteiro & Furness (1995: 10); Forrester *et al.* (1995: 5); Price *et al.* (2003: 188); Page *et al.* (2004: 643, 648); Boieiro *et al.* (2008: 268, 297); Hammer *et al.* (2010: 1113); Palma (2011b: 16).

Remarks: *Halipeurus theresae* is a “wing” louse parasitic on many gadfly petrels. The host given above as *Pterodroma deserta* has been variously referred to as “*Pterodroma feae*” (e.g. Monteiro & Furness 1995, from the Azores Islands) or as “*Pterodroma feae deserta*” (e.g. Zonfrillo 1993, from the Madeira Archipelago). However, the host named here as *Pterodroma feae* is from the Cape Verde Islands, and represents a new host record and locality record for *Halipeurus theresae* (voucher specimens in MONZ).

I have examined 4 males and 6 females of *Halipeurus theresae* misidentified by Ward & Downey (1973: 394) as “*Halipeurus leucophryna*”, and deposited in the USNM and BPBM.

### ***Halipeurus thompsoni* Edwards, 1961**

*Halipeurus (Halipeurus) thompsoni* Edwards, 1961: 147, figs 3k–7k.

*Halipeurus (Halipeurus) thompsoni* Edwards, 1961; Timmermann 1965: 145.

*Halipeurus thompsoni* Edwards, 1961; Pilgrim 1970: 75.

*Halipeurus (Halipeurus) thompsoni* Edwards, 1961; Wise 1977: 61.

*Halipeurus (Halipeurus) thompsoni* Edwards, 1961; Pilgrim & Palma 1982: 12.

*Halipeurus (H.) thompsoni* Edwards, 1961; Palma 2010: 408.

Holotype ♂ in AMNH.

Type host: *Puffinus bulleri* Salvin, 1888.

New Zealand host: *Puffinus bulleri* Salvin, 1888.

Other hosts: None.

New Zealand localities: ND, AK, CL, BP, WI, WN, NC, MC, SC, CH.

Geographic distribution: Pacific Ocean.

New Zealand references: Pilgrim (1970); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1371); Palma (2010).

Other significant references: Timmermann (1965); Palma (1996b: 186); Price *et al.* (2003: 188).

Remarks: *Halipeurus thompsoni* is a distinct “wing” louse species, exclusively parasitic on Buller’s shearwater, a host endemic to New Zealand, but which ranges widely over the Pacific Ocean (Checklist Committee 2010: 114).

### ***Halipeurus turtur* Edwards, 1961**

*Halipeurus (Halipeurus) turtur* Edwards, 1961: 149, figs 3o–7o.

*Halipeurus (Halipeurus) turtur* Edwards, 1961; Timmermann 1965: 149.

*Halipeurus turtur* Edwards, 1961; Pilgrim 1970: 75.

*Halipeurus (Halipeurus) turtur* Edwards, 1961; Wise 1977: 61.

*Halipeurus (Halipeurus) turtur* Edwards, 1961; Pilgrim & Palma 1982: 9, 30, note 15.

*Halipeurus (H.) turtur* Edwards, 1961; Palma 2010: 408.

Holotype ♂ in AMNH (Pilgrim 1970: 75).

Type host: “*Pachyptila turtur*”, in error (see Palma & Horning 2002: 20, note 4).

New Zealand host: *Pterodroma cookii* (G.R. Gray, 1843).

Other hosts: None.

New Zealand localities: ND, CL, AK, KA, SI.

Geographic distribution: Pacific Ocean.

New Zealand references: Pilgrim (1970); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1370); Palma (1999: 384); Page *et al.* (2004: 643, 648); Palma (2010).

Other significant references: Clay & Moreby (1967: 160, fig. 148); Timmermann (1965); Price *et al.* (2003: 188); Hammer *et al.* (2010: 1113).

Remarks: *Halipeurus turtur* is a “wing” louse exclusively parasitic on Cook’s petrel, a host endemic to New Zealand, but which ranges widely over the Pacific Ocean (Checklist Committee 2010: 96).

The record of “*Halipeurus turtur* Edwards, 1961” in Watson (1967: 72) —repeated in subsequent publications— is a misidentification of *H. diversus* (see Palma (1999: 384, note F). Furthermore, a record of “*Halipeurus turtur* Edwards, 1961” in Emerson (1971: 360) is a misidentification of *Naubates prioni* (Enderlein, 1908) (see Palma & Pilgrim 2002: 42).

### Genus *Harrisoniella* Bedford, 1929

*Harrisoniella* Bedford, 1929. *15th Annual Rep. Director Veterinary Services*: 529. Type species: *Lipeurus ferox* Giebel, 1867 = *Harrisoniella ferox* (Giebel, 1867) (by subsequent designation).

*Diomedicola* Kéler, 1957b. *Beitr. zur Entomol.* 7(3/4): 496. Type species: *Lipeurus ferox* Giebel, 1867 = *Harrisoniella ferox* (Giebel, 1867) (by original designation).

### *Harrisoniella ferox* (Giebel, 1867)

Figs 113–114

*Lipeurus ferox* Giebel, 1867: 195.

*Esthiopterum ferox* Giebel, 1867 [sic]; Harrison 1916: 134.

*Perineus ferox* (Giebel, 1867); Harrison 1937: 29.

*Harrisoniella ferox* (Giebel, 1867); Hopkins & Clay 1952: 165.

*Diomedicola ferox* (Giebel, 1867); Kéler 1957b: 502, figs 1a, 4, 8, 9.

*Diomedicola irroratae* Kéler, 1957b: 508, figs 3c, 10c.

*Harrisoniella chilensis* Carriker, 1964: 6, figs 4–7, 7a.

*Harrisoniella ferox* (Giebel, 1867); Timmermann 1965: 94, figs 27, 28a.

*Harrisoniella irroratae* (Kéler, 1957); Timmermann 1965: 95, figs 29c, 30c.

*Harrisoniella ferox* (Giebel, 1867); Pilgrim & Palma 1982: 6.

*Harrisoniella ferox* (Giebel, 1867); Palma & Pilgrim 1984: 149, figs 1–3, 7, 11, 15, 23.

*Harrisoniella ferox* (Giebel, 1867); Murray *et al.* 1990: 1368–1369.

*Harrisoniella ferox* (Giebel, 1867); Palma 2010: 408.

Neotype ♂ in SAIM (Palma & Pilgrim 1984: 150).

Type host: *Thalassarche melanophris* (Temminck, 1828).

New Zealand hosts: *Thalassarche melanophris* (Temminck, 1828); *Thalassarche impavida* Mathews, 1912; *Thalassarche bulleri bulleri* (Rothschild, 1893); *Thalassarche bulleri platei* (Reichenow, 1898); *Thalassarche cauta* (Gould, 1841); *Thalassarche cauta steadi* Falla, 1933; *Thalassarche eremita* Murphy, 1930; *Thalassarche salvini* (Rothschild, 1893).

Other hosts: *Thalassarche chlororhynchos* (J.F. Gmelin, 1789); *Phoebastria irrorata* (Salvin, 1883).

New Zealand localities: BP, WO, HB, WI, WN, KA, NC, MC, SC, WD, CH, SI, BO, AU, CA.

Geographic distribution: Southern Hemisphere.

New Zealand references: Pilgrim & Palma (1982); Palma & Pilgrim (1984: 150); Murray *et al.* (1990); Palma (1999: 375); Palma (2010).

Other significant references: Kéler (1957b); Timmermann (1965); Green & Palma (1991: 15, 25); Palma (1996b: 187); Price *et al.* (2003: 188); Page *et al.* (2004: 642, 648); Hänel & Palma (2007: 113, 124, 129); Palma & Peck (2013: 40).

Remarks: *Harrisoniella ferox* is mostly found under the wings of small albatrosses. Palma & Pilgrim (1984: 149) included a complete synonymy for *H. ferox*, which was misidentified for many years due to lack of type material.

### *Harrisoniella hopkinsi* Eichler, 1952

“*Philopterus diomedae*” Dufour, 1835: 671, pl. 21: figs 1–2 (not *Pediculus diomedae* J.C. Fabricius, 1775).

“*Lipeurus ferox*” Taschenberg, 1882: 145, pl. 5: figs 1, 1a (not *Lipeurus ferox* Giebel, 1867).

“*Lipeurus diomedae*” Osborn in Howard, 1890: 189 (not *Pediculus diomedae* J.C. Fabricius, 1775).

“*Lipeurus densus*” Waterston, 1914: 311 (not *Lipeurus densus* Kellogg, 1896).

“*Esthiopterum diomedae*” Harrison, 1916: 133 (not *Pediculus diomedae* J.C. Fabricius, 1775).

“*Perineus diomedae*” Harrison, 1937: 29 (not *Pediculus diomedae* J.C. Fabricius, 1775).

“*Harrisoniella diomedae*” Thompson, 1938b: 5, pl. 1 (not *Pediculus diomedae* J.C. Fabricius, 1775).

“? *Harrisoniella* [sic] *ferox*” Clay, 1940a: 298 (not *Lipeurus ferox* Giebel, 1867).

*Harrisoniella hopkinsi* Eichler, 1952a: 40, fig. 1.

*Harrisoniella thompsoni* Eichler, 1952a: 41, fig. 4.

*Harrisoniella hopkinsi* Eichler, 1952; Hopkins & Clay 1953: 438.

*Harrisoniella thompsoni* Eichler, 1952; Hopkins & Clay 1953: 438.

*Diomedicola hopkinsi* (Eichler, 1952); Kéler 1957b: 504, figs 1b, 3a,b, 7, 10b.

*Harrisoniella hopkinsi* Eichler, 1952; Timmermann 1965: 94, figs 28b, 29a,b, 30a,b, 31.

*Harrisoniella* sp.; Watson 1967: 72.

*Harrisoniella hopkinsi* Eichler, 1952; Wise 1977: 61.

*Harrisoniella hopkinsi* Eichler, 1952; Pilgrim & Palma 1982: 5.

*Harrisoniella hopkinsi* Eichler, 1952; Palma & Pilgrim 1984: 156, figs 5, 9, 13, 17, 21, 25, 27.

*Harrisoniella hopkinsi* Eichler, 1952; Palma 2001: 67, fig. 7.

*Harrisoniella hopkinsi* Eichler, 1952; Palma 2010: 408.

Holotype ♂ in ZMHU (Göllner-Scheiding 1973: 35). Holotype ♂ of *Harrisoniella thompsoni* in NHML (Palma & Pilgrim 1984: 156).

Type host: *Diomedea exulans* Linnaeus, 1758.

New Zealand hosts: *Diomedea exulans* Linnaeus, 1758; *Diomedea antipodensis antipodensis* Robertson & Warham, 1992; *Diomedea epomophora* Lesson, 1825; *Diomedea sanfordi* Murphy, 1917.

Other host: *Diomedea dabbenena* Mathews, 1929.

New Zealand localities: ND, HB, TK, WA, WN, SD, MB, KA, NC, MC, SC, WD, CO, DN, CH, AN, AU, CA, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Harrison (1937); Clay (1964a: 231); Gressitt (1964: 538); Watson (1967); Pilgrim (1974: 1035, fig. 4); Wise (1977); Pilgrim & Palma (1982); Palma & Pilgrim (1984); Murray *et al.* (1990: 1368); Marris (2000: 187); Palma (2001: 67, fig. 7); Palma & Horning (2002: 8, 15); Page *et al.* (2004: 642, 648); Palma (2010).

Other significant references: Thompson (1938b); Séguy (1944: 33, fig. 24); Séguy (1953: 595, figs 53–54); Kéler (1957b); Timmermann (1965; 1966: 86, figs 1c, 2b); Clay & Moreby (1967: 162, figs 84, 116, 126); Green & Palma (1991: 15, 25); Palma (1991c: 239); Palma (1996b: 187); Price *et al.* (2003: 188); Hänel & Palma (2007: 107, 113, 125, fig. 2a).

Remarks: *Harrisoniella hopkinsi* is mostly found under the wings of large albatrosses. Palma & Pilgrim (1984: 156) included a complete synonymy for *H. hopkinsi*, which was misidentified for 117 years, since Dufour (1835) until it was recognised as a different species by Eichler (1952a).

### Genus *Ibidoecus* Cummings, 1916

*Ibidoecus* Cummings, 1916b. *Proc. Zool. Soc. London* 1916: 663. Type species: *Philopterus plataleae* Denny, 1842 = *Ibidoecus plataleae* (Denny, 1842) (by original designation).

#### *Ibidoecus bisignatus* (Nitzsch [*in* Giebel], 1866)

*Docophorus bisignatus* Nitzsch [*in* Giebel], 1866: 362.

*Docophorus bisignatus* Nitzsch [*in* Giebel], 1874 [sic]; Piaget 1885: 11, pl. 2: fig. 1.

*Philopterus bisignatus* Nitzsch [*in* Giebel], 1866 [sic]; Harrison 1916: 89.

*Ibidoecus bisignatus* (Nitzsch, 1866) [sic]; Cummings 1916b: 663.

*Ibidoecus bisignatus* (Nitzsch [*in* Giebel], 1866); Hopkins & Clay 1952: 176.

*Ibidoecus bisignatus* (Nitzsch, 1866) [sic]; Tandan 1958b: 403, figs 6, 10, 19–21.

*Ibidoecus bisignatus* (Nitzsch, 1866) [sic]; Pilgrim & Palma 1982: 16.

*Ibidoecus bisignatus* (Nitzsch, 1866) [sic]; Murray *et al.* 1990: 1373.

*Ibidoecus bisignatus* (Nitzsch [*in* Giebel], 1866); Price *et al.* 2003: 191.

*Ibidoecus bisignatus* (Nitzsch [*in* Giebel], 1866); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Plegadis falcinellus* (Linnaeus, 1766).

New Zealand host: *Plegadis falcinellus* (Linnaeus, 1766).

Other host: *Plegadis chihi* (Vieillot, 1817).

New Zealand localities: MB, SL.

Geographic distribution: Eurasia; Africa; Americas; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Cummings (1916b); Dubinin (1938: fig. 15); Carriker (1947: 114); Tandan (1958b); Forrester *et al.* (1995: 12); Palma (1996b: 188); Price *et al.* (2003).

Remarks: *Ibidoecus bisignatus* lives mostly on the head and neck of the host. Although the glossy ibis is a regular vagrant to New Zealand (Checklist Committee 2010: 166), there are only two records of *I. bisignatus* from this country.

### ***Ibidoecus diana*e Tandan, 1958**

Figs 115–116

*Ibidoecus diana*e Tandan, 1958a: 151, figs 1–6.

*Ibidoecus diana*e Tandan, 1958; Pilgrim & Palma 1982: 16.

*Ibidoecus diana*e Tandan, 1958; Murray *et al.* 1990: 1373.

*Ibidoecus diana*e Tandan, 1958; Price *et al.* 2003: 191.

*Ibidoecus diana*e Tandan, 1958; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Threskiornis molucca pygmaeus* Mayr, 1931.

New Zealand host: *Threskiornis molucca strictipennis* (Gould, 1838).

Other hosts: None.

New Zealand locality: SL.

Geographic distribution: Australasia; Melanesia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Moreby (1976: 92); Clay (1976a: 5); Palma (1996b: 188); Price *et al.* (2003: 191).

Remarks: *Ibidoecus diana*e lives mostly on the head and neck of the host. The Australian white ibis is an occasional visitor to New Zealand (Checklist Committee 2010: 166), with only one record of *I. diana*e from this country.

### ***Ibidoecus platalea*e (Denny, 1842)**

*Philoater (Docophorus) platalea*e Denny, 1842: 46, 100, pl. 4: fig. 9.

*Philoater platalea*e Denny, 1842 [sic]; Harrison 1916: 102.

*Ibidoecus platalea*e (Denny, 1842); Cummings 1916b: 664, figs 15, 21.1.

*Ibidoecus platalea*e (Denny, 1842); Hopkins & Clay 1952: 177.

*Ibidoecus platalea*e (Denny, 1842); Tandan 1958b: 406, figs 5, 9, 22–26.

*Ibidoecus platalea*e (Denny, 1842); Pilgrim & Palma 1982: 16.

*Ibidoecus platalea*e (Denny, 1842); Murray *et al.* 1990: 1373.

*Ibidoecus platalea*e (Denny, 1842); Price *et al.* 2003: 191.

*Ibidoecus platalea*e (Denny, 1842); Palma 2010: 408.

Syntypes ♂♀ in NHML (Cummings 1916b: 663).

Type host: *Platalea leucorodia* Linnaeus, 1758.

New Zealand host: *Platalea regia* Gould, 1838.

Other hosts: None.

New Zealand localities: AK, HB, NN, WD.

Geographic distribution: Eurasia; Africa; Australasia; Melanesia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Cummings (1916b); Séguy (1944: 263, figs 398–400); Carriker (1947: 117); Tandan (1958b); Moreby (1976: 92); Butler & O'Connor (1994: 455); Martín-Mateo (1994: 114, figs 1f, 3c,d); Palma (1996b: 188); Price *et al.* (2003: 191); Martín-Mateo (2009: 219).

Remarks: Since the first record of the royal spoonbill in New Zealand in 1861, this host has become well established with many breeding localities in this country (Checklist Committee 2010: 167). *Ibidoecus plataleae* lives mostly on the head and neck of the host and is the most frequently collected louse species from New Zealand spoonbills.

### Genus *Incidifrons* Ewing, 1929

*Incidifrons* Ewing, 1929. *Manual External Parasites*: 111, 189. Type species: *Phlopterus pertusus* Nitzsch [sic] = *Incidifrons fulicae* (Linnaeus, 1758) (by original designation).

#### *Incidifrons fulicae* (Linnaeus, 1758)

“Pollino della folaga” Redi, 1668: pl. 4: fig. 3.

*Pediculus fulicae* Linnaeus, 1758: 613. *Nomen novum* for Redi’s “Pollino della folaga” Redi, 1668: pl. 4: fig. 3.

*Docophorus pertusus* Burmeister, 1838a: 436.

*Docophorus pertusus* Nitzsch [in Giebel], 1874: 108, pl. 11: figs 3, 12.

*Incidifrons pertusus* (Nitzsch) [sic]; Ewing, 1929: 190.

*Incidifrons fulicae* (Linnaeus, 1758); Hopkins 1940: 425.

*Incidifrons fulicae* (Linnaeus, 1758); Hopkins & Clay 1952: 178.

*Incidifrons fulicae* (Linnaeus, 1758); Pilgrim & Palma 1982: 19.

*Incidifrons fulicae* (Linnaeus, 1758); Murray *et al.* 1993: 961.

*Incidifrons fulicae* (Linnaeus, 1758); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1950: 257).

Type host: *Fulica atra* Linnaeus, 1758.

New Zealand host: *Fulica atra australis* Gould, 1845.

Other host: *Fulica cristata* J.F. Gmelin, 1789.

New Zealand locality: CO, DN.

Geographic distribution: Africa; Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Hopkins (1940); Clay & Hopkins (1950: 255, figs 48–51); Ledger (1980: 146); Castro & Cicchino (1983: 274, figs 4, 12, 16, 23, 27); Rékasi (1986: 124, figs 26–27); Green & Palma (1991: 15, 32); Palma (1996b: 189); Price *et al.* (2003: 192); Palma & Jensen (2005: 56, 63); Adam (2007: 181); Martín-Mateo (2009: 110, fig. 27).

Remarks: Hopkins (1940: 421) gives a complete account of the early taxonomic confusion among several species of lice described from *Fulica atra*, including a clarification of the synonymy of *Incidifrons fulicae*. Although the Australian coot is now widespread and increasing its range in New Zealand (Checklist Committee 2010: 190), there is only one record of *I. fulicae* from this country.

#### *Incidifrons porzanae* Blagoveshtchensky, 1951

#### New Record

Figs 117–118

*Incidifrons pertusus porzanae* Blagoveshtchensky, 1951: 294, fig. 10.

*Incidifrons porzanae* Blagoveshtchensky, 1951; Hopkins & Clay 1953: 439.

*Incidifrons* sp.; Pilgrim & Palma 1982: 19.

*Incidifrons* sp.; Murray *et al.* 1993: 961.

*Incidifrons porzanae* Blagoveshtchensky, 1951; Price *et al.* 2003: 192.

Holotype ♂, repository unknown.

Type host: *Porzana parva* (Scopoli, 1769).

New Zealand host: *Porzana pusilla affinis* (J.E. Gray, 1845).

Other hosts: None.

New Zealand localities: ND, WO, MC, SC, CO, SL.

Geographic distribution: Asia; New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993).

Other significant references: Hopkins & Clay (1953); Price *et al.* (2003).

Material examined and repository: 12♂, 25♀, 12N (7 samples, MONZ).



Remarks: This is the first record of *Incidifrons porzanae* for New Zealand, because the New Zealand references cited above reported this louse as “*Incidifrons* sp.” only.

### Genus *Lagopoecus* Waterston, 1922

*Lagopoecus* Waterston, 1922. *Entomol. Month. Mag.* 58: 159. Type species: *Nirmus cameratus* “Lyonet, (1830)” [sic] = *Lagopoecus lyrurus* Clay, 1938 (by original designation).

*Colinicola* Carriker, 1945b. *Rev. Acad. Colomb. Ciencias Exactas, Físicas y Naturales* 6: 360. Type species: *Goniodes numidianus* Denny, 1842 = *Lagopoecus numidianus* (Denny, 1842) (by original designation).

### *Lagopoecus docophoroides* (Piaget, 1880)

Figs 119–120

*Lipeurus docophoroides* Piaget, 1880: 357, pl. 28: fig. 9.

*Lipeurus docophoroides* Piaget, 1880; Harrison 1916: 83.

*Lagopoecus docophoroides* (Piaget, 1880); Clay 1938: 195: fig. 43d.

*Lagopoecus docophoroides* (Piaget, 1880); Hopkins & Clay 1952: 187.

*Lagopoecus docophoroides* (Piaget, 1880); Pilgrim & Palma 1982: 18.

*Lagopoecus docophoroides* (Piaget, 1880); Murray *et al.* 1993: 960.

*Colinicola docophoroides* (Piaget, 1880); Price *et al.* 2003: 163.

*Lagopoecus docophoroides* (Piaget, 1880); Palma 2010: 408.

Syntypes ♂♀ in NHML (Clay 1938: 195).

Type host: *Callipepla californica californica* (Shaw, 1798).

New Zealand host: *Callipepla californica brunnescens* (Ridgway, 1884).

Other hosts: None

New Zealand localities: SD, MB, NN, NC, MC, SC, CO, DN.

Geographic distribution: North & South America; Europe; Australasia, Hawaiian Islands.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Emerson (1949: 117, fig. 1); Price *et al.* (2003).

Remarks: *Lagopoecus docophoroides* was introduced to New Zealand and other countries with California quails by human agency (Checklist Committee 2010: 25). In agreement with Hopkins & Clay (1952: 73) and contrary to Price *et al.* (2003: 163), I do not recognise the genus *Colinicola* as separable from *Lagopoecus*.

### Genus *Lipeurus* Nitzsch, 1818

*Lipeurus* Nitzsch, 1818. *Germar's Mag. Entomol.* 3: 292. Type species: *Pediculus caponis* Linnaeus, 1758 = *Lipeurus caponis* (Linnaeus, 1758) (by subsequent designation).

### *Lipeurus caponis* (Linnaeus, 1758)

*Pediculus caponis* Linnaeus, 1758: 614.

*Ricinus caponis* (Linnaeus, 1758); Latreille 1804: 110.

*Lipeurus caponis* (Linnaeus, 1758); Clay 1938: 111, figs 1, 2a,b, 3a.

*Lipeurus (Lipeurus) caponis* (Linnaeus, 1758); Séguy 1944: 193, figs 278–281.

*Lipeurus caponis* (Linnaeus, 1758); Hopkins & Clay 1952: 192.

*Lipeurus caponis* (L.); Helson 1956: 13, 17.

*Lipeurus caponis*; Whitten 1971: 383.

*Lipeurus caponis* (Linnaeus, 1758); Wise 1977: 61.

*Lipeurus caponis* (Linnaeus, 1758); Murray *et al.* 1993: 960.

*Lipeurus caponis* (Linnaeus, 1758); Palma 1999: 384, note G.

*Lipeurus caponis* (Linnaeus, 1758); Palma 2010: 408.

Neotype ♂ in NHML (Clay & Hopkins 1950: 263).

Type host: *Gallus gallus* (Linnaeus, 1758).

New Zealand hosts: *Gallus gallus gallus* (Linnaeus, 1758); *Meleagris gallopavo* Linnaeus, 1758; *Numida meleagris* (Linnaeus, 1758) captive.

Other hosts: *Colinus virginianus* (Linnaeus, 1758); *Gallus sonneratii* Temminck, 1813; *Gallus lafayettii* Lesson, 1831; *Gallus varius* (Shaw, 1798); *Phasianus colchicus* Linnaeus, 1758.

New Zealand localities: AK, TO, WN, MB.

Geographic distribution: Eurasia; Americas; Australasia.

New Zealand references: Helson (1956); Whitten (1971); Wise (1977); Murray *et al.* (1993); Palma (1999); Palma (2010).

Other significant references: Clay (1938); Séguy (1944); Clay & Hopkins (1950: 263); Emerson (1956a: 69, pl. 5); Emerson & Ward (1958: 57); Mey (1986: 36, fig. 5, photo 3); Price (1987: 220, fig. 22.21); Palma (1996b: 190); Price *et al.* (2003: 194); Martín-Mateo (2009: 73, fig. 17); Palma & Peck (2013: 40).

Remarks: *Lipeurus caponis* was introduced to New Zealand and other countries with chickens by human agency. This louse species has spread onto a number of adventive hosts due to the human practice of mixing various species of game birds in captivity. *Gallus gallus gallus* has been listed for the first time in the latest edition of the New Zealand Checklist of Birds (Checklist Committee 2010: 27; see also Palma 1999: 383, note 5).

### ***Lipeurus maculosus maculosus* Clay, 1938**

Figs 121–122

*Lipeurus maculosus maculosus* Clay, 1938: 116, figs 5a, 6a, pl. 1: fig. 2.

*Lipeurus (Lipeurus) maculosus* Clay, 1938; Séguy 1944: 193, fig. 282.

*Lipeurus maculosus* Clay, 1938; Hopkins & Clay 1952: 196.

*Lipeurus maculosus maculosus* Clay, 1938; Pilgrim & Palma 1982: 18.

*Lipeurus m. maculosus* Clay, 1938; Murray *et al.* 1993: 960.

*Lipeurus maculosus maculosus* Clay, 1938; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Phasianus colchicus* Linnaeus, 1758.

New Zealand hosts: *Phasianus colchicus* Linnaeus, 1758; *Perdix perdix perdix* (Linnaeus, 1758).

Other hosts: None.

New Zealand localities: HB, WI, WN, NC, MC, SC.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Séguy (1944); Marconcini & Macchioni (1975: 106, figs 5–6); Rékasi (1986: 125, fig. 28); Modrzejewska & Złotorzycka (1987: 663, figs 5, 9); Kopociński *et al.* (1998: 81); Price *et al.* (2003: 195).

Remarks: *Lipeurus maculosus maculosus* was introduced to New Zealand with its primary host (pheasants) by human agency (Checklist Committee 2010: 28). *Perdix perdix perdix* appears to have died out in New Zealand (Checklist Committee 2010: 348).

### **Genus *Lunaceps* Clay & Meinertzhagen, 1939**

*Lunaceps* Clay & Meinertzhagen, 1939b. *Ann. Mag. Nat. Hist.* (Ser. 11) 4: 450. Type species: *Degeeriella actophila* (Kellogg & Chapman, 1899) = *Lunaceps actophilus* (Kellogg & Chapman, 1899) (by original designation).

### ***Lunaceps actophilus* (Kellogg & Chapman, 1899)**

*Nirmus actophilus* Kellogg & Chapman, 1899: 78, pl. 6: fig. 4.

*Degeeriella actophila* Kellogg & Chapman, 1899 [sic]; Harrison 1916: 107.

*Lunaceps actophilus* (Kellogg & Chapman, 1899); Hopkins & Clay 1952: 201.

*Lunaceps actophilus* (Kellogg & Chapman, 1899); Timmermann 1954b: 628.

*Lunaceps actophilus* (Kellogg & Chapman, 1899); Pilgrim & Palma 1982: 21.

*Lunaceps actophilus* (Kellogg & Chapman, 1899); Murray *et al.* 2006a: 1964.

*Lunaceps actophilus* (Kellogg & Chapman, 1899); Palma 2010: 408.

*Lunaceps actophilus* (Kellogg & Chapman, 1899); Gustafsson & Olsson 2012b: 11, figs 6a,b,c,d.

Lectotype ♀ in EMEC (Carriker 1957: 99; Emerson 1965: 49).

Type host: *Calidris alba* (Pallas, 1764).

New Zealand host: *Calidris alba* (Pallas, 1764).

Other hosts: None.

New Zealand locality: NC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010); Gustafsson & Olsson (2012b: 16).

Other significant references: Hackman & Nyholm (1968: 80); Forrester *et al.* (1995: 27); Price *et al.* (2003: 196); Palma & Jensen (2005: 56, 63); Gustafsson & Olsson (2012a: 94, 96, figs 1–2).

Remarks: The sanderling is an uncommon visitor to New Zealand (Checklist Committee 2010: 195), with a single record of *Lunaceps actophilus* from this country. The dunlin —*Calidris alpina* (Linnaeus, 1758)— previously regarded as a host of *Lunaceps actophilus* is now host to *Lunaceps schismatus* Gustafsson & Olsson, 2012b: 68.

#### ***Lunaceps drosti* Timmermann, 1954**

*Lunaceps drosti* Timmermann, 1954b: 627, fig. 1b.

*Lunaceps drosti* Timmermann, 1954; Pilgrim & Palma 1982: 21.

*Lunaceps drosti* Timmermann, 1954; Murray *et al.* 2006a: 1964.

*Lunaceps drosti* Timmermann, 1954; Palma 2010: 408.

*Lunaceps drosti* Timmermann, 1954; Gustafsson & Olsson 2012b: 19, figs 8a,b,c,d.

Holotype ♂ in NHML.

Type host: *Calidris canutus canutus* (Linnaeus, 1758).

New Zealand host: *Calidris canutus rogersi* (Mathews, 1913).

Other hosts: *Calidris canutus rufus* (Wilson, 1813); *Aphriza virgata* (J.F. Gmelin, 1789).

New Zealand localities: WN, MC, CA.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010); Gustafsson & Olsson (2012b: 21).

Other significant references: Forrester *et al.* (1995: 27); Palma (1996b: 191); Price *et al.* (2003: 196); Palma & Jensen (2005: 56, 63); Gustafsson & Olsson (2012a: 94, 96, figs 1–2).

Remarks: *Calidris canutus rogersi* is a regular and numerous annual visitor to New Zealand (Checklist Committee 2010: 195). The great knot —*Calidris tenuirostris* (Horsfield, 1821), previously regarded as a host of *Lunaceps drosti*— is now host to *Lunaceps mintoni* Gustafsson & Olsson, 2012b: 47.

#### ***Lunaceps falcinellus* Timmermann, 1954**

*Lunaceps falcinellus* Timmermann, 1954b: 627.

“*Lunaceps incoenis s. l.*” Pilgrim & Palma, 1982: 21 (not *Nirmus incoenis* Kellogg & Chapman, 1899).

*Lunaceps* sp.; Palma 1999: 380.

*Lunaceps* sp.; Murray *et al.* 2006a: 1964.

“*Lunaceps incoenis*” Murray *et al.*, 2006a: 1964 (not *Nirmus incoenis* Kellogg & Chapman, 1899).

“*Lunaceps incoenis*” Palma, 2010: 408 (not *Nirmus incoenis* Kellogg & Chapman, 1899).

*Lunaceps falcinellus* Timmermann, 1954; Gustafsson & Olsson 2012b: 24, figs 10a,b,c,d.

Holotype ♂ in NHML.

Type host: *Limicola falcinellus falcinellus* (Pontoppidan, 1763).

New Zealand hosts: *Calidris ferruginea* (Pontoppidan, 1763); *Calidris ruficollis* (Pallas, 1776).

Other hosts: *Calidris minuta* (Leisler, 1812); *Limicola falcinellus sibirica* Dresser, 1876.

New Zealand localities: WI, MC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Palma (1999); Murray *et al.* (2006a); Palma (2010); Gustafsson & Olsson (2012b: 27).

Other significant references: Price *et al.* (2003: 196); Gustafsson & Olsson (2012a: 94, 96, figs 1–2).

Remarks: Timmermann (1954b: 627) and Price *et al.* (2003: 196) listed *Tryngites subruficollis* (Vieillot, 1819) as a host for *Lunaceps falcinellus*, but Gustafsson & Olsson (2012b: 66) separated the population of *Lunaceps* from that host as the new species *L. rothkoi* Gustafsson & Olsson, 2012b: 66.

#### ***Lunaceps limosae* Bechet, 1968**

*Lunaceps limosella* Timmermann, 1954b: 629. In part.

*Lunaceps limosella limosae* Bechet, 1968: 127, fig. 5.

“*Lunaceps limosella*” Pilgrim & Palma, 1982: 20 (not *Lunaceps limosella* Timmermann, 1954).

*Lunaceps limosella* Timmermann, 1954; Price *et al.* 2003: 196. In part.

*Lunaceps limosella* Timmermann, 1954; Murray *et al.* 2006a: 1964. In part.

*Lunaceps limosae* Bechet, 1968; Gustafsson & Olsson 2012b: 40, figs 16 a,b,c,d.

Syntypes ♂♀ in Ion Bechet’s personal collection, Cluj-Napoca, Romania (Costica Adam pers. comm. July 2012).

Type host: *Limosa limosa limosa* (Linnaeus, 1758).

New Zealand host: *Limosa limosa melanuroides* Gould, 1846.

Other hosts: None.

New Zealand locality: AU.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Gustafsson & Olsson (2012b: 42).

Other significant references: Ledger (1980: 169); Butler & O’Connor (1994: 455); Price *et al.* (2003); Adam (2007: 180, figs 13b,c).

Remarks: *Lunaceps limosae* is an additional species to the New Zealand fauna because, with the exception of the original description by Bechet (1968), it had been regarded as a junior synonym of *L. limosella* in all subsequent publications until its resurrection by Gustafsson & Olsson (2012b: 40). The Asiatic black-tailed godwit is an uncommon annual visitor to New Zealand (Checklist Committee 2010: 203).

#### ***Lunaceps limosella* Timmermann, 1954**

*Lunaceps limosella* Timmermann, 1954b: 629. In part.

*Lunaceps limosella* Timmermann, 1954; Pilgrim & Palma 1982: 21.

*Lunaceps limosella* Timmermann, 1954; Murray *et al.* 2006a: 1964. In part.

*Lunaceps limosella* Timmermann, 1954; Palma 2010: 408.

*Lunaceps limosella* Timmermann, 1954; Gustafsson & Olsson 2012b: 42, figs 17 a,b,c,d.

Holotype ♂ in NHML.

Type host: *Limosa lapponica lapponica* (Linnaeus, 1758).

New Zealand host: *Limosa lapponica baueri* Naumann, 1836.

Other host: *Limosa lapponica menzbieri* Portenko, 1936.

New Zealand localities: NN, MB, KA, BR, MC, SL, SN, CA.

Geographic distribution: All continents, except the Americas and Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010); Gustafsson & Olsson (2012b: 45).

Other significant reference: Price *et al.* (2003: 196).

Remarks: Records of *Lunaceps limosella* from *Limosa limosa* are now referred to *Lunaceps limosae* (see above). The eastern bar-tailed godwit is the most numerous wader that visits New Zealand every year (Checklist Committee 2010: 203).

#### ***Lunaceps numenii madagascariensis* Gustafsson & Olsson, 2012**

Figs 123–124

*Lunaceps numenii* (Denny, 1842); Timmermann 1954: 631, fig. 5c. In part.

“*Lunaceps numenii numenii*” Pilgrim & Palma, 1982: 20 (not *Philopterus (Nirmus) numenii* Denny, 1842).

“*Lunaceps numenii numenii*” Price *et al.* 2003: 196 (not *Philopterus (Nirmus) numenii* Denny, 1842).

“*Lunaceps numenii numenii*” Murray *et al.* 2006a: 1964 (not *Philopterus (Nirmus) numenii* Denny, 1842).

“*Lunaceps numenii numenii*” Palma, 2010: 408 (not *Philopterus (Nirmus) numenii* Denny, 1842).

*Lunaceps numenii madagascariensis* Gustafsson & Olsson, 2012b: 55, figs 22a,b,c,d.

Holotype ♂ in NHML.

Type host: *Numenius madagascariensis* (Linnaeus, 1766).

New Zealand host: *Numenius madagascariensis* (Linnaeus, 1766).

Other hosts: None.

New Zealand localities: WD, SI.

Geographic distribution: Asia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010); Gustafsson & Olsson (2012b).

Other significant references: Timmermann (1954); Price *et al.* (2003).

Remarks: Although the eastern curlew is a regular annual visitor to New Zealand (Checklist Committee 2010: 200), there are only two records of *Lunaceps numenii madagascariensis* from this country.

### ***Lunaceps numenii oliveri* (Johnston & Harrison, 1912)**

*Degeeriella oliveri* Johnston & Harrison, 1912: 367, fig. 3.

*Degeeriella oliveri* Johnston & Harrison, 1912; Thompson 1939: 120.

*Lunaceps phaeopi* (Denny, 1842); Hopkins & Clay 1952: 202. In part.

"*Lunaceps phaeopi*" Emerson & Ward 1958: 57 (not *Philopterus (Nirmus) phaeopi* Denny, 1842).

"*Lunaceps phaeopi*" Watt, 1971: 236, 243 (not *Philopterus (Nirmus) phaeopi* Denny, 1842).

"*Lunaceps numenii phaeopi*" Wise, 1977: 61 (not *Philopterus (Nirmus) phaeopi* Denny, 1842).

*Lunaceps numenii oliveri* (Johnston & Harrison, 1912); Pilgrim & Palma 1982: 20, 31, note 22.

*Lunaceps numenii oliveri* (Johnston & Harrison, 1912); Price *et al.* 2003: 196.

*Lunaceps numenii oliveri* (Johnston & Harrison, 1912); Murray *et al.* 2006a: 1964.

*Lunaceps n. oliveri* (Johnston & Harrison, 1912); Palma 2010: 408.

*Lunaceps oliveri* (Johnston & Harrison, 1912) ("Clade 9"); Gustafsson & Olsson 2012a: 94, 96, figs 1–2.

*Lunaceps numenii phaeopi* (Denny, 1842); Gustafsson & Olsson 2012b: 57, figs 23a,b,c,d. In part.

Syntypes ♂♀ in MONZ (Palma *et al.* 1989: 45).

Type host: *Numenius phaeopus variegatus* (Scopoli, 1786).

New Zealand host: *Numenius phaeopus variegatus* (Scopoli, 1786).

Other hosts: None.

New Zealand localities: TK, KE.

Geographic distribution: Asia; Australasia.

New Zealand references: Johnston & Harrison (1912); Thompson (1939); Pilgrim (1970: 75); Wise (1977); Pilgrim & Palma (1982); Palma *et al.* (1989: 45); Murray *et al.* (2006a); Palma (2010); Gustafsson & Olsson (2012b: 60).

Other significant references: Emerson & Ward (1958); Moreby (1976: 93); Price *et al.* (2003); Gustafsson & Olsson (2012a).

Remarks: Contrary to Gustafsson & Olsson (2012b: 57, 59), I regard *Degeeriella oliveri* as a valid subspecies of *Lunaceps numenii*. Although the Asiatic whimbrel is a frequent visitor to New Zealand (Checklist Committee 2010: 201), its lice have been infrequently collected.

### ***Lunaceps superciliosus* Gustafsson & Olsson, 2012**

*Lunaceps* sp.; Palma 1999: 380.

*Lunaceps incoenis* (Kellogg & Chapman, 1899); Price *et al.* 2003: 196. In part.

*Lunaceps* sp.; Murray *et al.* 2006a: 1964.

*Lunaceps incoenis* (Kellogg & Chapman, 1899) ("Clade 3"); Gustafsson & Olsson 2012a: 94, 96, figs 1–2.

*Lunaceps* sp. ("Clade 3"); Gustafsson & Olsson 2012a: 94, 96, figs 1–2.

*Lunaceps superciliosus* Gustafsson & Olsson, 2012b: 71, figs 29a,b,c,d.

Holotype ♂ in MONZ.

Type host: *Calidris acuminata* (Horsfield, 1821).

New Zealand host: *Calidris acuminata* (Horsfield, 1821).

Other host: *Calidris subminuta* (Middendorff, 1853).

New Zealand localities: MC, KE.

Geographic distribution: Asia; Australasia.

New Zealand references: Palma (1999); Murray *et al.* (2006a); Gustafsson & Olsson (2012b: 73).

Other significant references: Price *et al.* (2003); Gustafsson & Olsson (2012a).

Remarks: Although the sharp-tailed sandpiper is a regular annual visitor to New Zealand (Checklist Committee 2010: 196), there are only two records of *Luniceps superciliosus* from this country.

### Genus *Melibrueelia* Valim & Palma, 2015

*Melibrueelia* Valim & Palma, 2015. *Zootaxa*: 3926: 483. Type species: *Melibrueelia novaeseelandiae* Valim & Palma, 2015 (by original designation). Endemic to New Zealand, but see Remarks below.

#### *Melibrueelia novaeseelandiae* Valim & Palma, 2015

Figs 125–126

*Brueelia* sp. nov.; Watt 1971: 235, 244.

*Brueelia* sp.; Pilgrim & Palma 1982: 27.

*Brueelia* sp.; Murray *et al.* 2001: 1263.

*Brueelia* sp.; Palma 2010: 408.

*Melibrueelia novaeseelandiae* Valim & Palma, 2015: 484, figs 1–3, 6a,b,c, 7a,b.

Holotype ♂ in MONZ.

Type host: *Prosthemadera novaeseelandiae novaeseelandiae* (J.F. Gmelin, 1788).

New Zealand hosts: *Anthornis melanura obscura* Falla, 1948; *Anthornis melanura oneho* Bartle & Sagar, 1987; *Anthornis melanura melanura* (Sparrman, 1786); *Prosthemadera novaeseelandiae novaeseelandiae* (J.F. Gmelin, 1788).

Other hosts: None.

New Zealand localities: ND, AK, CL, BP, WN, BR, WD, KE, AU.

Geographic distribution: New Zealand.

New Zealand references: Watt (1971); Pilgrim & Palma (1982); Murray *et al.* (2001); Palma (2010); Valim & Palma (2015).

Other significant references: None.

Remarks: *Melibrueelia novaeseelandiae* is an endemic species parasitic on members of the bird family Meliphagidae, the honeyeaters (Checklist Committee 2010: 288). Although *Melibrueelia* is at present endemic to New Zealand, there is a probability that species of this genus will be found on Australian species of honeyeaters.

### Genus *Naubates* Bedford, 1930

#### Subgenus *Guenterion* Palma & Pilgrim, 2002

*Naubates* Harrison, 1937. *Sci. Rep., Australasian Ant. Exped. 1911–14, series C, Zool. Bot.*: 30. Type species: *Naubates heteroproctus* Harrison, 1937 = *Naubates (Guenterion) heteroproctus* Harrison, 1937 (by original designation). Preoccupied by *Naubates* Bedford, 1930.

*Guenterion* Palma & Pilgrim, 2002. *Jour. Roy. Soc. New Zealand* 32: 29. Type species: *Lipeurus clypeatus* Giebel, 1874 = *Naubates (Guenterion) clypeatus* (Giebel, 1874) (by original designation).

#### *Naubates (Guenterion) clypeatus* (Giebel, 1874)

*Lipeurus clypeatus* Giebel, 1874: 236.

*Esthiopterum clypeatum* Giebel, 1874 [sic]; Harrison 1916: 132.

*Naubates clypeatus* (Giebel, 1874); Hopkins & Clay 1952: 234.

*Naubates clypeatus* (Giebel, 1874); Timmermann 1961d: 187, fig. 7.

*Naubates clypeatus* (Giebel, 1874); Timmermann 1965: 123, fig. 61.

*Naubates clypeatus* (Giebel, 1874); Wise 1977: 62.

*Naubates* sp.; Lowry *et al.* 1978: 138.

*Naubates clypeatus* (Giebel, 1874); Pilgrim & Palma 1982: 9.

*Naubates (Guenterion) clypeatus* (Giebel, 1874); Palma & Pilgrim 2002: 31, figs 5–6, 10, 17, 24, 40, 45, 52, 63.

*Naubates (Guenterion) clypeatus* (Giebel, 1874); Palma 2010: 408.

Syntypes ♂♀, presumed lost. See Palma & Pilgrim (2002: 31).

Type host: *Halobaena caerulea* (J.F. Gmelin, 1789).

New Zealand host: *Halobaena caerulea* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: AK, BP, TK, WI, WN, NC, MC, SC, Macquarie Island.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Wise (1977); Lowry *et al.* (1978); Pilgrim & Palma (1982); Palma (1996b: 191); Murray *et al.* (1990: 1370); Palma & Horning (2002: 9, 16); Palma & Pilgrim (2002); Palma (2010).

Other significant references: Giebel (1876: 389); Thompson (1935c: 487); Timmermann (1961d; 1965); Green & Palma (1991: 15, 26); Price *et al.* (2003: 199).

Remarks: *Naubates (Guenterion) clypeatus* is frequently collected from the wings of blue petrels. As discussed by Palma & Pilgrim (2002: 35, 42), *Naubates (Guenterion) clypeatus* has been reported from species of *Pachyptila*, but those records are likely to be misidentifications of *N. (G.) prioni* (see below).

### ***Naubates (Guenterion) damma* Timmermann, 1961**

*Naubates damma* Timmermann, 1961d: 185, fig. 6, 9 (bottom).

*Naubates pterodromi* Bedford, 1930; Timmermann 1961d: 183. In part.

*Naubates damma* Timmermann, 1961; Timmermann 1965: 123, fig. 60.

*Naubates damma* Timmermann, 1961; Pilgrim & Palma 1982: 9.

*Naubates (Guenterion) damma* Timmermann, 1961; Palma & Pilgrim 2002: 42, figs 3, 11, 18, 25, 42, 47, 53, 62, 66, 68–69.

*Naubates (G.) damma* Timmermann, 1961; Palma 2010: 408.

Holotype ♂ in NHML (Palma & Pilgrim 2002: 43).

Type host: *Pterodroma leucoptera* (Gould, 1844).

New Zealand hosts: *Pterodroma cervicalis* (Salvin, 1891); *Pterodroma cookii* (G.R. Gray, 1843); *Pterodroma leucoptera caledonica* Imber & Jenkins, 1981.

Other hosts: *Pterodroma externa* (Salvin, 1875); *Pterodroma arminjoniana* (Giglioli & Salvadori, 1869); *Pterodroma phaeopygia* (Salvin, 1876); *Pterodroma sandwichensis* (Ridgway, 1884); *Pterodroma hypoleuca* (Salvin, 1888).

New Zealand localities: ND, AK, CL, WI, KE, SI.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990: 1370); Palma & Pilgrim (2002); Palma (2010).

Other significant references: Timmermann (1965); Amerson & Emerson (1971: 4, 24); Palma (1996b: 192); Price *et al.* (2003: 199).

Remarks: *Naubates (Guenterion) damma* is frequently collected from the wings of its hosts, and it is morphologically intermediate between the two *Naubates* species from *Pachyptila* + *Halobaena*, and those from species of *Pterodroma* (see Palma & Pilgrim 2002: 44).

### ***Naubates (Guenterion) heteroproctus* Harrison, 1937**

*Naubates heteroproctus* Harrison, 1937: 30, pl. 2: figs 4–7.

*Naubates heteroproctus* Harrison, 1937; Hopkins & Clay 1952: 235.

*Naubates heteroproctus* Harrison, 1937; Timmermann 1961d: 184, fig. 9 (upper).

*Naubates heteroproctus* Harrison, 1937; Timmermann 1965: 122, fig. 63b, pl. 3: fig. 4.

*Naubates heteroproctus* Harrison, 1937; Wise 1977: 62.

*Naubates heteroproctus* Harrison, 1937; Pilgrim & Palma 1982: 8, 29.

*Naubates (Guenterion) heteroproctus* Harrison, 1937; Palma & Pilgrim 2002: 51, figs 55, 59.

*Naubates (G.) heteroproctus* Harrison, 1937; Palma 2010: 408.

Syntypes ♂♀ & nymph, presumed lost. See Palma & Pilgrim (2002: 52).

Type host: “*Pterodroma lessonii*”, in error (see Palma & Pilgrim 2002: 52).

New Zealand host: *Pterodroma macroptera gouldi* (Hutton, 1869).

Other hosts: *Pterodroma macroptera* (A. Smith, 1840).

New Zealand localities: ND, AK, CL, BP, WN, NC, MC, SC, SL, Macquarie Island.

Geographic distribution: Atlantic, Indian and Pacific Oceans.

New Zealand references: Harrison (1937); Thompson (1938d: 487); Gressitt (1970: 328); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1369); Palma & Horning (2002: 9, 20); Palma & Pilgrim (2002); Page *et al.* (2004: 638, 649); Palma (2010).

Other significant references: Séguy (1953: 572, figs 24–25); Timmermann (1961d; 1965); Green & Palma (1991: 15, 26); Palma (1996b: 192); Price *et al.* (2003: 199); Hänel & Palma (2007: 113, 125, 130).

Remarks: *Naubates (Guenterion) heteroproctus* is frequently collected from the wings of grey-faced petrels. At present, females of *N. (G.) heteroproctus* are indistinguishable from those of *N. (G.) pterodromi* and *N. (G.) lessonii* (see Palma & Pilgrim 2002: 31, 48).

### ***Naubates (Guenterion) lessonii* Palma & Pilgrim, 2002**

Figs 127–128

“*Naubates heteroproctus*” Watson, 1967: 72 (not *Naubates heteroproctus* Harrison, 1937).

“*Naubates fuliginosus*” Clay & Moreby, 1970: 217 (not *Lipeurus fuliginosus* Taschenberg, 1882).

“*Naubates heteroproctus*” Clay & Moreby, 1970: 218 (not *Naubates heteroproctus* Harrison, 1937).

*Naubates* sp.; Pilgrim & Palma 1982: 8, 30.

*Naubates* sp.; Murray *et al.* 1990: 1369.

*Naubates pterodromi sensu lato*; Green & Palma 1991: 15, 26.

*Naubates pterodromi*; Palma 1996b: 193. In part.

*Naubates pterodromi* Bedford, 1930 *sensu lato*; Palma & Horning 2002: 9, 16, 21.

*Naubates (Guenterion) lessonii* Palma & Pilgrim, 2002: 50, figs 43, 58, 70–71.

*Naubates (G.) lessonii* Palma & Pilgrim, 2002; Palma 2010: 408.

Holotype ♂ in MONZ.

Type host: *Pterodroma lessonii* (Garnot, 1826).

New Zealand host: *Pterodroma lessonii* (Garnot, 1826).

Other hosts: None.

New Zealand localities: AK, TK, WI, WN, WD, AU, Macquarie Island.

Geographic distribution: Southern Oceans.

New Zealand references: Watson (1967); Clay & Moreby (1970); Gressitt (1970: 327); Wise (1977: 62); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b); Palma & Horning (2002); Palma & Pilgrim (2002); Palma (2010).

Other significant references: Green & Palma (1991); Price *et al.* (2003: 199).

Remarks: *Naubates (Guenterion) lessonii* is frequently collected from the wings of white-headed petrels. At present, females of *N. (G.) lessonii* are indistinguishable from those of *N. (G.) pterodromi* and *N. (G.) heteroproctus* (see Palma & Pilgrim 2002: 31, 48).

### ***Naubates (Guenterion) prioni* (Enderlein, 1908)**

*Lipeurus prioni* Enderlein, 1908: 454, figs 194, 196–199.

*Esthiopterum clypeatum* Enderlein, 1908 [sic]; Harrison 1916: 140. In part.

“*Naubates (Micronaubates) clypeatus*” Pessôa & Guimarães, 1935: 112, fig. 12 (not *Lipeurus clypeatus* Giebel, 1874).

“*Naubates clypeatus*” Harrison, 1937: 31 (not *Lipeurus clypeatus* Giebel, 1874).

*Naubates prioni* (Enderlein, 1908); Hopkins & Clay 1952: 235.

*Naubates prioni* (Enderlein, 1908); Timmermann 1961d: 187, fig. 8.

*Naubates prioni* (Enderlein, 1908); Timmermann 1965: 123, figs 62a,b, pl. 3: fig. 3.

*Naubates prioni* (Enderlein, 1908); Watson 1967: 72.

*Naubates* sp.; Watson 1967: 72.

*Naubates prioni* (Enderlein, 1908); Wise 1977: 62.

*Naubates prioni* (Enderlein, 1908); Horning *et al.* 1980: 6, 10.

*Naubates prioni* (Enderlein, 1908); Pilgrim & Palma 1982: 10.

*Naubates prioni* (Enderlein, 1908); Mey 1994: 24, figs 9–10.

*Naubates (Guenterion) prioni* (Enderlein, 1908); Palma & Pilgrim 2002: 35, figs 16, 23, 41, 46, 51, 64.

*Naubates (G.) prioni* (Enderlein, 1908); Palma 2010: 408.

Lectotype ♂ in ZMHU (Göllner-Scheiding 1973: 41; Palma & Pilgrim 2002: 42).



Type host: *Pachyptila desolata* (J.F. Gmelin, 1789).

New Zealand hosts: *Pachyptila vittata* (G. Forster, 1777); *Pachyptila salvini salvini* (Mathews, 1912); *Pachyptila desolata* (J.F. Gmelin, 1789); *Pachyptila belcheri* (Mathews, 1912); *Pachyptila turtur* (Kuhl, 1820); *Pachyptila crassirostris crassirostris* (Mathews, 1912); *Pachyptila crassirostris pyramidalis* Fleming, 1939; *Pachyptila crassirostris flemingi* Tennyson & Bartle, 2005.

Other hosts: None.

New Zealand localities: ND, AK, CL, BP, WO, HB, TK, WI, WA, WN, SD, MB, NN, NC, MC, SC, WD, DN, SL, KE, CH, SI, BO, SN, AN, AU, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Harrison (1937); Watson (1967); Clay & Moreby (1970: 218); Gressitt (1970: 328); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990: 1370); Palma (1996b: 193); Paterson *et al.* (1999: 222); Marris (2000: 187); Palma & Horning (2002: 9, 16, 20); Palma & Pilgrim (2002); Page *et al.* (2004: 638, 649); Palma (2010).

Other significant references: Séguy (1953: 569, figs 21–23); Timmermann (1961d; 1965); Clay & Moreby (1967: 162, 168, fig. 89); Green & Palma (1991: 15, 26); Furness & Palma (1992: 35, 39); Mey (1994); Price *et al.* (2003: 199); Hänel & Palma (2007: 113, 125, 130).

Remarks: *Naubates (Guenterion) prioni* is both highly prevalent and abundant on all its hosts, and predominantly found on their wings. Records of *Naubates (Guenterion) clypeatus* from species of *Pachyptila* are likely to be misidentifications of *N. (G.) prioni* (see Palma & Pilgrim 2002: 35, 42).

#### ***Naubates (Guenterion) pterodromi* Bedford, 1930**

*Naubates pterodromi* Bedford, 1930: 170, figs 10, 13; *sensu* Kéler 1952.

*Naubates pterodromi* Bedford, 1930; Kéler 1952: 213, figs 6–10.

*Naubates pterodromi* Bedford, 1930; Timmermann 1961d: 183, figs 2, 5, 9 (middle), pl. 1: figs c,d. In part.

*Naubates pterodromi* Bedford, 1930; Timmermann 1965: 121, figs 54, 57, 59, 63, pl. 4: figs 3–4.

*Naubates pterodromi* Bedford, 1930; Palma & Pilgrim 1977: 290.

*Naubates pterodromi* Bedford, 1930; Horning *et al.* 1980: 6, 9.

*Naubates pterodromi* Bedford, 1930; Pilgrim & Palma 1982: 8, 30.

*Naubates (Guenterion) pterodromi* Bedford, 1930; Palma & Pilgrim 2002: 45, figs 4, 13, 15, 22, 49, 54, 57, 65.

*Naubates (G.) pterodromi* Bedford, 1930; Palma 2010: 408.

Holotype ♀ in SAMS, unidentifiable (Palma & Pilgrim 2002: 49). Pragmatype ♂ in SAIM (see Palma & Pilgrim 2002: 50)

Type host: “*Pterodroma macroptera*”, in error (see Palma & Pilgrim 2002: 48).

New Zealand hosts: *Pterodroma mollis* (Gould, 1844); *Pterodroma inexpectata* (J.R. Forster, 1844).

Other host: *Pterodroma incerta* (Schlegel, 1863).

New Zealand localities: ND, AK, WN, NC, MC, SC, SL, SI, SN, AN.

Geographic distribution: Atlantic, Indian and Pacific Oceans.

New Zealand references: Palma & Pilgrim (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990: 1370); Palma (1999: 378); Marris (2000: 187); Palma & Pilgrim (2002); Page *et al.* (2004: 638, 649); Palma (2010).

Other significant references: Kéler (1952); Timmermann (1961d; 1965); Green & Palma (1991: 15, 26); Furness & Palma (1992: 35, 40); Zonfrillo (1993: 327); Palma (1996b: 193); Price *et al.* (2003: 199); Hänel & Palma (2007: 113, 125, 130).

Remarks: *Naubates (Guenterion) pterodromi* is frequently collected from the wings of its hosts. Palma & Pilgrim (2002: 48) discussed the problems arising from the original concept of *N. (G.) pterodromi* based on females only, which are at present indistinguishable from those of *N. (G.) heteroproctus* and *N. (G.) lessonii* (see Palma & Pilgrim 2002: 31).

**Subgenus *Naubates* Bedford, 1930**

*Naubates* Bedford, 1930. *16th Report Director Vet. Services Animal Ind. Union of South Africa*: 167. Type species: *Esthiopterum fuliginosum* (Taschenberg, 1882) = *Naubates (Naubates) fuliginosus* (Taschenberg, 1882) (by original designation).

***Naubates (Naubates) fuliginosus* (Taschenberg, 1882)**

*Lipeurus fuliginosus* Taschenberg, 1882: 156, pl. 4: fig. 3.  
*Esthiopterum fuliginosum* Taschenberg, 1882 [sic]; Harrison 1916: 134.  
*Naubates fuliginosus* (Taschenberg, 1882); Bedford 1930: 168, figs 9, 11, 15, 16a.  
*Naubates fuliginosus* (Taschenberg, 1882); Hopkins & Clay 1952: 235.  
*Naubates fuliginosus* (Taschenberg, 1882); Timmermann 1961d: 177, figs 1, 3, pl. 1: figs a,b.  
*Naubates fuliginosus* (Taschenberg, 1882); Timmermann 1965: 118, figs 53, 55, pl. 4: figs 1–2.  
*Naubates fuliginosus* (Taschenberg, 1882); Wise 1977: 62.  
*Naubates fuliginosus* (Taschenberg, 1882); Pilgrim & Palma 1982: 11.  
*Naubates (Naubates) fuliginosus* (Taschenberg, 1882); Palma & Pilgrim 2002: 12, figs 7–8, 19, 26, 29, 31, 35.  
*Naubates (Naubates) fuliginosus* (Taschenberg, 1882); Palma 2010: 408.

Syntypes ♂♀, presumed lost. See Palma & Pilgrim (2002: 12).

Type hosts: *Diomedea exulans* Linnaeus, 1758 & *Thalassarche chlororhynchos* (J.F. Gmelin, 1789), both in error (see Palma & Pilgrim 2002: 18).

New Zealand hosts: *Procellaria aequinoctialis* Linnaeus, 1758; *Procellaria westlandica* Falla, 1946; *Procellaria parkinsoni* G.R. Gray, 1862; *Procellaria cinerea* J.F. Gmelin, 1789.

Other hosts: None.

New Zealand localities: ND, AK, CL, BP, WI, WN, NC, MC, SC, WD, SN, AN, AU, CA, Macquarie Island.

Geographic distribution: Southern Oceans.

New Zealand references: Gressitt (1970: 327); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1371); Palma (1996b: 192); Marris (2000: 187); Palma & Horning (2002: 9, 16); Palma & Pilgrim (2002); Page *et al.* (2004: 638, 648); Palma (2010).

Other significant references: Bedford (1930); Kéler (1957c: fig. 23b); Timmermann (1961d; 1965); Clay & Moreby (1967: 162, 168, fig. 88); Price *et al.* (2003: 199).

Remarks: *Naubates (Naubates) fuliginosus* is both highly prevalent and abundant on all its hosts, and predominantly found on their wings.

***Naubates (Naubates) harrisoni* Bedford, 1930**

*Naubates harrisoni* Bedford, 1930: 168, figs 12, 14, 16b.  
*Naubates harrisoni* Bedford, 1930; Hopkins & Clay 1952: 235.  
*Naubates harrisoni* Bedford, 1930; Timmermann 1961d: 180, fig. 4.  
*Naubates harrisoni* Bedford, 1930; Timmermann 1965: 119, figs 56, 58.  
*Naubates harrisoni* Bedford, 1930; Watt 1971: 236, 243, fig. 6.  
*Naubates harrisoni* Bedford, 1930; Pilgrim & Palma 1982: 11.  
*Naubates harrisoni* Bedford, 1930 *s. l.*; Pilgrim & Palma 1982: 12.  
*Naubates (Naubates) harrisoni* Bedford, 1930; Palma & Pilgrim 2002: 20, figs 1–2, 20, 27, 30, 32, 36, 38.  
*Naubates (N.) harrisoni* Bedford, 1930; Palma 2010: 408.

Holotype ♂ in SAMS (Palma & Pilgrim 2002: 20).

Type host: *Puffinus gravis* (O'Reilly, 1818).

New Zealand hosts: *Puffinus pacificus pacificus* (J.F. Gmelin, 1789); *Puffinus pacificus chlororhynchus* Lesson, 1831; *Puffinus bulleri* Salvin, 1888; *Puffinus carneipes* Gould, 1844; *Puffinus huttoni* Mathews, 1912.

Other hosts: *Puffinus nativitatis* Streets, 1877; *Puffinus puffinus* (Brünnich, 1764); *Puffinus opisthomelas* Coues, 1864; *Puffinus lherminieri lherminieri* Lesson, 1839; *Puffinus assimilis baroli* (Bonaparte, 1857); *Puffinus assimilis boydi* Mathews, 1912; *Puffinus assimilis tunneyi* Mathews, 1912; *Puffinus tenuirostris* (Temminck, 1835); *Puffinus creatopus* Coues, 1864.

New Zealand localities: ND, AK, CL, BP, WN, SD, MB, KA, NC, MC, SC, WD, CH, KE.

Geographic distribution: Atlantic, Indian and Pacific Oceans.

New Zealand references: Watt (1971); Wise (1977: 62); Pilgrim & Palma (1982); Murray *et al.* (1990: 1371); Paterson *et al.* (1999: 222); Palma & Pilgrim (2002); Galloway (2005: 16); Palma (2010).

Other significant references: Timmermann (1961d; 1965); Amerson & Emerson (1971: 6, 8, 24); Ward & Downey (1973: 394); Bourgeois & Threlfall (1979: 1356); Fowler & Shaw (1990: 15); Forrester *et al.* (1995: 6); Foster *et al.* (1996: 85); Palma (1996b: 192); Furness & Palma (1992: 35, 38); Price *et al.* (2003: 199); Page *et al.* (2004: 638, 648, 650); Hänel & Palma (2007: 113, 125, 130); Martín-Mateo (2009: 103, fig. 26).

Remarks: Pilgrim & Palma (1982: 12) regarded the population of *Naubates (Naubates) harrisoni* from *Puffinus huttoni* as somewhat different from those from several other hosts, and qualified it as *sensu lato*; however, my examination of more samples from all hosts shows that making such difference is not warranted (see also Palma & Pilgrim 2002: 20, 24). *Naubates (N.) harrisoni* is an infrequently collected species from all its hosts, and predominantly found on their wings.

### ***Naubates (Naubates) thieli* Timmermann, 1965**

Figs 129–130

*Naubates thieli* Timmermann, 1965: 121.

*Naubates thieli* Timmermann, 1965; Palma 1999: 378.

*Naubates (Naubates) thieli* Timmermann, 1965; Palma & Pilgrim 2002: 27, figs 9, 28, 33–34, 37, 39.

*Naubates (N.) thieli* Timmermann, 1965; Palma 2010: upper fig. p. 295, 408.

Holotype ♂ in ANIC (Palma & Pilgrim 2002: 27, 29).

Type host: *Pterodroma solandri* (Gould, 1844).

New Zealand host: *Pterodroma solandri* (Gould, 1844).

Other hosts: None.

New Zealand locality: ND.

Geographic distribution: Pacific Ocean.

New Zealand references: Palma (1999); Palma & Pilgrim (2002); Palma (2010).

Other significant references: Spratt (1983: 58); Palma (1996b: 193); Price *et al.* (2003: 199).

Remarks: *Pterodroma solandri* is an Australian petrel regarded as a “straggler to the northern North Island” by the Checklist Committee (2010: 89). *Naubates (Naubates) thieli* is a “wing” species and likely to be the result of a host-switch of *N. (N.) harrisoni*, or its ancestor, from a species of *Puffinus* onto *Pterodroma solandri* (see Palma & Pilgrim 2002: 28).

### **Genus *Neopsittaconirmus* Conci, 1942**

*Neopsittaconirmus* Conci, 1942c [20 May]. *Boll. Soc. Entomol. Italiana* 74: 37. Type species: *Neopsittaconirmus borgioli* Conci, 1942c (by original designation).

*Psittacicola* Guimarães, 1942 [28 November]. *Papéis Avulsos Dept. Zool. (São Paulo)* 2: 80. Type species: *Esthiopterum kea* (Kellogg, 1907) = *Neopsittaconirmus kea* (Kellogg, 1907) (by original designation).

### ***Neopsittaconirmus albus* (Le Souëf & Bullen, 1902)**

*Lipeurus albus* Le Souëf & Bullen, 1902a: 157, fig. 4.

*Esthiopterum album* Le Souëf & Bullen, 1902 [sic]; Harrison 1916: 130 (as junior synonym of *Esthiopterum capreolum* Gervais, 1847).

*Neopsittaconirmus albus* (Le Souëf & Bullen, 1902); Hopkins & Clay 1952: 237.

*Lipeurus albus* Le Souëf & Bullen, 1902; Guimarães 1974a: 187 (as “Species inquirendae”).

*Neopsittaconirmus albus* (Le Souëf & Bullen, 1902); Price & Emerson 1978: 33, figs 1–8.

*Neopsittaconirmus albus* (Le Souëf & Bullen, 1902); Palma 1999: 381.

*Neopsittaconirmus albus* (Le Souëf & Bullen, 1902); Palma 2010: 408.

Syntypes ♂♀, presumed lost. See Palma (1996b: 194).

Type host: *Cacatua galerita* (Latham, 1790).

New Zealand host: *Cacatua galerita* (Latham, 1790).

Other hosts: None.

New Zealand locality: WI.

Geographic distribution: Australasia.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Guimarães (1974a); Price & Emerson (1978); Green & Palma (1991: 15, 34); Palma (1996b: 194); Price *et al.* (2003: 200).

Remarks: It is not possible to assert if *Neopsittaconirmus albus* was introduced to New Zealand with sulphur-crested cockatoos by human agency, or if it was self-introduced with its host (Checklist Committee 2010: 252).

### ***Neopsittaconirmus kea* (Kellogg, 1907)**

Figs 131–132

*Lipeurus circumfasciatus* var. *kea* Kellogg, 1907: 122.

*Esthiopterum kea* Kellogg, 1907 [sic]; Harrison 1916: 136.

*Neopsittaconirmus kea* (Kellogg, 1907); Conci 1942c: 37.

*Psittacicola kea* (Kellogg, 1907); Guimarães 1942: 81, figs 1–5.

*Neopsittaconirmus kea* (Kellogg, 1907); Hopkins & Clay 1952: 238.

*Neopsittaconirmus kea* (Kellogg, 1907); Pilgrim 1970: 75.

*Neopsittaconirmus kea*; Miller 1971: 132.

*Neopsittaconirmus kea* (Kellogg, 1907); Guimarães 1974a: 181, figs 123–126.

*Neopsittaconirmus kea* (Kellogg, 1907); Wise 1977: 62.

*Neopsittaconirmus kea* (Kellogg, 1907); Pilgrim & Palma 1982: 24.

*Neopsittaconirmus kea* (Kellogg, 1907); Murray *et al.* 1999: 1241.

*Neopsittaconirmus kea* (Kellogg, 1907); Palma 2010: 408.

Syntypes ♂♀ in EMEC (Peter T. Oboyski pers. comm. October 2014).

Type host: *Nestor notabilis* Gould, 1856.

New Zealand hosts: *Nestor meridionalis septentrionalis* Lorenz, 1896; *Nestor meridionalis meridionalis* (J.F. Gmelin, 1788); *Nestor notabilis* Gould, 1856.

Other hosts: None.

New Zealand localities: ND, CL, HB, WA, WN, NN, NC, MC, WD, SI.

Geographic distribution: New Zealand.

New Zealand references: Kellogg (1907); Guimarães (1942: 83); Pilgrim (1970); Miller (1971: 132); Guimarães (1974a); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1999); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Conci (1942c); Price *et al.* (2003: 200); Sychra (2006: 65, fig. 8).

Remarks: *Neopsittaconirmus kea* is an endemic and “at risk” species (Buckley *et al.* 2012), exclusively parasitic on members of the parrot genus *Nestor*.

### **Genus *Nesiotinus* Kellogg, 1903**

*Nesiotinus* Kellogg, 1903. *Biol. Bull. Wood's Hole* 5(2): 89. Type species: *Nesiotinus demersa* [sic] Kellogg, 1903 = *Nesiotinus demersus* Kellogg, 1903 (by monotypy).

### ***Nesiotinus demersus* Kellogg, 1903**

Figs 133–134

*Nesiotinus demersa* [sic] Kellogg, 1903: 90, fig. 3.

*Nesiotinus demersus* Kellogg, 1903; Harrison 1916: 74. Emendation.

*Nesiotinus demersus* Kellogg, 1903; Hopkins & Clay 1952: 239.

*Nesiotinus demersus* Kellogg, 1903; Clay & Moreby 1967: 159, figs 76–77, 94–95.

*Nesiotinus demersus* Kellogg, 1903; Watson 1967: 72.

*Nesiotinus demersus* Kellogg, 1903; Wise 1977: 62.

*Nesiotinus* sp.; Lowry *et al.* 1978: 137.

*Nesiotinus demersus* Kellogg, 1903; Pilgrim & Palma 1982: 4.

*Nesiotinus demersus* Kellogg, 1903; Murray *et al.* 1990: 1367.

*Nesiotinus demersus* Kellogg, 1903; Banks & Paterson 2004: 96, figs 3e, 7a, 9–11.

*Nesiotinus demersus* Kellogg, 1903; Palma 2010: 408.

Holotype ♀ in EMEC.

Type host: *Aptenodytes patagonicus* Miller, 1778.

New Zealand host: *Aptenodytes patagonicus* Miller, 1778.

Other hosts: None.

New Zealand locality: Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Watson (1967); Gressitt (1970: 328); Wise (1977); Lowry *et al.* (1978); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b: 195); Palma & Horning (2002: 9, 17); Banks & Paterson (2004); Banks *et al.* (2006: 158); Palma (2010).

Other significant references: Cummings (1916c: 172); Kéler (1960b: 177, figs 1–6); Eichler (1941a: 361, fig. 26); Clay & Moreby (1967); Price *et al.* (2003: 200); Mey (2011: 79, figs 7, 11, 14, 16, 18–19, 22–23, 25–28).

Remarks: A second species of *Nesiotinus* described by Mey (2011) based on a single female louse from a Subantarctic diving petrel (*Pelecanoides urinatrix exsul* Salvin, 1896) needs confirmation of both its identity and its host association.

### Genus *Ornithobius* Denny, 1842

*Ornithobius* Denny, 1842. *Mon. Anopl. Brit.*: 183. Type species: *Ornithobius cygni* (Linnaeus, 1758) (by subsequent designation). As a subgenus of *Philopterus*.

#### *Ornithobius bucephalus* (Giebel, 1874)

*Lipeurus bucephalus* Giebel, 1874: 239.

*Ornithobius bucephalus* Giebel, 1874 [sic]; Harrison 1916: 128 (as junior synonym of *Ornithobius cygni* (Linnaeus, 1758)).

*Ornithobius bucephalus* (Giebel, 1874); Hopkins & Clay 1952: 253.

*Ornithobius bucephalus bucephalus* (Giebel, 1874); Timmermann 1962b: 136, fig. 5a.

*Ornithobius bucephalus tribulis* Timmermann, 1962b: 136, figs 1, 5b.

*Ornithobius bucephalus* (Giebel, 1874); Pilgrim & Palma 1982: 16.

*Ornithobius bucephalus* (Giebel, 1874); Murray *et al.* 1990: 1374.

*Ornithobius bucephalus* (Giebel, 1874); Arnold 2005: 159, figs 4, 12.

*Ornithobius bucephalus* (Giebel, 1874); Palma 2010: 408.

Holotype ♀, probably lost (Palma 1996b: 195).

Type host: *Cygnus olor* (J.F. Gmelin, 1789).

New Zealand host: *Cygnus olor* (J.F. Gmelin, 1789).

Other host: *Cygnus melancoryphus* (Molina, 1782).

New Zealand localities: WN, MC.

Geographic distribution: Eurasia; Africa; South America, New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Timmermann (1962b); Palma (1996b: 195); Price *et al.* (2003: 202); Arnold (2005).

Remarks: *Ornithobius bucephalus* was introduced to New Zealand with mute swans by human agency (Checklist Committee 2010: 32).

#### *Ornithobius fuscus* Le Souëf, 1902

Figs 135–136

*Ornithobius fuscus* Le Souëf, 1902b: 91.

*Ornithobius fuscus* Le Souëf, 1902; Hopkins & Clay 1952: 253.

*Ornithobius fuscus* Le Souëf, 1902; Timmermann 1962b: 137, figs 4, 5c.

*Ornithobius fuscus* Le Souëf, 1902; Pilgrim & Palma 1982: 16.

*Ornithobius fuscus* Le Souëf, 1902; Murray *et al.* 1990: 1374.

*Ornithobius fuscus* Le Souëf, 1902; Arnold 2005: 162, figs 7, 15, 23.

*Ornithobius fuscus* Le Souëf, 1902; Palma 2010: 408.

Holotype ♀, probably lost (Palma 1996b: 196).

Type host: *Cygnus atratus* (Latham, 1790).

New Zealand host: *Cygnus atratus* (Latham, 1790).

Other hosts: none.

New Zealand localities: AK, WO, WA, WN, SD, MB, NC, MC, SC.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Arnold (2005: 163); Palma (2010).

Other significant references: Timmermann (1962b); Green & Palma (1991: 16, 29); Palma (1996b: 196); Price *et al.* (2003: 202).

Remarks: The population of black swans living in New Zealand is probably the result of a mixture of self-introduced birds and others introduced by human agency (Checklist Committee 2010: 33). Therefore, it is not possible to determine if *Ornithobius fuscus* is a native or an introduced species.

### ***Ornithobius goniopleurus* (Denny, 1842)**

*Phlopterus (Ornithobius) goniopleurus* [sic] Denny, 1842: 60, 184, pl. 23: fig. 2.

*Ornithobius goniopleurus* Denny, 1842 [sic]; Harrison 1916: 129. Emendation.

*Ornithobius goniopleurus* Denny, 1842 [sic]; Hopkins & Clay 1952: 253.

*Ornithobius goniopleurus* Denny, 1842 [sic]; Pilgrim & Palma 1982: 16.

*Ornithobius goniopleurus* Denny, 1842 [sic]; Murray *et al.* 1990: 1374.

*Ornithobius goniopleurus* Denny, 1842 [sic]; Arnold 2005: 163, figs 1, 3, 8, 16, 24–25.

*Ornithobius goniopleurus* Denny, 1842 [sic]; Palma 2010: 408.

Syntypes ♂♀ in NHML (Vincent S. Smith pers. comm. December 2014).

Type host: *Branta canadensis* (Linnaeus, 1758).

New Zealand host: *Branta canadensis maxima* Delacour, 1951.

Other hosts: *Anser rossii* Cassin, 1861; *Branta canadensis hutchinsii* (Richardson, 1832); *Branta leucopsis* (Bechstein, 1803).

New Zealand localities: BP, MC.

Geographic distribution: North America; Europe; north-east Asia; New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Séguy (1944: 336, figs 502–504); Balát (1974: 3, 5, figs 1d, 2b, 3c, 4b); Price *et al.* (2003: 202); Arnold (2005).

Remarks: *Ornithobius goniopleurus* was introduced to New Zealand with Canada geese by human agency (Checklist Committee 2010: 36). Records of this louse species from *Anas strepera* Linnaeus, 1758 and *Cygnus atratus* in Arnold (2005: 164) are regarded as stragglers or contaminants.

Denny (1842: 183) clearly regarded *Ornithobius* as a subgenus of *Phlopterus*. However, subsequent authors ignored that fact and did not add parentheses around author and date when *Ornithobius* was used at the generic level.

### **Genus *Oxylipeurus* Mjöberg, 1910**

*Oxylipeurus* Mjöberg, 1910a. *Arkiv Zool.* 6(13): 91. Type species: *Lipeurus inaequalis* Piaget, 1880 = *Oxylipeurus inaequalis* (Piaget, 1880) (by original designation).

*Epicolinus* Carriker, 1945a. *Rev. Bras. Entomol.* 5(1): 104. Type species: *Lipeurus clavatus* McGregor, 1917 = *Oxylipeurus clavatus* (McGregor, 1917) (by original designation).

### ***Oxylipeurus clavatus* (McGregor, 1917)**

*Lipeurus clavatus* McGregor, 1917: 115, pl. 7: fig. 3

*Oxylipeurus clavatus* (McGregor, 1917); Clay 1938: 168, pl. 9: fig. 5.

*Epicolinus clavatus* (McGregor, 1917); Carriker 1945a: 105, figs 30–34.

*Oxylipeurus clavatus* (McGregor, 1917); Hopkins & Clay 1952: 256.

*Epicolinus clavatus* (McGregor, 1917); Kéler 1958b: 319, 321, figs 9–10.

*Oxylipeurus clavatus* (McGregor, 1917); Pilgrim & Palma 1982: 18.

*Oxylipeurus clavatus* (McGregor, 1917); Palma 2010: 408.

Holotype ♀ in USNM.

Type host: *Colinus virginianus texanus* (Lawrence, 1853).

New Zealand host: *Colinus virginianus taylori* Lincoln, 1915.

Other hosts: *Colinus virginianus virginianus* (Linnaeus, 1758); *Colinus virginianus floridanus* (Coues, 1872).

New Zealand locality: WN.

Geographic distribution: Central and North America.

New Zealand references: Pilgrim & Palma (1982); Palma (2010).

Other significant references: Clay (1938); Carriker (1945a); Kéler (1958b); Forrester *et al.* (1995: 22); Price *et al.* (2003: 203).

Remarks: *Oxylipeurus clavatus* was introduced to New Zealand with its host by human agency, but *Colinus virginianus* appears to have died out in New Zealand (Checklist Committee 2010: 349).

### ***Oxylipeurus ellipticus* (Kéler, 1958)**

*Epicolinus ellipticus* Kéler, 1958b: 322, figs 55–56.

*Oxylipeurus ellipticus* (Kéler, 1958); Pilgrim & Palma 1982: 18.

*Oxylipeurus ellipticus* (Kéler, 1958); Murray *et al.* 1993: 960.

*Oxylipeurus ellipticus* (Kéler, 1958); Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Callipepla gambelii* (Gambel, 1843).

New Zealand host: *Callipepla californica brunnescens* (Ridgway, 1884).

Other hosts: None.

New Zealand localities: AK, BP, SD, MB, NN, NC, MC, SC, CO, DN.

Geographic distribution: North & South America; Europe; Australasia, Hawaiian Islands.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Galloway (2005: 17); Palma (2010).

Other significant references: Price *et al.* (2003: 203).

Remarks: *Oxylipeurus ellipticus* was introduced to New Zealand with California quails by human agency (Checklist Committee 2010: 25).

### ***Oxylipeurus mesopelios colchicus* Clay, 1938**

*Oxylipeurus mesopelios colchicus* Clay, 1938: 177, pl. 11: fig. 3.

*Oxylipeurus colchicus* Clay, 1938; Hopkins & Clay 1952: 256.

*Oxylipeurus mesopelios* (ex *Phasianus colchicus*); Kéler 1958b: 327, 331, figs 3, 32.

*Oxylipeurus mesopelios colchicus* Clay, 1938; Pilgrim & Palma 1982: 18.

*Oxylipeurus mesopelios colchicus* Clay, 1938; Murray *et al.* 1993: 961.

*Oxylipeurus colchicus* Clay, 1938; Price *et al.* 2003: 203.

*Oxylipeurus mesopelios colchicus* Clay, 1938; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Phasianus colchicus* Linnaeus, 1758.

New Zealand host: *Phasianus colchicus* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: RI, WI, WN.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 219); Palma (2010).

Other significant references: Kéler (1958b); Modrzejewska & Złotorzycka (1987: 665, fig. 6); Kopociński *et al.* (1998: 82); Price *et al.* (2003).

Remarks: *Oxylipeurus mesopelios colchicus* was introduced to New Zealand and other countries with common pheasants by human agency (Checklist Committee 2010: 28). In agreement with Clay (1938: 177) and contrary to Price *et al.* (2003: 203), I regard this louse taxon as a subspecies.

***Oxylipeurus polytrapezius polytrapezius* (Burmeister, 1838)**

Figs 137–138

*Lipeurus polytrapezius* Burmeister, 1838a: 434.*Oxylipeurus polytrapezius polytrapezius* (Burmeister, 1838); Clay 1938: 181, figs 37a,c, 39b, pl. 12: fig. 4.*Lipeurus (Oxylipeurus) polytrapezius* (Nitzsch, 1818) [sic]; Séguy 1944: 189, figs 272–274.*Oxylipeurus polytrapezius* (Burmeister, 1838); Hopkins & Clay 1952: 259.*Oxylipeurus polytrapezius* (Burmeister, 1838); Kéler 1958b: 327, 330, figs 35–36, 47–49.*Oxylipeurus polytrapezius polytrapezius* (Burmeister, 1838); Pilgrim & Palma 1982: 18.*Oxylipeurus p. polytrapezius* (Burmeister, 1838); Murray *et al.* 1993: 960.*Oxylipeurus polytrapezius polytrapezius* (Burmeister, 1838); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma &amp; Pilgrim (1984: 150).

Type host: *Meleagris gallopavo* Linnaeus, 1758.New Zealand host: *Meleagris gallopavo* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: WO, HB, WA, WN.

Geographic distribution: North America; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 220); Palma (2010).Other significant references: Clay (1938); Séguy (1944); Emerson (1962: 201, figs 10–12); Forrester *et al.* (1995: 21);Palma (1996b: 196); Price *et al.* (2003: 204); Martín-Mateo (2009: 76).Remarks: *Oxylipeurus polytrapezius polytrapezius* was introduced to New Zealand and other countries with turkeys by human agency (Checklist Committee 2010: 28). In agreement with Clay (1938: 181) and contrary to Price *et al.* (2003: 204), I regard this louse taxon as a subspecies.**Genus *Paraclisis* Timmermann, 1965***Paraclisis* Timmermann, 1965. *Abhandl. Verhandl. Naturwiss. Ver. Hamburg, N.F. 8 (Supplement)*: 96. Type species: *Pediculus diomedeeae* J.C. Fabricius, 1775 = *Paraclisis diomedeeae* (J.C. Fabricius, 1775) (by original designation).“*Harrisoniella* Kéler, 1956”: 521 (not *Harrisoniella* Bedford, 1930) See Clay & Hopkins (1961); I.C.Z.N. (1963).***Paraclisis diomedeeae* (J.C. Fabricius, 1775)***Pediculus diomedeeae* J.C. Fabricius, 1775: 808.*Ricinus diomedeeae* (Linnaeus, 1758); Latreille 1804: 107.*Lipeurus lepturus* Enderlein, 1908: 453, figs 200–202, 209.*Esthiopterum diomedeeae* J.C. Fabricius, 1775 [sic]; Harrison 1916: 133.*Perineus diomedeeae* (J.C. Fabricius, 1775); Clay 1940a: 299, figs, 1–2, 4a, 5a, 6a.*Perineus diomedeeae diomedeeae* (J.C. Fabricius, 1775); Eichler, 1949b: 340.*Perineus diomedeeae enderleini* Eichler, 1949b: 342, figs 8–10.*Perineus diomedeeae* (J.C. Fabricius, 1775); Hopkins & Clay 1952: 277.*Perineus enderleini* Eichler, 1949; Hopkins & Clay 1952: 277.*Perineus diomedeeae lepturus* (Enderlein, 1908); Séguy 1953: 567, fig. 16.*Harrisoniella diomedeeae* (J.C. Fabricius, 1775); Kéler 1956: 522, figs 1–3a.*Harrisoniella diomedeeae* (J.C. Fabricius, 1775); Kéler 1957a: 285, figs 1d, 2a, 3, 9.*Perineus* (s. l.) *diomedeeae* (J.C. Fabricius, 1775); Clay 1964a: 231, fig. 1.*Perineus* (s. l.) sp.; Clay 1964a: 231.*Perineus diomedeeae* (J.C. Fabricius, 1775); Gressitt 1964: 538.*Paraclisis diomedeeae* (J.C. Fabricius, 1775); Timmermann 1965: 100, figs 32, 33a, 34a, 37a, 38, 39a, pl. 5: figs 1–2.*Paraclisis diomedeeae* (J.C. Fabricius, 1775); Clay & Moreby 1967: 162, 168, figs 117–119, 122, 143.*Paraclisis diomedeeae* (J.C. Fabricius, 1775); Wise 1977: 62.*Paraclisis diomedeeae* (J.C. Fabricius, 1775); Horning *et al.* 1980: 6, 9.*Paraclisis diomedeeae* (J.C. Fabricius, 1775); Pilgrim & Palma 1982: 6.*Paraclisis diomedeeae* (J.C. Fabricius, 1775) s. l.; Pilgrim & Palma 1982: 7.*Paraclisis diomedeeae* (J.C. Fabricius, 1775); Palma 2010: 408.

Neotype ♂ in NHML (Clay 1940a: 302).

Type host: “In Brasiliae diomedeis” = *Thalassarche melanophris* (Temminck, 1828) (see Clay 1940a: 299).



New Zealand hosts: *Thalassarche chlororhynchos* (J.F. Gmelin, 1789); *Thalassarche chrysostoma* (J.R. Forster, 1785); *Thalassarche melanophris* (Temminck, 1828); *Thalassarche impavida* Mathews, 1912; *Thalassarche bulleri bulleri* (Rothschild, 1893); *Thalassarche bulleri platei* (Reichenow, 1898); *Thalassarche cauta stadii* Falla, 1933; *Thalassarche eremita* Murphy, 1930; *Thalassarche salvini* (Rothschild, 1893); *Phoebetria palpebrata* (J.R. Forster, 1785).

Other hosts: *Thalassarche cauta cauta* (Gould, 1841); *Phoebetria fusca* (Hilsenberg, 1822).

New Zealand localities: ND, AK, BP, GB, HB, TK, WI, WA, WN, NN, KA, NC, MC, SC, WD, CO, DN, SL, SI, CH, BO, SN, AN, AU, CA, Macquarie Island.

Geographic distribution: Atlantic, Pacific, Indian and Southern Oceans.

New Zealand references: Clay (1964a); Gressitt (1964); Watson (1967: 73); Clay & Moreby (1970: 217); Gressitt (1970: 328); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990: 1368–1369); Palma (1996b: 197); Marris (2000: 187); Palma (2001: 67, fig. 6); Palma & Horning (2002: 10, 15); Page *et al.* (2004: 641, 649); Palma (2010).

Other significant references: Clay (1940a); Clay & Hopkins (1951: 34); Séguy (1953: 565, figs 11–15); Timmermann (1965); Clay & Moreby (1967); Green & Palma (1991: 16, 25); Price *et al.* (2003: 205); Palma & Jensen (2005: 56, 60); Hänel & Palma (2007: 113, 126, 129).

Remarks: Pilgrim & Palma (1982: 7) regarded the population of *Paraclisis diomedea* from *Phoebetria palpebrata* as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted. *Thalassarche chlororhynchos* is a new host record for *P. diomedea* in New Zealand (voucher specimens in MONZ). *Paraclisis diomedea* is a “wing” louse species and the most prevalent among all lice parasitic on small albatrosses.

### ***Paraclisis hyalina* (Neumann, 1911)**

*Lipeurus hyalinus* Neumann, 1911: 21, pl. 2: figs 1a,d.

*Esthiopterum hyalinum* Neumann, 1911 [sic]; Harrison 1916: 136.

*Perineus hyalinus* (Neumann, 1911); Clay 1940a: 305, figs 3c, 4b, 5d, 6c.

*Perineus hyalinus* (Neumann, 1911); Hopkins & Clay 1952: 277.

*Harrisoniella hyalina* (Neumann, 1911); Kéler 1957a: 290, figs 2e, 5–7.

*Harrisoniella hyalina* (Neumann, 1911); Kéler 1958a: 380.

*Paraclisis hyalina* (Neumann, 1911); Timmermann 1965: 103, figs 33e, 34c, 37b, 39d, 40, pl. 6: fig. 1.

*Paraclisis hyalina* (Neumann, 1911); Clay & Moreby 1967: 162, 168, figs 85, 120, 124–125, 144.

*Perineus hyalinus*; Watson 1967: 73.

*Paraclisis hyalina* (Neumann, 1911); Wise 1977: 62.

*Paraclisis hyalina* (Neumann, 1911); Pilgrim & Palma 1982: 5.

*Paraclisis hyalina* (Neumann, 1911); Palma 2010: 408.

Syntypes ♂♀, repository unknown (Palma 1996b: 197).

Type host: *Diomedea exulans* Linnaeus, 1758.

New Zealand hosts: *Diomedea exulans* Linnaeus, 1758; *Diomedea antipodensis antipodensis* Robertson & Warham, 1992; *Diomedea antipodensis gibsoni* Robertson & Warham, 1992; *Diomedea epomophora* Lesson, 1825; *Diomedea sanfordi* Murphy, 1917.

Other host: *Diomedea dabbenena* Mathews, 1929.

New Zealand localities: HB, WA, WN, SD, MB, NC, MC, SC, WD, AN, AU, CA, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Kéler (1957a: 291); Kéler (1958a: 381); Watson (1967); Clay & Moreby (1970: 217); Gressitt (1970: 328); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1368); Palma (1996b: 197); Marris (2000: 187); Palma & Horning (2002: 10, 15); Page *et al.* (2004: 641, 649); Palma (2010).

Other significant references: Clay (1940a); Timmermann (1965); Clay & Moreby (1967); Price *et al.* (2003: 205); Hänel & Palma (2007: 113, 126, 129).

Remarks: *Paraclisis hyalina* is a “wing” louse species, highly prevalent among the lice parasitic on large albatrosses.

***Paraclisis miriceps* (Kellogg & Kuwana, 1902)**

Figs 139–140

*Lipeurus miriceps* Kellogg & Kuwana, 1902: 480, pl. 30: fig. 4.*Esthiopterum miriceps* Kellogg & Kuwana, 1902 [sic]; Harrison 1916: 138 (as junior synonym of *Esthiopterum giganticola* (Kellogg, 1896a)).*Perineus miriceps* (Kellogg & Kuwana, 1902); Clay 1940a: 304, figs 3b, 5c, 6b.*Perineus miriceps* (Kellogg & Kuwana, 1902); Hopkins & Clay 1952: 278.*Harrisoniella miriceps* (Kellogg & Kuwana, 1902); Kéler 1957a: 290, figs 2d, 4.*Paraclisis miriceps* (Kellogg & Kuwana, 1902); Timmermann 1965: 103, figs 33d, 34b, 36, 39c.*Paraclisis miriceps* (Kellogg & Kuwana, 1902) s. l.; Pilgrim & Palma 1982: 7.*Paraclisis miriceps* (Kellogg & Kuwana, 1902); Price *et al.* 2003: 205.*Paraclisis miriceps* (Kellogg & Kuwana, 1902); Palma 2010: 408.

Holotype ♂, presumed lost. See Palma &amp; Peck (2013: 41).

Type host: “*Geospiza fuliginosa*”, in error (see Hopkins & Clay 1952: 278).New Zealand host: *Macronectes giganteus* (J.F. Gmelin, 1789).Other host: *Phoebastria irrorata* (Salvin, 1883).

New Zealand localities: TK, NN, MC.

Geographic distribution: Pacific Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990: 1369); Paterson *et al.* (1999: 222); Palma (2010).Other significant references: Clay (1940a); Kéler (1957a); Timmermann (1965); Price *et al.* (2003); Page *et al.* (2004: 641, 649); Palma & Peck (2013: 41).Remarks: The primary host of *P. miriceps* is *Phoebastria irrorata*, the Galápagos albatross (Palma & Peck 2013: 41).

Although *Macronectes giganteus* is regularly parasitised by *Paraclisis obscura* (see below), there is a population of *P. miriceps* also living on *M. giganteus*, but with a much reduced prevalence of infestation. Pilgrim & Palma (1982: 7) qualified that population as *sensu lato* due to the absence of comparison material from the primary host; however, my examination of extensive samples from *Phoebastria irrorata* shows that that qualification is not warranted.

***Paraclisis obscura* (Rudow, 1869)***Lipeurus obscurus* Rudow, 1869b: 30.*Esthiopterum obscurum* Rudow, 1870 [sic]; Harrison 1916: 139.*Perineus obscurus* Rudow, 1870 [sic]; Harrison 1937: 29.*Perineus obscurus* (Rudow, 1869); Clay 1940a: 307, figs 3d, 4d, 5c, 6d.*Perineus obscurus* (Rudow, 1869); Hopkins & Clay 1952: 278.*Harrisoniella obscura* (Rudow, 1869); Kéler 1957a: 281, figs 1a,b,c, 2b.*Perineus* (s. l.) *obscurus* (Rudow, 1869); Clay 1964a: 231.*Perineus obscurus* (Rudow, 1869); Gressitt 1964: 538.*Paraclisis obscura* (Rudow, 1869); Timmermann 1965: 100, figs 33b, 34d, 35, 37d, 39e, pl. 5: figs 3–4.*Paraclisis obscura* (Rudow, 1869); Clay & Moreby 1967: 162, 168, figs 121, 123, 127, 153.*Paraclisis obscura* (Rudow, 1869); Wise 1977: 62.*Paraclisis obscura* (Rudow, 1869); Horning *et al.* 1980: 6, 9.*Paraclisis obscura* (Rudow, 1869); Pilgrim & Palma 1982: 6.*Paraclisis obscura* (Rudow, 1869); Murray *et al.* 1990: 1369.*Paraclisis obscura* (Rudow, 1869); Palma 2010: 408.

Neotype ♂ in NHML (Clay 1940a: 308).

Type host: *Macronectes giganteus* (J.F. Gmelin, 1789).New Zealand hosts: *Macronectes giganteus* (J.F. Gmelin, 1789); *Macronectes halli* Mathews, 1912.

Other hosts: None.

New Zealand localities: ND, BP, WO, TK, WA, WN, NN, KA, NC, MC, SC, WD, CH, SN, AN, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Clay (1964a); Gressitt (1964); Watson (1967: 73); Clay & Moreby (1970: 217); Gressitt (1970: 328); Wise (1977); Lowry *et al.* (1978: 138); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990);

Palma (1996b: 197); Paterson *et al.* (1999: 222); Marris (2000: 187); Palma & Horning (2002: 10, 16); Palma (2010).

Other significant references: Clay (1940a); Kéler (1957a); Timmermann (1965); Clay & Moreby (1967); Green & Palma (1991: 16, 25); Price *et al.* (2003: 205); Page *et al.* (2004: 641, 649); Hänel & Palma (2007: 113, 126, 129).

Remarks: *Paraclisis obscura* is a “wing” louse species, highly prevalent among the lice parasitic on giant petrels.

### Genus *Pectinopygus* Mjöberg, 1910

*Pectinopygus* Mjöberg, 1910a. *Arkiv Zool.* 6(13): 95. Type species *Lipeurus pullatus* Nitzsch [*in* Giebel], 1866 = *Pectinopygus bassani* (O. Fabricius, 1780) (by original designation).

*Epipelicanus* Thompson, 1935b. *Ann. Mag. Nat. Hist.* (Ser. 10) 16: 149. Type species: *Lipeurus forficulatus* Nitzsch [*in* Giebel], 1866 = *Pectinopygus forficulatus* (Nitzsch [*in* Giebel], 1866) (by original designation).

*Epifregata* Thompson, 1935b. *Ann. Mag. Nat. Hist.* (Ser. 10) 16: 150. Type species: *Lipeurus gracilicornis* Piaget, 1880 = *Pectinopygus gracilicornis* (Piaget, 1880) (by original designation).

*Philichthyophaga* Thompson, 1935b. *Ann. Mag. Nat. Hist.* (Ser. 10) 16: 150. Type species: *Lipeurus brevicornis* (Denny, 1842) = *Pectinopygus brevicornis* (Denny, 1842) (by original designation).

### *Pectinopygus annulatus* (Piaget, 1880)

*Lipeurus annulatus* Piaget, 1880: 340, pl. 27: fig. 10.

*Lipeurus potens* Kellogg & Kuwana, 1902: 477, pl. 30: fig. 1.

*Esthiopterum annulatum* Piaget, 1880 [sic]; Harrison 1916: 130.

*Pectinopygus jamaicensis* Thompson, 1948b: 348, figs 1–8.

*Pectinopygus annulatus* (Piaget, 1880); Hopkins & Clay 1952: 268.

*Pectinopygus potens* (Kellogg & Kuwana, 1902); Hopkins & Clay 1952: 270.

*Pectinopygus annulatus* (Piaget, 1880); Watt 1971: 236, 243.

*Pectinopygus annulatus* (Piaget, 1880); Pilgrim & Palma 1982: 14.

*Pectinopygus annulatus* (Piaget, 1880); Murray *et al.* 1990: 1372.

*Pectinopygus annulatus* (Piaget, 1880); Palma 2010: 408.

Lectotype ♂ in NHML (Clay 1973: 219).

Type host: *Sula leucogaster* (Boddaert, 1783).

New Zealand hosts: *Sula leucogaster plotus* (J.R. Forster, 1844); *Sula dactylatra tasmani* van Tets, *et al.* 1988.

Other hosts: *Sula variegata* (Tschudi, 1843); *Sula granti* Rothschild, 1902.

New Zealand localities: ND, WO, WD, KE.

Geographic distribution: Tropical and subtropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Watt (1971); Wise (1977: 62); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999: 378); Palma (2010).

Other significant references: Thompson (1948b); Clay (1964b: 15, fig. 1); Amerson & Emerson (1971: 10, 24); Ward & Downey (1973: 394); Clay (1973: 219); Moreby (1976: 92); Forrester *et al.* (1995: 6); Palma (1996b: 198); Price *et al.* (2003: 207); Hughes *et al.* (2007: 238, 249); Palma & Peck (2013: 41); Rivera-Parra *et al.* (2014: 571).

Remarks: *Pectinopygus annulatus* is a “wing” louse species, highly prevalent among the lice parasitic on boobies.

### *Pectinopygus australis* Thompson, 1948

*Pectinopygus* (*Epipelicanus*) *australis* Thompson, 1948a: 323, figs 10–15, pl. 10: figs 1–2.

*Pectinopygus australis* Thompson, 1948; Pilgrim & Palma 1982: 14.

*Pectinopygus australis* Thompson, 1948; Murray *et al.* 1990: 1372.

*Pectinopygus australis* Thompson, 1948; Price *et al.* 2003: 207.

*Pectinopygus australis* Thompson, 1948; Palma 2010: 408.

Holotype ♀ in NHML (Palma 1996b: 198, in error as “♂”).

Type host: *Pelecanus conspicillatus* Temminck, 1824.

New Zealand host: *Pelecanus conspicillatus* Temminck, 1824.

Other hosts: None.

New Zealand locality: SL.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Moreby (1976: 91); Green & Palma (1991: 16, 28); Palma (1996b: 198); Price *et al.* (2003).

Remarks: *Pelecanus conspicillatus* breeds in Australia and is a straggler to New Zealand (Checklist Committee (2010: 138), with only one record of *Pectinopygus australis* from this country.

### ***Pectinopygus bassani* (O. Fabricius, 1780)**

*Pediculus bassani* O. Fabricius, 1780: 218.

*Pectinopygus bassanae* [sic] O. Fabricius, 1780 [sic]; Harrison 1916: 143.

*Pectinopygus (Pectinopygus) bassanae* [sic] O. Fabricius, 1780 [sic]; Harrison 1937: 32.

*Pectinopygus (Pectinopygus) bassani* (O. Fabricius, 1780); Thompson 1940a: 372, figs 1–9, pl. 8.

*Pectinopygus (Pectinopygus) bassani serrator* Thompson, 1940a: 381.

*Pectinopygus (Pectinopygus) bassanae* [sic] serrator Thompson, 1940; Thompson 1940b: 498.

*Pectinopygus (Pectinopygus) bassanus* [sic] (O. Fabricius, 1780); Séguy 1944: 379, figs 557–562.

*Pectinopygus bassani* (O. Fabricius, 1780); Hopkins & Clay 1952: 268.

*Pectinopygus bassani* (O. Fabricius, 1780); Pilgrim & Palma 1982: 14.

*Pectinopygus bassani* (O. Fabricius, 1780); Murray *et al.* 1990: 1372.

*Pectinopygus bassani* (O. Fabricius, 1780); Palma 2010: 408.

Neotype ♂ in NHML (Thompson 1940a: 380). Syntypes ♂♀ of *Pectinopygus (Pectinopygus) bassani serrator* in NHML.

Type host: *Morus bassanus* (Linnaeus, 1758).

New Zealand host: *Morus serrator* (G.R. Gray, 1843).

Other host: *Morus capensis* (Lichtenstein, 1823).

New Zealand localities: ND, AK, CL, WO, BP, HB, TK, WN, SD, MB, NC, MC, SC, CO, DN.

Geographic distribution: North Atlantic Ocean; Indian Ocean; Pacific Ocean; eastern North America; western Europe; southern Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Hughes *et al.* (2007: 238, 249); Palma (2010).

Other significant references: Harrison (1937); Thompson (1940a; 1940b); Séguy (1944); Clay & Hopkins (1954: 247); Clay (1973: 217); Green & Palma (1991: 16, 28); Butler & O'Connor (1994: 455); Forrester *et al.* (1995: 7); Martín-Mateo (1992a: 43, figs 10–16); Palma (1996b: 198); Price *et al.* (2003: 207); Martín-Mateo (2009: 99, fig. 24).

Remarks: *Pectinopygus bassani* is a “wing” louse species, highly prevalent among the lice parasitic on gannets.

### ***Pectinopygus carunculatus* Timmermann, 1964**

*Pectinopygus carunculatus* Timmermann, 1964: 280, fig. 6, pl. 9: figs 2–3.

*Pectinopygus carunculatus* Timmermann, 1964; Wise 1977: 62.

*Pectinopygus carunculatus* Timmermann, 1964; Pilgrim & Palma 1982: 15.

*Pectinopygus carunculatus* Timmermann, 1964; Murray *et al.* 1990: 1373.

*Pectinopygus carunculatus* Timmermann, 1964; Palma 2010: 408.

Holotype in NHML.

Type host: *Leucocarbo colensoi* (Buller, 1888).

New Zealand host: *Leucocarbo colensoi* (Buller, 1888).

Other hosts: None.

New Zealand locality: AU.

Geographic distribution: New Zealand.

New Zealand references: Timmermann (1964); Pilgrim (1970: 74); Weidner (1977: 102); Pilgrim & Palma (1982); Wise (1977); Murray *et al.* (1990); Palma (2010).

Other significant references: Price *et al.* (2003: 208).

Remarks: *Pectinopygus carunculatus* is an endemic but infrequently collected species. Although Timmermann (1964: 280) regarded the population of *Pectinopygus* from *Leucocarbo campbelli* as conspecific with that from

*Leucocarbo colensoi*, I regard the former —together with other *Pectinopygus* populations from other species of *Leucocarbo*— as *P. carunculatus sensu lato* (see below).

***Pectinopygus carunculatus* Timmermann, 1964 *sensu lato***

*Pectinopygus* spp.; Gressitt 1964: 539.

*Pectinopygus carunculatus* Timmermann, 1964: 280. In part.

*Pectinopygus* sp.; Pilgrim & Palma 1982: 14–15.

*Pectinopygus carunculatus* Timmermann, 1964 s. l.; Pilgrim & Palma 1982: 15.

*Pectinopygus* sp.; Murray *et al.* 1990: 1373.

New Zealand hosts: *Leucocarbo carunculatus* (J.F. Gmelin, 1789); *Leucocarbo chalconotus* (G.R. Gray, 1845); *Leucocarbo onslowi* Forbes, 1893; *Leucocarbo ranfurlyi* (Ogilvie-Grant, 1901); *Leucocarbo campbelli* (Filhol, 1888).

Other hosts: None.

New Zealand localities: SD, MB, KA, SC, DN, SL, CH, BO, CA.

Geographic distribution: New Zealand.

New Zealand references: Gressitt (1964); Timmermann (1964); Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 222).

Other significant references: None.

Remarks: The endemic populations of *Pectinopygus* from the hosts listed above are morphologically close to *Pectinopygus carunculatus sensu stricto*. However, they exhibit subtle but consistent differences which may warrant treating them as separate taxa. Detailed morphological and molecular studies of all these populations are needed to determine their correct taxonomic position.

***Pectinopygus dispar* (Piaget, 1880)**

*Nirmus dispar* Piaget, 1880: 174, pl. 14: fig. 7.

*Lipeurus subsetosus* Piaget, 1880: 336, pl. 27: fig. 5.

*Esthiopterum dispar* Piaget, 1880 [sic]; Harrison 1916: 133 (as junior synonym of *Esthiopterum acutifrons* (Rudow, 1869)).

*Pectinopygus dispar* (Piaget, 1880); Hopkins & Clay 1952: 269.

*Pectinopygus subsetosus* (Piaget, 1880); Hopkins & Clay 1952: 270.

*Pectinopygus dispar* (Piaget, 1880); Clay 1973: 219, fig. 6, pl. 1: fig. 1, pl. 2: fig. 7, pl. 3: fig. 20.

*Pectinopygus subsetosus* (Piaget, 1880); Moreby 1976: 92.

*Pectinopygus dispar* (Piaget, 1880); Horning *et al.* 1980: 6, 11.

*Pectinopygus dispar* (Piaget, 1880); Pilgrim & Palma 1982: 14.

*Pectinopygus dispar* (Piaget, 1880); Murray *et al.* 1990: 1373.

*Pectinopygus dispar* (Piaget, 1880); Palma 2010: 408.

Lectotype ♀ in NHML (Clay 1973: 219).

Type host: “*Phalacrocorax sulcirostris*”, in error (see Clay 1973: 220).

New Zealand host: *Phalacrocorax melanoleucos brevirostris* Gould, 1837.

Other hosts: *Phalacrocorax melanoleucos melanoleucos* (Vieillot, 1817); *Phalacrocorax melanoleucos brevicauda* Mayr, 1931.

New Zealand localities: AK, BP, WO, WA, WN, NC, MC, SC, CO, DN, SN.

Geographic distribution: Southeast Asia; Australasia; western Pacific Ocean.

New Zealand references: Clay (1958c: 145); Clay (1973: 207); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Moreby (1976); Green & Palma (1991: 16, 28); Palma (1996b: 199); Price *et al.* (2003: 208).

Remarks: *Pectinopygus dispar* is a “wing” louse species, frequently found on the New Zealand little shag.

***Pectinopygus garbei* (Pessôa & Guimarães, 1935)**

*Naubates (Micronaubates) garbei* Pessôa & Guimarães, 1935b: 109, figs 6–11.

*Naubates garbei* Pessôa & Guimarães, 1935; Thompson 1938b: 486.

*Pectinopygus (Pectinopygus) garbei* (Pessôa & Guimarães, 1935); Guimarães 1945: 184, figs 6–13.

*Pectinopygus garbei* (Pessôa & Guimarães, 1935); Hopkins & Clay 1952: 269.

*Pectinopygus garbei* (Pessôa & Guimarães, 1935); Pilgrim & Palma 1982: 14.

*Pectinopygus garbei* (Pessôa & Guimarães, 1935); Murray *et al.* 1990: 1372.

*Pectinopygus garbei* (Pessôa & Guimarães, 1935); Palma 2010: 408.

Holotype ♂ in Laboratório de Parasitologia, Faculdade de Medicina da Universidade de São Paulo, Brazil (Michel P. Valim pers. comm. December 2014).

Type host: *Sula leucogaster leucogaster* (Boddaert, 1783) (see Guimarães 1945: 184).

New Zealand host: *Sula leucogaster plotus* (J.R. Forster, 1844).

Other host: *Sula sula* (Linnaeus, 1766).

New Zealand locality: KE.

Geographic distribution: Tropical Central and South America; Africa; Asia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Clay (1940a: 310); Guimarães (1945); Clay (1964b: 15, fig. 1); Clay (1976b: 540); Palma (1996b: 199); Price *et al.* (2003: 208); Silva *et al.* (2014: 942).

Remarks: Palma (1996b: 199) incorrectly stated “MZUSP” as the repository of the holotype of *Pectinopygus garbei*, but only paratypes are held at that museum (Valim 2009: 208). *Sula leucogaster plotus* is a vagrant to New Zealand (Checklist Committee 2010: 141), with a single record of *Pectinopygus garbei* from this country.

#### ***Pectinopygus gyricornis* (Denny, 1842)**

*Philopterus (Lipeurus) gyricornis* Denny, 1842: 58, 167, pl. 15: fig. 1.

*Esthiopterum gyricorne* Denny, 1842 [sic]; Harrison 1916: 135.

*Pectinopygus (Philichthyophaga) gyricornis* (Denny, 1842); Séguy 1944: 375, fig. 553.

*Pectinopygus gyricornis* (Denny, 1842); Hopkins & Clay 1952: 269.

*Pectinopygus gyricornis* (Denny, 1842); Timmermann 1964: 273, fig. 1.

*Pectinopygus gyricornis* (Denny, 1842); Horning *et al.* 1980: 6, 11.

*Pectinopygus gyricornis* (Denny, 1842); Pilgrim & Palma 1982: 14.

*Pectinopygus gyricornis* (Denny, 1842); Murray *et al.* 1990: 1372.

*Philichthyophaga gyricornis* (Denny, 1842); Mey 1994: 27: figs 11–12.

*Pectinopygus gyricornis* (Denny, 1842); Palma 2010: 408.

Holotype ♂ in NHML (Clay 1973: 221).

Type host: “*Sterna hirundo*”, in error (see Hopkins & Clay 1952: 269).

New Zealand host: *Phalacrocorax carbo novaehollandiae* Stephens, 1826.

Other hosts: *Phalacrocorax carbo carbo* (Linnaeus, 1758); *Phalacrocorax carbo sinensis* (Blumenbach, 1798).

New Zealand localities: WA, WN, SD, MB, NN, NC, MC, SC, CO, DN, SN.

Geographic distribution: Eurasia; Africa; North America; Australasia.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Galloway (2005: 17); Palma (2010).

Other significant references: Séguy (1944); Brelih & Tovornik (1961: 104); Timmermann (1964); Clay (1973: 215, 221); Green & Palma (1991: 16, 28); Mey (1994); Butler & O’Connor (1994: 455); Palma (1996b: 200); Price *et al.* (2003: 208); Adam (2007: 174); Hughes *et al.* (2007: 238, 249).

Remarks: *Pectinopygus gyricornis* is a “wing” louse species, frequently found on all its hosts.

#### ***Pectinopygus punctatus* Timmermann, 1964**

*Pectinopygus punctatus* Timmermann, 1964: 277, fig. 5.

*Pectinopygus punctatus* Timmermann, 1964; Wise 1977: 62.

*Pectinopygus punctatus* Timmermann, 1964; Pilgrim & Palma 1982: 15.

*Pectinopygus punctatus* Timmermann, 1964 s. l.; Pilgrim & Palma 1982: 15.

*Pectinopygus punctatus* Timmermann, 1964; Murray *et al.* 1990: 1372.

*Pectinopygus punctatus* Timmermann, 1964; Palma 2010: 408.

Holotype ♂ in CMNZ (Nicholls *et al.* 1998: 30).

Type host: *Stictocarbo punctatus punctatus* (Sparrman, 1786).

New Zealand hosts: *Stictocarbo punctatus punctatus* (Sparrman, 1786); *Stictocarbo punctatus oliveri* Mathews, 1931.

Other hosts: None.

New Zealand localities: HB, WI, WN, KA, NN, MC, BR, DN.

Geographic distribution: New Zealand.

New Zealand references: Timmermann (1964); Pilgrim (1970: 75); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Nicholls *et al.* (1998: 30); Paterson *et al.* (1999: 223); Hughes *et al.* (2007: 238, 249); Palma (2010).

Other significant references: Clay (1973: 215); Price *et al.* (2003: 208).

Remarks: *Pectinopygus punctatus* is an endemic species, exclusively parasitic on spotted and blue shags. Pilgrim & Palma (1982: 15) regarded the population of *Pectinopygus punctatus* from *Stictocarbo punctatus oliveri* (as *Stictocarbo punctatus steadi*) as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted.

### ***Pectinopygus setosus* (Piaget, 1880)**

*Lipeurus setosus* Piaget, 1880: 335, pl. 27: fig. 4.

*Esthiopterum setosum* Piaget, 1880 [sic]; Harrison 1916: 141.

*Pectinopygus setosus* (Piaget, 1880); Hopkins & Clay 1952: 270.

*Pectinopygus setosus* (Piaget, 1880); Timmermann 1964: 278.

*Pectinopygus setosus* (Piaget, 1880); Clay 1973: 215, 217, 221.

*Pectinopygus setosus* (Piaget, 1880); Pilgrim & Palma 1982: 14.

*Pectinopygus setosus* (Piaget, 1880); Murray *et al.* 1990: 1372.

*Pectinopygus setosus* (Piaget, 1880); Palma 2010: 408.

Lectotype ♂ in NHML (Clay 1973: 221).

Type host: *Phalacrocorax sulcirostris* (J.F. Brandt, 1837).

New Zealand host: *Phalacrocorax sulcirostris* (J.F. Brandt, 1837).

Other hosts: None

New Zealand localities: BP, HB, WA, WN.

Geographic distribution: Australasia; Indonesia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Hughes *et al.* (2007: 238, 249); Palma (2010).

Other significant references: Kéler (1957c: fig. 22b); Timmermann (1964); Clay (1973); Green & Palma (1991: 16, 28); Palma (1996b: 200); Price *et al.* (2003: 208).

Remarks: *Pectinopygus setosus* is a frequently found “wing” louse species on all its hosts.

### ***Pectinopygus turbinatus* (Piaget, 1890)**

*Oncophorus turbinatus* Piaget, 1890a: 233, pl. 8: fig. 10.

*Pectinopygus (Philichthyophaga) macquariensis* Harrison, 1937: 34, fig. 6.

*Pectinopygus macquariensis* Harrison, 1937; Hopkins & Clay 1952: 269.

*Pectinopygus turbinatus* (Piaget, 1890); Hopkins & Clay 1952: 270.

*Pectinopygus turbinatus* (Piaget, 1890); Timmermann 1964: 280, fig. 7, pl. 9: fig. 4.

*Pectinopygus turbinatus* (Piaget, 1890); Clay & Moreby 1967: 162, 169, figs 78, 115.

*Pectinopygus turbinatus* (Piaget, 1890); Wise 1977: 62.

*Pectinopygus turbinatus* (Piaget, 1890); Pilgrim & Palma 1982: 15.

*Pectinopygus turbinatus* (Piaget, 1890); Murray *et al.* 1990: 1373.

*Pectinopygus turbinatus* (Piaget, 1890); Palma 2010: 408.

Holotype ♂ in NHML (Clay 1973: 222). Holotype ♀ of *Pectinopygus (Philichthyophaga) macquariensis* in AMSA (Palma 1996b: 200).

Type host: “*Mycteria senegalensis*”, in error (see Hopkins & Clay 1952: 270; Clay 1973: 222).

New Zealand host: *Leucocarbo purpurascens* (J.F. Brandt, 1837).

Other hosts: *Leucocarbo atriceps* (King, 1828); *Leucocarbo albiventer* (Lesson, 1831).

New Zealand locality: Macquarie Island.

Geographic distribution: Pacific, Atlantic and Indian Oceans; South America; Antarctica.

New Zealand references: Harrison (1937); Watson (1967: 73); Gressitt (1970: 328); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b: 200); Paterson *et al.* (1999: 222); Palma & Horning (2002: 10, 17); Palma (2010).

Other significant references: Timmermann (1964); Clay & Moreby (1967); Clay (1973: 217, 222); Price *et al.* (2003: 209).

Remarks: *Pectinopygus macquariensis* was described from the endemic Macquarie Island shag, *Leucocarbo purpurascens*, but later synonymised under *P. turbinatus* by Timmermann (1964: 280).

### ***Pectinopygus varius* Timmermann, 1964**

Figs 141–142

*Pectinopygus varius* Timmermann, 1964: 273, fig. 2.

*Pectinopygus varius* Timmermann, 1964; Clay 1973: 215, fig. 2, pl. 2: figs 8–9, pl. 3: figs 13–14

*Pectinopygus varius* Timmermann, 1964; Wise 1977: 63.

*Pectinopygus varius* Timmermann, 1964; Pilgrim & Palma 1982: 14.

*Pectinopygus varius* Timmermann, 1964; Murray *et al.* 1990: 1372.

*Pectinopygus varius* Timmermann, 1964; Palma 2010: 408.

Holotype ♂ in CMNZ (Nicholls *et al.* 1998: 30).

Type host: *Phalacrocorax varius varius* (J.F. Gmelin, 1789).

New Zealand host: *Phalacrocorax varius varius* (J.F. Gmelin, 1789).

Other host: *Phalacrocorax varius hypoleucos* (J.F. Brandt, 1837).

New Zealand localities: ND, AK, CL, BP, GB, SD, MB, SI.

Geographic distribution: Australasia.

New Zealand references: Timmermann (1964); Pilgrim (1970: 75); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Nicholls *et al.* (1998: 30); Palma (2010).

Other significant references: Clay (1973); Palma (1996b: 201); Price *et al.* (2003: 209); Hughes *et al.* (2007: 238, 249).

Remarks: *Pectinopygus varius* is a “wing” louse species, frequently found on both subspecies of the pied shag.

### ***Pectinopygus* species**

*Pectinopygus* sp.; Pilgrim & Palma 1982: 15.

*Pectinopygus* sp.; Murray *et al.* 1990: 1372.

New Zealand host: *Stictocarbo featherstoni* (Buller, 1873).

Other hosts: None.

New Zealand locality: CH.

Geographic distribution: Chatham Islands, New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990).

Other significant references: None.

Remarks: At present, this endemic population of *Pectinopygus* is not identifiable because the available samples are inadequate.

### **Genus *Pelmatocerandra* Enderlein, 1908**

*Pelmatocerandra* Enderlein, 1908. *Deutsche Südpolar-Exped. 10 (Zoologie) 2*: 449. Type species: *Pelmatocerandra setosa* (Giebel, 1876) (by monotypy).

### ***Pelmatocerandra setosa* (Giebel, 1876)**

Figs 143–144

*Nirmus setosus* Giebel, 1876: 388.

*Lipeurus setosus* (Giebel, 1876); Kellogg 1914: 81, 87.

*Lipeurus eatoni* Kellogg, 1914: 81, 86. Unnecessary *nomen novum* for *Nirmus setosus* Giebel, 1876 on transfer to *Lipeurus*.

*Pelmatocerandra eatoni* Kellogg, 1914 [sic]; Harrison 1916: 144 (as junior synonym of *P. setosa* Giebel, 1876)

*Pelmatocerandra setosa* Giebel, 1876 [sic]; Harrison 1916: 144.



*Pelmatocerandra setosa* (Giebel, 1876); Hopkins & Clay 1952: 274.

*Pelmatocerandra setosa* (Giebel, 1876); Clay 1958a: 252, figs 1, 4, 7, pl. 4: fig. 1.

*Pelmatocerandra setosa* (Giebel, 1876); Timmermann 1965: 162, figs 99–100, 102, pl. 9: figs 2–3.

*Pelmatocerandra setosa* (Giebel, 1876); Wise 1977: 63.

*Pelmatocerandra setosa* (Giebel, 1876); Horning *et al.* 1980: 6, 11.

*Pelmatocerandra setosa* (Giebel, 1876); Pilgrim & Palma 1982: 13.

*Pelmatocerandra setosa* (Giebel, 1876); Murray *et al.* 1990: 1372.

*Pelmatocerandra setosa* (Giebel, 1876); Palma 2010: 408.

Lectotype ♂ in NHML (Clay 1958a: 252).

Type host: *Pelecanoides urinatrix* (J.F. Gmelin, 1789).

New Zealand hosts: *Pelecanoides urinatrix urinatrix* (J.F. Gmelin, 1789); *Pelecanoides urinatrix chathamensis* Murphy & Harper, 1916; *Pelecanoides urinatrix exsul* Salvin, 1896; *Pelecanoides georgicus* Murphy & Harper, 1916.

Other host: *Pelecanoides urinatrix dacunhae* Nicholl, 1906.

New Zealand localities: ND, CL, BP, GB, TO, WO, WI, WA, WN, WD, CO, DN, CH, SI, SN, AN, AU, CA, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Clay (1958a); Clay (1964a: 232); Gressitt (1964: 538); Watson (1967: 73); Clay & Moreby (1970: 219); Gressitt (1970: 328); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b: 201); Marris (2000: 187); Palma & Horning (2002: 11, 15); Palma (2010).

Other significant references: Thompson (1940c: 104, figs 1–10, pl. 2); Kéler (1952: 216, figs 11–13); Paulian (1953: 199); Séguy (1953: 559, figs 3–4); Timmermann (1965); Clay & Moreby (1967: 160, fig. 147); Payne & Prince (1979: 316); Green & Palma (1991: 16, 28); Furness & Palma (1992: 35, 42); Price *et al.* (2003: 209); Page *et al.* (2004: 638, 649); Hänel & Palma (2007: 113, 126, 131).

Remarks: The natural, regular species of *Pelmatocerandra* parasitic on *Pelecanoides georgicus* in the Atlantic and Indian Oceans is *Pelmatocerandra enderleini* Eichler, 1949b. However, from the evidence of two independent samples taken in 1978 and 2003 respectively, the New Zealand population of the South Georgian diving petrel appears to be regularly parasitised by *Pelmatocerandra setosa*.

### Genus *Penenirmus* Clay & Meinertzhagen, 1938

*Penenirmus* Clay & Meinertzhagen, 1938a. *Entomologist* 71: 73. Type species: *Pediculus albiventris* Scopoli, 1763 = *Penenirmus albiventris* (Scopoli, 1763) (by original designation).

#### *Penenirmus* species

Fig. 145

*Penenirmus* sp.; Palma 1999: 382.

*Penenirmus* sp.; Murray *et al.* 2006b: 1959.

New Zealand host: *Bowdleria punctata wilsoni* Stead, 1936.

New Zealand locality: SI.

Geographic distribution: Codfish Island, Stewart Islands, New Zealand.

New Zealand references: Palma (1999); Murray *et al.* (2006b); Buckley *et al.* (2012: App. 2).

Other significant references: Emerson & Johnson (1961); Dalglish (1972); Price *et al.* (2003: 209).

Remarks: The single available record of *Penenirmus* could not be identified to species because the sample contains females only (voucher specimens in MONZ).

### Genus *Perineus* Thompson, 1936

*Perineus* Thompson, 1936. *Ann. Mag. Nat. Hist. (Ser. 10)* 18: 41. Type species: *Lipeurus nigrolimbatus* Giebel, 1874 = *Perineus nigrolimbatus* (Giebel, 1874) (by original designation).

#### *Perineus circumfasciatus* Kéler, 1957

“*Perineus concinnus*” Harrison, 1937: 29 (not *Lipeurus concinnus* Kellogg & Chapman, 1899).

*Perineus circumfasciatus* Kéler, 1957b: 525, figs 29–30.

*Perineus* sens. str.; Clay 1964a: 231. In part.

*Perineus* sp.; Gressitt 1964: 538. In part.

*Perineus circumfasciatus* Kéler, 1957; Timmermann 1965: 107, fig. 46.

*Perineus circumfasciatus* Kéler, 1957; Watson 1967: 73.

*Perineus* sp.; Watson 1967: 73.

*Perineus circumfasciatus* Kéler, 1957; Wise 1977: 63. In part.

“*Perineus concinnus*” Wise, 1977: 63 (not *Lipeurus concinnus* Kellogg & Chapman, 1899).

*Perineus circumfasciatus* Kéler, 1957; Horning *et al.* 1980: 6, 9.

*Perineus circumfasciatus* Kéler, 1957; Pilgrim & Palma 1982: 6.

*Perineus circumfasciatus* Kéler, 1957; Palma & Pilgrim 1988: 580, figs 7, 14, 22, 27, 35, 41, 48, 52, 54.

*Perineus circumfasciatus* Kéler, 1957; Palma 2010: 408.

Holotype ♂ in NHML (Palma & Pilgrim 1988: 580).

Type host: *Thalassarche melanophris* (Temminck, 1828).

New Zealand hosts: *Thalassarche chlororhynchos* (J.F. Gmelin, 1789); *Thalassarche chrysostoma* (J.R. Forster, 1785); *Thalassarche melanophris* (Temminck, 1828); *Thalassarche impavida* Mathews, 1912; *Thalassarche bulleri bulleri* (Rothschild, 1893); *Thalassarche bulleri platei* (Reichenow, 1898); *Thalassarche cauta steadi* Falla, 1933; *Thalassarche eremita* Murphy, 1930; *Thalassarche salvini* (Rothschild, 1893); *Phoebetria fusca* (Hilsenberg, 1822); *Phoebetria palpebrata* (J.R. Forster, 1785).

Other host: *Thalassarche cauta cauta* (Gould, 1841).

New Zealand localities: AK, BP, HB, WA, WN, KA, NC, MC, SC, SL, CH, BO, SN, AU, CA, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Harrison (1937); Clay (1964a); Gressitt (1964); Watson (1967); Gressitt (1970: 328); Pilgrim (1970: 74); Wise (1977); Pilgrim & Palma (1979: 177); Horning *et al.* (1980); Pilgrim & Palma (1982); Palma & Pilgrim (1988); Murray *et al.* (1990: 1368–1369); Green & Palma (1991: 16, 25); Palma (1996b: 201); Palma (1999: 375, 383, note 1); Marris (2000: 188); Palma & Horning (2002: 11, 15); Page *et al.* (2004: 642, 649); Palma (2010).

Other significant references: Timmermann (1965); Clay & Moreby (1967: 163, 168, fig. 129); Price *et al.* (2003: 211); Hänel & Palma (2007: 113, 126, 129).

Remarks: *Perineus circumfasciatus* is a “wing” louse species frequently collected from small albatrosses.

### ***Perineus concinnoides* Kéler, 1957**

Figs 146–147

*Perineus concinnoides* Kéler, 1957b: 521, fig. 26.

*Perineus* sens. str.; Clay 1964a: 231. In part.

*Perineus* sp.; Gressitt 1964: 538. In part.

*Perineus concinnoides* Kéler, 1957; Timmermann 1965: 109, fig. 48.

*Perineus concinnoides* Kéler, 1957; Wise 1977: 63.

*Perineus concinnoides* Kéler, 1957; Pilgrim & Palma 1982: 5.

*Perineus concinnoides* Kéler, 1957; Palma & Pilgrim 1988: 578, figs 3, 10, 20, 26, 29, 36, 42, 44.

*Perineus concinnoides* Kéler, 1957; Palma 2010: 408.

Holotype ♂ in NHML (Palma & Pilgrim 1988: 579).

Type host: *Diomedea exulans* Linnaeus, 1758.

New Zealand hosts: *Diomedea exulans* Linnaeus, 1758; *Diomedea epomophora* Lesson, 1825; *Diomedea sanfordi* Murphy, 1917.

Other host: *Diomedea dabbenena* Mathews, 1929.

New Zealand localities: WA, KA, NC, MC, SC, WD, CO, DN, CH.

Geographic distribution: Southern Hemisphere.

New Zealand references: Pilgrim (1970: 74); Wise (1977); Pilgrim & Palma (1982); Palma & Pilgrim (1988); Murray *et al.* (1990: 1368); Palma (2010).

Other significant references: Timmermann (1965); Clay & Moreby (1967: 162, 168, fig. 130); Green & Palma (1991: 16, 25); Palma (1996b: 202); Page *et al.* (2004: 642, 649); Price *et al.* (2003: 211).

Remarks: *Perineus concinnoides* is a “wing” louse species infrequently collected from large albatrosses.

### ***Perineus macronecti* Palma & Pilgrim, 1988**

*Perineus circumfasciatus* Kéler, 1957b: 525. In part.

*Perineus* sp. nov.; Watson 1967: 73.

“*Perineus circumfasciatus*” Clay & Moreby, 1970: 217 (not *Perineus circumfasciatus* Kéler, 1957).

*Perineus circumfasciatus* Kéler, 1957; Wise 1977: 63. In part.

*Perineus* sp.; Horning *et al.* 1980: 6, 9.

*Perineus* sp.; Pilgrim & Palma 1982: 7.

*Perineus macronecti* Palma & Pilgrim, 1988: 584, figs 5, 12, 21, 28, 30, 37, 43, 49, 53, 55.

*Perineus macronecti* Palma & Pilgrim, 1988; Palma 1999: 377.

*Perineus macronecti* Palma & Pilgrim, 1988; Palma 2010: 408.

Holotype ♂ in MONZ (Palma *et al.* 1989: 45).

Type host: *Macronectes halli* Mathews, 1912.

New Zealand hosts: *Macronectes giganteus* (J.F. Gmelin, 1789); *Macronectes halli* Mathews, 1912.

Other hosts: None.

New Zealand localities: ND, TK, WN, KA, MC, SN, AN, Macquarie Island.

Geographic distribution: Southern Hemisphere.

New Zealand references: Watson (1967); Clay & Moreby (1970); Gressitt (1970: 328); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Palma & Pilgrim (1988); Palma *et al.* (1989: 45); Murray *et al.* (1990: 1369); Palma (1996b: 202); Palma (1999); Paterson *et al.* (1999: 222); Marris (2000: 188); Palma & Horning (2002: 11, 16); Palma (2010).

Other significant references: Price *et al.* (2003: 212); Hänel & Palma (2007: 113, 126, 129).

Remarks: *Perineus macronecti* is a “wing” louse species infrequently collected from giant petrels.

### ***Perineus nigrolimbatus* (Giebel, 1874)**

*Lipeurus nigrolimbatus* Giebel, 1874: 233.

*Esthiopterum nigrolimbatum* Giebel, 1874 [sic]; Harrison 1916: 138.

*Perineus nigrolimbatus* (Giebel, 1874); Hopkins & Clay 1952: 278.

*Perineus nigrolimbatus* (Giebel, 1874); Kéler 1957b: 512, figs 13–25.

*Perineus nigrolimbatus* (Giebel, 1874); Timmermann 1965: 108, figs 42–45, pl. 9: fig. 4.

*Perineus nigrolimbatus* (Giebel, 1874); Clay & Moreby 1967: 163, 168, fig. 128.

*Perineus nigrolimbatus* (Giebel, 1874); Pilgrim & Palma 1982: 7.

*Perineus nigrolimbatus* (Giebel, 1874); Palma & Pilgrim 1988: 569, figs 1–2, 8–9, 15–17, 25, 31, 33, 40, 45.

*Perineus nigrolimbatus* (Giebel, 1874); Murray *et al.* 1990: 1369.

*Perineus nigrolimbatus* (Giebel, 1874); Palma 2010: 408.

Neotype ♂ in SDEI (Kéler 1957b: 517).

Type host: *Fulmarus glacialis glacialis* (Linnaeus, 1761).

New Zealand host: *Fulmarus glacialis* (A. Smith, 1840).

Other hosts: *Fulmarus glacialis auduboni* Bonaparte, 1857; *Fulmarus glacialis rodgersii* Cassin, 1862.

New Zealand localities: AK, TK, WI, WN, NN, KA, NC, MC, SC, SI, Macquarie Island, RO.

Geographic distribution: Northern Pacific and Atlantic Oceans; Southern Oceans.

New Zealand references: Pilgrim & Palma (1982); Palma & Pilgrim (1988); Murray *et al.* (1990); Palma (1996b: 202); Palma (2001: 66, fig. 5); Palma & Horning (2002: 11, 16); Palma (2010).

Other significant references: Thompson (1936: 42); Harrison (1937: 30); Clay (1940a: 299); Séguy (1944: 367, figs 546); Kéler (1957b); Timmermann (1965); Clay & Moreby (1967); Hackman & Nyholm (1968: 80); Fowler & Miller (1984: 24, figs 1–2c); Green & Palma (1991: 17, 26); Butler & O’Connor (1994: 455); Price *et al.* (2003: 212); Page *et al.* (2004: 642, 649, 650); Palma & Jensen (2005: 56, 60).

Remarks: *Perineus nigrolimbatus* is a highly prevalent “wing” louse species on all its hosts. *Fulmarus glacialis* breeds on the coast of Antarctica and is a regular straggler to New Zealand seas (Checklist Committee 2010: 80).

### Genus *Philoceanus* Kellogg, 1903

*Philoceanus* Kellogg, 1903. *Biol. Bull. Wood's Hole* 5(2): 87. Type species: *Philoceanus becki* Kellogg, 1903 (by monotypy).

#### *Philoceanus fasciatus* (Carriker, 1958)

Figs 148–149

*Naubates fasciatus* Carriker, 1958: 184, figs 1–6.

*Philoceanus wolfherrei* Timmermann, 1961b: 534.

*Philoceanus fasciatus* (Carriker, 1958); Timmermann 1965: 159.

*Philoceanus fasciatus* (Carriker, 1958); Clay & Moreby 1967: 163, 169, fig. 86.

*Philoceanus fasciatus* (Carriker, 1958); Pilgrim & Palma 1982: 13.

*Philoceanus fasciatus* (Carriker, 1958); Murray *et al.* 1990: 1372.

*Philoceanus fasciatus* (Carriker, 1958); Palma 1999: 378.

*Philoceanus fasciatus* (Carriker, 1958); Palma 2010: 408.

Holotype ♂ in FMLA. Holotype ♀ of *Philoceanus wolfherrei* in NHML.

Type host: “*Oceanites oceanicus oceanicus*”, in error (see Clay & Moreby 1967: 163).

New Zealand hosts: *Pealeornis maoriana* Mathews, 1932; *Fregetta tropica* (Gould, 1844); *Fregetta grallaria* (Vieillot, 1817).

Other hosts: None.

New Zealand localities: ND, CL, KE, AN, AU, CA.

Geographic distribution: Southern Hemisphere.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999); Marris (2000: 188); Stephenson *et al.* (2009: 198, 201, fig. 5); Palma (2010).

Other significant references: Timmermann (1961b; 1965); Clay & Moreby (1967); Furness & Palma (1992: 35, 41); Price *et al.* (2003: 212); Hänel & Palma (2007: 113, 126, 130).

Remarks: *Philoceanus fasciatus* is a “wing” louse species infrequently collected from storm petrels. *Philoceanus wolfherrei* was described from *Fregetta tropica*, but later synonymised under *Ph. fasciatus* by Clay & Moreby (1967: 163).

#### *Philoceanus garrodiae* (Clay, 1940)

*Naubates garrodiae* Clay, 1940a: 310, figs 7, 8, 9a,d, 11a.

*Philoceanus garrodiae* (Clay, 1940); Hopkins & Clay 1952: 279.

*Philoceanus garrodiae* (Clay, 1940); Timmermann 1961b: 532, fig. 1.

*Philoceanus garrodiae* (Clay, 1940); Clay 1964a: 231.

*Philoceanus garrodiae* (Clay, 1940); Timmermann 1965: 157, pl. 8: figs 1–2.

*Philoceanus garrodiae* (Clay, 1940); Wise 1977: 63

*Philoceanus garrodiae* (Clay, 1940); Pilgrim & Palma 1982: 13.

*Philoceanus garrodiae* (Clay, 1940); Murray *et al.* 1990: 1372.

*Philoceanus garrodiae* (Clay, 1940); Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Garrodia nereis* (Gould, 1841).

New Zealand host: *Garrodia nereis* (Gould, 1841).

Other hosts: None.

New Zealand localities: WN, NC, MC, SC, WD, CO, DN, CH, AN, AU, CA.

Geographic distribution: Southern Oceans.

New Zealand references: Clay (1964a); Gressitt (1964: 538); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Marris (2000: 188); Page *et al.* (2004: 638, 650); Palma (2010).

Other significant references: Timmermann (1961b; 1965); Green & Palma (1991: 17, 27); Palma (1996b: 203); Price *et al.* (2003: 212).

Remarks: *Philoceanus garrodiae* is a highly prevalent “wing” louse species, known from the grey-backed storm petrel only.

***Philoceanus robertsi* (Clay, 1940)**

*Naubates robertsi* Clay, 1940a: 313, figs 9c, 10a, 11b.

*Philoceanus robertsi* (Clay, 1940); Hopkins & Clay 1952: 279.

*Philoceanus robertsi* (Clay, 1940); Timmermann 1961b: 533, fig. 4.

*Philoceanus robertsi* (Clay, 1940); Timmermann 1965: 158, fig. 97.

*Philoceanus robertsi* (Clay, 1940); Clay & Moreby 1967: 163, 169, fig. 149.

*Philoceanus robertsi* (Clay, 1940); Pilgrim & Palma 1982: 13.

*Philoceanus robertsi* (Clay, 1940); Murray *et al.* 1990: 1372.

*Philoceanus robertsi* (Clay, 1940); Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Oceanites oceanicus exasperatus* Mathews, 1912.

New Zealand host: *Oceanites oceanicus exasperatus* Mathews, 1912.

Other hosts: *Oceanites oceanicus oceanicus* (Kuhl, 1820); *Oceanites pincoyae* Harrison *et al.*, 2013; *Hydrobates pelagicus* (Linnaeus, 1758).

New Zealand localities: ND, RO.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Séguy (1953: 572, figs 26–29); Timmermann (1961b; 1965); Clay & Moreby (1967); Fowler & Miller (1984: 24); Fowler & Price (1987: 44, fig. 1); Palma (1996b: 203); Price *et al.* (2003: 212); Page *et al.* (2004: 638, 650); Harrison *et al.* (2013: 186).

Remarks: *Philoceanus robertsi* is an infrequently collected “wing” louse species.

**Genus *Philopteroides* Mey, 2004**

*Philopteroides* Mey, 2004. *Ornithol. Anz.* 43: 173. Type species: *Philopteroides novaezelandiae* Mey, 2004 (by original designation).

***Philopteroides fuliginosus* Valim & Palma, 2013**

Figs 150–151

*Philopterus* sp.; Pilgrim & Palma 1982: 26.

*Philopterus* sp.; Paterson *et al.* 1999: 221.

*Philopterus* sp.; Galloway 2005: 17.

*Philopterus* sp.; Murray *et al.* 2006b: 1956.

*Philopteroides fuliginosus* Valim & Palma, 2013: 81, figs 5–6, 11, 21–24, 34–35.

Holotype ♀ in MONZ.

Type host: *Rhipidura fuliginosa placabilis* Bangs, 1921.

New Zealand hosts: *Rhipidura fuliginosa fuliginosa* (Sparrman, 1787); *Rhipidura fuliginosa placabilis* Bangs, 1921.

Other hosts: None.

New Zealand localities: CL, WN, NN, MC, WD.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999); Galloway (2005); Murray *et al.* (2006b); Valim & Palma (2013); Najer *et al.* (2016: 527).

Other significant reference: Mey (2004).

Remarks: *Philopteroides fuliginosus* is an endemic species, although it is possible that some Australian species of *Rhipidura* closely related to *Rhipidura fuliginosa* (see Checklist Committee 2010: 298) may harbour this species of *Philopteroides*.

***Philopteroides macrocephalus* Valim & Palma, 2013**

*Philopterus* sp.; Horning *et al.* 1980: 6, 12.

*Philopterus* sp.; Pilgrim & Palma 1982: 26.

*Philopterus* sp.; Murray *et al.* 2002: 1216.

*Philopteroides macrocephalus* Valim & Palma, 2013: 83, figs 7–8, 12, 25–28, 36–37.

Holotype ♀ in MONZ.

Type host: *Petroica macrocephala macrocephala* (J.F. Gmelin, 1789).

New Zealand hosts: *Petroica macrocephala toitoi* (Lesson, 1828); *Petroica macrocephala macrocephala* (J.F. Gmelin, 1789); *Petroica macrocephala damnefaerdi* (Rothschild, 1894).

Other hosts: None.

New Zealand localities: WN, BR, WD, CH, SN.

Geographic distribution: New Zealand.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (2002); Valim & Palma (2013); Najer *et al.* (2016: 527).

Other significant reference: Mey (2004).

Remarks: *Philopteroides macrocephalus* is an endemic species, exclusively parasitic on New Zealand tomtits.

***Philopteroides novaezelandiae* Mey, 2004**

*Philopteroides novaezelandiae* Mey, 2004: 174, figs 21, 22c,d.

*Philopteroides novaezelandiae* Mey, 2004; Palma 2010: 408.

*Philopteroides novaezelandiae* Mey, 2004; Valim & Palma, 2013: 80.

Holotype ♂ in NHMR.

Type host: *Acanthisitta chloris chloris* (Sparrman, 1787).

New Zealand host: *Acanthisitta chloris chloris* (Sparrman, 1787).

Other hosts: None.

New Zealand locality: WD.

Geographic distribution: South Island, New Zealand.

New Zealand references: Mey (2004); Palma (2010); Buckley *et al.* (2012: App. 2); Valim & Palma (2013); Najer *et al.* (2016: 527).

Other significant references: None.

Remarks: *Philopteroides novaezelandiae* is an endemic species, known from the two type specimens only (Mey 2004: 174).

***Philopteroides pilgrimi* Valim & Palma, 2013**

*Philopterus* sp.; Gill 1980: 246.

*Philopterus* sp.; Pilgrim & Palma 1982: 26.

*Philopterus* sp.; Murray *et al.* 2002: 1215.

*Philopteroides pilgrimi* Valim & Palma, 2013: 77, figs 3–4, 10, 17–20, 32–33.

Holotype ♀ in MONZ.

Type host: *Gerygone igata* (Quoy & Gaimard, 1830).

New Zealand host: *Gerygone igata* (Quoy & Gaimard, 1830).

Other hosts: None.

New Zealand localities: WN, KA.

Geographic distribution: New Zealand.

New Zealand references: Gill (1980); Pilgrim & Palma (1982); Murray *et al.* (2002); Valim & Palma (2013); Najer *et al.* (2016: 527, 540).

Other significant reference: Mey (2004).

Remarks: *Philopteroides pilgrimi* is an endemic species, exclusively parasitic on New Zealand grey warblers.

***Philopteroides xenicus* Mey, 2004**

*Philopteroides xenicus* Mey, 2004: 176, figs 22a,b,e.

*Philopteroides xenicus* Mey, 2004; Mey 2005: 214.

*Philopteroides xenicus* Mey, 2004; Palma 2010: 408.

*Philopteroides xenicus* Mey, 2004; Valim & Palma, 2013: 80.

Holotype ♂ in NHMR.

Type host: *Xenicus longipes longipes* (J.F. Gmelin, 1789).

New Zealand host: *Xenicus longipes longipes* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand locality: WD.

Geographic distribution: New Zealand: South Island.

New Zealand references: Mey (2004; 2005); Palma (2010); Buckley *et al.* (2012: 140); Valim & Palma (2013); Rózsa & Vas (2015b: 108); Najer *et al.* (2016: 527).

Other significant references: None.

Remarks: *Philopteroides xenicus* is an endemic species, known from the holotype only. Considering that the host of this louse—the South Island bush wren—is extinct (Checklist Committee 2010: 277), its louse is also regarded as extinct (Buckley *et al.* 2012; Rózsa & Vas 2015b: 108).

**Genus *Philopterus* Nitzsch, 1818**

*Philopterus* Nitzsch, 1818. *Germer's Mag. Entomol.* 3: 281, 288. Type species: *Philopterus (Docophorus) ocellatus* "Nitzsch" = *Philopterus ocellatus* (Scopoli, 1763) (by subsequent designation).

*Docophorus* Nitzsch, 1818. *Germer's Mag. Entomol.* 3: 289. Type species: *Philopterus (Docophorus) ocellatus* (Scopoli, 1763) (by subsequent designation).

*Docophorus* Eichler, 1944b. *Stettin. Entomol. Zeit.* 105: 80. Type species: *Docophorus communis* Nitzsch var. *passeris* Piaget, 1880 = *Philopterus fringillae* (Scopoli, 1772) (by original designation).

***Philopterus irkutensis* Fedorenko, 1985****New Record**

*Philopterus* sp.; Pilgrim & Palma 1982: 25.

*Philopterus irkutensis* Fedorenko, 1985: 14, fig. 3.

*Philopterus* sp.; Paterson *et al.* 1999: 223.

*Philopterus* species; Marris 2000: 188.

*Philopterus irkutensis* Fedorenko, 1985; Price *et al.* 2003: 214.

*Philopterus* sp.; Murray *et al.* 2006b: 1959.

Syntypes ♂♀ in the Schmalhausen Institute of Zoology, Kiev, Ukraine (Fedorenko 1985: 15).

Type host: *Anthus richardi* Vieillot, 1818.

New Zealand hosts: *Anthus novaeseelandiae novaeseelandiae* (J.F. Gmelin, 1789); *Anthus novaeseelandiae steindachneri* Reischek, 1889.

Other host: *Anthus richardi* Vieillot, 1818.

New Zealand localities: GB, MC, WD, AN.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999); Marris (2000); Murray *et al.* (2006b).

Other significant reference: Price *et al.* (2003).

Material examined and repository: 22♂, 25♀ (6 samples, MONZ).

Remarks: This is the first record of *Philopterus irkutensis* for New Zealand because the New Zealand references cited above reported this louse as "*Philopterus* sp." only.

***Philopterus modularis* Denny, 1842****New Record**

*Philopterus (Docophorus) modularis* Denny, 1842: 47, 107, pl. 3: fig. 3.

*Philopterus modularis* Denny, 1842; Harrison 1916: 99.

*Philopterus modularis* (Denny, 1842) [sic]; Hopkins & Clay 1952: 286.

*Docophorus modularis* (Denny, 1842); Złotorzycka 1964b: 417, fig. 5c.

*Philopterus* sp.; Pilgrim & Palma 1982: 26.  
*Philopterus* sp.; Paterson *et al.* 1999: 220.  
*Philopterus* sp.; Murray *et al.* 2006b: 1958.  
*Philopterus modularis* (Denny, 1842) [sic]; Price *et al.* 2003: 215.

Syntypes ♀♀ in NHML (Vincent S. Smith pers. comm. December 2014).

Type host: *Prunella modularis occidentalis* (Hartert, 1910).

New Zealand host: *Prunella modularis* (Linnaeus, 1758).

Other hosts: None.

New Zealand localities: WN, MC, WD, CH, CA.

Geographic distribution: Europe, Eastern Asia.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999); Murray *et al.* (2006b).

Other significant references: Złotorzycka (1964b); Price *et al.* (2003).

Material examined and repository: 11♂, 18♀ (7 samples, MONZ).

Remarks: This is the first record of *Philopterus modularis* for New Zealand, because the New Zealand references cited above reported this louse as “*Philopterus* sp.” only. *Philopterus modularis* was introduced to New Zealand with dunnocks by human agency (Checklist Committee 2010: 318), and it is the only species that has become established from a total of six species which parasitise dunnocks in their home range (Price *et al.* 2003: 354).

Denny (1842: 63) included *Docophorus* as a subgenus of *Philopterus*; therefore, there is no reason to place author and date within parentheses when this species is combined with the genus *Philopterus*, as in Hopkins & Clay (1952: 286) and other authors who followed them.

#### ***Philopterus novaezealandiae* Palma & Price, 2000**

Figs 152–153

*Philopterus* sp.; Pilgrim & Palma 1982: 28.  
*Philopterus novaezealandiae* Palma & Price, 2000: 293, figs 1–4.  
*Philopterus novaezealandiae* Palma & Price, 2000; Murray *et al.* 2006b: 1958.  
*Philopterus novaezealandiae* Palma & Price, 2000; Palma 2010: 408.

Holotype ♂ in MONZ.

Type host: *Callaeas wilsoni* (Bonaparte, 1851).

New Zealand hosts: *Callaeas wilsoni* (Bonaparte, 1851); *Callaeas cinerea* (J.F. Gmelin, 1788).

Other hosts: None.

New Zealand localities: BP, WI, WN, FD.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Palma & Price (2000); Price *et al.* (2003: 215); Murray *et al.* (2006b); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: None.

Remarks: *Philopterus novaezealandiae* is an endemic and vulnerable species (Buckley *et al.* 2012), exclusively parasitic on the two species of kokako.

#### ***Philopterus stadleri* (Eichler, 1959)**

**New record**

*Docophorus stadleri* Eichler, 1959: 1173, fig. 3.  
*Philopterus stadleri* (Eichler, 1959); Fedorenko 1978: 57.  
*Philopterus* sp.; Pilgrim & Palma 1982: 25.  
*Philopterus stadleri* (Eichler, 1959); Price *et al.* 2003: 216.  
*Philopterus* sp.; Murray *et al.* 2006b: 1958.

Holotype ♀, repository unknown.

Type host: *Alauda arvensis arvensis* Linnaeus, 1758.

New Zealand host: *Alauda arvensis* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: BR, MC.

Geographic distribution: Eurasia; Australasia.



New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006b).

Other significant references: Fedorenko (1978); Price *et al.* (2003).

Material examined and repository: 1♂, 2♀, 4N (3 samples, MONZ).

Remarks: This is the first record of *Philopterus stadleri* for New Zealand, because the New Zealand references cited above reported this louse as “*Philopterus sp.*” only. *Philopterus stadleri* was introduced to New Zealand with skylarks by human agency (Checklist Committee 2010: 306).

### ***Philopterus turdi* Denny, 1842**

*Philopterus (Docophorus) turdi* Denny, 1842: 43, 76, pl. 4: fig. 5.

*Philopterus (Docophorus) merulae* Denny, 1842: 47, 106, pl. 3: fig. 1.

*Philopterus turdi* Denny, 1842; Harrison 1916: 106.

*Philopterus merulae* (Denny, 1842) [sic]; Hopkins & Clay 1952: 286.

*Philopterus turdi* (Denny, 1842) [sic]; Hopkins & Clay 1952: 289.

*Docophorus merulae* (Denny, 1842); Złotorzycka 1964b: 415, figs 5a, 6a.

*Docophorus turdi* (Denny, 1842); Złotorzycka 1964b: 423, figs 5l, 8a.

*Philopterus merulae* (Denny, 1842); Baum 1968: 143, fig. 7.

*Philopterus turdi* (Denny, 1842) [sic]; Nelson 1969: 199.

*Philopterus turdi* (Denny, 1842) [sic]; Watt 1971: 238, 244.

*Philopterus turdi* (Denny, 1842) [sic]; Wise 1977: 63.

*Philopterus turdi* (Denny, 1842) [sic]; Horning *et al.* 1980: 6, 12.

*Philopterus turdi* (Denny, 1842) [sic]; Pilgrim & Palma 1982: 26.

*Philopterus turdi* (Denny, 1842) [sic]; Murray *et al.* 2006b: 1959.

*Philopterus turdi* (Denny, 1842) [sic]; Palma 2010: 408.

Syntypes ♀♀ in NHML (Palma 1996b: 204).

Type host: *Turdus philomelos* Brehm, 1831.

New Zealand hosts: *Turdus merula merula* Linnaeus, 1758; *Turdus philomelos* Brehm, 1831.

Other hosts: None.

New Zealand localities: TK, WI, WN, NN, KA, BR, NC, MC, SC, WD, CO, DN, KE, Norfolk Island, CH, SN.

Geographic distribution: Eurasia; north Africa; Australasia.

New Zealand references: Nelson (1969); Watt (1971); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Galloway (2005: 15); Murray *et al.* (2006b); Palma (2010).

Other significant references: Złotorzycka (1964b); Baum (1968); Jiménez-González *et al.* (1980: 213, figs 21–25); Palma (1996b: 204); Price *et al.* (2003: 217); Palma & Jensen (2005: 57, 67); Martín-Mateo (2009: 148).

Remarks: *Philopterus turdi* was introduced to New Zealand with blackbirds and song thrushes by human agency (Checklist Committee 2010: 313). Denny (1842: 63) included *Docophorus* as a subgenus of *Philopterus*; therefore, there is no reason to place author and date within parentheses when this species is combined with the genus *Philopterus*, as in Hopkins & Clay (1952: 289) and other authors who followed them.

### ***Philopterus* species 1**

*Philopterus* sp.; Pilgrim & Palma 1982: 28.

*Philopterus* sp.; Hughes 1984a: 459.

*Philopterus* sp.; Paterson *et al.* 1999: 221.

*Philopterus* sp.; Murray *et al.* 2006b: 1957.

New Zealand host: *Gymnorhina tibicen* (Latham, 1802).

Other hosts: None.

New Zealand localities: BP, GB, HB.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999: 221); Murray *et al.* (2006b).

Other significant references: Hughes (1984a); Hughes (1984b: 467); Toon & Hughes (2008: 128).

Remarks: This unidentified and probably new species of *Philopterus* was introduced to New Zealand with Australian magpies by human agency (Checklist Committee 201: 297).

***Philopterus* species 2**

*Philopterus* sp.; Pilgrim & Palma 1982: 26.

*Philopterus* sp.; Murray *et al.* 2002: 1216.

New Zealand host: *Mohoua albicilla* (Lesson, 1830).

Other hosts: None.

New Zealand locality: WN.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2002).

Other significant references: None.

Remarks: The available sample of *Philopterus* from *Mohoua albicilla* is not identifiable at present because it contains a single nymph (voucher specimen in MONZ).

**Genus *Pseudonirmus* Mjöberg, 1910**

*Pseudonirmus* Mjöberg, 1910a. *Arkiv Zool.* 6(13): 149. Type species: *Degeeriella charcoti* Neumann, 1907a = *Pseudonirmus charcoti* (Neumann, 1907) (by original designation).

***Pseudonirmus charcoti* (Neumann, 1907)**

*Degeeriella charcoti* Neumann, 1907a: 15, figs 2–3.

*Ricinus charcoti* (Neumann, 1907); Enderlein 1908: 477.

*Nirmus antarcticus* Valette, 1913: 298, pl. 5: figs 1–5.

*Esthiopterum charcoti* Neumann, 1907 [sic]; Harrison 1916: 132.

*Pseudonirmus charcoti* (Neumann, 1907); Hopkins & Clay 1952: 303.

*Pseudonirmus antarcticus* (Valette, 1913); Hopkins & Clay 1955: 186 (as a junior synonym).

*Pseudonirmus charcoti* (Neumann, 1907); Timmermann 1961c: 33, figs 1, 5a, 6.

*Pseudonirmus charcoti* (Neumann, 1907); Timmermann 1965: 113, figs 51, 52a, pl. 10: fig. 1.

*Pseudonirmus charcoti* (Neumann, 1907); Clay & Moreby 1967: 163, 168, figs 131, 135, 138.

*Pseudonirmus charcoti* (Neumann, 1907); Pilgrim & Palma 1982: 8.

*Pseudonirmus charcoti* (Neumann, 1907); Murray *et al.* 1990: 1369.

*Pseudonirmus charcoti* (Neumann, 1907); Palma 2010: 408.

Syntypes ♂♀, repository unknown (Palma 1996b: 206), probably in the Ecole Vétérinaire de Toulouse, France.

Type host: *Pagodroma nivea* (G. Forster, 1777).

New Zealand hosts: *Pagodroma nivea nivea* (G. Forster, 1777); *Pagodroma nivea major* (Schlegel, 1863).

Other hosts: None.

New Zealand locality: RO.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 222); Palma (2010).

Other significant references: Thompson (1935a: 555, 557); Harrison (1937: 26, fig. 2c); Timmermann (1961c; 1965); Clay & Moreby (1967); Steele *et al.* (1997: 292); Price *et al.* (2003: 221).

Remarks: *Pseudonirmus charcoti* is an infrequently collected “wing” louse species, exclusively parasitic on snow petrels.

***Pseudonirmus gurlti* (Taschenberg, 1882)**

*Lipeurus gurlti* Taschenberg, 1882: 151, pl. 5: fig. 6.

*Esthiopterum gurlti* Taschenberg, 1882 [sic]; Harrison 1916: 135.

*Pseudonirmus gurlti* (Taschenberg, 1882); Hopkins & Clay 1952: 303.

*Pseudonirmus gurlti* (Taschenberg, 1882); Timmermann 1961c: 35, figs 3, 5c.

*Pseudonirmus gurlti* (Taschenberg, 1882); Timmermann 1965: 115, fig. 52c, pl. 10: fig. 3.

*Pseudonirmus gurlti* (Taschenberg, 1882); Clay & Moreby 1967: 163, 168, figs 133, 137, 140.

*Pseudonirmus gurlti* (Taschenberg, 1882); Horning *et al.* 1980: 7, 9.

*Pseudonirmus gurlti* (Taschenberg, 1882); Pilgrim & Palma 1982: 7.

*Pseudonirmus gurlti* (Taschenberg, 1882); Murray *et al.* 1990: 1369.

*Pseudonirmus gurlti* (Taschenberg, 1882); Palma & Horning 2002: 4 (figs), 11, 16.

*Pseudonirmus gurlti* (Taschenberg, 1882); Palma 2010: 408.

Syntypes ♂♀, repository unknown, presumed lost. See Palma (1996b: 206).

Type host: *Daption capense* (Linnaeus, 1758).

New Zealand hosts: *Daption capense capense* (Linnaeus, 1758); *Daption capense australe* Mathews, 1913.

Other hosts: None.

New Zealand localities: ND, AK, HB, TK, WI, WN, NC, MC, SC, WD, SL, SI, BO, SN, Macquarie Island, RO.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Watson (1967: 73); Gressitt (1970: 328); Wise (1977: 63); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b: 206); Paterson *et al.* (1999: 222); Palma & Horning (2002); Greenslade (2006: figs 20.3–20.4); Palma (2010).

Other significant references: Thompson (1935c: 486, fig. 2); Harrison (1937: 26, fig. 2a); Clay (1940a: 298); Guimarães (1943: 425, fig. 1); Séguy (1953: 563, figs 9–10); Timmermann (1961c; 1965); Clay & Moreby (1967); Green & Palma (1991: 17, 26); Price *et al.* (2003: 221); Page *et al.* (2004: 638, 650).

Remarks: *Pseudonirmus gurlti* is a frequently collected “wing” louse species, exclusively parasitic on Cape petrels.

### ***Pseudonirmus lugubris* (Taschenberg, 1882)**

Figs 154–155

*Lipeurus lugubris* Taschenberg, 1882: 153, pl. 6: fig. 9.

*Esthiopterum lugubre* Taschenberg, 1882 [sic]; Harrison 1916: 137 (as junior synonym of *Esthiopterum gurlti* Taschenberg, 1882 [sic]).

*Pseudonirmus antarcticus* Harrison, 1937: 26, fig. 2b. Preoccupied by *Pseudonirmus antarcticus* (Valette, 1913).

*Pseudonirmus antarcticus* Harrison, 1937; Hopkins & Clay 1952: 303.

*Pseudonirmus lugubris* (Taschenberg, 1882); Hopkins & Clay 1952: 303 (as junior synonym of *Pseudonirmus gurlti* (Taschenberg, 1882)).

*Pseudonirmus lugubris* (Taschenberg, 1882); Clay & Hopkins 1955: 65.

*Pseudonirmus antarcticus* Harrison, 1937; Timmermann 1961c: 34, figs 2, 5b.

*Pseudonirmus lugubris* (Taschenberg, 1882); Timmermann 1965: 113, figs 50, 52b, pl. 10: fig. 2.

*Pseudonirmus lugubris* (Taschenberg, 1882); Clay & Moreby 1967: 163, 168, figs 132, 136, 139.

*Pseudonirmus lugubris* (Taschenberg, 1882); Pilgrim & Palma 1982: 7.

*Pseudonirmus lugubris* (Taschenberg, 1882); Murray *et al.* 1990: 1369.

*Pseudonirmus lugubris* (Taschenberg, 1882); Palma 2010: 408.

Lectotype ♂ in ZMHG (Clay & Hopkins 1955: 65; Weidner 1966: 258). Holotype ♂ of *Pseudonirmus antarcticus* in AMSA (Palma 1996b: 207).

Type host: *Thalassoica antarctica* (J.F. Gmelin, 1789).

New Zealand host: *Thalassoica antarctica* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: ND, AK, WI, RO.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Harrison (1937); Clay & Hopkins (1955); Timmermann (1961c; 1965); Weidner (1966); Clay & Moreby (1967); Palma (1996b: 206); Price *et al.* (2003: 221).

Remarks: *Pseudonirmus lugubris* is a “wing” louse species, exclusively parasitic on Antarctic petrels. *Thalassoica antarctica* breeds on the coast of Antarctica and straggles to New Zealand seas (Checklist Committee 2010: 82).

### **Genus *Psittoecus* Conci, 1942**

*Psittoecus* Conci, 1942c. *Boll. Soc. Entomol. Italiana* 74: 40. Type species: *Philopterus waterstoni* Fresca, 1923 = *Psittoecus waterstoni* (Fresca, 1923) (by original designation).

### ***Psittoecus vanzolinii* Guimarães, 1974**

Figs 156–157

*Psittoecus vanzolinii* Guimarães, 1974b: 168, figs 9–11.

*Psittoecus vanzolinii* Guimarães, 1974; Palma 1999: 381.

*Psittoecus vanzolinii* Guimarães, 1974; Price *et al.* 2003: 222.

*Psittoecus vanzolinii* Guimarães, 1974; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Cacatua galerita* (Latham, 1790).

New Zealand host: *Cacatua galerita* (Latham, 1790).

Other hosts: None.

New Zealand locality: WI.

Geographic distribution: Australasia.

New Zealand references: Palma (1999; 2010).

Other significant references: Green & Palma (1991: 18, 34); Palma (1996b: 208); Price *et al.* (2003).

Remarks: It is not possible to assert if *Psittoecus vanzolinii* was introduced to New Zealand with sulphur-crested cockatoos by human agency, or if it was self-introduced with its host (Checklist Committee 2010: 252).

### Genus *Quadriceps* Clay & Meinertzhagen, 1939

*Quadriceps* Clay & Meinertzhagen, 1939b. *Ann. Mag. Nat. Hist.* (Ser. 11) 4: 453. Type species: *Degeeriella vanelli* (Denny, 1842) = *Quadriceps charadrii hospes* (Nitzsch [in Giebel], 1866) (by original designation).

*Koeniginirmus* Eichler, 1940b. *Zool. Anz.* 130: 101. Type species: “*Koeniginirmus punctatus* (Nitzsch in Burmeister, 1838a)” = *Quadriceps punctatus* (Burmeister, 1838a) (by original designation).

*Glareolites* Eichler, 1944b. *Stettin. Entomol. Zeit.* 105: 80. Type species: *Nirmus ellipticus* Nitzsch [in Giebel], 1866 = *Quadriceps ellipticus* (Nitzsch [in Giebel], 1866) (by original designation).

*Haematophagus* Timmermann, 1950. *Fauna Islandica* 2: 1, 2. Type species: *Quadriceps haematopi* (Denny, 1842 = *Quadriceps auratus* (Haan, 1829) (by original designation).

*Proneptis* Timmermann, 1953a. *Bombus* 78/79: 331. Type species: *Proneptis semifissa* (Nitzsch [in Giebel], 1866) = *Quadriceps semifissus* (Nitzsch [in Giebel], 1866) (by original designation).

*Chadriceps* Złotorzycka, 1967. *Polskie Pismo Entomol.* 37(4): 728. Type species: *Pediculus hiaticulae* O. Fabricius, 1780 = *Quadriceps hiaticulae* (O. Fabricius, 1780) (by original designation).

*Anousceps* Złotorzycka, 1967. *Polskie Pismo Entomol.* 37(4): 730. Type species: *Nirmus separatus* Kellogg & Kuwana, 1902 = *Quadriceps separatus* (Kellogg & Kuwana, 1902) (by original designation).

*Himantophagus* Złotorzycka, 1967. *Polskie Pismo Entomol.* 37(4): 733. Type species: *Nirmus hemichrous* Nitzsch [in Giebel], 1866 = *Quadriceps hemichrous* (Nitzsch [in Giebel], 1866) (by original designation).

*Laminonirmus* Złotorzycka, 1967. *Polskie Pismo Entomol.* 37(4): 754. Type species: *Koeniginirmus ornatus* (Grube, 1851) = *Quadriceps ornatus ornatus* (Grube, 1851) (by original designation). As subgenus of *Koeniginirmus* Eichler, 1940.

### *Quadriceps assimilis* (Piaget, 1890)

*Nirmus assimilis* Piaget, 1890b: xxiii, fig. 1.

*Degeeriella assimilis* Piaget, 1890 [sic]; Harrison 1916: 108.

*Quadriceps assimilis* (Piaget, 1890); Hopkins & Clay 1952: 308.

*Quadriceps assimilis assimilis* (Piaget, 1890); Timmermann 1953b: 179.

*Quadriceps assimilis assimilis* (Piaget, 1890); Timmermann 1969b: 250.

*Quadriceps assimilis* (Piaget, 1890); Palma 1999: 380.

*Quadriceps assimilis* (Piaget, 1890); Palma 2010: 408.

Syntypes ♂♀, repository unknown.

Type host: *Charadrius asiaticus* Pallas, 1773.

New Zealand host: *Charadrius veredus* Gould, 1848.

Other hosts: *Charadrius peronii* (Schlegel, 1865); *Charadrius montanus* Townsend, 1837.

New Zealand locality: KE.

Geographic distribution: Eurasia; Africa; North America; Australasia.

New Zealand references: Palma (1999; 2010).

Other significant references: Timmermann (1953b; 1957a: 147; 1969); Price *et al.* (2003: 223).

Remarks: The oriental dotterel is a rare visitor to New Zealand (Checklist Committee 2010: 219), with only a single record of *Quadriceps assimilis* from this country.

***Quadriceps auratus* (Haan, 1829)**

- Philopterus* (*Docophorus*) *auratus* Haan, 1829: 310, pl. 5: fig. 9.  
*Degeeriella aurata* Lyonet [sic], 1829 [sic]; Harrison 1916: 109.  
*Quadriceps auratus* (Haan, 1829); Hopkins 1949a: 29, pl. 1: figs 1–2.  
*Quadriceps auratus* (Haan, 1829); Hopkins & Clay 1952: 309.  
*Haematophagus auratus* (Haan, 1829); Złotorzycka 1967: 748, pl. 10: figs 1–4.  
*Quadriceps auratus*; Baker 1974: 20.  
*Quadriceps auratus* (Haan, 1829); Clay 1981b: 936, figs 4, 12, 14, 17–18, 20, 22.  
*Quadriceps auratus* (Haan, 1829); Pilgrim & Palma 1982: 19.  
*Cummingsiella auratus* (de Haan, 1829) [sic]; Butler & O'Connor 1994: 454.  
*Quadriceps auratus* (Haan, 1829); Murray *et al.* 1993: 961.  
*Quadriceps auratus* (Haan, 1829); Palma 2010: 408.

Neotype ♀ in NHML (Hopkins 1949a: 31, pl 1, fig. 2).

Type host: *Haematopus ostralegus* Linnaeus, 1758.

New Zealand hosts: *Haematopus finschi* Martens, 1897; *Haematopus unicolor* J.R. Forster, 1844; *Haematopus chathamensis* Hartert, 1927.

Other hosts: *Haematopus ater* Vieillot & Oudart, 1825; *Haematopus bachmani* Audubon, 1838; *Haematopus palliatus* Temminck, 1820; *Haematopus moquini* Bonaparte, 1856; *Haematopus longirostris* Vieillot, 1817; *Haematopus fuliginosus* Gould, 1845.

New Zealand localities: AK, WI, SD, MB, NN, NC, MC, SC, WD, CO, DN, SL, CH.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Baker (1974); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 223); Palma (2010).

Other significant references: Hopkins (1949a); Złotorzycka (1967); Clay (1981b); Green & Palma (1991: 18, 32); Butler & O'Connor (1994); Forrester *et al.* (1995: 25); Palma (1995: 219); Palma (1996b: 209); Price *et al.* (2003: 223); Palma & Peck (2013: 49).

Remarks: *Quadriceps auratus* is the most widespread and frequently collected species of *Quadriceps* from oystercatchers.

***Quadriceps birostris* (Giebel, 1874)**

- Nirmus birostris* Giebel, 1874: 174.  
*Nirmus gloriosus* Kellogg & Kuwana, 1902: 467, pl. 29: fig. 1.  
*Degeeriella birostris* Giebel, 1874 [sic]; Harrison 1916: 109.  
*Degeeriella gloriosa* Kellogg & Kuwana, 1902 [sic]; Harrison 1916: 114.  
*Degeeriella gloriosa* (Kellogg & Kuwana, 1902); Ferris 1932a: 68, figs 18a,b,c,d,e.  
*Koeniginirmus nychthemerus birostris* Giebel, 1874 [sic]; Eichler 1951b: 127.  
*Quadriceps birostris* (Giebel, 1874); Hopkins & Clay 1952: 309.  
*Quadriceps birostris* (Giebel, 1874); Timmermann 1952a: 76, figs 4–5.  
*Quadriceps birostris* (Giebel, 1874); Timmermann 1957a: 71, pl. 9: figs a,b.  
*Koeniginirmus (Laminonirmus) nychthemerus birostris* (Giebel, 1874); Złotorzycka 1967: 761.  
*Quadriceps birostris* (Giebel, 1874); Nelson 1969: 199.  
*Quadriceps birostris* (Giebel, 1874); Watt 1971: 238, 243.  
*Quadriceps birostris* (Giebel, 1874); Pilgrim & Palma 1982: 23.  
*Quadriceps birostris* (Giebel, 1874); Murray *et al.* 2006a: 1965.  
*Quadriceps birostris* (Giebel, 1874); Palma 2010: 408.

Syntypes ♂♂ presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: “*Sterna n. sp. probably Sterna fuliginosa*” [= *Onychoprion fuscatus* (Linnaeus, 1766)] (Hopkins & Clay 1952: 309).

New Zealand host: *Onychoprion fuscatus serratus* (J.R. Forster, 1830).

Other hosts: *Onychoprion fuscatus fuscatus* (Linnaeus, 1766); *Onychoprion fuscatus oahuensis* Bloxham, 1826; *Onychoprion fuscatus crissalis* (Lawrence, 1872).

New Zealand localities: WO, KE, Norfolk Island.

Geographic distribution: Tropical and subtropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Nelson (1969); Watt (1971); Wise (1977: 63); Pilgrim & Palma (1982); Palma (1994b: 268); Murray *et al.* (2006a); Palma (2010).

Other significant references: Ferris (1932a); Eichler (1951b); Timmermann (1952a; 1957a); Carriker (1957: 100); Złotorzycka (1967); Amerson & Emerson (1971: 17, 25); Ward & Downey (1973: 394); Forrester *et al.* (1995: 31); Palma (1996b: 210); Price *et al.* (2003: 223); Palma & Peck (2013: 50).

Remarks: *Quadriceps birostris* appears to be restricted to the subspecies of the sooty tern.

### ***Quadriceps caspius* (Giebel, 1874)**

Figs 158–159

*Nirmus caspius* Giebel, 1874: 174.

*Degeeriella caspia* Giebel, 1866 [sic]; Harrison 1916: 110.

*Koeniginirmus sellatus caspius* Giebel, 1874 [sic]; Eichler 1951b: 134.

*Quadriceps caspius* (Giebel, 1874); Timmermann 1952a: 83.

*Quadriceps caspius* (Giebel, 1874); Hopkins & Clay 1952: 309.

*Quadriceps caspius* (Giebel, 1874); Timmermann 1957a: 72, pl. 11: figs a,b.

*Koeniginirmus (Laminonirmus) caspius caspius* (Giebel, 1874); Złotorzycka 1967: 756, pl. 17: fig. 3.

*Quadriceps caspius* (Giebel, 1874); Pilgrim & Palma 1982: 22.

*Quadriceps caspius* (Giebel, 1874); Murray *et al.* 2006a: 1965.

*Quadriceps caspius* (Giebel, 1874); Palma 2010: 408.

Syntypes ♂♀, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Hydroprogne caspia* (Pallas, 1770).

New Zealand host: *Hydroprogne caspia* (Pallas, 1770).

Other hosts: None.

New Zealand localities: ND, AK, SC.

Geographic distribution: Eurasia; Africa; North America; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Eichler (1951b); Timmermann (1952a; 1957a); Złotorzycka (1967); Forrester *et al.* (1995: 30); Price *et al.* (2003: 223).

Remarks: *Quadriceps caspius* is frequently collected from Caspian terns.

### ***Quadriceps cedemajori* Timmermann, 1969**

*Quadriceps assimilis cedemajori* Timmermann, 1969b: 250, fig. 6b.

*Quadriceps assimilis cedemajori* Timmermann, 1969; Wise 1977: 63.

*Quadriceps cedemajori* Timmermann, 1969; Martens & Palma 1981: 83.

*Quadriceps cedemajori* Timmermann, 1969; Pilgrim & Palma 1982: 20.

*Quadriceps cedemajori* Timmermann, 1969; Murray *et al.* 1993: 962.

*Quadriceps cedemajori* Timmermann, 1969; Palma 2010: 407.

Holotype ♂ in BPBM (Tenorio 1979: 12).

Type host: *Charadrius bicinctus* Jardine & Selby, 1827.

New Zealand hosts: *Charadrius bicinctus bicinctus* Jardine & Selby, 1827; *Charadrius bicinctus exilis* Falla, 1978; *Anarhynchus frontalis* Quoy & Gaimard, 1830.

Other hosts: None

New Zealand localities: AK, CL, TK, WN, NN, KA, MC, WD, OL, CH, AU.

Geographic distribution: Australasia; Oceania.

New Zealand references: Wise (1977); Weidner (1977: 102); Martens & Palma (1981); Pilgrim & Palma (1982); Murray *et al.* (1993); Davies (1995: 46); Paterson *et al.* (1999: 223); Palma (2010).

Other significant reference: Price *et al.* (2003: 223).

Remarks: Remarks: *Quadriceps cedemajori* has an unusual host distribution on three New Zealand endemic plovers (Martens & Palma 1981: 84).

***Quadriceps charadrii orarius* (Kellogg, 1896)**

*Nirmus orarius* Kellogg, 1896a: 104, pl. 5: fig. 5.

*Degeeriella oraria* Kellogg [sic]; Johnston & Harrison 1912: 368.

*Degeeriella oraria* (Kellogg, 1896); Thompson 1939: 120.

*Quadriceps orarius* (Kellogg, 1896); Hopkins & Clay 1952: 314.

*Quadriceps charadrii orarius* (Kellogg, 1896); Timmermann 1953b: 184.

*Quadriceps orarius* (Kellogg, 1896); Złotorzycka 1967: 720.

*Quadriceps charadrii orarius* (Kellogg, 1896); Emerson 1972a: 137.

*Quadriceps orarius* (Kellogg, 1896); Wise 1977: 64.

“*Quadriceps charadrii*” Pilgrim & Palma, 1982: 20, 31, note 21 (not *Pediculus charadrii* Linnaeus, 1758).

“*Quadriceps charadrii*” Murray *et al.*, 1993: 962 (not *Pediculus charadrii* Linnaeus, 1758).

*Quadriceps orarius* (Kellogg, 1896); Price *et al.* 2003: 226.

“*Quadriceps charadrii charadrii*” Palma, 2010: 408 (not *Pediculus charadrii* Linnaeus, 1758).

Holotype ♀ in EMEC (Carriker 1957: 101).

Type host: *Pluvialis dominicus* (Statius Müller, 1776).

New Zealand host: *Pluvialis fulva* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: AK, KE.

Geographic distribution: Americas; Asia; Australasia; Pacific Ocean.

New Zealand references: Johnston & Harrison (1912); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Thompson (1939); Timmermann (1953b); Carriker (1957: 101); Złotorzycka (1967); Amerson & Emerson (1971: 13, 25); Emerson (1972a); Moreby (1976: 93); Price *et al.* (2003).

Remarks: In agreement with Timmermann (1953b: 184) and contrary to Price *et al.* (2003: 226), I regard this louse taxon as a subspecies of *Quadriceps charadrii* (Linnaeus, 1758). I also regard these taxa as subspecies: *Q. charadrii hospes* (Nitzsch [*in* Giebel], 1866) and *Q. charadrii punctifer* Hopkins, 1949c (see Timmermann 1953b: 183; Emerson 1972a: 136).

***Quadriceps coenocoryphae* Timmermann, 1955**

*Quadriceps coenocoryphae* Timmermann, 1955: 523, fig. 9.

*Quadriceps coenocoryphae* Timmermann, 1955; Złotorzycka 1967: 713.

*Quadriceps coenocoryphae* Timmermann, 1955; Wise 1977: 63.

*Quadriceps coenocoryphae* Timmermann, 1955; Horning *et al.* 1980: 7, 11.

*Quadriceps coenocoryphae* Timmermann, 1955; Pilgrim & Palma 1982: 21.

*Quadriceps coenocoryphae* Timmermann, 1955; Murray *et al.* 2006a: 1964.

*Quadriceps coenocoryphae* Timmermann, 1955; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Coenocorypha aucklandica aucklandica* (G.R. Gray, 1845).

New Zealand hosts: *Coenocorypha pusilla* (Buller, 1869); *Coenocorypha huegeli* (Tristram, 1893); *Coenocorypha aucklandica aucklandica* (G.R. Gray, 1845); *Coenocorypha aucklandica perseverance* Miskelly & Baker, 2010.

Other hosts: None.

New Zealand localities: CH, SN, AU, CA.

Geographic distribution: New Zealand.

New Zealand references: Timmermann (1955); Złotorzycka (1967); Pilgrim (1970: 74); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003: 223).

Remarks: *Quadriceps coenocoryphae* is an endemic and “at risk” species (Buckley *et al.* 2012), exclusively parasitic on several species and subspecies of New Zealand snipes. *Coenocorypha aucklandica perseverance* is a new host record for *Quadriceps coenocoryphae* (voucher specimens in MONZ).

***Quadriceps dominella* Timmermann, 1953**

- Quadriceps dominella* Timmermann, 1953b: 186.  
*Quadriceps dominella* Timmermann, 1953; Hopkins & Clay 1955: 186.  
*Quadriceps dominella* Timmermann, 1953; Wise 1977: 63.  
*Quadriceps dominella* Timmermann, 1953; Martens 1980: 350, figs 1–2.  
*Quadriceps dominella* Timmermann, 1953; Martens & Palma 1981: 83.  
*Quadriceps dominella* Timmermann, 1953; Pilgrim & Palma 1982: 20.  
*Quadriceps dominella* Timmermann, 1953; Murray *et al.* 1993: 962.  
*Quadriceps dominella* Timmermann, 1953; Palma 2010: 408.

Holotype ♀ in NHML.

Type host: *Charadrius obscurus obscurus* (J.F. Gmelin, 1789).

New Zealand hosts: *Charadrius obscurus obscurus* (J.F. Gmelin, 1789); *Charadrius obscurus aquilonius* Dowding, 1994.

Other hosts: None.

New Zealand localities: ND, AK, MC, SI.

Geographic distribution: New Zealand.

New Zealand references: Timmermann (1953b); Złotorzycka (1967: 714); Pilgrim (1970: 74); Wise (1977); Martens (1980); Martens & Palma (1981); Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003: 224).

Remarks: *Quadriceps dominella* is an endemic and vulnerable species (Buckley *et al.* 2012), exclusively parasitic on New Zealand dotterels. *Charadrius obscurus aquilonius* is a new host record for *Quadriceps dominella* (voucher specimens in MONZ).

***Quadriceps ellipticus* (Nitzsch [*in* Giebel], 1866) *sensu lato***

- Nirmus ellipticus* Nitzsch [*in* Giebel], 1866: 371.  
*Degeeriella elliptica* Nitzsch [*in* Giebel], 1866 [sic]; Harrison 1916: 112.  
*Glareolites ellipticus* Nitzsch [*in* Giebel], 1866 [sic]; Eichler 1944b: 80, fig. 1.  
*Quadriceps ellipticus* (Nitzsch [*in* Giebel], 1866); Timmermann 1952d: 1030.  
*Glareolites ellipticus* (Nitzsch, 1866) [sic]; Złotorzycka 1967: 741.  
*Quadriceps ellipticus* (Nitzsch, 1866) [sic] *s. l.*; Pilgrim & Palma 1982: 22.  
*Quadriceps ellipticus* (Nitzsch [*in* Giebel], 1866); Martín-Mateo 1992b: 410, figs 1–3, 7, 12, 17.  
*Quadriceps ellipticus* (Nitzsch [*in* Giebel], 1866); Murray *et al.* 2006a: 1964  
*Quadriceps ellipticus* (Nitzsch [*in* Giebel], 1866); Palma 2010: 408.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Glareola pratincola* (Linnaeus, 1766).

New Zealand host: *Glareola maldivarum* J.R. Forster, 1795.

Other host: *Glareola lactea* Temminck, 1820.

New Zealand locality: NN.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Eichler (1944b); Timmermann (1952d); Tendeiro (1964: 201, fig. 19, pl. 10: fig. 30, pl. 11: fig. 32, pl. 12: fig. 34); Złotorzycka (1967); Martín-Mateo (1992b); Price *et al.* (2003: 224).

Remarks: In agreement with Pilgrim & Palma (1982: 22), I regard the population of *Quadriceps ellipticus* from *Glareola maldivarum* as not completely identical to that from the type host, but not sufficiently distinct to warrant its taxonomic separation. The oriental pratincole is a rare visitor to New Zealand (Checklist Committee 2010: 223), with only a single record of *Quadriceps ellipticus* from this country.

***Quadriceps hemichrous* (Nitzsch [*in* Giebel], 1866)**

- Nirmus hemichrous* Nitzsch [*in* Giebel], 1866: 372.  
*Degeeriella hemichroa* Nitzsch [*in* Giebel], 1866 [sic]; Harrison 1916: 114.



- Quadriceps hemichrous* (Nitzsch [*in* Giebel], 1866); Hopkins & Clay 1952: 311.  
*Quadriceps hemichrous* (Nitzsch [*in* Giebel], 1866); Timmermann 1954f: 169, fig. 8.  
*Himantophagus hemichrous* (Nitzsch [*in* Giebel], 1866); Złotorzycka 1967: 734.  
*Quadriceps hemichrous* (Nitzsch, 1866) [*sic*]; Pilgrim & Palma 1982: 21.  
*Quadriceps hemichrous* (Nitzsch, 1866) [*sic*]; Murray *et al.* 1993: 962.  
*Quadriceps hemichrous* (Nitzsch *in* Giebel, 1866); Palma 2010: 408.

Neotype ♂ in NHML (Timmermann 1954f: 170).

Type host: *Himantopus himantopus himantopus* (Linnaeus, 1758).

New Zealand hosts: *Himantopus himantopus leucocephalus* Gould, 1837; *Himantopus novaezelandiae* Gould, 1841.

Other host: *Himantopus himantopus mexicanus* (Stadius Müller, 1776).

New Zealand localities: AK, BP, HB, WI, WN, KA, NC, MC, SC, WD.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Timmermann (1954f: 170); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 221, 223); Palma (2010).

Other significant references: Timmermann (1957a: 65); Złotorzycka (1967); Hinojos & Canaris (1988: 328); Palma (1995: 219); Forrester *et al.* (1995: 26); Palma (1996b: 211); Price *et al.* (2003: 224); Palma & Peck (2013: 51).

Remarks: In addition to *Quadriceps hemichrous*, the species of the host genus *Himantopus* are parasitised by *Quadriceps semifissus* (see below).

#### ***Quadriceps hopkinsi apophoretus* Timmermann, 1969**

- Quadriceps hopkinsi apophoretus* Timmermann, 1969c: 198, fig. 4b.  
 “*Quadriceps hopkinsi*” Nelson, 1969: 199 (not *Quadriceps hopkinsi* Timmermann, 1952a).  
*Quadriceps hopkinsi apophoretus* Timmermann, 1969; Watt 1971: 238, 243, fig. 9.  
*Quadriceps hopkinsi apophoretus* Timmermann, 1969; Pilgrim & Palma 1982: 23.  
*Quadriceps apophoretus* Timmermann, 1969; Price *et al.* 2003: 223.  
*Quadriceps hopkinsi apophoretus* Timmermann, 1969; Murray *et al.* 2006a: 1965.  
*Quadriceps hopkinsi apophoretus* Timmermann, 1969; Palma 2010: 408.

Holotype ♂ in NZAC.

Type host: *Procelsterna cerulea albivitta* Bonaparte, 1856.

New Zealand host: *Procelsterna cerulea albivitta* Bonaparte, 1856.

Other host: *Procelsterna cerulea cerulea* (Bennett, 1840).

New Zealand localities: ND, BP, MC, KE, Norfolk Island.

Geographic distribution: Oceania; New Zealand.

New Zealand references: Timmermann (1969c: 200); Nelson (1969); Watt (1971); Wise (1977: 63); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Palma (1996b: 212); Price *et al.* (2003: 223).

Remarks: In agreement with Timmermann (1969c: 198) and contrary to Price *et al.* (2003: 223), I regard this louse taxon as a subspecies of *Quadriceps hopkinsi* Timmermann, 1952 (see below).

#### ***Quadriceps hopkinsi hopkinsi* Timmermann, 1952**

- Quadriceps hopkinsi* Timmermann, 1952a: 74, figs 1–3.  
*Quadriceps hopkinsi* Timmermann, 1952; Hopkins & Clay 1953: 443.  
*Anousceps hopkinsi* (Timmermann, 1952); Złotorzycka 1967: 730.  
*Quadriceps hopkinsi hopkinsi* Timmermann, 1952; Timmermann 1969c: 198, fig. 4a.  
*Quadriceps hopkinsi hopkinsi* Timmermann, 1952; Watt 1971: 238, 243.  
*Quadriceps hopkinsi hopkinsi* Timmermann, 1952; Pilgrim & Palma 1982: 23.  
*Quadriceps hopkinsi* Timmermann, 1952; Price *et al.* 2003: 224.  
*Quadriceps hopkinsi hopkinsi* Timmermann, 1952; Murray *et al.* 2006a: 1965.  
*Quadriceps h. hopkinsi* Timmermann, 1952; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Anous minutus melanogenys* G.R. Gray, 1846.

New Zealand host: *Anous minutus minutus* Boie, 1844.

Other hosts: *Anous tenuirostris* (Temminck, 1823); *Anous minutus atlanticus* (Mathews, 1912).

New Zealand locality: KE.

Geographic distribution: Australasia; Oceania; Central & South America.

New Zealand references: Watt (1971); Wise (1977: 63); Pilgrim & Palma (1982); Palma (1994b: 268); Murray *et al.* (2006a); Palma (2010).

Other significant references: Złotorzycka (1967); Timmermann (1969c); Amerson & Emerson (1971: 19, 25); Ward & Downey (1973: 394); Benoit (1976: 233); Moyer & Wagenbach (1995: 1073); Palma (1996b: 211); Price *et al.* (2003: 224); Silva *et al.* (2014: 942).

Remarks: In agreement with Timmermann (1969c: 198) and contrary to Price *et al.* (2003: 224), I regard this louse taxon as a subspecies (see above).

#### ***Quadriceps normifer alpha* (Kellogg, 1914)**

*Nirmus triangulatus* Nitzsch, var. *alpha* Kellogg, 1914: 84.

*Degeeriella alpha* Kellogg, 1914 [sic]; Harrison 1916: 107.

*Koeniginirmus normifer alpha* Kellogg, 1914 [sic]; Eichler 1951b: 127.

*Quadriceps alpha* (Kellogg, 1914); Hopkins & Clay 1952: 308.

*Quadriceps alpha* (Kellogg, 1914); Clay & Moreby 1967: 164, 169, fig. 93.

*Koeniginirmus (Laminonirmus) normifer alpha* (Kellogg, 1914); Złotorzycka 1967: 759.

*Quadriceps normifer alpha* (Kellogg, 1914); Emerson 1972a: 141.

*Quadriceps alpha* (Kellogg, 1914); Pilgrim & Palma 1982: 22.

*Quadriceps alpha* (Kellogg, 1914); Price *et al.* 2003: 223.

*Quadriceps alpha* (Kellogg, 1914); Murray *et al.* 2006a: 1965.

*Quadriceps normifer alpha* (Kellogg, 1914); Palma 2010: 408.

*Quadriceps normifer alpha* (Kellogg, 1914); Palma 2015b: 153, figs 1–3.

Lectotype ♂ in EMEC (Palma 2015b: 151, figs 1, 3).

Type host: *Catharacta chilensis* (Bonaparte, 1857) (see Palma 2015b: 151).

New Zealand host: *Catharacta maccormicki* (Saunders, 1893).

Other host: *Catharacta antarctica antarctica* (Lesson, 1831) (not confirmed, see Palma 2015b: 154).

New Zealand localities: RO.

Geographic distribution: South America; Subantarctic Islands; Antarctica; southern Atlantic, Indian and Pacific Oceans.

New Zealand references: Clay & Moreby (1967); Złotorzycka (1967); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010); Palma (2015b: 153).

Other significant references: Eichler (1951b); Emerson (1972a); Cohen *et al.* (1997: 186); Price *et al.* (2003).

Remarks: In agreement with Eichler (1951b: 127) and Emerson (1972a: 141), and contrary to Pilgrim & Palma (1982: 22) and Price *et al.* (2003: 223), I regard this louse taxon as a subspecies of *Quadriceps normifer* (Grube, 1851). Palma (2015b) summarised the taxonomic history and host records of *Quadriceps normifer alpha*, clarified the identity of the type host, and designated a lectotype.

#### ***Quadriceps normifer normifer* (Grube, 1851)**

*Nirmus normifer* Grube, 1851: 478, pl. 31: fig. 8.

*Degeeriella normifer* Grube, 1851 [sic]; Harrison 1916: 119.

*Koeniginirmus normifer* (Grube, 1851); Timmermann 1949b: 88, fig. 3.

*Koeniginirmus normifer normifer* Grube, 1851 [sic]; Eichler 1951b: 126.

*Quadriceps normifer* (Grube, 1851); Hopkins & Clay 1952: 313.

*Koeniginirmus (Laminonirmus) normifer normifer* (Grube, 1851); Złotorzycka 1967: 760, pl. 15: fig. 1.

*Quadriceps normifer normifer* (Grube, 1851); Emerson 1972a: 141.

*Quadriceps normifer* (Grube, 1851); Pilgrim & Palma 1982: 22.

*Quadriceps normifer normifer*; Crossland 1993: 305.

*Quadriceps normifer* (Grube, 1851); Price *et al.* 2003: 225.

*Quadriceps normifer normifer* (Grube, 1851); Murray *et al.* 2006a: 1965.

*Quadriceps n. normifer* (Grube, 1851); Palma 2010: 408.

Syntypes ♂♀, repository unknown.

Type host: *Stercorarius parasiticus* (Linnaeus, 1758).

New Zealand host: *Stercorarius parasiticus* (Linnaeus, 1758).

Other hosts: None.

New Zealand localities: HB, WN, KA, MC.

Geographic distribution: Arctic Region; South America; South Africa; Atlantic, Indian and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Crossland (1993); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1949b); Eichler (1951b); Złotorzycka (1967); Hackman & Nyholm (1968: 79); Emerson (1972a); Cohen *et al.* (1997: 186); Price *et al.* (2003); Palma & Jensen (2005: 57, 64).

Remarks: In agreement with Eichler (1951b: 126) and Emerson (1972a: 141), and contrary to Pilgrim & Palma (1982: 22) and Price *et al.* (2003: 225), I regard this louse taxon as a subspecies.

#### ***Quadriceps normifer parvopallidus* (Eichler, 1951)**

*Nirmus triangulatus* variety Piaget, 1880: 202.

*Koeniginirmus normifer parvopallidus* Eichler, 1951b: 127. *Nomen novum* for *Nirmus triangulatus* variety Piaget, 1880: 202.

*Quadriceps parvopallidus* (Eichler, 1951); Hopkins & Clay 1953: 444.

*Koeniginirmus (Laminonirmus) normifer parvopallidus* Eichler, 1951; Złotorzycka 1967: 760.

*Quadriceps normifer parvopallidus* (Eichler, 1951); Emerson 1972a: 142.

*Quadriceps* sp.; Melville 1985: 67.

*Quadriceps normifer parvopallidus* (Eichler, 1951); Palma 1999: 380.

*Quadriceps parvopallidus* (Eichler, 1951); Price *et al.* 2003: 226.

*Quadriceps normifer parvopallidus* (Eichler, 1951); Murray *et al.* 2006a: 1965.

*Quadriceps n. parvopallidus* (Eichler, 1951); Palma 2010: 408.

Syntypes ♂♀ in NHML (Thompson 1937–1939: 421).

Type host: *Stercorarius longicaudus* Vieillot, 1819.

New Zealand host: *Stercorarius longicaudus* Vieillot, 1819.

Other hosts: None.

New Zealand localities: ND, WO, WI, WN.

Geographic distribution: Cold and temperate regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Melville (1985); Palma (1999); Murray *et al.* (2006a); Palma (2010).

Other significant references: Złotorzycka (1967); Emerson (1972a); Cohen *et al.* (1997: 186); Price *et al.* (2003); Palma & Jensen (2005: 57, 64).

Remarks: In agreement with Eichler (1951b: 127) and Emerson (1972a: 142), and contrary to Price *et al.* (2003: 226), I regard this louse taxon as a subspecies of *Quadriceps normifer*.

#### ***Quadriceps novaeseelandiae* Timmermann, 1953**

*Quadriceps novaeseelandiae* Timmermann, 1953b: 185, fig. 1.

*Quadriceps novaeseelandiae* Timmermann, 1953; Hopkins & Clay 1955: 187.

*Quadriceps novaeseelandiae* Timmermann, 1953; Wise 1977: 63.

*Quadriceps novaeseelandiae* Timmermann, 1953; Martens 1980: 351, figs 3–4.

*Quadriceps novaeseelandiae* Timmermann, 1953; Martens & Palma 1981: 83.

*Quadriceps novaeseelandiae* Timmermann, 1953; Pilgrim & Palma 1982: 20.

*Quadriceps novaeseelandiae* Timmermann, 1953; Murray *et al.* 1993: 962.

*Quadriceps novaeseelandiae* Timmermann, 1953; Palma 2010: 407.

Holotype ♂ in NHML.

Type host: *Thinornis novaeseelandiae* (J.F. Gmelin, 1789).

New Zealand hosts: *Charadrius bicinctus exilis* Falla, 1978; *Thinornis novaeseelandiae* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: CH, AU.

Geographic distribution: New Zealand.

New Zealand references: Timmermann (1953b); Pilgrim (1970: 74); Wise (1977); Martens (1980); Martens & Palma (1981); Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010); Buckley *et al.* (2012, App. 2).

Other significant reference: Price *et al.* (2003: 225).

Remarks: *Quadriceps novaeseelandiae* is an endemic and “at risk” species (Buckley *et al.* 2012), with an unusual host distribution on two species of New Zealand endemic plovers (Martens & Palma 1981: 84).

### ***Quadriceps nycthemerus* (Burmeister, 1838)**

*Nirmus nycthemerus* Burmeister, 1838a: 428.

*Degeeriella nycthemera* Nitzsch in Burmeister, 1838 [sic]; Harrison 1916: 119.

*Koeniginirmus nycthemerus nycthemerus* Nitzsch in Burmeister, 1838 [sic]; Eichler 1951b: 127. Unjustified emendation.

*Quadriceps nycthemerus* (Burmeister, 1838); Timmermann 1952a: 78, fig. 6 left.

*Quadriceps nycthemerus* (Burmeister, 1838); Hopkins & Clay 1952: 313. Unjustified emendation.

*Quadriceps nycthemerus* (Burmeister, 1838); Timmermann 1957a: 71, pl. 8: figs a,b.

*Koeniginirmus (Laminonirmus) nycthemerus nycthemerus* (Burmeister, 1838); Złotorzycka 1967: 762, pl. 17: fig. 4.

Unjustified emendation.

*Quadriceps nycthemerus* (Burmeister, 1838); Pilgrim & Palma 1982: 23. Unjustified emendation.

*Quadriceps nycthemerus* (Burmeister, 1838); Price *et al.* 2003: 225.

*Quadriceps nycthemerus* (Burmeister, 1838); Murray *et al.* 2006a: 1965. Unjustified emendation.

*Quadriceps nycthemerus* (Burmeister, 1838); Palma 2010: 408. Unjustified emendation.

Neotype ♂ in NHML (Timmermann 1952a: 78).

Type host: *Sternula albifrons albifrons* (Pallas, 1764).

New Zealand host: *Sternula albifrons sinensis* (J.F. Gmelin, 1789).

Other host: *Sterna nereis* Gould, 1843.

New Zealand locality: ND.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Séguy (1944: 289, fig. 439); Eichler (1951b); Timmermann (1952a); Timmermann (1952d: 1033); Timmermann (1957a); Złotorzycka (1967); Palma (1996b: 213); Price *et al.* (2003).

Remarks: I agree with Timmermann (1952a: 78) and Price *et al.* (2003: 225) in that there is no valid reason to change Burmeister’s (1838) original spelling of the species epithet “*nycthemerus*”, as done by Hopkins & Clay (1952: 313) and several other authors who followed them (see synonymy above).

### ***Quadriceps ornatus fuscolaminulatus* (Enderlein, 1908)**

*Ricinus fuscolaminulatus* Enderlein, 1908: 447, figs 193, 195.

*Degeeriella fuscolaminulata* Enderlein, 1908 [sic]; Harrison 1916: 113.

*Koeniginirmus ornatus fuscolaminulatus* Enderlein, 1908 [sic]; Eichler 1951b: 130.

*Quadriceps ornatus fuscolaminulatus* (Enderlein, 1908); Timmermann 1952b: 218.

*Koeniginirmus fuscolaminulatus* (Enderlein, 1908); Séguy 1953: 580, figs 39–40.

*Quadriceps fuscolaminulatus* (Enderlein, 1908); Clay 1964a: 232.

*Koeniginirmus (Laminonirmus) ornatus fuscolaminulatus* (Enderlein, 1908); Złotorzycka 1967: 764.

*Quadriceps ornatus fuscolaminulatus* (Enderlein, 1908); Wise 1977: 64.

*Quadriceps ornatus fuscolaminulatus* (Enderlein, 1908); Pilgrim & Palma 1982: 22.

*Quadriceps ornatus fuscolaminulatus* (Enderlein, 1908); Murray *et al.* 2006a: 1965.

*Quadriceps ornatus fuscolaminulatus* (Enderlein, 1908); Palma 2010: 408.

Holotype ♀ in ZMHU (Göllner-Scheiding 1973: 35).

Type host: *Larus dominicanus* Lichtenstein, 1823

New Zealand host: *Larus dominicanus dominicanus* Lichtenstein, 1823

Other hosts: None.

New Zealand localities: CA, Macquarie Island.

Geographic distribution: Southern Oceans.

New Zealand references: Clay (1964a); Gressitt (1964: 539); Clay & Moreby (1970: 220); Gressitt (1970: 328); Wise (1977); Pilgrim & Palma (1982); Palma (1996b: 213); Palma & Horning (2002: 12, 17); Murray *et al.* (2006a); Palma (2010).

Other significant references: Eichler (1951b); Timmermann (1952b,c); Séguy (1953); Clay & Moreby (1967: 164, 169, fig. 91); Timmermann (1971: 143, fig. 1); Price *et al.* (2003: 226); Yamagishi *et al.* (2014: 384).

Remarks: Within the New Zealand Subregion, *Quadriceps ornatus fuscolaminulatus* is the only species/subspecies of *Quadriceps* parasitising the Subantarctic populations of *Larus dominicanus* (also see below under *Quadriceps punctatus sublingulatus*).

#### ***Quadriceps punctatus lingulatus* (Waterston, 1914)**

*Nirmus punctatus lingulatus* Waterston, 1914: 285.

*Degeeriella lingulata* Waterston, 1914 [sic]; Harrison 1916: 116.

*Koeniginirmus lingulatus* Waterston, 1914 [sic]; Eichler 1951b: 126.

*Quadriceps lingulatus* (Waterston, 1914); Hopkins & Clay 1952: 313.

*Quadriceps punctatus lingulatus* (Waterston, 1914); Timmermann 1952b: 214, fig. 2c.

*Quadriceps lingulatus* (Waterston, 1914); Clay 1964a: 232.

*Koeniginirmus (Koeniginirmus) lingulatus lingulatus* (Waterston, 1914); Złotorzycka 1967: 750.

*Quadriceps lingulatus* (Waterston, 1914); Wise 1977: 63.

*Quadriceps punctatus* (Burmeister, 1838a) *s. l.*; Horning *et al.* 1980: 7, 11.

*Quadriceps punctatus* (Burmeister, 1838) *s. l.*; Pilgrim & Palma 1982: 22. In part.

*Quadriceps punctatus lingulatus* (Waterston, 1914); Price *et al.* 2003: 227.

*Quadriceps punctatus lingulatus* (Waterston, 1914); Murray *et al.* 2006a: 1965.

*Quadriceps punctatus* (Burmeister, 1838) *sensu lato*; Murray *et al.* 2006a: 1965. In part.

*Quadriceps punctatus* (Burmeister, 1838); Palma 2010: 408. In part.

Syntypes ♂♀ in SAMS (Palma 1996b: 213).

Type host: *Larus hartlaubii* Bruch, 1853.

New Zealand hosts: *Larus novaehollandiae scopulinus* J.R. Forster, 1843; *Larus bulleri* Hutton, 1871.

Other hosts: *Larus belcheri* Vigors, 1829; *Larus modestus* Tschudi, 1843; *Larus novaehollandiae novaehollandiae* Stephens, 1826; *Larus pipixcan* Wagler, 1831.

New Zealand localities: AK, HB, WN, SD, MB, NC, MC, SC, WD, SI, SN, AU.

Geographic distribution: Americas; Australasia.

New Zealand references: Clay (1964a); Gressitt (1964: 539); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Galloway (2005: 17); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1952b,c); Timmermann (1971: 143, fig. 1); Palma (1995: 220); Palma (1996b: 213); Price *et al.* (2003); Palma & Peck (2013: 53).

Remarks: Price *et al.* (2003: 227, 291) incorrectly listed *Larus bulleri* as a host of *Quadriceps punctatus punctatus* (Burmeister, 1838a).

#### ***Quadriceps punctatus sublingulatus* Timmermann, 1952**

#### **New Record**

*Quadriceps punctatus sublingulatus* Timmermann, 1952b: 215.

*Quadriceps sublingulatus* Timmermann, 1952; Hopkins & Clay 1953: 444.

*Quadriceps punctatus sublingulatus* Timmermann, 1952; Clay & Moreby 1967: 164, 169, fig. 90.

*Koeniginirmus (Koeniginirmus) lingulatus sublingulatus* (Timmermann, 1952); Złotorzycka 1967: 750.

*Quadriceps punctatus* (Burmeister, 1838a) *s. l.*; Pilgrim & Palma 1982: 22. In part.

*Quadriceps punctatus sublingulatus* Timmermann, 1952; Price *et al.* 2003: 227.

*Quadriceps punctatus* (Burmeister, 1838) *sensu lato*; Murray *et al.* 2006a: 1965. In part.

*Quadriceps punctatus* (Burmeister, 1838); Palma 2010: 408. In part.

Holotype ♂ in NHML.

Type host: *Larus delawarensis* Ord, 1815.

New Zealand host: *Larus dominicanus dominicanus* Lichtenstein, 1823.

Other hosts: *Larus californicus* Lawrence, 1854; *Larus occidentalis* Audubon, 1839; *Larus cirrocephalus* Vieillot, 1818; *Larus philadelphia* (Ord, 1815).

New Zealand localities: AK, WN, NC, MC, SC, WD.

Geographic distribution: Americas; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1952b,c); Clay & Moreby (1967); Złotorzycka (1967); Timmermann (1971: 143, fig. 1); Forrester *et al.* (1995: 29); Price *et al.* (2003: 227); González-Acuña *et al.* (2011: 300); Yamagishi *et al.* (2014: 384).

Material examined and repository: 57♂, 60♀ (15 samples, MONZ).

Remarks: This is the first record of *Quadriceps punctatus sublingulatus* for New Zealand, because the New Zealand references cited above reported this louse as “*Quadriceps punctatus* (Burmeister, 1838) *sensu lato*” or as “*Quadriceps punctatus* (Burmeister, 1838)”. Within the New Zealand Subregion, *Quadriceps punctatus sublingulatus* is the only species/subspecies of *Quadriceps* parasitising the mainland populations of *Larus dominicanus* (also see above under *Quadriceps ornatus fuscoluminulatus*).

### ***Quadriceps renschi* Timmermann, 1954**

*Quadriceps renschi* Timmermann, 1954e: 206, fig. 13.

*Quadriceps renschi* Timmermann, 1954; Hopkins & Clay 1955: 187.

*Quadriceps renschi* Timmermann, 1954; Wise 1977: 64.

*Quadriceps renschi* Timmermann, 1954; Pilgrim & Palma 1982: 20.

*Quadriceps renschi* Timmermann, 1954; Murray *et al.* 1993: 962.

*Quadriceps renschi* Timmermann, 1954; Palma 2010: 408.

Holotype ♂ in NHML.

Type host: *Vanellus miles novaehollandiae* Stephens, 1819.

New Zealand host: *Vanellus miles novaehollandiae* Stephens, 1819.

Other hosts: None.

New Zealand localities: NN, BR, MC, SC, DN, SL.

Geographic distribution: Australasia.

New Zealand references: Pilgrim (1970: 75); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 221); Palma (2010).

Other significant references: Złotorzycka (1967: 722); Green & Palma (1991: 18, 32); Palma (1996b: 214); Price *et al.* (2003: 227).

Remarks: *Quadriceps renschi* is a frequently collected “wing” louse species from spur-winged plovers.

### ***Quadriceps ridgwayi* (Kellogg, 1906)**

*Nirmus ridgwayi* Kellogg, 1906: 317.

*Degeeriella ridgwayi* Kellogg, 1906 [sic]; Harrison 1916: 122.

*Quadriceps ridgwayi* (Kellogg, 1906); Hopkins & Clay 1952: 316.

*Quadriceps ridgwayi*; Baker 1974: 20.

*Quadriceps ridgwayi* (Kellogg, 1906); Clay 1976b: 540, 546.

*Quadriceps ridgwayi* (Kellogg, 1906); Clay 1981b: 936, 938.

*Quadriceps ridgwayi* (Kellogg, 1906); Pilgrim & Palma 1982: 19.

*Quadriceps ridgwayi* (Kellogg, 1906); Murray *et al.* 1993: 961.

*Quadriceps ridgwayi* (Kellogg, 1906); Palma 2010: 408.

Lectotype ♂ in EMEC (Palma 1996b: 214).

Type host: *Haematopus palliatus galapagensis* Ridgway, 1886.

New Zealand host: *Haematopus unicolor* J.R. Forster, 1844.

Other hosts: *Haematopus ater* Vieillot & Oudart, 1825; *Haematopus longirostris* Vieillot, 1817; *Haematopus fuliginosus* Gould, 1845.

New Zealand localities: AK, WD, SL.

Geographic distribution: North and South America; Australasia; Oceania.

New Zealand references: Baker (1974); Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Złotorzycka (1967: 748); Timmermann (1971: 158); Clay (1976b; 1981b); Green & Palma (1991: 18, 32); Palma (1996b: 214); Price *et al.* (2003: 227); Palma & Peck (2013: 55).

Remarks: *Quadriceps ridgwayi* has a reduced host distribution (Clay 1981: 938, table 1), if compared with the other species of *Quadriceps* parasitic on oystercatchers, i.e. *Q. auratus* (see above).

#### ***Quadriceps sellatus houri* Hopkins, 1949**

*Quadriceps houri* Hopkins, 1949d: 52, pl. 3: figs 9–10.

*Quadriceps houri* Hopkins, 1949; Hopkins 1951a: 373.

*Quadriceps sellatus houri* Hopkins, 1949; Timmermann 1952a: 79.

*Quadriceps houri* Hopkins, 1949; Hopkins & Clay 1952: 312.

*Quadriceps houri* Hopkins, 1949; Timmermann 1957a: 71, pl. 7: figs a,b.

*Koeniginirmus (Laminonirmus) houri* (Hopkins, 1949); Złotorzycka 1967: 758.

*Quadriceps houri* Hopkins, 1949; Clay & Moreby 1967: 164, 169, fig. 146.

*Quadriceps houri* Hopkins, 1949; Pilgrim & Palma 1982: 23.

*Quadriceps houri* Hopkins, 1949; Palma 1999: 380.

*Quadriceps houri* Hopkins, 1949; Murray *et al.* 2006a: 1965.

*Quadriceps houri* Hopkins, 1949; Palma 2010: 408.

Holotype ♀ in NHML.

Type host: *Sterna paradisaea* Pontoppidan, 1763.

New Zealand hosts: *Sterna paradisaea* Pontoppidan, 1763; *Sterna vittata bethunei* Buller, 1896.

Other host: *Sterna vittata vittata* J.F. Gmelin, 1789.

New Zealand localities: WN, SL, SI, Macquarie Island.

Geographic distribution: Arctic and Subantarctic regions; Atlantic and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Palma (1996b: 212); Palma (1999); Palma & Horning (2002: 12, 18); Murray *et al.* (2006a); Palma (2010).

Other significant references: Hopkins (1951a); Timmermann (1952a; 1957a); Clay & Moreby (1967); Złotorzycka (1967); Price *et al.* (2003: 224); Palma & Jensen (2005: 57, 65); Hänel & Palma (2007: 113, 127, 131).

Remarks: In agreement with Timmermann (1952a: 79), and contrary to Pilgrim & Palma (1982: 23), Price *et al.* (2003: 224) and Palma (2010: 408), I regard this louse taxon as a subspecies of *Quadriceps sellatus*. The population of *Sterna vittata bethunei* from Macquarie Island is parasitised by *Quadriceps sellatus houri*, while the population of the same host from the Snares Islands is parasitised by *Quadriceps sellatus sellatus* (see below).

#### ***Quadriceps sellatus sellatus* (Burmeister, 1838)**

*Nirmus sellatus* Burmeister, 1838a: 428.

*Degeeriella sellata* Burmeister, 1838 [sic]; Harrison 1916: 122.

*Koeniginirmus sellatus* (Burmeister, 1838); Timmermann 1949b: 87, fig. 2.

*Quadriceps sellatus* (Burmeister, 1838); Hopkins 1949d: 52, pl. 3: figs 11–12.

*Koeniginirmus sellatus sellatus* Burmeister, 1838 [sic]; Eichler 1951b: 134.

*Quadriceps sellatus sellatus* (Burmeister, 1838); Timmermann 1952a: 80.

*Koeniginirmus sellatus* (Burmeister, 1839) [sic]; Séguy 1953: 581, fig. 41.

*Quadriceps sellatus* (Burmeister, 1838); Timmermann 1957a: 71, pl. 7: figs c,d.

*Koeniginirmus (Laminonirmus) sellatus sellatus* (Burmeister, 1838); Złotorzycka 1967: 767.

*Quadriceps sellatus* (Burmeister, 1838); Clay & Moreby 1967: 164, 169, fig. 145.

*Quadriceps sellatus* (Burmeister, 1838) *s. l.*; Horning *et al.* 1980: 7, 11.

*Quadriceps sellatus* (Burmeister, 1838) *s. l.*; Pilgrim & Palma 1982: 23.

*Quadriceps sellatus* (Burmeister, 1838); Price *et al.* 2003: 227.

*Quadriceps sellatus* (Burmeister, 1838); Murray *et al.* 2006a: 1965.

*Quadriceps sellatus* (Burmeister, 1838); Palma 2010: 408.

Neotype ♂ in NHML (Hopkins 1949d: 52; Palma 1996b: 214).

Type host: *Sterna hirundo hirundo* Linnaeus, 1758.

New Zealand hosts: *Sterna striata* J.F. Gmelin, 1789; *Sterna vittata bethunei* Buller, 1896.

Other hosts: *Sterna bergii* Lichtenstein, 1823; *Sterna bengalensis* Lesson, 1831.

New Zealand localities: AK, KA, NC, MC, SC, WD, CO, DN, SL, SN, AU.

Geographic distribution: Cosmopolitan.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Hopkins (1949d); Timmermann (1949b; 1952a; 1957a); Eichler (1951b); Séguy (1953); Złotorzycka (1967); Clay & Moreby (1967); Moreby (1976: 94); Green & Palma (1991: 18, 33); Forrester *et al.* (1995: 30); Palma (1996b: 214); Price *et al.* (2003).

Remarks: In agreement with Timmermann (1952a: 80), and contrary to Price *et al.* (2003: 227) and Palma (2010: 408), I regard this louse taxon as a subspecies. The population of *Sterna vittata bethunei* from Snares Islands is parasitised by *Quadraceps sellatus sellatus*, while the population of the same host from Macquarie Island is parasitised by *Quadraceps sellatus houri* (see above).

### ***Quadraceps semifissus* (Nitzsch [*in* Giebel], 1866)**

*Nirmus semifissus* Nitzsch [*in* Giebel], 1866: 372.

*Degeeriella semifissa* Nitzsch *in* Giebel, 1866 [sic]; Harrison 1916: 123.

*Quadraceps semifissa mexicana* [sic] Carriker, 1944: 99, pl. 5: figs 8–9.

*Quadraceps mexicanus* Carriker, 1944; Hopkins & Clay 1952: 313. Emendation.

*Quadraceps semifissus* (Nitzsch [*in* Giebel], 1866); Hopkins & Clay 1952: 316.

*Proneptis semifissa* (Nitzsch [*in* Giebel], 1866); Timmermann 1953a: 333, fig. right.

*Proneptis semifissa* (Nitzsch [*in* Giebel], 1866); Timmermann 1954f: 166, figs 3–5.

*Proneptis semifissa mexicana* (Carriker, 1944); Złotorzycka 1967: 732.

*Proneptis semifissa semifissa* (Nitzsch [*in* Giebel], 1866); Złotorzycka 1967: 733, pl. 7: figs 1, 3, pl. 8: fig. 1.

*Quadraceps semifissus* (Nitzsch, 1866) [sic]; Pilgrim & Palma 1982: 21.

*Quadraceps semifissus mexicanus* Carriker, 1944; Hinojos & Canaris 1988: 328.

*Quadraceps semifissus* (Nitzsch, 1866) [sic]; Murray *et al.* 1993: 962.

*Quadraceps semifissus* (Nitzsch [*in* Giebel], 1866); Palma 1999: 380.

*Quadraceps semifissus* (Nitzsch [*in* Giebel], 1866); Palma 2010: 408.

Syntypes ♂♀, presumed lost. See Clay (1949a: 1), Palma & Pilgrim (1984: 150) and Palma (1996b: 215).

Type host: *Himantopus himantopus himantopus* (Linnaeus, 1758).

New Zealand hosts: *Himantopus himantopus leucocephalus* Gould, 1837; *Himantopus novaezelandiae* Gould, 1841.

Other hosts: *Himantopus himantopus mexicanus* (Stadius Müller, 1776); *Recurvirostra americana* J.F. Gmelin, 1789; *Recurvirostra andina* Philippi & Landbeck, 1861; *Recurvirostra avosetta* Linnaeus, 1758; *Recurvirostra novaehollandiae* Vieillot, 1816.

New Zealand localities: AK, BP, HB, WI, WN, NN, KA, NC, MC, SC, CO, DN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Palma (1999); Murray *et al.* (1993); Paterson *et al.* (1999: 221, 223); Palma (2010).

Other significant references: Timmermann (1953a; 1954f); Timmermann (1957a: 76, figs 48–50); Złotorzycka (1967); Timmermann (1971: 165, fig. 10); Hinojos & Canaris (1988); Palma (1995: 220); Forrester *et al.* (1995: 26); Palma (1996b: 215); Price *et al.* (2003: 227); Martín-Mateo (2009: 131); Palma & Peck (2013: 55).

Remarks: In addition to *Quadraceps semifissus*, the species of the host genus *Himantopus* are parasitised by *Quadraceps hemichrous* (see above).

### ***Quadraceps separatus* (Kellogg & Kuwana, 1902)**

*Nirmus separatus* Kellogg & Kuwana, 1902: 472, pl. 29: fig. 6.

*Nirmus gloriosus emarginatus* Kellogg & Chapman, 1902: 159.

*Degeeriella emarginata* Kellogg & Chapman, 1902 [sic]; Harrison 1916: 112.

*Degeeriella separata* Kellogg & Kuwana, 1902 [sic]; Harrison 1916: 123.

*Quadraceps separatus* (Kellogg & Kuwana, 1902); Hopkins & Clay 1952: 316.

*Quadraceps separatus* (Kellogg & Kuwana, 1902); Timmermann 1952a: 73, figs 1–2.

*Anousceps separatus* (Kellogg & Kuwana, 1902); Złotorzycka 1967: 731.



*Quadriceps separatus* (Kellogg & Kuwana, 1902); Palma 1994b: 268.

*Quadriceps separatus* (Kellogg & Kuwana, 1902); Palma 1999: 380.

*Quadriceps separatus* (Kellogg & Kuwana, 1902); Murray *et al.* 2006a: 1965.

*Quadriceps separatus* (Kellogg & Kuwana, 1902); Palma 2010: 408.

Lectotype ♂ in EMEC (Palma 1996b: 215).

Type host: “*Geospiza conirostris*”, in error (see Palma 1994b: 270).

New Zealand host: *Anous stolidus pileatus* (Scopoli, 1786).

Other host: *Anous stolidus galapagensis* Sharpe, 1879.

New Zealand locality: KE.

Geographic distribution: Tropical and subtropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Palma (1994b); Palma (1999); Murray *et al.* (2006a); Palma (2010).

Other significant references: Ferris (1932a: 69, figs 19a,b,c,d,e); Timmermann (1952a); Timmermann (1957a: 70, pl. 9: figs c,d); Złotorzycka (1967); Amerson & Emerson (1971: 18, 25); Ward & Downey (1973: 395); Moreby (1976: 94); Forrester *et al.* (1995: 32); Palma (1994b; 1996b: 215); Price *et al.* (2003: 227); Palma & Peck (2013: 55); Silva *et al.* (2014: 942).

Remarks: Within the New Zealand Subregion, *Anous stolidus pileatus* breeds on Norfolk island and, more recently, on Curtis Island at the Kermadec Islands (Checklist Committee 2010: 231), with only one record of *Q. separatus* from this country.

#### ***Quadriceps strepsilaris* (Denny, 1842)**

*Philopterus (Nirmus) strepsilaris* Denny, 1842: 52, 135, pl. 11: fig. 4.

*Nirmus lepidus* Kellogg & Kuwana, 1902: 473, pl. 29: fig. 7.

*Degeeriella lepida* Kellogg & Kuwana, 1902 [sic]; Harrison 1916: 116.

*Degeeriella strepsilaris* Denny, 1842 [sic]; Harrison 1916: 124.

*Quadriceps strepsilaris* (Denny, 1842); Timmermann 1950: 2, fig. 2.

*Quadriceps lepidus* (Kellogg & Kuwana, 1902); Hopkins & Clay 1952: 312.

*Quadriceps strepsilaris* (Denny, 1842); Hopkins & Clay 1952: 317.

*Quadriceps strepsilaris* (Denny, 1842); Złotorzycka 1967: 724, pl. 1: fig. 3, pl. 2: fig. 2, pl. 3: fig. 4.

*Quadriceps strepsilaris* (Denny, 1842); Pilgrim & Palma 1982: 21.

*Quadriceps strepsilaris* (Denny, 1842); Palma 1994b: 269.

*Cummingsiella strepsilaris* (Denny, 1842); Butler & O'Connor 1994: 454.

*Quadriceps strepsilaris* (Denny, 1842); Price *et al.* 2003: 228.

*Quadriceps strepsilaris* (Denny, 1842); Murray *et al.* 2006a: 1964.

*Quadriceps strepsilaris* (Denny, 1842); Palma 2010: 408.

Holotype ♀ in NHML (Thompson 1937a: 80).

Type host: *Arenaria interpres* (Linnaeus, 1758).

New Zealand host: *Arenaria interpres* (Linnaeus, 1758).

Other host: *Arenaria melanocephala* (Vigors, 1829).

New Zealand localities: AK, MC, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Palma (1994b); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1950); Emerson & Ward (1958: 58); Złotorzycka (1967); Hackman & Nyholm (1968: 79); Amerson & Emerson (1971: 15, 20); Butler & O'Connor (1994); Hunter & Colwell (1994: 402); Forrester *et al.* (1995: 26); Price *et al.* (2003); Palma & Jensen (2005: 58, 64); Palma & Peck (2013: 56).

Remarks: The ruddy turnstone is a very common summer visitor to New Zealand (Checklist Committee 2010: 207).

#### ***Quadriceps* species**

*Quadriceps* sp.; Pilgrim & Palma 1982: 22.

*Quadriceps* sp.; Murray *et al.* 2006a: 1965.

New Zealand host: *Chlidonias albobristatus* (G.R. Gray, 1845).

New Zealand locality: MC.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a).

Other significant references: Timmermann (1952a-d); Price *et al.* (2003: 22).

Remarks: The two available samples of *Quadriceps* from *Chlidonias albostrigatus* could not be identified to species because each contains one female only (voucher specimens in MONZ).

### Genus *Rallicola* Johnston & Harrison, 1911

*Rallicola* Johnston & Harrison, 1911. *Proc. Linn. Soc. New South Wales* 36(2): 324. Type species: *Oncophorus attenuatus* N. [sic] = *Rallicola (Rallicola) ortygometrae* (Schrank, 1781) (by original designation).

### Subgenus *Aptericola* Harrison, 1915

*Aptericola* Harrison, 1915b. *Parasitology* 8: 90. Type species: *Rallicola (Aptericola) gadowi* Harrison, 1915b (by original designation).

### *Rallicola (Aptericola) gadowi* Harrison, 1915

*Rallicola (Aptericola) gadowi* Harrison, 1915b: 90, figs 1–2.

*Aptericola gadowi* (Harrison, 1915); Tillyard 1926: 134, fig. O2.

*Rallicola gadowi* Harrison, 1915; Hopkins & Clay 1952: 319.

*Rallicola gadowi* Harrison, 1915; Clay 1953a: 568, figs 9–11, 45–46.

*Aptericola gadowi*; Miller 1971: 132.

*Rallicola gadowi* Harrison, 1915; Clay 1972: 71, figs 1–3, 5–7, 10, 13. In part.

*Rallicola gadowi* Harrison, 1915; Wise 1977: 64. In part.

*Rallicola (Aptericola) gadowi* Harrison, 1915; Pilgrim & Palma 1982: 3.

*Rallicola (Aptericola) gadowi* Harrison, 1915; Palma 2010: 409.

Lectotype ♀ in NHML (Clay 1972: 72).

Type host: *Apteryx australis australis* Shaw, 1813.

New Zealand host: *Apteryx australis australis* Shaw, 1813.

Other hosts: None.

New Zealand localities: WD, FD.

Geographic distribution: New Zealand.

New Zealand references: Harrison (1915); Tillyard (1926); Clay (1953a); Miller (1971); Clay (1972); Reid & Williams (1975: 324); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1367); Palma (1991a: 317); Mey (1994: 21, figs 7–8); Baker *et al.* (1995: 8256); Burbidge *et al.* (2003: 174); Sales (2005: 15); Heath (2010: 151); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Hopkins & Clay (1952); Price *et al.* (2003: 229).

Remarks: *Rallicola (Aptericola) gadowi* is an endemic and “at risk” species (Buckley *et al.* 2012). Samples published by Clay (1972: 73) as originating from “*Apteryx australis mantelli*” are actually from *Apteryx australis australis* (see Palma 1991a: 317). Also, see below under Remarks for *Rallicola (Aptericola) gadowi* Harrison, 1915 *sensu lato*.

### *Rallicola (Aptericola) gadowi* Harrison, 1915 *sensu lato*

*Rallicola (Aptericola) novae-zealandiae* Harrison, 1915b: 92, figs 3–5.

*Rallicola novaezealandiae* Harrison, 1915; Clay 1953a: 579, figs 47–48.

*Rallicola gadowi* Harrison, 1915; Clay 1972: 71, figs 1–3, 5–7, 10, 13. In part.

*Rallicola gadowi* Harrison, 1915; Wise 1977: 64. In part.

*Rallicola (Aptericola) gadowi* Harrison, 1915 *s. l.*; Pilgrim & Palma 1982: 3.

*Aptericola gadowi* ssp.; Mey 1994: 21: figs 7–8.

*Rallicola (Aptericola) gadowi* Harrison, 1915 *sensu lato*; Palma & Price 2004: 71.

Lectotype of *Rallicola (Aptericola) novaezealandiae* Harrison, 1915: ♀ in NHML (Clay 1972: 72).

New Zealand hosts: *Apteryx australis lawryi* Rothschild, 1893; *Apteryx owenii* Gould, 1847; *Apteryx rowi* Tennyson *et al.* 2003.

Other hosts: None.

New Zealand localities: WN (Kapiti Island), WD, SI.

Geographic distribution: New Zealand.

New Zealand references: Harrison (1915); Clay (1953a); Clay (1972); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1367); Mey (1994); Palma & Price (2004); Heath (2010: 151).

Other significant references: Clay (1951b: 188, fig. 12); Hopkins & Clay (1952); Price *et al.* (2003: 229).

Remarks: The populations of *Rallicola (Aptericola) gadowi* from its four hosts show a remarkable degree of morphological variability, both among and within populations, and especially those from the subspecies of *Apteryx australis*, within which I have identified several male and female morphs. Clay (1972: 73) concluded that the best option was to consider them as one species under one name. However, the material she had for study was limited in number of samples and of specimens. I have examined over 2,000 specimens from all four hosts and from all the localities where kiwis occur today, and conclude that a molecular study is necessary to attempt to elucidate the complex morphology shown by these lice (see Palma & Price 2004: 71).

### ***Rallicola (Aptericola) gracilentus* Clay, 1953**

*Rallicola (Aptericola) gracilis* Harrison, 1915b: 93, fig. 6. Preoccupied by *Docophorus gracilis* Piaget, 1871: 120 = *Rallicola gracilis* (Piaget, 1871).

*Rallicola gracilentus* Clay, 1953a: 584, figs 43–44. *Nomen novum* for *Rallicola gracilis* Harrison, 1915.

*Rallicola gracilentus* Clay, 1953; Hopkins & Clay 1955: 187.

*Rallicola gracilentus* Clay, 1953; Clay 1972: 73, figs 8, 11, 14.

*Rallicola gracilentus* Clay, 1953; Wise 1977: 64.

*Rallicola (Aptericola) gracilentus* Clay, 1953; Pilgrim & Palma 1982: 3.

*Rallicola (A.) gracilentus* Clay, 1953; Palma 2010: 409.

Lectotype ♀ in NHML (Clay 1972: 74).

Type host: *Apteryx haastii* Potts, 1872.

New Zealand host: *Apteryx haastii* Potts, 1872.

Other hosts: None, but see Remarks, below.

New Zealand localities: NN, BR, WD.

Geographic distribution: South Island, New Zealand.

New Zealand references: Harrison (1915); Clay (1953a); Pilgrim (1970: 75); Clay (1972); Reid & Williams (1975: 324); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1367); Baker *et al.* (1995: 8256); Burbidge *et al.* (2003: 174); Green & Turner (2003a: 97); Sales (2005: 15); Heath (2010: 151); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant references: Hopkins & Clay (1955); Price *et al.* (2003: 229).

Remarks: *Rallicola (Aptericola) gracilentus* is an endemic and vulnerable species (Buckley *et al.* 2012), exclusively parasitic on great spotted kiwis. Samples of *Rallicola (Aptericola)* from some kiwis identified as possible hybrids between *Apteryx owenii* and *Apteryx haastii* collected in the South Island during the past 60 years (Shepherd *et al.* 2012: 7) are *R. (Aptericola) gracilentus*.

### ***Rallicola (Aptericola) pilgrimi* Clay, 1972**

*Rallicola pilgrimi* Clay, 1972: 74, figs 9, 12, 15.

*Rallicola pilgrimi* Clay, 1972; Wise 1977: 64.

*Rallicola (Aptericola) pilgrimi* Clay, 1972; Pilgrim & Palma 1982: 3.

*Rallicola (A.) pilgrimi* Clay, 1972; Palma 2010: 409.

Holotype ♀ in NHML.

Type host: *Apteryx owenii* Gould, 1847.

New Zealand host: *Apteryx owenii* Gould, 1847.

Other hosts: None.

New Zealand localities: WA, NN.

Geographic distribution: New Zealand.

New Zealand references: Clay (1972); Reid & Williams (1975: 324); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990: 1367); Baker *et al.* (1995: 8256); Burbidge *et al.* (2003: 174); Sales (2005: 15); Heath (2010: 151); Palma (2010); Buckley *et al.* (2012: 140); Rózsa & Vas (2015b: 109).

Other significant reference: Price *et al.* (2003: 230).

Remarks: Although there is a viable population of *Apteryx owenii* on Kapiti Island (Checklist Committee 2010: 22), *Rallicola (Aptericola) pilgrimi* has not been found there yet. All records of this louse are from mainland North and South Islands and, considering that *A. owenii* is almost certainly extinct on those islands (Checklist Committee 2010: 22), *Rallicola (Aptericola) pilgrimi* is also regarded as extinct (Buckley *et al.* 2012; Rózsa & Vas 2015b: 109).

### ***Rallicola (Aptericola) rodericki* Palma, 1991**

Figs 160–161

*Rallicola (Aptericola)* sp.; Pilgrim & Palma 1982: 3.

*Rallicola (Aptericola)* sp.; Murray *et al.* 1990: 1367.

*Rallicola (Aptericola) rodericki* Palma, 1991a: 314, figs 1–4.

*Rallicola (A.) rodericki* Palma, 1991; Palma 2010: 409.

Holotype ♂ in MONZ.

Type host: *Apteryx mantelli* Bartlett, 1852.

New Zealand host: *Apteryx mantelli* Bartlett, 1852.

Other hosts: None.

New Zealand localities: AK (Ponui Island), CL (Little Barrier Island).

Geographic distribution: Hauraki Gulf, New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1991a); Baker *et al.* (1995: 8257); Palma (1999: 375); Sales (2005: 15); Heath (2010: 151); Palma (2010) Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003: 231).

Remarks: *Rallicola (Aptericola) rodericki* is an endemic and “at risk” species (Buckley *et al.* 2012), exclusively parasitic on North Island brown kiwis. Despite a search for lice on over 100 dead and live specimens of *Apteryx mantelli* from all the mainland populations of North Island kiwi (Palma 1991a: 316), *Rallicola (Aptericola) rodericki* has not been found on mainland kiwis.

### **Subgenus *Huiacola* Mey, 1990**

*Huiacola* Mey, 1990. *Zool. Anz.* 224: 52. Type species: *Huiacola extinctus* Mey, 1990 = *Rallicola (Huiacola) extinctus* (Mey, 1990) (by original designation).

### ***Rallicola (Huiacola) extinctus* (Mey, 1990)**

Figs 162–163

*Rallicola* sp.; Pilgrim & Palma 1982: 28.

*Huiacola extinctus* Mey, 1990: 54, figs 2–13.

*Rallicola (Huiacola) extinctus* (Mey, 1990); Palma 1999: 382.

*Rallicola (Huiacola) extinctus* (Mey, 1990); Murray *et al.* 2006b: 1958.

*Rallicola (Huiacola) extinctus* (Mey, 1990); Palma 2010: 409.

Holotype ♂ in NHMR.

Type host: *Heteralocha acutirostris* (Gould, 1837).

New Zealand host: *Heteralocha acutirostris* (Gould, 1837).

Other hosts: None.

New Zealand localities: WA, WN.

Geographic distribution: North Island, New Zealand.

New Zealand references: Mason (1921: 359); Pilgrim & Palma (1982); Mey (1990); Palma (1999); Mey (2005: 213); Murray *et al.* (2006b); Palma (2010: 295); Buckley *et al.* (2012: 140); Mey (2014: 98, fig. right); Rózsa & Vas (2015b: 109).

Other significant reference: Price *et al.* (2003: 229).

Remarks: *Rallicola (Huiacola) extinctus* is an endemic and extinct species (Buckley *et al.* 2012; Rózsa & Vas 2015b: 109), because its only host is extinct (Checklist Committee 2010: 283).

### Subgenus *Rallicola* Johnston & Harrison, 1911

*Rallicola* Johnston & Harrison, 1911. *Proc. Linn. Soc. New South Wales* 36(2): 324. Type species: *Oncophorus attenuatus* N. [sic] = *Rallicola (Rallicola) ortygometae* (Schrank, 1781) (by original designation).

#### *Rallicola (Rallicola) fulicae* (Denny, 1842)

*Philopterus (Nirmus) fulicae* Denny, 1842: 50, 125, pl. 9: fig. 2.

*Rallicola fulicae* Denny, 1842 [sic]; Harrison 1916: 126 (as junior synonym of *Rallicola cuspidata* Scopoli).

*Rallicola fulicae* (Denny, 1842); Hopkins 1940: 425.

*Rallicola fulicae* (Denny, 1842); Hopkins & Clay 1952: 319.

*Rallicola fulicae* (Denny, 1842); Clay 1953a: 573, figs 18, 25.

*Rallicola fulicae* (Denny, 1842); Pilgrim & Palma 1982: 19.

*Rallicola fulicae* (Denny, 1842); Murray *et al.* 1993: 961.

*Rallicola (Rallicola) fulicae* (Denny, 1842); Price *et al.* 2003: 229.

*Rallicola (Rallicola) fulicae* (Denny, 1842); Palma 2010: 408.

Holotype ♂ in NHML (Palma 1996b: 216).

Type host: *Fulica atra* Linnaeus, 1758.

New Zealand host: *Fulica atra australis* Gould, 1845.

Other host: *Fulica cristata* J.F. Gmelin, 1789.

New Zealand localities: SD, MB, NC, MC, SC, CO, DN.

Geographic distribution: Africa; Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Hopkins (1940); Clay (1953a); Emerson (1955: 285, fig. 1); Tendeiro (1965b: 114); Green & Palma (1991: 18, 32); Palma (1996b: 216); Price *et al.* (2003); Palma & Jensen (2005: 58, 63); Adam (2007: 178); Martín-Mateo (2009: 109).

Remarks: Hopkins (1940: 421) gives a complete account of the early taxonomic confusion among several species of lice described from *Fulica atra*, including a clarification of the synonymy of *Rallicola (R.) fulicae*.

#### *Rallicola (Rallicola) harrisoni* Emerson, 1955

*Rallicola harrisoni* Emerson, 1955: 288, figs 29–30.

*Rallicola harrisoni* Emerson, 1955; Wise 1977: 64.

*Rallicola* sp.; Lowry *et al.* 1978: 139.

*Rallicola harrisoni* Emerson, 1955; Pilgrim & Palma 1982: 18.

*Rallicola harrisoni* Emerson, 1955 *s. l.*; Pilgrim & Palma 1982: 19.

*Rallicola harrisoni* Emerson, 1955; Murray *et al.* 1993: 961.

*Rallicola harrisoni* Emerson, 1955; Palma & Horning 2002: 12, 17.

*Rallicola (Rallicola) harrisoni* Emerson, 1955; Price *et al.* 2003: 229.

*Rallicola (R.) harrisoni* Emerson, 1955; Palma 2010: 408.

Holotype ♂, originally in the M.A. Carriker Collection, now deposited in USNM.

Type host: *Gallirallus australis australis* (Sparrman, 1786).

New Zealand hosts: *Gallirallus australis greyi* (Buller, 1888); *Gallirallus australis hectori* (Hutton, 1873); *Gallirallus australis scotti* (Ogilvie-Grant, 1905).

Other hosts: None.

New Zealand localities: HB, NN, BR, WD, CH, SI, Macquarie Island.

Geographic distribution: New Zealand.

New Zealand references: Emerson (1955); Pilgrim (1970: 75); Wise (1977: 64); Lowry *et al.* (1978); Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (1996b: 216); Paterson *et al.* (1999: 223); Palma & Horning (2002); Palma (2010).

Other significant reference: Price *et al.* (2003: 229).

Remarks: *Rallicola (Rallicola) harrisoni* is an endemic species, exclusively parasitic on wekas, and frequently collected.

Pilgrim & Palma (1982: 19) regarded the population of *R. harrisoni* from *Gallirallus australis scotti* as somewhat different from that of the type host, and qualified it as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted.

### ***Rallicola (Rallicola) lugens* (Giebel, 1874)**

*Nirmus lugens* Giebel, 1874: 170.

*Oncophorus fallax* Piaget, 1880: 220, pl. 18: fig. 6.

*Rallicola lugens* Giebel, 1874 [sic]; Harrison 1916: 127.

*Rallicola fallax* (Piaget, 1880); Hopkins & Clay 1952: 319.

*Rallicola lugens* (Giebel, 1874); Hopkins & Clay 1952: 320.

*Rallicola lugens* (Giebel, 1874); Clay 1953a: 570, figs 14, 20, 41.

*Rallicola lugens* (Giebel, 1874); Watt 1971: 238, 243.

*Rallicola lugens* (Giebel, 1874); Pilgrim & Palma 1982: 19.

*Rallicola lugens* (Giebel, 1874); Murray *et al.* 1993: 961.

*Rallicola (Rallicola) lugens* (Giebel, 1874); Price *et al.* 2003: 230.

*Rallicola (R.) lugens* (Giebel, 1874); Palma 2010: 408.

Syntypes ♂♀, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150). Syntypes of *Oncophorus fallax* in NHML (Palma 1996b: 217).

Type host: *Porphyrio poliocephalus* (Latham, 1802).

New Zealand host: *Porphyrio melanotus melanotus* Temminck, 1820.

Other hosts: *Porphyrio melanotus bellus* Gould, 1841; *Porphyrio porphyrio* (Linnaeus, 1758); *Porphyrio indicus* Horsfield, 1821.

New Zealand localities: AK, BP, TK, SD, MB, NC, MC, SC, WD, KE.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Watt (1971); Wise (1977: 64); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 221, 223); Palma (2010).

Other significant references: Clay (1953a); Emerson (1955: 296, fig. 24); Tendeiro (1965b: 132, figs 79–81); Green & Palma (1991: 18, 32); Palma (1996b: 217); Price *et al.* (2003).

Remarks: *Rallicola (R.) lugens* is a widespread louse species, frequently collected from New Zealand pukekos.

### ***Rallicola (Rallicola) ortyometrae philippensis* Emerson, 1966**

*Rallicola ortyometrae philippensis* Emerson, 1966: 337.

*Rallicola ortyometrae* (Schrank, 1781) *s. l.*; Pilgrim & Palma 1982: 18.

*Rallicola ortyometrae philippensis* Emerson, 1966; Murray *et al.* 1993: 961.

*Rallicola (Rallicola) philippensis* Emerson, 1966; Price *et al.* 2003: 230.

“*Rallicola (R.) ortyometrae*”; Palma 2010: 408 (not *Pediculus ortyometrae* Schrank, 1781).

Holotype ♂ in NHML.

Type host: *Gallirallus philippensis philippensis* (Linnaeus, 1766).

New Zealand host: *Gallirallus philippensis assimilis* (G.R. Gray, 1843).

Other hosts: *Gallirallus philippensis mellori* (Mathews, 1912); *Gallirallus philippensis lesouefi* (Mathews, 1911).

New Zealand localities: ND, AK, BP, GB, SD, NN.

Geographic distribution: Australasia; Oceania.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).

Other significant references: Palma (1996b: 218); Price *et al.* (2003).

Remarks: In agreement with Emerson (1966: 337), and contrary to Price *et al.* (2003: 230) and Palma (2010: 408), I regard this louse taxon as a subspecies of *Rallicola ortyometrae*.

***Rallicola (Rallicola) tabuensis* Emerson, 1966**

Figs 164–165

*Rallicola tabuensis* Emerson, 1966: 336, fig. 1.*Rallicola* sp.; Pilgrim & Palma 1982: 19.*Rallicola tabuensis* Emerson, 1966; Pilgrim & Palma 1982: 19.*Rallicola* sp.; Murray *et al.* 1993: 961.*Rallicola tabuensis* Emerson, 1966; Murray *et al.* 1993: 961.*Rallicola (Rallicola) tabuensis* Emerson, 1966; Price *et al.* 2003: 231.*Rallicola (R.) tabuensis* Emerson, 1966; Palma 2010: 409.

Holotype ♂ in BPBM (Tenorio 1979: 13).

Type host: *Porzana tabuensis tabuensis* (J.F. Gmelin, 1789).New Zealand hosts: *Porzana pusilla affinis* (J.E. Gray, 1845); *Porzana tabuensis tabuensis* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: ND, BP, TH, CH.

Geographic distribution: Oceania.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010).Other significant references: Clay (1953a: 572); Moreby (1976: 93); Tenorio (1979: 13); Green & Palma (1991: 19, 31); Palma (1996b: 218); Price *et al.* (2003).Remarks: This is the first record of *Rallicola (R.) tabuensis* from *Porzana pusilla affinis* because the references cited above reported this louse under that host as “*Rallicola* sp.” only.

The illustration of the male genitalia of *Rallicola (R.) tabuensis* in Emerson (1966: 336, fig. 1) is highly inaccurate. Besides the basal plate and the parameres, it shows a “Y” shaped median structure which is the result of an misinterpreted fusion of the internal mesosomal ring with the external and heavily pigmented distal process arising from the last ventral abdominal segment (see Clay 1953a: 572). Secondly, Emerson’ figure (1966: 336, fig. 1) shows thin, pointed parameres when, in fact, each paramere has a triangular, hyaline lateral expansion with a wide distal end.

***Rallicola (Rallicola) takahe* Holloway, 1956***Rallicola takahe* Holloway, 1956: 113, figs 1–5.*Rallicola takahe*; Miller 1971: 132, fig. 346.*Rallicola takahe* Holloway, 1956; Wise 1977: 64.*Rallicola takahe* Holloway, 1956; Pilgrim & Palma 1982: 19.*Rallicola takahe* Holloway, 1956; Murray *et al.* 1993: 961.*Rallicola (Rallicola) takahe* Holloway, 1956; Price *et al.* 2003: 231.*Rallicola (R.) takahe* Holloway, 1956; Palma 2010: 409.Holotype ♂ in MONZ (Palma *et al.* 1989: 45).Type host: *Porphyrio hochstetteri* (A.B. Meyer, 1883).New Zealand host: *Porphyrio hochstetteri* (A.B. Meyer, 1883).

Other hosts: None.

New Zealand localities: WA, WN, FD.

Geographic distribution: New Zealand.

New Zealand references: Holloway (1956); Pilgrim (1970: 75); Miller (1971: 132); Wise (1977); Pilgrim & Palma (1982); Palma *et al.* (1989: 45); Murray *et al.* (1993); Palma (2010); Buckley *et al.* (2012: 137, App. 2).Other significant reference: Price *et al.* (2003).

Remarks: *Rallicola takahe* is an endemic and critically threatened species (Buckley *et al.* 2012), exclusively parasitic on the South Island takahe, *Porphyrio hochstetteri*. The name of the type host used in the original description of this louse is “*Notornis mantelli* Owen, 1848”. However, the present name of this bird species is *Porphyrio hochstetteri* (A.B. Meyer, 1883), the South Island takahe, while the name *Porphyrio mantelli* (Owen, 1848) is now applied to the extinct North Island takahe (Checklist Committee 2010: 188).

**Genus *Saemundssonina* Timmermann, 1936**

*Saemundssonina* Timmermann, 1936 [April]. *Zool. Anz.* 114: 97. Type species: *Philopterus gonothorax* (Giebel, 1874) = *Saemundssonina (Saemundssonina) lari* (O. Fabricius, 1780) (by original designation).

**Subgenus *Puffinoecus* Eichler, 1949**

*Puffinoecus* Eichler, 1949a. *Boll. Soc. Entomol. Italiana* 79: 12. Type species: *Puffinoecus peusi* Eichler, 1949a = *Saemundssonina (Puffinoecus) peusi* (Eichler, 1949) (by original designation).

***Saemundssonina (Puffinoecus) enderleini* (Eichler, 1949)**

Figs 166–167

“*Docophorus Schillingi*” Enderlein, 1908: 444, figs 192, 213 (not *Trabeculus schillingi* Rudow, 1866b).

*Puffinoecus enderleini* Eichler, 1949a: 13. *Nomen novum* for *Docophorus schillingi* Enderlein, 1908.

*Saemundssonina enderleini* (Eichler, 1949); Hopkins & Clay 1952: 330.

*Puffinoecus Enderleini* Eichler, 1946 [sic]; Séguy 1953: 598, figs 55–56.

*Saemundssonina enderleini* (Eichler, 1949); Timmermann 1965: 81.

*Saemundssonina* sp. [ex *Pterodroma lessonii*]; Watson 1967: 74.

*Saemundssonina* sp.; Clay & Moreby 1970: 218.

*Saemundssonina* sp.; Gressitt 1970: 329.

*Saemundssonina* sp.; Pilgrim & Palma 1982: 8.

*Saemundssonina* sp.; Murray *et al.* 1990: 1369.

*Saemundssonina (Puffinoecus)* sp.; Green & Palma 1991: 20, 26.

*Puffinoecus enderleini* Eichler, 1949; Martín-Mateo 1996: 61.

*Saemundssonina (Puffinoecus) enderleini* (Eichler, 1949); Palma 1999: 378.

*Saemundssonina (Puffinoecus)* species; Marris 2000: 188.

*Saemundssonina (Puffinoecus)* sp.; Palma & Horning 2002: 13, 16.

*Saemundssonina (Puffinoecus) enderleini* (Eichler, 1949); Price *et al.* 2003: 233.

*Saemundssonina (Puffinoecus) enderleini* (Eichler, 1949); Palma 2010: 409.

Syntypes ♂♀ in ZMHU, but syntypes ♂ now lost (Jürgen Deckert pers. comm. September 2013).

Type host: *Pterodroma mollis* (Gould, 1844).

New Zealand hosts: *Pterodroma macroptera gouldi* (Hutton, 1869); *Pterodroma lessonii* (Garnot, 1826); *Pterodroma mollis* (Gould, 1844).

Other hosts: None.

New Zealand localities: AK, BP, TK, WN, MC, CH, AN, Macquarie Island.

Geographic distribution: Atlantic, Indian and Pacific Oceans.

New Zealand references: Clay & Moreby (1970); Gressitt (1970); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999); Marris (2000); Palma & Horning (2002); Palma (2010).

Other significant references: Séguy (1953); Timmermann (1965); Martín-Mateo (1996); Price *et al.* (2003).

Remarks: *Saemundssonina (P.) enderleini* is a frequently collected “head & neck” louse parasitic on three gadfly petrel species.

***Saemundssonina (Puffinoecus) jamaicensis* Timmermann, 1962****New Record**

*Saemundssonina jamaicensis* Timmermann, 1962a: 430.

*Saemundssonina jamaicensis* Timmermann, 1962; Timmermann 1965: 80.

*Puffinoecus jamaicensis* Timmermann, 1961 [sic]; Martín-Mateo 1996: 51, figs 1c, 3d–5d, 6a.

*Saemundssonina (Puffinoecus)* sp.; Palma 1999: 376.

*Saemundssonina (Puffinoecus) jamaicensis* Timmermann, 1962; Price *et al.* 2003: 234.

Holotype ♂ in NHML.

Type host: *Pterodroma hasitata caribbaea* Carte, 1866.

New Zealand host: *Pseudobulweria rostrata* (Peale, 1848).

Other host: *Pterodroma hasitata hasitata* (Kuhl, 1820).

New Zealand locality: ND.

Geographic distribution: Pacific and Atlantic Oceans.



New Zealand reference: Palma (1999).

Other significant references: Timmermann (1965); Zonfrillo (1993: 327); Forrester *et al.* (1995: 5); Martín-Mateo (1996); Price *et al.* (2003).

Material examined and repository: 4♂, 3♀, 1N (1 sample, MONZ).

Remarks: *Saemundssonina (P.) jamaicensis* is an infrequently collected “head & neck” louse parasitic on gadfly petrels. Also, it is a new louse species for New Zealand because the New Zealand reference cited above reported this louse as “*Saemundssonina (Puffinoecus)* sp.” only. *Pseudobulweria rostrata* is a rare vagrant to New Zealand (Checklist Committee 2010: 110), and a new host record for *S. (P.) jamaicensis* (voucher specimens in MONZ).

#### ***Saemundssonina (Puffinoecus) orientalis* (Uchida, 1949)**

*Philopterus validus* var *orientalis* Uchida, 1949: 541, fig. 11.

*Saemundssonina orientalis* (Uchida, 1949); Hopkins & Clay 1952: 334.

*Saemundssonina orientalis* (Uchida, 1948) [sic]; Timmermann 1965: 80.

*Puffinoecus orientalis* (Uchida, 1948) [sic]; Martín-Mateo 1996: 55, figs 1d, 3c–5c, 6d.

*Saemundssonina (Puffinoecus) orientalis* (Uchida, 1949); Price *et al.* 2003: 236.

*Saemundssonina (Puffinoecus) orientalis* (Uchida, 1949); Scofield *et al.* 2011: 214.

Syntypes ♂♀, not located in NSMJ (Takuya Kiyoshi pers. comm. May 2013).

Type host: *Calonectris leucomelas* (Temminck, 1836).

New Zealand host: *Calonectris leucomelas* (Temminck, 1836).

Other hosts: None.

New Zealand locality: WO.

Geographic distribution: Japan; Korea; eastern China; Pacific Ocean.

New Zealand reference: Scofield *et al.* (2011).

Other significant references: Timmermann (1965); Martín-Mateo (1996); Price *et al.* (2003).

Remarks: *Saemundssonina (P.) orientalis* is an infrequently collected “head & neck” louse exclusively parasitic on streaked shearwaters. *Calonectris leucomelas* has been recorded only once in New Zealand (Checklist Committee 2010: 111; Scofield *et al.* 2011).

#### ***Saemundssonina (Puffinoecus) puellula* Timmermann, 1965**

*Saemundssonina puellula* Timmermann, 1965: 82.

*Saemundssonina puellula* Timmermann, 1965; Watt 1971: 238, 243.

*Saemundssonina puellula* Timmermann, 1965; Pilgrim & Palma 1982: 11.

*Puffinoecus nadleri* Mey, 1989: 54, figs 1–3.

*Saemundssonina puellula* Timmermann, 1965; Murray *et al.* 1990: 1371.

*Saemundssonina (Puffinoecus) puellula* Timmermann, 1965; Palma 1994a: 67.

*Puffinoecus puellulus* (Timmermann, 1965); Martín-Mateo 1996: 59, figs 1e,f, 3g–5g, 6f, 8.

*Saemundssonina (Puffinoecus) puellula* Timmermann, 1965; Price *et al.* 2003: 237.

*Saemundssonina (P.) puellula* Timmermann, 1965; Palma 2010: 409.

Holotype ♂ in NHML.

Type host: *Puffinus pacificus cuneatus* Salvin, 1888.

New Zealand host: *Puffinus pacificus pacificus* (J.F. Gmelin, 1789).

Other hosts: *Puffinus creatopus* Coues, 1864; *Puffinus gravis* (O'Reilly, 1818).

New Zealand locality: KE.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Watt (1971); Wise (1977: 65); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1994a); Paterson *et al.* (1999: 222); Palma (2010).

Other significant references: Amerson & Emerson (1971: 7, 26); Mey (1989); Martín-Mateo (1996); Palma (1996b: 227); Price *et al.* (2003).

Remarks: *Saemundssonina (P.) puellula* is an infrequently collected “head & neck” louse parasitic on the larger species of shearwaters.

**Saemundssonina (*Puffinoecus*) *valida* (Kellogg & Chapman, 1899)**

- Docophorus validus* Kellogg & Chapman, 1899: 56, pl. 5: fig. 2.  
*Philopterus validus* Kellogg & Chapman, 1899 [sic]; Harrison 1916: 106.  
*Saemundssonina valida* (Kellogg & Chapman, 1899); Hopkins & Clay 1952: 336.  
*Saemundssonina valida* (Kellogg & Chapman, 1899); Timmermann 1965: 81.  
*Saemundssonina* sp.; Pilgrim & Palma 1982: 12.  
*Saemundssonina* sp.; Murray *et al.* 1990: 1371.  
*Puffinoecus validus* (Kellogg & Chapman, 1899); Martín-Mateo 1996: 61.  
*Saemundssonina (Puffinoecus) valida* (Kellogg & Chapman, 1899); Palma 1999: 376.  
*Saemundssonina (Puffinoecus) valida* (Kellogg & Chapman, 1899); Price *et al.* 2003: 238.  
*Saemundssonina (P.) valida* (Kellogg & Chapman, 1899); Palma 2010: 409.

Holotype ♀ in EMEC (Carriker 1957: 99; Palma 1996b: 227).

Type host: *Puffinus opisthomelas* Coues, 1864.

New Zealand host: *Puffinus gavia* (J.R. Forster, 1844).

Other hosts: None.

New Zealand localities: AK, CL, WN, NN, TH.

Geographic distribution: Pacific Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999); Palma (2010).

Other significant references: Kellogg & Kuwana (1902: 461); Uchida (1949: 541); Timmermann (1965); Martín-Mateo (1996); Palma (1996b: 227); Price *et al.* (2003).

Remarks: *Saemundssonina (P.) valida* is a frequently collected “head & neck” louse parasitic on the smaller species of sheawaters. Kellogg & Kuwana (1902: 461) and Uchida (1949: 541) stated that the type host of *Docophorus validus* Kellogg & Chapman, 1899 was *Puffinus gavia*. However, that statement is incorrect, as clarified by Timmermann (1965: 81).

***Saemundssonina (Puffinoecus)* species**

- Saemundssonina* sp.; Pilgrim & Palma 1982: 11.  
*Saemundssonina* sp.; Murray *et al.* 1990: 1371.  
*Saemundssonina (Puffinoecus)* sp.; Palma 1999: 378.  
*Saemundssonina (Puffinoecus)* sp.; Palma & Imber 2000: 230.

New Zealand hosts: *Pterodroma magentae* (Giglioli & Salvadori, 1869); *Puffinus carneipes* Gould, 1844.

New Zealand localities: ND, CL, BP, CH.

Geographic distribution: Pacific Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999); Palma & Imber (2000); Palma (2010); Buckley *et al.* (2012: App. 2).

Other significant reference: Martín-Mateo (1996).

Remarks: Available records of *Saemundssonina (Puffinoecus)* from the two hosts listed above could not be identified to species because the samples contain females only (voucher specimens in MONZ) (see Palma 2000: 127).

**Subgenus *Saemundssonina* Timmermann, 1936**

- Saemundssonina* Timmermann, 1936 [April]. *Zool. Anz.* 114: 97. Type species: *Philopterus gonothorax* (Giebel, 1874) = *Saemundssonina (Saemundssonina) lari* (O. Fabricius, 1780) (by original designation).  
*Hastaephorus* Kéler, 1936 [November]. *Arb. Morph. tax. Entomol. Berlin-Dahlem* 3: 261. Type species: *Docophorus alpinus* Giebel, 1874 = *Saemundssonina (Saemundssonina) tringae* (O. Fabricius, 1780) (by original designation).

***Saemundssonina (Saemundssonina) albatrossa* Palma, 2012**

Figs 168–169

- Saemundssonina* sp.; Pilgrim & Palma 1982: 6.  
*Saemundssonina* sp.; Murray *et al.* 1990: 1368–1369.  
*Saemundssonina (Saemundssonina)* species 2; Marris 2000: 188.  
*Saemundssonina* sp. M; Palma 2010: 409.  
*Saemundssonina (Saemundssonina) albatrossa* Palma, 2012: 39, figs 1–4, 11, 18.

Holotype ♂ in MONZ.

Type host: *Phoebetria palpebrata* (J.R. Forster, 1785).

New Zealand hosts: *Phoebetria palpebrata* (J.R. Forster, 1785); *Thalassarche chrysostoma* (J.R. Forster, 1785); *Thalassarche impavida* Mathews, 1912.

Other hosts: None.

New Zealand localities: ND, AK, TK, WI, CA.

Geographic distribution: Southern Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Marris (2000); Palma (2010; 2012).

Other significant references: None.

Remarks: *Saemundssonina* (*S.*) *albatrossa* is an infrequently collected “head & neck” louse parasitic on small albatrosses.

### ***Saemundssonina* (*Saemundssonina*) *albemarlensis* (Kellogg & Kuwana, 1902)**

*Docophorus albemarlensis* Kellogg & Kuwana, 1902: 465, pl. 28: fig. 5.

*Philopterus albemarlensis* Kellogg & Kuwana, 1902 [sic]; Harrison 1916: 87.

*Saemundssonina albemarlensis* (Kellogg & Kuwana, 1902); Hopkins & Clay 1952: 329 (as junior synonym of *Saemundssonina phaetona* (Osborn, 1890)).

*Saemundssonina petersi* Ward, 1955: 90, figs 1b,d,f.

*Saemundssonina albemarlensis* (Kellogg & Kuwana, 1902); Emerson 1972a: 154.

*Saemundssonina albemarlensis* (Kellogg & Kuwana, 1902); Pilgrim & Palma 1982: 23.

*Saemundssonina* (*Saemundssonina*) *albemarlensis* (Kellogg & Kuwana, 1902); Price *et al.* 2003: 232.

*Saemundssonina* (*Saemundssonina*) *albemarlensis* (Kellogg & Kuwana, 1902); Murray *et al.* 2006a: 1965.

*Saemundssonina* (*Saemundssonina*) *albemarlensis* (Kellogg & Kuwana, 1902); Palma 2010: 409.

Lectotype ♂ in EMEC (Palma & Peck 2013: 58).

Type host: “*Camarhynchus psittacula affinis*”, in error (see Emerson 1972a: 154).

New Zealand host: *Onychoprion fuscatus serratus* (J.R. Forster [in Wagler], 1830).

Other hosts: *Onychoprion fuscatus fuscatus* (Linnaeus, 1766); *Onychoprion fuscatus oahuensis* (Bloxxham, 1826); *Onychoprion fuscatus crissalis* (Lawrence, 1872).

New Zealand localities: WO, KE.

Geographic distribution: Pacific, Atlantic and Indian Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Ward (1955: 90); Amerson & Emerson (1971: 17, 26); Emerson (1972a); Ward & Downey (1973: 395); Forrester *et al.* (1995: 31); Price *et al.* (2003: 232); Palma & Peck (2013: 57).

Remarks: *Saemundssonina* (*S.*) *albemarlensis* is an infrequently collected “head & neck” louse parasitic on sooty terns. *Saemundssonina petersi* Ward, 1955 was demoted to a junior synonym of *S.* (*S.*) *albemarlensis* by Emerson (1972a: 154).

### ***Saemundssonina* (*Saemundssonina*) *antarctica* (Wood, 1937)**

“*Philopterus melanocephalus*” Neumann, 1907a: 14 (not *Docophorus melanocephalus* Burmeister, 1838).

*Philopterus antarcticus* Wood, 1937 in Harrison, 1937: 22, fig. 1.

*Saemundssonina antarctica* (Wood, 1937); Hopkins & Clay 1952: 329.

*Saemundssonina nivea* Timmermann, 1956: 190, fig. 5.

*Saemundssonina nivea* Timmermann, 1956; Timmermann 1959a: 150, fig. 1.

*Saemundssonina nivea* Timmermann, 1956; Timmermann 1965: 76, fig. 14.

*Saemundssonina antarctica* (Wood, 1937); Timmermann 1965: 77.

“*Saemundssonina lari*” Clay & Moreby, 1967: 164 (not *Pediculus lari* O. Fabricius, 1780).

*Saemundssonina nivea* Timmermann, 1956; Clay & Moreby 1967: 165, 168, figs 154, 159, 175.

*Saemundssonina antarctica* (Wood, 1937); Pilgrim & Palma 1982: 7.

*Saemundssonina* (*Saemundssonina*) *antarctica* (Wood, 1937); Pilgrim & Palma 1994: 241, figs 1–3.

*Saemundssonina* (*S.*) *antarctica* (Wood, 1937); Palma 2010: 409.

Holotype ♂ in AMSA (Pilgrim & Palma 1994: 241). Holotype ♂ of *Saemundssonina nivea* Timmermann, 1956 in NHML (Pilgrim & Palma 1994: 241).

Type host: *Pagodroma nivea nivea* (G. Forster, 1777).

New Zealand hosts: *Thalassoica antarctica* (J.F. Gmelin, 1789); *Pagodroma nivea major* (Schlegel, 1863).

Other hosts: None.

New Zealand localities: ND, AK, WI, RO.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990: 1369); Pilgrim & Palma (1994); Paterson *et al.* (1999: 222); Palma (2010).

Other significant references: Harrison (1937); Timmermann (1956; 1959a; 1965); Clay & Moreby (1967); Steele *et al.* (1997: 292); Price *et al.* (2003: 232).

Remarks: The record of *Saemundssonina (S.) antarctica* from Macquarie Island in Palma (1996b: 219) and in Palma & Horning (2002: 12, 16) is not included here because it is the result of straggling or contamination (see Pilgrim & Palma 1994: 242). The record of this louse species from *Pagodroma nivea nivea* in Pilgrim & Palma (1982: 8) is erroneous; it is actually from *Pagodroma nivea major*, due to a confusion on the identity of the host (Checklist Committee 2010: 84).

### ***Saemundssonina (Saemundssonina) bicolor* (Rudow, 1870)**

*Docophorus bicolor* Rudow, 1870: 459.

? *Philopterus bicolor* (Rudow, 1870); Clay 1940a: 297.

*Saemundssonina bicolor* (Rudow, 1870); Hopkins & Clay 1952: 329.

*Saemundssonina creatopae* Carriker, 1964: 14, figs 11–13.

*Docophorus bicolor* Rudow, 1870; Timmermann 1965: 76 (as a *nomen dubium*).

*Saemundssonina bicolor* (Rudow, 1870); Clay & Moreby 1967: 165, 168, figs 161, 178.

*Saemundssonina* sp.; Lowry *et al.* 1978: 139.

*Saemundssonina bicolor* (Rudow, 1870); Pilgrim & Palma 1982: 7.

*Saemundssonina bicolor* (Rudow, 1870); Murray *et al.* 1990: 1369.

*Saemundssonina (Saemundssonina) bicolor* (Rudow, 1870); Palma 1994a: 67.

*Saemundssonina (S.) bicolor* (Rudow, 1870); Palma 2010: 409.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma (1994a: 67).

Type host: *Fulmarus glacialis* (A. Smith, 1840).

New Zealand host: *Fulmarus glacialis* (A. Smith, 1840).

Other hosts: None.

New Zealand localities: AK, TK, WI, WN, NN, KA, NC, MC, SC, CO, DN, SL, Macquarie Island.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Lowry *et al.* (1978); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b: 219); Palma & Horning (2002: 12, 16); Palma (2010).

Other significant references: Clay (1940a); Carriker (1964); Clay & Moreby (1967); Green & Palma (1991: 19, 26); Palma (1994a); Price *et al.* (2003: 233).

Remarks: *Saemundssonina (S.) bicolor* is a frequently collected “head & neck” louse parasitic on Antarctic fulmars. *Fulmarus glacialis* breeds on the coast of Antarctica and is a regular straggler to New Zealand seas (Checklist Committee 2010: 80).

### ***Saemundssonina (Saemundssonina) cephalus* (Denny, 1842)**

*Philopterus (Docophorus) cephalus* Denny, 1842: 44, 81, pl. 2: fig. 8.

*Docophorus pustulosus* Nitzsch [*in* Giebel], 1866: 363.

*Docophorus lari magna* Piaget, 1880: 112.

*Docophorus atlanticus* Kellogg, 1914: 81, pl. 16: fig. 1.

*Philopterus cephalus* Denny, 1842; Harrison 1916: 91.

*Saemundssonina pustulosa* (Nitzsch *in* Giebel), 1866 [sic]; Timmermann 1949a: 10, figs 4, 6.

*Saemundssonina cephalus* (Denny, 1842); Hopkins & Clay 1952: 330.

*Saemundssonina cephalus* (Denny, 1842); Pilgrim & Palma 1982: 22.

*Saemundssonina (Saemundssonina) cephalus* (Denny, 1842); Palma 2000: 122, fig. 1.

*Saemundssonina (Saemundssonina) cephalus* (Denny, 1842); Murray *et al.* 2006a: 1965.

*Saemundssonina (S.) cephalus* (Denny, 1842); Palma 2010: 409.

Lectotype ♀ in NHML (Palma 2000: 122). Lectotype ♂ of *Docophorus atlanticus* in USNM (Palma 2004: 16, fig. 1).

Type host: *Stercorarius parasiticus* (Linnaeus, 1758).

New Zealand host: *Stercorarius parasiticus* (Linnaeus, 1758).

Other hosts: None.

New Zealand localities: HB, KA, MC.

Geographic distribution: Eurasia; North America; southern South America, southern Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Palma (2000); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1949a; 1957a: 47); Hackman & Nyholm (1968: 77); Cohen *et al.* (1997: 186); Ramli *et al.* (2000: 71); Price *et al.* (2003: 233); Palma (2004: 13); Palma & Jensen (2005: 58, 64).

Remarks: *Saemundssonina (S.) cephalus* is an infrequently collected “head & neck” louse parasitic on Arctic skuas. *Stercorarius parasiticus* is a summer migrant to New Zealand (Checklist Committee 2010: 226), with only two records of *Saemundssonina (S.) cephalus* from this country.

### ***Saemundssonina (Saemundssonina) chathamensis* Timmermann, 1977**

*Saemundssonina chathamensis* Timmermann, 1977: 135.

*Saemundssonina chathamensis* Timmermann, 1977; Pilgrim & Palma 1982: 20.

*Saemundssonina chathamensis* Timmermann, 1977; Murray *et al.* 1993: 962.

*Saemundssonina (Saemundssonina) chathamensis* Timmermann, 1977; Price *et al.* 2003: 233.

*Saemundssonina (S.) chathamensis* Timmermann, 1977; Palma 2010: 409.

Holotype ♂ in NZAC.

Type host: *Thinornis novaeseelandiae* (J.F. Gmelin, 1789).

New Zealand host: *Thinornis novaeseelandiae* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand locality: CH.

Geographic distribution: Chatham Islands, New Zealand.

New Zealand references: Timmermann (1977); Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (2010); Buckley *et al.* (2012: 137, App. 2).

Other significant reference: Price *et al.* (2003).

Remarks: *Saemundssonina (S.) chathamensis* is an infrequently collected “head & neck” louse parasitic on New Zealand shore birds. Both the host and *Saemundssonina (S.) chathamensis* are endemic and critically threatened species (Buckley *et al.* 2012).

### ***Saemundssonina (Saemundssonina) conica conica* (Denny, 1842)**

*Philopterus (Docophorus) conicus* Denny, 1842: 45, 90, pl. 5: fig. 2.

*Docophorus fuliginosus hawaiiensis* Kellogg & Chapman, 1902: 157.

*Philopterus wallacei* Johnston & Harrison, 1912: 369, figs 5–6.

*Philopterus numeniicola* Johnston & Harrison, 1912: 372, figs 11–12.

*Philopterus conicus* (Denny, 1842) [sic]; Thompson 1939: 16.

*Saemundssonina conica* (Denny, 1842); Timmermann 1949a: 18, figs 10–11.

*Saemundssonina conica* (Denny, 1842); Hopkins & Clay 1952: 330.

*Saemundssonina hawaiiensis* (Kellogg & Chapman, 1902); Hopkins & Clay 1952: 332.

*Saemundssonina numeniicola* (Johnston & Harrison, 1912); Hopkins & Clay 1952: 334.

*Saemundssonina hawaiiensis* (Kellogg & Chapman, 1902); Pilgrim 1970: 75.

*Saemundssonina numeniicola* (Johnston & Harrison, 1912); Pilgrim 1970: 75.

*Saemundssonina hawaiiensis* (Kellogg & Chapman, 1902); Watt 1971: 238, 243.

*Saemundssonina numeniicola* (Johnston & Harrison, 1912); Watt 1971: 238, 243.

*Saemundssonina conica conica* (Denny, 1842); Emerson 1972a: 155.

*Saemundssonina conica conica* (Denny, 1842); Wise 1977: 64.

*Saemundssonina conica conica* (Denny, 1842); Pilgrim & Palma 1982: 20.

*Saemundssonina conica conica* (Denny, 1842); Murray *et al.* 1993: 962.

*Saemundssonina (Saemundssonina) conica conica* (Denny, 1842); Price *et al.* 2003: 233.

*Saemundssonina (S.) conica conica* (Denny, 1842); Palma 2010: 409.

Holotype nymph, probably lost (Vincent S. Smith pers. comm. December 2014). Both, the holotype ♂ of *Philopterus wallacei* and the lectotype ♀ of *Philopterus numeniicola* in MONZ (Palma *et al.* 1989: 45; Palma 1996b: 220).

Type host: *Pluvialis apricaria* (Linnaeus, 1758).

New Zealand host: *Pluvialis fulva* (J.F. Gmelin, 1789).

Other hosts: *Pluvialis dominicus* (Stadius Müller, 1776); *Charadrius vociferus* Linnaeus, 1758.

New Zealand localities: AK, SL, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Johnston & Harrison (1912); Thompson (1939: 71); Pilgrim (1970); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Palma *et al.* (1989: 45); Murray *et al.* (1993); Palma (1996b: 220); Palma (2010).

Other significant references: Timmermann (1949a; 1951c: 401); Amerson & Emerson (1971: 13, 26); Emerson (1972a); Moreby (1976: 93); Timmermann (1977: 137); Forrester *et al.* (1995: 25); Price *et al.* (2003).

Remarks: *Saemundssonina (S.) conica conica* is an infrequently collected “head & neck” louse parasitic on several plover species.

### ***Saemundssonina (Saemundssonina) desolata* Timmermann, 1959**

*Saemundssonina desolata* Timmermann, 1959a: 151, figs 3, 3a.

*Saemundssonina desolata* Timmermann, 1959; Timmermann 1965: 73, fig. 8.

*Saemundssonina desolata* Timmermann, 1959; Clay & Moreby 1967: 165, 168, figs 156–157.

*Saemundssonina desolata* Timmermann, 1959; Horning *et al.* 1980: 7, 10.

*Saemundssonina desolata* Timmermann, 1959; Pilgrim & Palma 1982: 10.

*Saemundssonina (Saemundssonina) desolata* Timmermann, 1959; Price *et al.* 2003: 233.

*Saemundssonina (S.) desolata* Timmermann, 1959; Palma 2010: 409.

Holotype ♂ in NHML.

Type host: *Pachyptila desolata* (J.F. Gmelin, 1789).

New Zealand hosts: *Pachyptila vittata* (G. Forster, 1777); *Pachyptila salvini salvini* (Mathews, 1912); *Pachyptila desolata* (J.F. Gmelin, 1789); *Pachyptila belcheri* (Mathews, 1912); *Pachyptila turtur* (Kuhl, 1820); *Pachyptila crassirostris crassirostris* (Mathews, 1912); *Pachyptila crassirostris pyramidalis* Fleming, 1939.

Other hosts: None.

New Zealand localities: ND, AK, BP, HB, WI, WA, WN, NN, NC, MC, SC, WD, CH, BO, SN, AN.

Geographic distribution: Southern Hemisphere.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990: 1370); Paterson *et al.* (1999: 222); Marris (2000: 188); Palma (2010).

Other significant references: Timmermann (1965); Clay & Moreby (1967); Green & Palma (1991: 19, 26); Palma (1996b: 220); Furness & Palma (1992: 35, 39); Price *et al.* (2003); Hänel & Palma (2007: 113, 127, 130).

Remarks: *Saemundssonina (S.) desolata* is a frequently collected “head & neck” louse parasitic on all the species and subspecies of prions.

### ***Saemundssonina (Saemundssonina) euryrhyncha* (Giebel, 1874)**

*Docophorus euryrhynchus* Giebel, 1874: 112.

*Philopterus euryrhynchus* Giebel, 1874 [sic]; Harrison 1916: 93 (as junior synonym of *Philopterus pustulosus* (Nitzsch, [in Giebel], 1866)).

*Saemundssonina stresemanni* Timmermann, 1949: 13, fig. 7, 9.

*Saemundssonina euryrhyncha* (Giebel, 1874); Hopkins & Clay 1952: 330.

*Saemundssonina stresemanni* Timmermann, 1949; Hopkins & Clay 1952: 336.

*Saemundssonina stresemanni* Timmermann, 1949; Timmermann 1957a: 47, fig. 19.

*Saemundssonina stresemanni* Timmermann, 1949; Clay 1964a: 232.

*Saemundssonina* sp. [ex “*Catharacta skua lonnbergi*”]; Watson 1967: 74.

*Saemundssonina stresemanni* Timmermann, 1949; Clay & Moreby 1967: 165, 169, fig. 162.

*Saemundssonina stresemanni* Timmermann, 1949; Spellerberg 1971: 19.

*Saemundssonina stresemanni* Timmermann, 1949; Wise 1977: 65.  
*Saemundssonina stresemanni* Timmermann, 1949; Horning *et al.* 1980: 7, 11.  
*Saemundssonina stresemanni* Timmermann, 1949; Pilgrim & Palma 1982: 22.  
*Saemundssonina stresemanni* ssp.; Mey 1994: 41, figs 20–21.  
*Saemundssonina euryrhyncha* (Giebel, 1874); Butler & O'Connor 1994: 456.  
*Saemundssonina (Saemundssonina) stresemanni* Timmermann, 1949; Palma 1996b: 225.  
*Saemundssonina stresemanni* Timmermann, 1949; Cohen *et al.* 1997: 186.  
*Saemundssonina (Saemundssonina) euryrhyncha* (Giebel, 1874); Palma 2000: 124, fig. 2.  
*Saemundssonina (Saemundssonina) euryrhyncha* (Giebel, 1874); Palma & Horning 2002: 12, 17, 22.  
*Saemundssonina (Saemundssonina) euryrhyncha* (Giebel, 1874); Murray *et al.* 2006a: 1965.  
*Saemundssonina (S.) euryrhyncha* (Giebel, 1874); Palma 2010: 409.

Syntypes ♂♀, presumed lost. See Palma (2000: 124). Holotype ♂ of *Saemundssonina stresemanni* in NHML (Palma 2000: 124).

Type host: *Coprotheres pomarinus* (Temminck, 1815).

New Zealand hosts: *Catharacta antarctica lonnbergi* Mathews, 1912; *Catharacta maccormicki* (Saunders, 1893).

Other hosts: *Catharacta antarctica antarctica* (Lesson, 1831); *Catharacta skua* Brünnich, 1764; *Catharacta chilensis* (Bonaparte, 1857); *Catharacta hamiltoni* Hagen, 1952.

New Zealand localities: SD, MB, SI, CH, SN, AU, CA, Macquarie Island, RO.

Geographic distribution: Antarctica; Atlantic, Indian and Pacific Oceans.

New Zealand references: Clay (1964a); Gressitt (1964: 539); Clay & Moreby (1967); Watson (1967); Gressitt (1970: 329); Schaefer & Strandtmann (1971: 16); Spellerberg (1971); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Palma (1996b); Palma (2000); Palma & Horning (2002); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1949a; 1957a); Amerson & Emerson (1971: 15, 26); Butler & O'Connor (1994); Mey (1994); Cohen *et al.* (1997); Ramli *et al.* (2000: 71); Price *et al.* (2003: 233); Palma & Jensen (2005: 58, 64); Hänel & Palma (2007: 113, 127, 131).

Remarks: *Saemundssonina (S.) euryrhyncha* is a frequently collected “head & neck” louse parasitic on large skuas. Records of *Saemundssonina* from species of *Catharacta* published before Palma (2000) are given as *S. stresemanni*.

### ***Saemundssonina (Saemundssonina) gaini* (Neumann, 1913)**

*Philopterus gaini* Neumann, 1913: 189, figs 1–3.  
*Philopterus gaini* Neumann, 1913; Harrison 1916: 95.  
*Saemundssonina gaini* (Neumann, 1913); Hopkins & Clay 1952: 331.  
*Saemundssonina gaini* (Neumann, 1913); Timmermann 1962a: 435.  
*Saemundssonina gaini* (Neumann, 1913); Timmermann 1965: 75, fig. 12, pl. 1: fig. 4.  
*Saemundssonina gaini* (Neumann, 1913); Clay & Moreby 1967: 165, 168, figs 160, 177.  
*Saemundssonina gaini* (Neumann, 1913); Horning *et al.* 1980: 7, 9.  
*Saemundssonina gaini* (Neumann, 1913); Pilgrim & Palma 1982: 7.  
*Saemundssonina (Saemundssonina) gaini* (Neumann, 1913); Price *et al.* 2003: 234.  
*Saemundssonina (S.) gaini* (Neumann, 1913); Palma 2010: 409.  
*Saemundssonina (Saemundssonina) gaini* (Neumann, 1913); Palma 2012: 40, figs 5–6, 12.

Lectotype ♂ in NHML (Timmermann 1962a: 435).

Type host: *Macronectes giganteus* (J.F. Gmelin, 1789).

New Zealand hosts: *Macronectes giganteus* (J.F. Gmelin, 1789); *Macronectes halli* Mathews, 1912.

Other hosts: None.

New Zealand localities: WN, SN.

Geographic distribution: Southern Hemisphere.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990: 1369); Paterson *et al.* (1999: 222); Palma (2010; 2012).

Other significant references: Timmermann (1962a; 1965); Green & Palma (1991: 19, 25); Palma (1996b: 221); Price *et al.* (2003).

Remarks: *Saemundssonina gaini* is an infrequently collected “head & neck” louse parasitic on both giant petrel species.

**Saemundssonina (Saemundssonina) gygisa Palma, 2012**

*Saemundssonina* sp.; Nelson 1969: 199.

*Saemundssonina* sp.; Amerson & Emerson 1971: 19, 27.

*Saemundssonina (Saemundssonina)* sp.; Palma 1999: 381.

*Saemundssonina (Saemundssonina)* sp.; Murray *et al.* 2006a: 1965.

*Saemundssonina* sp. M; Palma 2010: 409.

*Saemundssonina (Saemundssonina) gygisa* Palma, 2012: 44, figs 8, 15–17, 20.

Holotype ♂ in MONZ.

Type host: *Gygis alba candida* (J.F. Gmelin, 1789).

New Zealand host: *Gygis alba candida* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: AK, WN, KE, Norfolk Island.

Geographic distribution: Tropical Pacific Ocean.

New Zealand references: Nelson (1969); Palma (1999); Murray *et al.* (2006a); Palma (2010); Palma (2012).

Other significant reference: Amerson & Emerson (1971).

Remarks: *Saemundssonina (S.) gygisa* is an infrequently collected “head & neck” louse parasitic on white terns only.

**Saemundssonina (Saemundssonina) haematopi (Linnaeus, 1758)**

*Pediculus haematopi* Linnaeus, 1758: 613.

*Ricinus haematopi* (Linnaeus, 1758); Latreille 1804: 108.

*Docophorus acanthus* Giebel, 1874: 101.

*Philopterus acanthus* (Giebel, 1874); Cummings 1916b: 677, fig. 24.

*Hastaephorus acanthus* Giebel [sic]; Kéler 1936: 263, figs 2b,d.

*Saemundssonina haematopi* (Linnaeus, 1758); Hopkins & Clay 1952: 331.

*Saemundssonina haematopi*; Baker 1974: 20.

*Saemundssonina haematopi* (Linnaeus, 1758); Clay 1981b: 933, figs 2, 6, 8, 10.

*Saemundssonina haematopi* (Linnaeus, 1758); Pilgrim & Palma 1982: 19.

*Saemundssonina haematopi* (Linnaeus, 1758); Murray *et al.* 1993: 961.

*Saemundssonina (Saemundssonina) haematopi* (Linnaeus, 1758); Price *et al.* 2003: 234.

*Saemundssonina (S.) haematopi* (Linnaeus, 1758); Palma 2010: 409.

Neotype ♂ in NHML (Clay & Hopkins 1950: 260).

Type host: *Haematopus ostralegus ostralegus* Linnaeus, 1758.

New Zealand hosts: *Haematopus finschi* Martens, 1897; *Haematopus unicolor* J.R. Forster, 1844.

Other hosts: *Haematopus ater* Vieillot & Oudart, 1825; *Haematopus bachmani* Audubon, 1838; *Haematopus palliatus* Temminck, 1820; *Haematopus moquini* Bonaparte, 1856; *Haematopus longirostris* Vieillot, 1817; *Haematopus fuliginosus* Gould, 1845.

New Zealand localities: WI, WN, NN, KA, NC, MC, SC, WD, CO, DN, SL.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Baker (1974); Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999: 223); Palma (2010).

Other significant references: Cummings (1916b); Kéler (1936); Timmermann (1949a: 23, fig. 14 right); Clay & Hopkins (1950: 259); Timmermann (1957a: 36, fig. 5); Timmermann (1971: 158); Clay (1981b); Green & Palma (1991: 19, 32); Forrester *et al.* (1995: 26); Palma (1996b: 221); Price *et al.* (2003); Palma & Jensen (2005: 58, 63); Palma & Peck (2013: 59).

Remarks: *Saemundssonina (Saemundssonina) haematopi* is a frequently collected “head & neck” louse parasitic on most species of oystercatchers (Clay 1981b: 938).

**Saemundssonina (Saemundssonina) hexagona (Giebel, 1874)**

*Docophorus hexagonus* (Giebel, 1874): 116.

*Saemundssonina hexagonus* [sic] (Giebel, 1874); Thompson 1938c: 461, figs 1, 3b.

*Saemundssonina hexagona* (Giebel, 1874); Hopkins & Clay 1952: 332.

*Saemundssonina hexagona* (Giebel, 1874); Timmermann 1955: 516.



*Saemundssonina hexagona* (Giebel, 1874); Watt 1971: 238, 243, fig. 10.

*Saemundssonina hexagona* (Giebel, 1874); Pilgrim & Palma 1982: 14.

*Saemundssonina* (*Saemundssonina*) *hexagona* (Giebel, 1874); Price *et al.* 2003: 234.

*Saemundssonina* (*S.*) *hexagona* (Giebel, 1874); Palma 2010: 409.

Holotype probably ♀, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Phaethon rubricauda* Boddaert, 1783.

New Zealand host: *Phaethon rubricauda* Boddaert, 1783.

Other hosts: None.

New Zealand localities: ND, AK, KE.

Geographic distribution: Tropical Pacific and Indian Oceans.

New Zealand references: Watt (1971); Wise (1977: 64); Pilgrim & Palma (1982); Murray *et al.* (1990: 1373); Miller & Miller (1986: 50); Palma (2010).

Other significant references: Thompson (1938c); Timmermann (1955); Amerson & Emerson (1971: 9, 26); Ward & Downey (1973: 395); Palma (1996b: 221); Price *et al.* (2003).

Remarks: *Saemundssonina* (*S.*) *hexagona* is an infrequently collected “head & neck” louse parasitic on red-tailed tropicbirds.

### ***Saemundssonina* (*Saemundssonina*) *inexpectata* Timmermann, 1951**

*Saemundssonina inexpectata* Timmermann, 1951a: 9, fig. 1.

*Saemundssonina inexpectata* Timmermann, 1951; Hopkins & Clay 1953: 445.

*Saemundssonina inexpectata* Timmermann, 1951; Timmermann 1957a: 44, 47, fig. 14.

*Saemundssonina* sp.; Melville 1985: 67.

*Saemundssonina* (*Saemundssonina*) *inexpectata* Timmermann, 1951; Palma 1999: 380.

*Saemundssonina* (*Saemundssonina*) *inexpectata* Timmermann, 1951; Palma 2000: 125, fig. 3.

*Saemundssonina* (*Saemundssonina*) *inexpectata* Timmermann, 1951; Murray *et al.* 2006a: 1965.

*Saemundssonina* (*S.*) *inexpectata* Timmermann, 1951; Palma 2010: 409.

Holotype ♂ in NHML (Palma 2000: 125).

Type host: “*Larus minutus*”, in error (see Palma 2000: 126).

New Zealand host: *Stercorarius longicaudus* Vieillot, 1819.

Other hosts: None

New Zealand localities: WN, WI.

Geographic distribution: Cold and temperate regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Melville (1985); Palma (1999); Palma (2000); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1957a); Cohen *et al.* (1997: 186); Ramli *et al.* (2000: 71); Price *et al.* (2003: 234); Palma & Jensen (2005: 58, 64).

Remarks: *Saemundssonina* (*S.*) *inexpectata* is an infrequently collected “head & neck” louse parasitic on long-tailed skuas. *Stercorarius longicaudus* is a rare visitor to New Zealand (Checklist Committee 2010: 226), with only two records of *Saemundssonina* (*S.*) *inexpectata* from this country.

### ***Saemundssonina* (*Saemundssonina*) *lari* (O. Fabricius, 1780) *sensu lato***

*Pediculus lari* O. Fabricius, 1780: 219.

*Docophorus gonothorax* Giebel, 1874: 450.

*Philoapterus lari* O. Fabricius, 1780 [sic]; Harrison 1916: 97 (as junior synonym of *Philoapterus gonothorax* Giebel, 1871 [sic]).

*Philoapterus gonothorax* Giebel, 1871 [sic]; Harrison 1937: 21.

*Saemundssonina gonothorax* Giebel, 1871 [sic]; Timmermann 1949a: 4, figs 1–3.

*Saemundssonina lari fallai* Timmermann, 1951a: 7.

*Saemundssonina lari gonothorax* (Giebel, 1874); Timmermann 1951a: 7.

*Saemundssonina gonothorax* (Giebel, 1874); Hopkins & Clay 1952: 331.

*Saemundssonina lari* (O. Fabricius, 1780); Hopkins & Clay 1952: 332.

*Saemundssonina fallai* Timmermann, 1951; Hopkins & Clay 1953: 445.

*Saemundssonina lari lari* (O. Fabricius, 1780); Clay & Hopkins 1954: 249, figs 37–39.

*Saemundssonina lari* (O. Fabricius, 1780); Clay 1964a: 232.

*Saemundssonina* sp. [ex "*Larus dominicanus*"]; Watson 1967: 74.  
*Saemundssonina fallai* Timmermann, 1951; Pilgrim 1970: 75.  
*Saemundssonina lari* (O. Fabricius, 1780); Wise 1977: 64.  
*Saemundssonina lari fallai* Timmermann, 1951; Wise 1977: 64.  
*Saemundssonina lari gonothorax* (Giebel, 1874); Wise 1977: 65.  
*Saemundssonina lari* (O. Fabricius, 1780) *s. l.*; Horning *et al.* 1980: 7, 11.  
*Saemundssonina lari* (O. Fabricius, 1780) *s. l.*; Pilgrim & Palma 1982: 22.  
*Saemundssonina (Saemundssonina) lari* (O. Fabricius, 1780); Murray *et al.* 2006a: 1965.  
*Saemundssonina (S.) lari* (O. Fabricius, 1780); Palma 2010: 409.

Neotype ♂ in NHML (Clay & Hopkins 1954: 249, pl. 11: fig. 5). Holotype ♂ of *Saemundssonina lari fallai* in NHML.

Type host: *Larus hyperboreus* Gunnerus, 1767.

New Zealand hosts: *Larus dominicanus* Lichtenstein, 1823; *Larus novaehollandiae scopulinus* J.R. Forster, 1844; *Larus bulleri* Hutton, 1871.

Other hosts: At least 28 other species of *Larus* (see Price *et al.* 2003: 234); *Gabianus pacificus* (Latham, 1802); *Pagophila eburnea* (Phipps, 1774); *Rissa brevirostris* (Bruch, 1853); *Rissa tridactyla* (Linnaeus, 1758); *Xema sabini* (Sabine, 1819).

New Zealand localities: AK, HB, WA, WN, SD, MB, NN, KA, NC, MC, SC, WD, CO, DN, SL, SI, CH, SN, AU, CA, Macquarie Island.

Geographic distribution: Cosmopolitan.

New Zealand references: Harrison (1937); Clay (1964a); Gressitt (1964: 539); Watson (1967); Pilgrim (1970); Gressitt (1970: 329); Wise (1977); Weidner (1977: 103); Horning *et al.* (1980); Pilgrim & Palma (1982); Palma (1996b: 222); Palma & Horning (2002: 13, 18); Galloway (2005: 17); Murray *et al.* (2006a); Palma (2010; 2012: 44, fig. 14).

Other significant references: Timmermann (1949a); Timmermann (1951a); Timmermann (1957a: 42, figs 9, 12); Clay & Hopkins (1954); Clay & Moreby (1967: 165, 169, figs 163–167); Choe & Kim (1987: 3000; 1988: 988); Green & Palma (1991: 19, 33); Forrester *et al.* (1995: 29); Price *et al.* (2003: 234); Palma & Jensen (2005: 58, 64); Adam (2007: 179); Martín-Mateo (2009: 117, fig. 29); González-Acuña *et al.* (2011: 300); Palma & Peck (2013: 60); Yamagishi *et al.* (2014: 384).

Remarks: *Saemundssonina (Saemundssonina) lari* is a morphologically variable “head & neck” louse frequently collected from many species of gulls. Timmermann (1949a; 1951) subdivided this taxon into several subspecies which, without justification, have not been recognised as valid by Price *et al.* (2003: 238). Therefore, I qualify this louse species as “*sensu lato*”, recognising that, while it shows a range of intraspecific variation, it needs a detailed study to determine the validity of subdividing it into subspecies.

### ***Saemundssonina (Saemundssonina) limosae* (Denny, 1842)**

*Philoater (Docophorus) limosae* Denny, 1842: 44, 86, pl. 4: fig. 2.  
*Philoater limosae* Denny, 1842; Harrison 1916: 98.  
*Saemundssonina limosae* (Denny, 1842); Timmermann 1951c: 395, figs 3–4, 7 upper.  
*Saemundssonina limosae* (Denny, 1842); Hopkins & Clay 1952: 333.  
*Saemundssonina limosae* (Denny, 1842); Watt 1971: 238, 243, fig. 10.  
*Saemundssonina limosae* (Denny, 1842); Pilgrim & Palma 1982: 21.  
*Saemundssonina (Saemundssonina) limosae* (Denny, 1842); Murray *et al.* 2006a: 1964.  
*Saemundssonina (S.) limosae* (Denny, 1842); Palma 2010: 409.

Holotype ♀ in NHML (Vincent S. Smith pers. comm. December 2014).

Type host: *Limosa lapponica lapponica* (Linnaeus, 1758).

New Zealand host: *Limosa lapponica baueri* Naumann, 1836.

Other hosts: None.

New Zealand localities: ND, NN, KA, KE.

Geographic distribution: Eurasia; North America; Australasia.

New Zealand references: Watt (1971); Wise (1977: 65); Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1951c); Price *et al.* (2003: 235).

Remarks: *Saemundssonina (S.) limosae* is an infrequently collected “head & neck” louse parasitic on bar-tailed godwits. The eastern bar-tailed godwit is the most numerous wader that visits New Zealand every year (Checklist Committee 2010: 203).

#### ***Saemundssonina (Saemundssonina) lockleyi* Clay, 1949**

“*Philopterus melanocephalus*” Harrison, 1937: 22 (not *Docophorus melanocephalus* Burmeister, 1838a).

*Saemundssonina lockleyi* Clay, 1949a: 11, figs 17, 24–25.

*Saemundssonina* sp.; Clay 1964a: 232.

*Saemundssonina* sp. [ex *Sterna vittata bethunei*]; Watson 1967: 74.

*Saemundssonina lockleyi* Clay, 1949; Clay & Moreby 1967: 165, 169, figs 170, 173–174.

“*Saemundssonina melanocephalus*” Wise, 1977: 65 (not *Docophorus melanocephalus* Burmeister, 1838).

*Saemundssonina lockleyi* Clay, 1949 s. l.; Horning *et al.* 1980: 7, 12.

*Saemundssonina lockleyi* Clay, 1949 s. l.; Pilgrim & Palma 1982: 23, 31, note 24.

*Saemundssonina (Saemundssonina) lockleyi* Clay, 1949; Palma & Horning 2002: 13, 18, 21.

*Saemundssonina (Saemundssonina) lockleyi* Clay, 1949; Murray *et al.* 2006a: 1965.

*Saemundssonina (S.) lockleyi* (O. Fabricius, 1780); Palma 2010: 409.

Holotype ♂ in NHML.

Type host: *Sterna vittata georgiae* Reichenow, 1904.

New Zealand hosts: *Chlidonias albostrigatus* (G.R. Gray, 1845); *Sterna vittata bethunei* Travers, 1896; *Sterna paradisaea* Pontoppidan, 1763.

Other hosts: *Onychoprion anaethetus* (Scopoli, 1786); *Sterna virgata* Cabanis, 1875.

New Zealand localities: WN, CO, DN, SL, SN, AU, CA, Macquarie Island.

Geographic distribution: Cosmopolitan.

New Zealand references: Harrison (1937); Clay (1949a); Ward (1955: 87); Clay (1964a); Gressitt (1964: 539); Watson (1967); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Palma (1996b: 223); Palma & Horning (2002); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay & Moreby (1967; 1970: 220); Moreby (1976: 93); Forrester *et al.* (1995: 31); Price *et al.* (2003: 235); Palma & Jensen (2005: 59, 65); Hänel & Palma (2007: 113, 127, 131).

Remarks: *Saemundssonina (S.) lockleyi* is a frequently collected “head & neck” louse parasitic on several tern species. Horning *et al.* (1980) and Pilgrim & Palma (1982) regarded the populations of *Saemundssonina (S.) lockleyi* from *Sterna vittata bethunei* and *St. paradisaea* as somewhat different from that of the type host, and qualified them as *sensu lato*; however, my examination of more samples shows that making such difference is not warranted. *Chlidonias albostrigatus* is a new host record for *S. (S.) lockleyi* (voucher specimens in MONZ).

#### ***Saemundssonina (Saemundssonina) marina* Timmermann, 1956**

*Saemundssonina marina* Timmermann, 1956: 191, fig. 6.

*Saemundssonina marina* Timmermann, 1956; Timmermann 1965: 73, fig. 10.

*Saemundssonina marina* Timmermann, 1956; Pilgrim & Palma 1982: 13.

*Saemundssonina marina* Timmermann, 1956; Murray *et al.* 1990: 1372.

*Saemundssonina (Saemundssonina) marina* Timmermann, 1956; Palma 1996b: 223.

*Saemundssonina (Saemundssonina) marina* Timmermann, 1956; Hänel & Palma 2007: 107, 113, 127, 130, fig. 2b.

*Saemundssonina (S.) marina* Timmermann, 1956; Palma 2010: 409.

Holotype ♂ in NHML (Palma 1996b: 223; Vincent S. Smith pers. comm. February 2015).

Type host: *Pelagodroma marina* (Latham, 1790).

New Zealand host: *Pelagodroma marina maoriana* Mathews, 1912.

Other hosts: *Pelagodroma marina marina* (Latham, 1790); *Pelagodroma marina hypoleuca* (Moquin-Tandon, 1841); *Pelagodroma marina dulciae* Mathews, 1912.

New Zealand localities: ND, WA, NC, DN, CH, SN.

Geographic distribution: Australasia; Atlantic and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 222); Palma (2010).

Other significant references: Timmermann (1965); Furness & Palma (1992: 35, 42); Palma (1996b); Price *et al.* (2003: 235); Hänel & Palma (2007).

Remarks: *Saemundssonina (S.) marina* is a frequently collected “head & neck” louse parasitic on white-faced storm petrels. *Pelagodroma marina hypoleuca* and *P. m. dulciae* are new host records for *Saemundssonina (S.) marina* (voucher specimens in NHML and in MONZ, respectively).

***Saemundssonina (Saemundssonina) melanocephalus* (Burmeister, 1838)**

*Docophorus melanocephalus* Burmeister, 1838a: 426.

*Philopterus melanocephalus* Nitzsch, in Burmeister [sic], 1838a; Harrison 1916: 99.

*Saemundssonina melanocephalus* (Burmeister, 1838); Clay 1949a: 11, figs 8, 11, 19, 28–29.

*Saemundssonina melanocephalus* (Burmeister, 1838); Hopkins & Clay 1952: 333.

*Saemundssonina melanocephalus* (Burmeister, 1838); Timmermann 1957a: 44, 46, fig. 16a, pl. 10: fig. b.

*Saemundssonina melanocephalus* (Burmeister, 1838); Pilgrim & Palma 1982: 23.

*Saemundssonina (Saemundssonina) melanocephalus* (Burmeister, 1838); Murray *et al.* 2006a: 1965.

*Saemundssonina (S.) melanocephalus* (Burmeister, 1838); Palma 2010: 409.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Sternula albifrons albifrons* (Pallas, 1764) (see Clay 1949a: 11).

New Zealand host: *Sternula albifrons sinensis* (J.F. Gmelin, 1789).

Other host: *Sternula nereis nereis* Gould, 1843.

New Zealand localities: WI, WN.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay (1949a); Ward (1955: 87); Timmermann (1957a); Forrester *et al.* (1995: 31); Palma (1996b: 223); Price *et al.* (2003: 235); Martín-Mateo (2009: 116).

Remarks: *Saemundssonina (S.) melanocephalus* is an infrequently collected “head & neck” louse parasitic on small terns. Wise (1977: 65) listed “*Saemundssonina melanocephalus* (Burmeister, 1838)” for New Zealand, following Harrison’s (1937: 22) record of “*Philopterus melanocephalus*” from Macquarie Island, which is a misidentification of *Saemundssonina (Saemundssonina) lockleyi* Clay, 1949 (see above).

***Saemundssonina (Saemundssonina) nereis* Timmermann, 1956**

“? *Philopterus platycephalus*” Clay, 1940a: 298 (not *Docophorus platycephalus* Kellogg & Kuwana, 1902).

*Saemundssonina nereis* Timmermann, 1956: 189, fig. 4.

*Saemundssonina nereis* Timmermann, 1956; Timmermann 1965: 75, fig. 11.

*Saemundssonina nereis* Timmermann, 1956; Pilgrim & Palma 1982: 13.

*Saemundssonina nereis* Timmermann, 1956; Murray *et al.* 1990: 1372.

*Saemundssonina (Saemundssonina) nereis* Timmermann, 1956; Price *et al.* 2003: 235.

*Saemundssonina (S.) nereis* Timmermann, 1956; Palma 2010: 409.

Holotype ♂ in NHML.

Type host: *Garrodia nereis* (Gould, 1841).

New Zealand host: *Garrodia nereis* (Gould, 1841).

Other hosts: None.

New Zealand localities: WN, MC, CH, AN, AU.

Geographic distribution: Subantarctic Islands; Southern Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Marris (2000: 188); Palma (2010).

Other significant references: Timmermann (1965); Price *et al.* (2003).

Remarks: *Saemundssonina (S.) nereis* is a frequently collected “head & neck” louse exclusively parasitic on grey-backed storm petrels.

***Saemundssonina (Saemundssonina) platycephalus* (Kellogg & Kuwana, 1902)      New Record**

*Docophorus platycephalus* Kellogg & Kuwana, 1902: 461, pl. 28: fig. 1.

*Philopterus platycephalus* Kellogg & Kuwana, 1902 [sic]; Harrison 1916: 102.

- Saemundssonina platycephalus* (Kellogg & Kuwana, 1902); Hopkins & Clay 1952: 335.  
*Saemundssonina platycephalus* (Kellogg & Kuwana, 1902); Timmermann 1965: 73, fig. 9.  
*Saemundssonina platycephalus* (Kellogg & Kuwana, 1902); Clay in Linsley & Usinger 1966: 132.  
 “*Saemundssonina ?marina*” Clay & Moreby, 1967: 165, fig. 158 (not *Saemundssonina marina* Timmermann, 1956).  
*Saemundssonina* sp.; Pilgrim & Palma 1982: 13.  
 “*Saemundssonina ?marina*” Murray *et al.* 1990: 1372 (not *Saemundssonina marina* Timmermann, 1956).  
*Saemundssonina* sp.; Murray *et al.* 1990: 1372.  
*Saemundssonina* (*Saemundssonina*) species 1; Marris 2000: 188.  
*Saemundssonina* (*Saemundssonina*) *platycephalus* (Kellogg & Kuwana, 1902); Price *et al.* 2003: 236.  
*Saemundssonina* (*Saemundssonina*) *platycephalus* (Kellogg & Kuwana, 1902); Palma & Peck 2013: 60.

Lectotype ♂ in EMEC (Palma & Peck 2013: 61).

Type host: *Oceanites gracilis galapagoensis* Lowe, 1921.

New Zealand host: *Fregetta tropica* (Gould, 1844).

Other hosts: *Oceanites oceanicus oceanicus* (Kuhl, 1820); *Oceanites oceanicus exasperatus* Mathews, 1912; *Oceanites gracilis gracilis* (Elliot, 1859).

New Zealand locality: AN.

Geographic distribution: Antarctica; Southern Oceans; Pacific Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Marris (2000).

Other significant references: Timmermann (1965); Clay & Moreby (1967); Price *et al.* (2003); Palma & Peck (2013).

Material examined and repository: 1♂, 2♀ (1 sample, MONZ).

Remarks: *Saemundssonina* (*S.*) *platycephalus* is an infrequently collected “head & neck” louse parasitic on several storm petrel species. This is the first record of *Saemundssonina* (*S.*) *platycephalus* for New Zealand because the New Zealand references cited above reported this louse as “*Saemundssonina* sp.” only. Furthermore, *Fregetta tropica* is new host record for *Saemundssonina* (*S.*) *platycephalus* (voucher specimens in MONZ).

### ***Saemundssonina* (*Saemundssonina*) *platygaster* (Denny, 1842) *sensu lato***

- Philopterus* (*Docophorus*) *platygaster* Denny, 1842: 44, 83, pl. 2: fig. 5.  
*Philopterus platygaster* Denny, 1842; Harrison 1916: 102.  
*Saemundssonina platygaster* (Denny, 1842); Hopkins & Clay 1952: 335.  
*Saemundssonina platygaster* (Denny, 1842) *s. l.*; Pilgrim & Palma 1982: 21.  
*Saemundssonina platygaster* (Denny, 1842); Murray *et al.* 1993: 962. In part.  
*Saemundssonina platygaster* (Denny, 1842); Paterson *et al.* 1999: 221, 223.  
*Saemundssonina* (*Saemundssonina*) *platygaster* (Denny, 1842) *sensu lato*; Palma 1999: 380.  
*Saemundssonina* (*Saemundssonina*) *platygaster* (Denny, 1842); Price *et al.* 2003: 236.  
*Saemundssonina* (*S.*) *platygaster* (Denny, 1842) *s. l.*; Palma 2010: 409.

Syntypes ♀♀, probably lost (Thompson 1937a: 78).

Type host: *Charadrius hiaticula* Linnaeus, 1758.

New Zealand hosts: *Tringa incana* (J.F. Gmelin, 1789); *Himantopus himantopus leucocephalus* Gould, 1837; *Himantopus novaehollandiae* Gould, 1841; *Charadrius obscurus aquilonius* Dowding, 1994; *Charadrius obscurus obscurus* (J.F. Gmelin, 1789).

Other hosts: *Tringa cinerea* (Güldenstaedt, 1774); *Charadrius dubius* Scopoli, 1786; *Charadrius marginatus* Vieillot, 1818.

New Zealand localities: AK, WI, NN, NC, MC, SC, CO, DN, SL, SI.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Paterson *et al.* (1999); Palma (1999); Palma (2010).

Other significant references: Timmermann (1969a: 233); Green & Palma (1991: 19, 33); Palma (1996b: 224); Price *et al.* (2003: 236); Martín-Mateo (2009: 120); Palma & Peck (2013: 61).

Remarks: *Saemundssonina* (*Saemundssonina*) *platygaster* is a morphologically variable and frequently collected “head & neck” louse parasitic on a large range of shore birds. Timmermann (1969a: 240) subdivided it into several subspecies, which are recognised as valid by Price *et al.* (2003: 236). The only subspecies recorded in New

Zealand is *S. (S.) platygaster balati* from *Charadrius bicinctus* (see below). However, I regard all samples from other New Zealand hosts as *S. (S.) platygaster sensu lato* until a thorough revision of these populations is made. *Tringa incana*, *Himantopus novaezelandiae* and *Charadrius obscurus aquilonius* are new host records for *S. (S.) platygaster s. l.* in New Zealand (voucher specimens in MONZ).

***Saemundssonina (Saemundssonina) platygaster balati* Timmermann, 1969**

*Saemundssonina (Saemundssonina) platygaster balati* [sic] Timmermann, 1969a: 241.

*Saemundssonina platygaster* (Denny, 1842) *s. l.*; Pilgrim & Palma 1982: 20.

*Saemundssonina platygaster* (Denny, 1842); Murray *et al.* 1993: 962. In part.

*Saemundssonina (Saemundssonina) platygaster balati* Timmermann, 1969; Palma 1999: 380.

*Saemundssonina (Saemundssonina) platygaster balati* Timmermann, 1969; Price *et al.* 2003: 236.

*Saemundssonina (S.) p. balati* Timmermann, 1969; Palma 2010: 409.

Holotype ♂ in ANIC (Palma 1996b: 224).

Type host: *Charadrius bicinctus bicinctus* Jardine & Selby, 1827.

New Zealand host: *Charadrius bicinctus bicinctus* Jardine & Selby, 1827.

Other hosts: None.

New Zealand localities: AK, MC.

Geographic distribution: Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1993); Palma (1999); Palma (2010).

Other significant references: Palma (1996b: 224); Price *et al.* (2003).

Remarks: Although the type host of *Saemundssonina (Saemundssonina) platygaster balati* only breeds in New Zealand, the type locality of this louse species is in Australia (Palma 1996b: 224), where most birds migrate to in winter (Checklist Committee 2010: 217).

***Saemundssonina (Saemundssonina) pterodromae* Timmermann, 1959**

*Saemundssonina pterodromae* Timmermann, 1959a: 153, fig. 4.

*Saemundssonina halobaenae* Timmermann, 1965: 77. New synonymy.

*Saemundssonina pterodromae* Timmermann, 1959; Timmermann 1965: 78, fig. 16.

*Saemundssonina pterodromae* Timmermann, 1959; Wise 1977: 65.

*Saemundssonina pterodromae* Timmermann, 1959; Pilgrim & Palma 1982: 8.

*Saemundssonina pterodromae* Timmermann, 1959; Palma & Pilgrim 1983: 149.

*Saemundssonina (Saemundssonina) pterodromae* Timmermann, 1959; Palma & Horning 2002: 13, 16.

*Saemundssonina (S.) pterodromae* Timmermann, 1959; Palma 2010: 409.

Holotype ♂ in NHML.

Type host: *Lugensa brevirostris* (Lesson, 1833).

New Zealand hosts: *Lugensa brevirostris* (Lesson, 1833); *Halobaena caerulea* (J.F. Gmelin, 1789).

Other hosts: None.

New Zealand localities: AK, TK, WI, WN, NC, MC, SC, WD, Macquarie Island.

Geographic distribution: Southern Oceans; Atlantic, Indian and Pacific Oceans.

New Zealand references: Clay & Moreby (1970: 217); Gressitt (1970: 328); Wise (1977); Pilgrim & Palma (1982); Palma & Pilgrim (1983); Murray *et al.* (1990: 1369–1370); Palma (1996b: 225); Palma & Horning (2002); Palma (2010).

Other significant references: Timmermann (1965); Green & Palma (1991: 19, 26); Price *et al.* (2003: 237).

Remarks: *Saemundssonina (S.) pterodromae* is a frequently collected “head & neck” louse parasitic on two quite distinct petrel species. Timmermann (1965: 77) recorded “*Saemundssonina halobaenae* Edwards (MS)”, with *Halobaena caerulea* as the type host, and included a brief description of the louse. To the best of my knowledge, Edwards has not published any description of such species; therefore, the authorship must be given to Timmermann. Furthermore, the *Saemundssonina* population from *Halobaena caerulea* belongs to the species *Saemundssonina (S.) pterodromae* (see Pilgrim & Palma 1982: 9; Palma & Pilgrim 1983: 149), hence *Saemundssonina halobaenae* becomes a junior synonym of the former species.

**Saemundssonina (Saemundssonina) remota Timmermann, 1951**

- Saemundssonina lobaticeps remota* Timmermann, 1951b: 142.  
*Saemundssonina remota* Timmermann, 1951; Hopkins & Clay 1953: 445.  
*Saemundssonina remota* Timmermann, 1951; Ward & Downey 1973: 395.  
*Saemundssonina lobaticeps remota* Timmermann, 1951; Pilgrim & Palma 1982: 23.  
*Saemundssonina (Saemundssonina) remota* Timmermann, 1951; Price *et al.* 2003: 237.  
*Saemundssonina (Saemundssonina) lobaticeps remota* Timmermann, 1951; Murray *et al.* 2006a: 1965.  
*Saemundssonina (S.) lobaticeps remota* Timmermann, 1951; Palma 2010: 409.  
*Saemundssonina (Saemundssonina) remota* Timmermann, 1951; Palma & Peck 2013: 61.

Holotype ♂ in NHML.

Type host: *Anous stolidus stolidus* (Linnaeus, 1758).

New Zealand host: *Anous minutus minutus* Boie, 1844.

Other hosts: None.

New Zealand locality: KE.

Geographic distribution: Temperate and tropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Amerson & Emerson (1971: 19, 26); Ward & Downey (1973); Moreby (1976: 94); Forrester *et al.* (1995: 32); Palma (1996b: 223); Price *et al.* (2003); Palma & Peck (2013); Silva *et al.* (2014: 942).

Remarks: *Saemundssonina (S.) remota* is an infrequently collected “head & neck” louse parasitic on noddies. In agreement with Ward & Downey (1973: 395) and Price *et al.* (2003: 237), and contrary to Pilgrim & Palma (1982: 23) and Palma (2010: 409), I regard the populations of *Saemundssonina* from species of *Anous* as a full species, distinct from *Saemundssonina (S.) lobaticeps* (Giebel, 1874), a louse parasitic on species of *Chlidonias*.

**Saemundssonina (Saemundssonina) scolopacisphaeopodis scolopacisphaeopodis (Schrank, 1803)**

- Pediculus scolopacis phaeopodis* Schrank, 1803: 191.  
*Philoapterus armatus* Johnston & Harrison, 1912: 370, figs 7–10.  
*Pediculus scolopacisphaeopodis* Schrank, 1803; Harrison 1916: 18 (as junior synonym of “*Docophorus rotundatus* Nitzsch” [sic])  
*Saemundssonina scolopacis-phaeopodis* (Schrank, 1803); Timmermann 1949a: 20, figs 12–14.  
*Saemundssonina sc.-ph. scolopacis-phaeopodis* (Schrank, 1803); Timmermann 1951c: 392, figs 1–2.  
*Saemundssonina scolopacisphaeopodis* (Schrank, 1803); Hopkins & Clay 1952: 335.  
*Saemundssonina scolopacisphaeopodis* (Schrank, 1803); Clay & Hopkins 1960: 15, figs 19–26, pl. 3: figs 1, 3.  
*Saemundssonina scolopacis-phaeopodis scolopacis-phaeopodis* (Schrank, 1803); Timmermann 1969a: 233, fig. 1b.  
*Saemundssonina scolopacisphaeopodis* (Schrank, 1803); Watt 1971: 238, 243.  
*Saemundssonina scolopacisphaeopodis* (Schrank, 1803); Wise 1977: 65.  
*Saemundssonina scolopacisphaeopodis scolopacisphaeopodis* (Schrank, 1803); Pilgrim & Palma 1982: 20.  
*Saemundssonina (Saemundssonina) scolopacisphaeopodis scolopacisphaeopodis* (Schrank, 1803); Murray *et al.* 2006a: 1964.  
*Saemundssonina (S.) scolopacisphaeopodis scolopacisphaeopodis* (Schrank, 1803); Palma 2010: 409.

Neotype ♂ in NHML (Clay & Hopkins 1960: 18, pl. 3: fig. 1). Syntypes ♂♀ of *Philoapterus armatus* in MONZ (Palma *et al.* 1989: 45).

Type host: *Numenius phaeopus phaeopus* (Linnaeus, 1758).

New Zealand host: *Numenius phaeopus variegatus* (Scopoli, 1786).

Other host: *Numenius borealis* (J.R. Forster, 1772).

New Zealand localities: TK, KE.

Geographic distribution: Eurasia; Africa; Australasia.

New Zealand references: Johnston & Harrison (1912); Thompson (1939: 16); Watt (1971); Wise (1977); Pilgrim (1970: 75); Pilgrim & Palma (1982); Palma *et al.* (1989: 45); Murray *et al.* (2006a); Palma (2010).

Other significant references: Timmermann (1949a; 1951c); Emerson & Ward (1958: 59); Clay & Hopkins (1960); Hackman & Nyholm (1968: 77); Timmermann (1969a; 1971: 139); Ward & Downey (1973: 395); Moreby (1976: 93); Forrester *et al.* (1995: 26); Price *et al.* (2003: 237); Palma & Jensen (2005: 59, 64); Martín-Mateo (2009: 119).

Remarks: *Saemundssonina (S.) scolopacisphaeopodis scolopacisphaeopodis* is an infrequently collected “head & neck” louse parasitic on curlews and whimbrels. Clay & Hopkins (1960: 16) discussed the status of some possible junior

synonyms of *Saemundssonina (S.) scolopacisphaeopodis scolopacisphaeopodis*. Although the Asiatic whimbrel is a frequent visitor to New Zealand (Checklist Committee 2010: 201), its lice have been infrequently collected.

***Saemundssonina (Saemundssonina) stammeri* Timmermann, 1959**

*Saemundssonina stammeri* Timmermann, 1959a: 149, fig. 2.

*Saemundssonina stammeri* Timmermann, 1959; Timmermann 1965: 77, fig. 15.

*Saemundssonina stammeri* Timmermann, 1959; Clay & Moreby 1967: 165, 168, figs 155, 176.

*Saemundssonina stammeri* Timmermann, 1959; Horning *et al.* 1980: 7, 9.

*Saemundssonina stammeri* Timmermann, 1959; Pilgrim & Palma 1982: 7.

*Saemundssonina (Saemundssonina) stammeri* Timmermann, 1959; Price *et al.* 2003: 237.

*Saemundssonina (S.) stammeri* Timmermann, 1959; Palma 2010: 409.

Holotype ♂ in NHML.

Type host: *Daption capense* (Linnaeus, 1758).

New Zealand hosts: *Daption capense capense* (Linnaeus, 1758); *Daption capense australe* Mathews, 1913.

Other hosts: None.

New Zealand localities: ND, AK, HB, TK, WI, WN, NC, MC, SC, WD, SL, SN, AN.

Geographic distribution: Antarctica; Subantarctic Islands; Southern Oceans.

New Zealand references: Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990: 1369); Paterson *et al.* (1999: 222); Palma (2010).

Other significant references: Timmermann (1965); Clay & Moreby (1967); Green & Palma (1991: 19, 26); Palma (1996b: 225); Price *et al.* (2003).

Remarks: *Saemundssonina (S.) stammeri* is a frequently collected “head & neck” louse exclusively parasitic on Cape petrels.

***Saemundssonina (Saemundssonina) sterna* (Linnaeus, 1758)**

*Pediculus sterna* Linnaeus, 1758: 612.

*Ricinus sterna* (Linnaeus, 1758); Latreille 1804: 107.

*Philopterus sterna* Linnaeus, 1758 [sic]; Harrison 1916: 104 (qualified as “Identity uncertain”).

*Saemundssonina sterna* (Linnaeus, 1758); Clay 1949a: 4, figs 1–6, 16, 22–23.

*Saemundssonina sterna* (Linnaeus, 1758); Hopkins & Clay 1952: 336.

*Saemundssonina sterna* (Linnaeus, 1758); Clay & Moreby 1967: 164, figs 168–169, 171–172.

*Saemundssonina sterna* (Linnaeus, 1758); Pilgrim & Palma 1982: 23.

*Saemundssonina (Saemundssonina) sterna* (Linnaeus, 1758); Price *et al.* 2003: 237.

*Saemundssonina (Saemundssonina) sterna* (Linnaeus, 1758); Murray *et al.* 2006a: 1965.

*Saemundssonina (S.) sterna* (Linnaeus, 1758); Palma 2010: 409.

Neotype ♂ in NHML (Clay 1949a: 10).

Type host: *Sterna hirundo hirundo* Linnaeus, 1758.

New Zealand host: *Sterna striata* J.F. Gmelin, 1789.

Other hosts: *Sterna dougallii* Montagu, 1813; *Sterna vittata* J.F. Gmelin, 1789.

New Zealand localities: AK, WN, NN, KA, NC, MC, SC, WD, CO, DN, SL.

Geographic distribution: Eurasia; Africa; North America; Australasia; Southern Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay (1949a); Clay & Hopkins (1950: 245); Ward (1955: 85); Clay & Moreby (1967); Hackman & Nyholm (1968: 78); Green & Palma (1991: 19, 33); Forrester *et al.* (1995: 31); Palma (1996b: 225); Price *et al.* (2003); Hänel & Palma (2007: 113, 127, 131); Martín-Mateo (2009: 117).

Remarks: *Saemundssonina (S.) sterna* is a frequently collected “head & neck” louse parasitic on several tern species.

***Saemundssonina (Saemundssonina) thalassidromae incisa* Timmermann, 1950**

*Saemundssonina incisa* Timmermann, 1950: 7.

*Saemundssonina incisa* Timmermann, 1950; Timmermann 1951b: 142, fig. 1, right.

*Saemundssonina incisa* Timmermann, 1950; Hopkins & Clay 1952: 332.



- Saemundssonina incisa* Timmermann, 1950; Timmermann 1957a: 52, fig. 25.  
*Saemundssonina incisa* Timmermann, 1950; Timmermann 1965: 73, fig. 7.  
*Saemundssonina incisa* Timmermann, 1950; Palma 1999: 378.  
*Saemundssonina (Saemundssonina) incisa* Timmermann, 1950; Price *et al.* 2003: 234.  
*Saemundssonina (Saemundssonina) thalassidromae incisa* Timmermann, 1950; Palma & Jensen 2005: 59, 61.  
*Saemundssonina (S.) incisa* Timmermann, 1950; Palma 2010: 409.

Holotype ♂ in NHML (Vincent S. Smith pers. comm. February 2015).

Type host: *Oceanodroma leucorhoa* (Vieillot, 1818).

New Zealand host: *Oceanodroma leucorhoa leucorhoa* (Vieillot, 1818).

Other host: *Oceanodroma tethys* (Bonaparte, 1852).

New Zealand locality: ND.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Palma (1999); Palma (2010).

Other significant references: Timmermann (1951b; 1957a; 1965); Fowler & Hodson (1988: 47); Price *et al.* (2003); Palma & Jensen (2005: 59, 61).

Remarks: *Saemundssonina (S.) incisa* is an infrequently collected “head & neck” louse parasitic on some storm petrels. In agreement with Palma & Jensen (2005: 59) and contrary to Palma (1999: 378) and Price *et al.* (2003: 234), I regard this louse taxon as a subspecies of *Saemundssonina (Saemundssonina) thalassidromae*.

#### ***Saemundssonina (Saemundssonina) thompsoni* Timmermann, 1951**

- Saemundssonina thompsoni* Timmermann, 1951c: 396, figs 5–7 below.  
*Saemundssonina thompsoni* Timmermann, 1951; Hopkins & Clay 1953: 445.  
*Saemundssonina thompsoni* Timmermann, 1951; Tendeiro 1963: 96, photos 77–79.  
*Saemundssonina thompsoni* Timmermann, 1951; Pilgrim & Palma 1982: 20.  
*Saemundssonina (Saemundssonina) thompsoni* Timmermann, 1951; Price *et al.* 2003: 237.  
*Saemundssonina (Saemundssonina) thompsoni* Timmermann, 1951; Murray *et al.* 2006a: 1964.  
*Saemundssonina (S.) thompsoni* Timmermann, 1951; Palma 2010: 409.

Holotype ♂ in NHML (originally in the Thompson Collection, see George 1981: 88).

Type host: *Limosa limosa* (Linnaeus, 1758).

New Zealand host: *Limosa limosa melanuroides* Gould, 1846.

Other host: *Limosa limosa limosa* (Linnaeus, 1758).

New Zealand locality: AU.

Geographic distribution: Eurasia; Australasia.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (2006a); Palma (2010).

Other significant references: Tendeiro (1963); Price *et al.* (2003); Adam (2007: 180).

Remarks: *Saemundssonina (S.) thompsoni* is an infrequently collected “head & neck” louse parasitic on black-tailed godwits. *Limosa limosa melanuroides* is an uncommon annual visitor to New Zealand (Checklist Committee 2010: 203), with only a single record of *Saemundssonina (S.) thompsoni* from this country.

#### ***Saemundssonina (Saemundssonina) tringae* (O. Fabricius, 1780) *sensu lato***

- Pediculus tringae* O. Fabricius, 1780: 219.  
*Philoaterus (Docophorus) variabilis* Denny, 1842: 42, 71, pl. 3: fig. 4.  
*Docophorus alpinus* Giebel, 1874: 105.  
*Pediculus tringae* O. Fabricius, 1780; Harrison 1916: 19 (as “Not recognisable”).  
*Hastaeophorus alpinus* Giebel [sic]; Kéler 1936: 262, figs 2a,c.  
“*Philoaterus limosae*” Harrison, 1937: 21, 46 (not *Philoaterus (Docophorus) limosae* Denny, 1842).  
*Saemundssonina tringae* (O. Fabricius, 1780); Hopkins & Clay 1952: 336.  
*Saemundssonina variabilis* (Denny, 1842); Hopkins & Clay 1952: 336.  
*Saemundssonina tringae* (O. Fabricius, 1780); Clay & Hopkins 1954: 249, figs 40–44.  
*Saemundssonina* sp. [ex *Calidris canutus rogersi*]; Watson 1967: 74.  
*Saemundssonina* sp.; Gressitt 1970: 329.  
“*Saemundssonina limosae*” Wise, 1977: 65 (not *Philoaterus (Docophorus) limosae* Denny, 1842).

*Saemundssonina tringae* (O. Fabricius, 1780) s. l.; Pilgrim & Palma 1982: 21, 31, note 23.

*Saemundssonina* (*Saemundssonina*) sp.; Palma & Horning 2002: 13, 17, 22.

*Saemundssonina* (*Saemundssonina*) *tringae* (O. Fabricius, 1780); Price *et al.* 2003: 237.

*Saemundssonina* (*Saemundssonina*) *tringae tringae* (O. Fabricius, 1780); Murray *et al.* 2006a: 1964.

*Saemundssonina* (*S.*) *tringae* (O. Fabricius, 1780); Palma 2010: 409.

Neotype ♂ in NHML (Clay & Hopkins (1954: 250).

Type host: *Calidris maritima* (Brünnich, 1764).

New Zealand host: *Calidris canutus rogersi* (Mathews, 1913).

Other hosts: At least 14 other species of *Calidris* (Price *et al.* 2003: 237); *Arenaria interpres* (Linnaeus, 1758); *Arenaria melanocephala* (Vigors, 1829); *Eurynorhynchus pygmaeus* (Linnaeus, 1758); *Limnodromus griseus* (J.F. Gmelin, 1789); *Limnodromus scolopaceus* (Say, 1823); *Phalaropus lobatus* (Linnaeus, 1758); *Philomachus pugnax* (Linnaeus, 1758).

New Zealand localities: MC, CA, Macquarie Island.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Harrison (1937); Watson (1967); Gressitt (1970); Wise (1977); Pilgrim & Palma (1982); Palma & Horning (2002); Murray *et al.* (2006a); Palma (2010).

Other significant references: Clay & Hopkins (1954); Hackman & Nyholm (1968: 78); Martens (1974: 142); Hunter & Colwell (1994: 402); Palma (1996b: 226); Price *et al.* (2003).

Remarks: *Saemundssonina* (*S.*) *tringae* is a morphologically variable and infrequently collected “head & neck” louse parasitic on many shore bird species. Martens (1974: 142) divided this taxon into several subspecies which, without justification, have not been recognised as valid by Price *et al.* (2003: 238). Therefore, I qualify the material from *Calidris canutus rogersi* as “*sensu lato*”, recognising that there is a range of intraspecific variation that needs detailed study to determine the validity of the subspecies.

### ***Saemundssonina* (*Saemundssonina*) *uppalensis* (Rudow, 1870)**

*Docophorus uppalensis* Rudow, 1870: 455.

*Philopterus uppalensis* Rudow, 1870 [sic]; Harrison 1916: 106.

*Saemundssonina uppalensis* (Rudow, 1870); Hopkins 1951b: 374.

*Saemundssonina upoluensis* (Rudow, 1870); Hopkins & Clay 1952: 336. Unjustified emendation.

*Saemundssonina upolensis* [sic] (Rudow, 1870); Timmermann 1955: 516. Unjustified emendation.

*Saemundssonina upoluensis* (Rudow, 1870); Pilgrim & Palma 1982: 14. Unjustified emendation.

*Saemundssonina upoluensis* (Rudow, 1870); Murray *et al.* 1990: 1373. Unjustified emendation.

*Saemundssonina uppalensis* (Rudow, 1870); Forrester *et al.* 1995: 6.

*Saemundssonina* (*Saemundssonina*) *uppalensis* (Rudow, 1870); Price *et al.* 2003: 238.

*Saemundssonina* (*S.*) *uppalensis* (Rudow, 1870); Palma 2010: 409.

Type presumed lost. See Hopkins (1951: 374).

Type host: “*Phaethon aethereus*”, in error (see Hopkins 1951: 374).

New Zealand host: *Phaethon lepturus dorotheae* Mathews, 1913.

Other hosts: *Phaethon lepturus catesbyi* J.F. Brandt, 1838; *Phaethon lepturus lepturus* Daudin, 1802; *Phaethon lepturus fulvus* J.F. Brandt, 1838.

New Zealand locality: ND.

Geographic distribution: Tropical and subtropical regions of the Atlantic, Indian and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (2010).

Other significant references: Hopkins (1951b); Timmermann (1955); Amerson & Emerson (1971: 10, 27); Martín-Mateo (1992a: 49, figs 26–28); Forrester *et al.* (1995); Palma (1996b: 226); Price *et al.* (2003).

Remarks: *Saemundssonina* (*S.*) *uppalensis* is an infrequently collected “head & neck” louse parasitic on white-tailed tropicbirds. *Phaethon lepturus dorotheae* is a regular straggler to northern New Zealand (Checklist Committee 2010: 137).

Hopkins & Clay (1952: 336) emended the species epithet of this louse following Hopkins (1951: 374) who believed that the name referred to Upolu Island in the Samoan Archipelago, and not to “Uppala” as Rudow (1870:

455) had spelt it. However, regardless of the spelling of the type locality, the original species name must remain unaltered (see Palma 1999: 384, note C).

**Saemundssonina (Saemundssonina) species**

**New Record**

New Zealand host: *Pealeornis maoriana* Mathews, 1932.

New Zealand locality: CL.

Geographic distribution: New Zealand seas.

New Zealand reference: This paper.

Material examined and repository: 1♀ (1 sample, MONZ).

Remarks: The only available specimen could not be identified to species because it is a female (see Palma 2000: 127).

**Genus *Strigiphilus* Mjöberg, 1910**

*Strigiphilus* Mjöberg, 1910a. *Arkiv Zool.* 6(13): 132. Type species: *Docophorus heterocerus* Nitzsch [*in* Giebel], 1861a = *Strigiphilus goniodicerus* Eichler, 1949 (by original designation).

***Strigiphilus aitkeni* Clay, 1966**

*Strigiphilus aitkeni* Clay, 1966b: 12, figs 5, 7, 9, 11–12, pl. 2.

*Strigiphilus aitkeni* Clay, 1966; Wise 1977: 65.

*Strigiphilus aitkeni* Clay, 1966; Pilgrim & Palma 1982: 25.

*Strigiphilus aitkeni* Clay, 1966; Murray *et al.* 1999: 1241.

*Strigiphilus aitkeni* Clay, 1966; Palma 2010: 409.

Holotype ♂ in NHML.

Type host: *Tyto alba hellmayri* Griscom & Greenway, 1937.

New Zealand host: *Tyto alba delicatula* (Gould, 1837).

Other hosts: *Tyto alba javanica* (J.F. Gmelin, 1788); *Tyto alba tuidara* (J.E. Gray, 1829); *Tyto novaehollandiae novaehollandiae* (Stephens, 1826); *Tyto novaehollandiae castanops* (Gould, 1837); *Tyto capensis longimembris* (Jerdon, 1839).

New Zealand localities: MC, WD.

Geographic distribution: Americas; Asia; Australasia.

New Zealand references: Clay (1966b: 17, fig. 12); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1999); Palma (2010).

Other significant references: Clay (1976b: 540, 546); Green & Palma (1991: 20; 36); Palma (1996b: 227); Price *et al.* (2003: 239).

Remarks: *Strigiphilus aitkeni* is a “head & neck” louse parasitic on many species of barn owls. *Tyto alba delicatula* is a vagrant to New Zealand (Checklist Committee 2010: 268), with only two records of *Strigiphilus aitkeni* from this country.

***Strigiphilus cursitans* (Nitzsch [*in* Giebel], 1861)**

*Docophorus cursitans* Nitzsch [*in* Giebel], 1861a: 529.

*Philopterus cursitans* Nitzsch, [*in* Giebel], 1861 [sic]; Harrison 1916: 92.

*Philopterus cursitans* (Nitzsch, 1861) [sic]; Marples 1942: 245.

*Strigiphilus cursitans* (Nitzsch, 1861) [sic]; Hopkins & Clay 1952: 339.

*Strigiphilus cursitans* (Nitzsch, 1861) [sic]; Wise 1977: 65.

*Strigiphilus cursitans* (Nitzsch, 1861) [sic]; Pilgrim & Palma 1982: 25.

*Strigiphilus cursitans* (Nitzsch, 1861) [sic]; Murray *et al.* 1999: 1241.

*Strigiphilus cursitans* (Nitzsch [*in* Giebel], 1861); Price *et al.* 2003: 239.

*Strigiphilus cursitans* (Nitzsch, 1861) [sic]; Palma 2010: 409.

Syntypes ♂♂ presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Athene noctua* (Scopoli, 1769).

New Zealand host: *Athene noctua* (Scopoli, 1769).

Other hosts: *Athene noctua bactriana* Blyth, 1847; *Athene noctua glaux* (Savigny, 1809); *Athene noctua saharae* (Kleinschmidt, 1909); *Athene noctua somaliensis* Reichenow, 1905; *Athene noctua vidalii* Brehm, 1857; *Strix butleri* (Hume, 1878).

New Zealand localities: NC, MC, SC, WD, CO, DN, SL.

Geographic distribution: Europe; Asia; Africa; New Zealand.

New Zealand references: Marples (1942); Wise (1977); Pilgrim & Palma (1982); Clayton & Price (1984: 343, figs 1–2, 7, 11, 32, 52, 54); Murray *et al.* (1999); Paterson *et al.* (1999: 220); Palma (2010).

Other significant references: Clayton (1990: 260); Price *et al.* (2003: 239); Adam (2007: 181); Martín-Mateo (2009: 169).

Remarks: *Strigiphilus cursitans* is a “head & neck” louse parasitic on several small species of owls. *Strigiphilus cursitans* was introduced to New Zealand with little owls by human agency (Checklist Committee 2010: 267).

### ***Strigiphilus vapidus* Clay, 1977**

Figs 170–171

*Strigiphilus vapidus* Clay, 1977: 1, figs 1, 3, 5–6, 8.

*Strigiphilus vapidus* Clay, 1977; Pilgrim & Palma 1982: 25.

*Strigiphilus vapidus* Clay, 1977; Clayton & Price 1984: 345, figs 12, 33, 55.

*Strigiphilus vapidus* Clay, 1977; Murray *et al.* 1999: 1241.

*Strigiphilus vapidus* Clay, 1977; Palma 2010: 409.

Holotype ♂ in ANIC.

Type host: *Ninox boobook ocellata* (Bonaparte, 1850).

New Zealand host: *Ninox novaeseelandiae novaeseelandiae* (J.F. Gmelin, 1788).

Other host: *Ninox novaeseelandiae leucopsis* (Gould, 1838).

New Zealand localities: WO, BP, BP, TO, HB, WA, WN, SD, MB, BR, WD.

Geographic distribution: Australasia.

New Zealand references: Clay (1977: 2); Pilgrim & Palma (1982); Clayton & Price (1984); Murray *et al.* (1999); Palma (2010).

Other significant references: Clayton (1990: 260); Green & Palma (1991: 20, 36); Palma (1996b: 227); Price *et al.* (2003: 240).

Remarks: *Strigiphilus vapidus* is a “head & neck” louse belonging to the “*Strigiphilus cursitans* group” of species (Clayton & Price 1984) and restricted to Australasian owls.

### ***Strigiphilus* species**

*Strigiphilus* sp.; Pilgrim & Palma 1982: 25.

*Strigiphilus*; Murray *et al.* 1999: 1241.

New Zealand host: *Sceloglaux albifacies albifacies* (Gray, 1844).

Other hosts: None

New Zealand locality: CO, DN.

Geographic distribution: New Zealand.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1999).

Other significant references: None.

Remarks: This species is represented by a single male louse belonging to the “*Strigiphilus cursitans* group” as defined by Clayton & Price (1984). It is morphologically close to *St. vapidus* Clay, 1977, but it exhibits some differences in head shape and details of the genitalia. However, considering that the host is almost certainly extinct (Checklist Committee 2010: 266), the probability of finding more specimens to confirm its identity is extremely low.

### **Genus *Sturnidoecus* Eichler, 1944**

*Sturnidoecus* Eichler, 1944b. *Stettin. Entomol. Zeit.* 105: 81. Type species: “*Docophorus leontodon* Nitzsch, *sensu* Piaget, 1880” = *Sturnidoecus sturni* (Schrank, 1776) (by original designation).

***Sturnidoecus sturni* (Schrank, 1776)**

Figs 172–173

- Pediculus sturni* Schrank, 1776: 118, pl. 5: figs 11–14.  
*Philopterus (Docophorus) leontodon* Nitzsch, 1818: 290.  
*Philopterus leontodon* Nitzsch, 1818; Johnston & Harrison 1912: 368.  
*Philopterus sturni* (Schrank, 1776); Thompson 1939: 71.  
*Sturnidoecus sturni* (Schrank, 1776); Hopkins & Clay 1952: 345.  
*Sturnidoecus sturni* (Schrank, 1776); Watt 1971: 238, 244, fig. 12.  
*Sturnidoecus sturni* (Schrank, 1776); Pilgrim & Palma 1982: 27.  
*Sturnidoecus sturni* (Schrank, 1776); Murray *et al.* 2006b: 1960.  
*Sturnidoecus sturni* (Schrank, 1776); Palma 2010: 409.

Neotype ♂ in NHML (Clay &amp; Hopkins 1954: 232, pl. 10: fig. 5).

Type host: *Sturnus vulgaris vulgaris* Linnaeus, 1758.New Zealand host: *Sturnus vulgaris vulgaris* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: HB, WI, WN, NC, MC, SC, WD, KE, Norfolk Island.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Johnston & Harrison (1912); Nelson (1969: 199); Watt (1971); Wise (1977: 65); Pilgrim & Palma (1982); Paterson *et al.* (1999: 220); Murray *et al.* (2006b); Palma (2010).Other significant references: Thompson (1939); Eichler (1944b: 81); Séguy (1944: 253, figs 381–384); Clay & Hopkins (1954: 232, fig. 12, pl. 10: fig. 5); Kettle (1983: 403); Green & Palma (1991: 20, 41); Butler & O'Connor (1994: 457); Palma (1996b: 228); Price *et al.* (2003: 243); Johnson & Clayton (2003: 462); Palma & Jensen (2005: 59, 68); Adam (2007: 185); Martín-Mateo (2009: 205); Bartlow *et al.* (2016: 222).Remarks: *Sturnidoecus sturni* was introduced to New Zealand with starlings by human agency (Checklist Committee 2010: 315).***Sturnidoecus* species**

- Sturnidoecus* sp.; Pilgrim & Palma 1982: 27.  
*Sturnidoecus* sp.; Paterson *et al.* 1999: 220.  
*Sturnidoecus* sp.; Murray *et al.* 2006b: 1960.

New Zealand host: *Acridotheres tristis* (Linnaeus, 1766).

Other hosts: None.

New Zealand localities: ND, CL.

Geographic distribution: Asia; Australia; South Africa; islands of the Pacific and Indian Oceans.

New Zealand references: Pilgrim & Palma (1982); Paterson *et al.* (1999); Murray *et al.* (2006b).Remarks: The two available records of *Sturnidoecus* from *A. tristis* could not be identified to species because the samples contain females only (voucher specimens in MONZ). *Acridotheres tristis* and its lice were introduced to New Zealand by human agency (Checklist Committee 2010: 315)**Genus *Trabeculus* Rudow, 1866**

- Trabeculus* Rudow, 1866b. *Z. ges. NatWiss.* 27: 466. Type species: *Trabeculus schillingii* Rudow, 1866 (by monotypy).  
*Oncophorus* Rudow, 1870. *Z. ges. NatWiss.* 35: 475. Type species: *Oncophorus schillingii* Rudow, 1870 = *Trabeculus schillingii* Rudow, 1866b (by monotypy).  
*Giebelia* Kellogg, 1896a. *Proc. Calif. Acad. Sci.* 6: 137. Type species: *Giebelia mirabilis* Kellogg, 1896a = *Trabeculus mirabilis* (Kellogg, 1896) (by monotypy).  
*Mackayia* Waterston, 1912a. *Scott. Nat.* 11: 252. Type species: *Mackayia dimorpha* Waterston, 1912a = *Trabeculus aviator* (Evans, 1912) (by monotypy).  
*Cecalymenus* Enderlein, 1917. *Zool. Anz.* 49: 242. Type species: *Cecalymenus oestrelatae* Enderlein, 1917 = *Trabeculus schillingii* Rudow, 1866b (by original designation).

***Trabeculus aviator* (Evans, 1912)**

- Docophorus*, ? sp. n. Evans, 1912 (Oct.): 268, 270, 276, fig. 4.  
*Docophorus aviator* Evans, 1912 (Oct.): 268, 270, 276, fig. 4.  
*Mackayia dimorpha* Waterston, 1912a (Nov.): 252, figs 1–6.  
*Trabeculus aviator* Evans, 1912 [sic]; Harrison 1916: 144. Listed as *nomen nudum*.  
*Trabeculus dimorphus* Waterston, 1912 [sic]; Harrison 1916: 144.  
*Trabeculus dimorphus* (Waterston, 1912); Timmermann 1951b: 142, fig. 1, left.  
*Trabeculus aviator* (Evans, 1912); Hopkins & Clay 1952: 348.  
*Trabeculus aviator* (Evans, 1912); Timmermann 1959b: 498, fig. 10.  
*Trabeculus aviator* (Evans, 1912); Timmermann 1965: 133, fig. 74, pl. 2: figs 3–4.  
*Trabeculus aviator* (Evans, 1912); Pilgrim & Palma 1982: 12.  
*Trabeculus aviator* (Evans, 1912); Price *et al.* 2003: 244.  
*Trabeculus aviator* (Evans, 1912); Palma 2010: 409.

Syntypes ♀♀, probably lost (Robert McGowan pers. comm. February 2014).

Type host: *Puffinus puffinus* (Brünnich, 1764).

New Zealand host: *Puffinus puffinus* (Brünnich, 1764).

Other hosts: None

New Zealand localities: WN, WI.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990: 1371); Palma (2010).

Other significant references: Timmermann (1951b); Timmermann (1959b; 1965); Fowler & Shaw (1990: 15); Butler & O'Connor (1994: 457); Price *et al.* (2003); Palma & Jensen (2005: 59–60).

Remarks: *Puffinus puffinus* is a straggler to Australasia, recorded in New Zealand three times (Checklist Committee 2010: 119), but with only two records of *Trabeculus aviator* from this country.

***Trabeculus flemingi* Timmermann, 1959**

- Trabeculus flemingi* Timmermann, 1959b: 497, fig. 9.  
*Trabeculus flemingi* Timmermann, 1959; Timmermann 1965: 133, fig. 73.  
*Trabeculus flemingi* Timmermann, 1959; Wise 1977: 65.  
*Trabeculus flemingi* Timmermann, 1959; Pilgrim & Palma 1982: 12.  
*Trabeculus flemingi* Timmermann, 1959; Murray *et al.* 1990: 1371.  
*Trabeculus flemingi* Timmermann, 1959; Price *et al.* 2003: 244.  
*Trabeculus flemingi* Timmermann, 1959; Palma 2010: 409.

Holotype ♂ in CMNZ (Nicholls *et al.* 1998: 30).

Type host: *Puffinus huttoni* Mathews, 1912.

New Zealand host: *Puffinus huttoni* Mathews, 1912.

Other hosts: None.

New Zealand localities: AK, WN, KA, NC, MC, SC.

Geographic distribution: Pacific Ocean.

New Zealand references: Timmermann (1959; 1965); Pilgrim (1970: 75); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Nicholls *et al.* (1998: 30); Page *et al.* (2004: 634, 638, 650); Galloway (2005: 16); Buckley *et al.* (2012: App. 2).

Other significant reference: Price *et al.* (2003).

Remarks: *Trabeculus flemingi* is an endemic and “at risk” species (Buckley *et al.* 2012), exclusively parasitic on Hutton’s shearwaters.

***Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912)**

Figs 174–175

- Philopterus fuscoclypeatus* Johnston & Harrison, 1912: 368, fig. 4.  
*Giebelia fuscoclypeata* Johnston & Harrison, 1912 [sic]; Harrison 1916: 144.  
*Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912); Hopkins & Clay 1952: 349.  
*Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912); Timmermann 1959b: 491, fig. 5.  
*Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912); Timmermann 1965: 129, fig. 69.

*Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912); Watt 1971: 238, 242, fig. 11.

*Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912); Wise 1977: 65.

*Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912); Pilgrim & Palma 1982: 9.

*Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912); Palma 2010: 409.

Holotype nymph in MONZ (Palma *et al.* 1989: 45).

Type host: *Pterodroma neglecta neglecta* (Schlegel, 1863).

New Zealand host: *Pterodroma neglecta neglecta* (Schlegel, 1863).

Other hosts: *Pterodroma hasitata hasitata* (Kuhl, 1820); *Pterodroma arminjoniana* (Giglioli & Salvadori, 1869); *Pterodroma heraldica* (Salvin, 1888); *Pterodroma atrata* (Mathews, 1912); *Pterodroma alba* (J.F. Gmelin, 1789); *Pterodroma phaeopygia* (Salvin, 1876).

New Zealand localities: ND, KE.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Johnston & Harrison (1912); Thompson (1939: 213); Pilgrim (1970: 75); Watt (1971); Wise (1977); Pilgrim & Palma (1982); Palma *et al.* (1989: 45); Murray *et al.* (1990: 1369); Palma (2010).

Other significant references: Timmermann (1959; 1965); Zonfrillo (1993: 327); Price *et al.* (2003: 244); Palma & Peck (2013: 63).

Remarks: The population of *Trabeculus* from *Pterodroma phaeopygia* has been regarded as *Trabeculus fuscoclypeatus sensu lato* (see Palma & Peck 2013: 63). *Pterodroma heraldica* and *Pt. atrata* are new host records for this louse species (voucher specimens in MONZ).

#### ***Trabeculus hexakon* (Waterston, 1914)**

*Giebelia hexakon* Waterston, 1914: 291, pl. 25: figs 7, 11, pl. 26: fig. 14.

*Giebelia hexakon* Waterston, 1914; Harrison 1916: 144.

*Trabeculus hexakon* (Waterston, 1914); Hopkins & Clay 1952: 349. Unjustified emendation.

*Trabeculus hexakon* (Waterston, 1914); Timmermann 1959b: 487, figs 1, 4. In part.

*Trabeculus hexakon* (Waterston, 1914); Timmermann 1965: 128, fig. 68, pl. 2: figs 1–2.

*Trabeculus hexakon* (Waterston, 1914); Clay & Moreby 1967: 166, 168, figs 80, 152. Unjustified emendation.

*Trabeculus hexakon* (Waterston, 1914); Wise 1977: 65. In part. Unjustified emendation.

*Trabeculus hexakon* (Waterston, 1914); Pilgrim & Palma 1982: 11.

*Trabeculus hexakon* (Waterston, 1914); Murray *et al.* 1990: 1371.

*Trabeculus hexakon* (Waterston, 1914); Marris 2000: 188.

*Trabeculus hexakon* (Waterston, 1914) *sensu lato*; Palma & Horning 2002: 13, 16. In part.

*Trabeculus hexakon* (Waterston, 1914); Price *et al.* 2003: 244. In part.

*Trabeculus hexakon* (Waterston, 1914); Palma 2010: 409. In part.

Syntypes ♂♀ in SAMS (see Remarks).

Type host: *Procellaria aequinoctialis* Linnaeus, 1758.

New Zealand hosts: *Procellaria aequinoctialis* Linnaeus, 1758; *Procellaria westlandica* Falla, 1946; *Procellaria parkinsoni* G.R. Gray, 1862; *Procellaria cinerea* J.F. Gmelin, 1789.

Other hosts: None.

New Zealand localities: ND, AK, CL, WO, WN, WD, SN, AN, AU, CA, Macquarie Island.

Geographic distribution: Pacific and Atlantic Oceans.

New Zealand references: Gressitt (1970: 328); Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Paterson *et al.* (1999: 222); Marris (2000); Palma & Horning (2002); Palma (2010).

Other significant references: Séguy (1953: 587, fig. 44); Timmermann (1959; 1965); Clay & Moreby (1967); Palma (1996b: 229); Price *et al.* (2003: 244); Page *et al.* (2004: 637, 650).

Remarks: The original syntypes of *Trabeculus hexakon* were one male and four females (Waterston 1914: 291), of which I have examined the male (slide 200) and one female (slide 202). Although the male is labelled as “TYPE”, it can not be the holotype because Waterston (1914) did not mention any type, and I am not aware of a lectotype having been designated.

A record of “*Giebelia hexakon*” from *Pachyptila desolata* in Harrison (1937: 37) was fully discussed and listed as “*Trabeculus* sp.” by Palma & Horning (2002: 14, 22, note 24).

***Trabeculus hexakon* (Waterston, 1914) *sensu lato***

*Giebelia hexakon* Waterston, 1914; Guimarães 1943: 430, fig. 4.

*Giebelia hexakon* Waterston, 1914; Kéler 1952: 205, figs 1–3.

*Trabeculus hexakon* (Waterston, 1914); Timmermann 1959b: 488. In part

*Trabeculus hexacon* [sic] (Waterston, 1914) *sensu lato*; Clay 1964a: 232.

*Trabeculus hexacon* [sic] (Waterston, 1914); Gressitt 1964: 538.

“*Saemundssonina* sp.” Watt, 1971: 238, 242 (not *Saemundssonina* Timmermann, 1936).

*Trabeculus hexacon* [sic] (Waterston, 1914); Watt, 1971: 238, 242.

*Trabeculus hexacon* [sic] (Waterston, 1914); Wise 1977: 65. In part.

*Trabeculus hexakon* (Waterston, 1914) *s. l.*; Horning *et al.* 1980: 7, 10.

*Trabeculus hexakon* (Waterston, 1914) *s. l.*; Pilgrim & Palma 1982: 8–9, 11–12.

*Trabeculus hexakon* (Waterston, 1914); Murray *et al.* 1990: 1370–1371.

*Trabeculus hexakon* (Waterston, 1914) *sensu lato*; Palma & Horning 2002: 13, 15. In part.

*Trabeculus hexakon* (Waterston, 1914); Price *et al.* 2003: 244. In part.

*Trabeculus hexakon* (Waterston, 1914); Palma 2010: 409. In part.

New Zealand hosts: *Pterodroma externa* (Salvin, 1875); *Pterodroma nigripennis* (Rothschild, 1893); *Pterodroma axillaris* (Salvin, 1893); *Pterodroma cookii* (G.R. Gray, 1843); *Pterodroma longirostris* (Stejneger, 1888); *Pterodroma pycrofti* Falla, 1933; *Pterodroma leucoptera caledonica* Imber & Jenkins, 1981; *Puffinus pacificus pacificus* (J.F. Gmelin, 1789); *Puffinus pacificus chlororhynchus* Lesson, 1831; *Puffinus carneipes* Gould, 1844; *Puffinus griseus* (J.F. Gmelin, 1789); *Puffinus tenuirostris* (Temminck, 1835).

Other hosts: *Pterodroma incerta* (Schlegel, 1863); *Puffinus creatopus* Coues, 1864; *Puffinus gravis* (O’Reilly, 1818).

New Zealand localities: ND, AK, CL, BP, WA, WN, NC, MC, SC, KA, WD, CO, DN, SI, KE, Norfolk Island, CH, SN, AN, Macquarie Island.

Geographic distribution: Pacific and Atlantic Oceans.

New Zealand references: Clay (1964a); Gressitt (1964); Nelson (1969: 199); Clay & Moreby (1970: 218); Gressitt (1970: 329); Watt (1971); Wise (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b: 229); Palma & Horning (2002); Page *et al.* (2004: 638, 650); Palma (2010).

Other significant references: Guimarães (1943); Kéler (1952); Séguy (1953: 587); Ward & Downey (1973: 395); Bourgeois & Threlfall (1979: 1356); Green & Palma (1991: 20, 27); Furness & Palma (1992: 35, 39); Forrester *et al.* (1995: 6); Foster *et al.* (1996: 85); Price *et al.* (2003); Palma & Jensen (2005: 59–60); Hänel & Palma (2007: 113, 128, 130); Martín-Mateo (2009: 139, fig. 31).

Remarks: The populations of *Trabeculus* from species of *Pterodroma* and *Puffinus* listed above are qualified as *sensu lato* because they have subtle but clear and consistent differences from the population of *Trabeculus hexakon* from the type host, and from the other species of *Procellaria*. The entire genus *Trabeculus* is in need of a systematic revision to evaluate those differences and reflect them into a proper nomenclatorial structure.

A record of “4 nymphs” of “*Saemundssonina* sp.” from *Pterodroma nigripennis* in Watt (1971: 238, 242) is a misidentification of *Trabeculus hexakon sensu lato* (see Pilgrim & Palma 1982: 30, note 13).

***Trabeculus mirabilis* (Kellogg, 1896)**

*Giebelia mirabilis* Kellogg, 1896a: 138, pl. 11: figs 7–8.

*Giebelia mirabilis* Kellogg, 1896; Kellogg 1908: 34, fig. 5.

*Giebelia mirabilis* Kellogg, 1896; Harrison 1916: 144.

*Trabeculus mirabilis* (Kellogg, 1896); Hopkins & Clay 1952: 349.

*Trabeculus mirabilis* (Kellogg, 1896); Timmermann 1959b: 496, fig. 8.

*Trabeculus mirabilis* (Kellogg, 1896); Timmermann 1965: 132, fig. 72.

*Trabeculus mirabilis* Kellogg, 1896 *s. l.*; Pilgrim & Palma 1982: 12.

*Trabeculus mirabilis* (Kellogg, 1896); Murray *et al.* 1990: 1371.

*Trabeculus mirabilis* (Kellogg, 1896); Marris 2000: 188.

*Trabeculus mirabilis* (Kellogg, 1896); Price *et al.* 2003: 245.

*Trabeculus mirabilis* (Kellogg, 1896); Palma 2010: 409.

Lectotype ♂ in EMEC (Palma & Peck 2013: 64).

Type host: *Puffinus opisthomelas* Coues, 1864.



New Zealand hosts: *Puffinus newelli* Henshaw, 1900; *Puffinus assimilis kermadecensis* Murphy, 1927; *Puffinus assimilis haurakiensis* Fleming & Serventy, 1943; *Puffinus elegans* Giglioli & Salvadori, 1869.

Other hosts: *Puffinus lherminieri bannermani* Mathews & Iredale, 1915; *Puffinus lherminieri lherminieri* Lesson, 1839; *Puffinus lherminieri subalaris* Ridgway, 1897; *Puffinus assimilis baroli* (Bonaparte, 1857); *Puffinus assimilis boydi* Mathews, 1912.

New Zealand localities: ND, AK, CL, BP, WI, WA, WN, WD, KE, CH, AN.

Geographic distribution: Atlantic and Pacific Oceans.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1999: 376, 383, note 2); Marris (2000); Palma (2010).

Other significant references: Timmermann (1959b; 1965); Palma (1996b: 229); Furness & Palma (1992: 35, 39); Forrester *et al.* (1995: 6); Price *et al.* (2003); Page *et al.* (2004: 638, 650); Hänel & Palma (2007: 113, 128, 129); Palma & Peck (2013: 64).

Remarks: Pilgrim & Palma (1982: 12) regarded the populations of *Trabeculus mirabilis* from three subspecies of *Puffinus assimilis* as somewhat different from that of the type host, and qualified them as *sensu lato*; however, my examination of more samples from those and other hosts, including the type host, shows that making such difference is not warranted.

### ***Trabeculus schillingi* Rudow, 1866**

*Trabeculus Schillingi* [sic] Rudow, 1866b: 467.

*Oncophorus Schillingii* [sic] Rudow, 1870: 476.

*Docophorus Schillingi* [sic] (Rudow, 1866); Giebel 1874: 121.

*Oncophorus Schillingi* [sic] Rudow, 1870; Piaget 1880: 221.

*Mackayia heteracanthus* Waterston, 1912b: 258.

*Mackayia heteracanthus* Waterston, 1912; Waterston 1914: 292, pl. 25: fig. 8, pl. 26: figs 13, 16, 18.

*Trabeculus schillingi* Rudow, 1866; Harrison 1916: 144.

*Cecalymenus oestrelatae* Enderlein, 1917: 242, figs 2–4.

*Trabeculus heteracanthus* (Waterston, 1912); Hopkins & Clay 1952: 349.

*Trabeculus schillingi* Rudow, 1866; Hopkins & Clay 1952: 349.

*Trabeculus schillingi* Rudow, 1866; Timmermann 1959b: 492, fig. 7c. In part.

*Trabeculus schillingi* Rudow, 1866; Timmermann 1965: 130, fig. 71c. In part.

*Trabeculus schillingi* Rudow, 1866; Clay & Moreby 1970: 218. In part.

*Trabeculus schillingi* Rudow, 1866; Wise 1977: 66. In part.

*Trabeculus schillingi* Rudow, 1866; Pilgrim & Palma 1982: 8.

*Trabeculus schillingi* Rudow, 1866; Murray *et al.* 1990: 1369. In part.

*Trabeculus schillingi* Rudow, 1866; Marris 2000: 188. In part.

*Trabeculus schillingi* Rudow, 1866; Price *et al.* 2003: 245. In part.

*Trabeculus schillingi* Rudow, 1866; Palma 2010: 409. In part.

Syntypes ♂♀, presumed lost. See Palma (1996b: 229).

Type host: *Pterodroma mollis* (Gould, 1844).

New Zealand host: *Pterodroma mollis* (Gould, 1844).

Other hosts: *Pterodroma madeira* Mathews, 1934; *Pterodroma deserta* Mathews, 1934; *Pterodroma hasitata caribbaea* Carte, 1866.

New Zealand localities: WN, AN.

Geographic distribution: Pacific, Atlantic and Indian Oceans.

New Zealand references: Wise (1977); Pilgrim & Palma (1982); Murray *et al.* (1990); Marris (2000); Palma (2010).

Other significant references: Waterston (1914: 292, pl. 25: fig. 8, pl. 26: figs 13, 16, 18); Harrison (1916); Thompson (1935a: 558); Eichler (1941a: 354, fig. 16); Kéler (1957c: fig. 29); Timmermann (1959; 1965); Clay & Moreby (1967); Clay & Moreby (1970); Zonfrillo (1993: 327); Palma (1996b: 229); Furness & Palma (1992: 35, 41); Price *et al.* (2003); Hänel & Palma (2007: 113, 128, 130).

Remarks: See under *Trabeculus schillingi* Rudow, 1866 *sensu lato* below.

***Trabeculus schillingi* Rudow, 1866 *sensu lato***

*Trabeculus schillingi* Rudow, 1866b; Harrison 1937: 37.

*Trabeculus schillingi* Rudow, 1866; Timmermann 1959: 492, figs 2, 6, 7a,b. In part.

*Trabeculus schillingi* Rudow, 1866; Timmermann 1965: 130, figs 4, 67a, 70, 71a,b. In part.

*Trabeculus heteracanthus* (Waterston, 1912); Clay & Moreby 1967: 166, 168.

*Trabeculus schillingi* Rudow, 1866 *sens. lat.*; Clay & Moreby 1967: 166, 168, 191, figs 150–151.

*Trabeculus schillingi* Rudow, 1866; Clay & Moreby 1970: 218. In part.

*Trabeculus schillingi* Rudow, 1866; Wise 1977: 66. In part.

*Trabeculus schillingi* Rudow, 1866; Palma & Pilgrim 1977: 290.

*Trabeculus schillingi* Rudow, 1866b *s. l.*; Horning *et al.* 1980: 7, 9.

*Trabeculus schillingi* Rudow, 1866 *s. l.*; Pilgrim & Palma 1982: 8.

*Trabeculus schillingi* Rudow, 1866; Murray *et al.* 1990: 1369. In part.

*Trabeculus schillingi* Rudow, 1866; Palma 1999: 378.

*Trabeculus schillingi* Rudow, 1866; Marris 2000: 188. In part.

*Trabeculus schillingi* Rudow, 1866; Palma & Horning 2002: 13, 16.

*Trabeculus schillingi* Rudow, 1866; Price *et al.* 2003: 245. In part.

*Trabeculus schillingi* Rudow, 1866; Palma 2010: 409. In part.

New Zealand hosts: *Pterodroma macroptera gouldi* (Hutton, 1869); *Pterodroma lessonii* (Garnot, 1826); *Pterodroma solandri* (Gould, 1844); *Pterodroma inexpectata* (J.R. Forster, 1844).

Other hosts: *Pterodroma incerta* (Schlegel, 1863); *Pterodroma ultima* Murphy, 1949.

New Zealand localities: ND, AK, BP, TK, WI, WN, NC, MC, SC, WD, SI, Norfolk Island, SN, AU, AN, Macquarie Island.

Geographic distribution: Pacific and Atlantic Oceans.

New Zealand references: Harrison (1937); Clay & Moreby (1970); Gressitt (1970: 329); Wise (1977); Palma & Pilgrim (1977); Horning *et al.* (1980); Pilgrim & Palma (1982); Murray *et al.* (1990); Palma (1996b: 229); Palma (1999); Marris (2000); Palma & Horning (2002); Page *et al.* (2004: 638, 650); Palma (2010).

Other significant references: Timmermann (1959; 1965); Clay & Moreby (1967); Green & Palma (1991: 20, 26); Furness & Palma (1992: 35, 40); Price *et al.* (2003); Hänel & Palma (2007: 113, 128, 130).

Remarks: The populations of *Trabeculus* from the species of *Pterodroma* listed above are qualified as *sensu lato* because they have subtle but clear and consistent differences from the population of *Trabeculus schillingi* from the type host (e.g. Timmermann 1965: 130, fig. 71). The entire genus *Trabeculus* is in need of a systematic revision to evaluate those differences and reflect them into a proper nomenclatorial structure.

*Pterodroma ultima* is a new host record for *Trabeculus schillingi sensu lato* (voucher specimens in MONZ).

***Trabeculus* species 1**

*Trabeculus* sp.; Pilgrim & Palma 1982: 8.

*Trabeculus* sp.; Murray *et al.* 1990: 1370.

New Zealand host: *Pterodroma cervicalis* (Salvin, 1891).

New Zealand localities: BP, KE.

Geographic distribution: Pacific Ocean.

New Zealand references: Pilgrim & Palma (1982); Murray *et al.* (1990).

Other significant reference: Timmermann (1959).

Remarks: The population of *Trabeculus* from *Pterodroma cervicalis* represents an undescribed, unnamed species (voucher specimens in MONZ).

***Trabeculus* species 2**

*Trabeculus* undescribed species; Taylor & Tennyson 1994: 288.

*Trabeculus* sp.; Palma 1999: 375.

New Zealand host: *Puffinus nativitatis* Streets, 1877.

New Zealand locality: KE.

Geographic distribution: Pacific Ocean.

New Zealand references: Taylor & Tennyson (1994); Palma (1999).

Other significant reference: Timmermann (1959).

Remarks: The population of *Trabeculus* from *Puffinus nativitatis* represents an undescribed, unnamed species (voucher specimens in MONZ). This petrel is a rare vagrant to New Zealand (Checklist Committee 2010: 119), with a single record of *Trabeculus* from this country. Records of “*Trabeculus hexakon*” on *Puffinus nativitatis* from other Pacific Ocean localities in Amerson & Emerson (1971: 8, 27) and in Ward & Downey (1973: 395) are misidentifications.

## Family TRICHODECTIDAE Kellogg, 1896

Trichodectidae Kellogg, 1896a. *Proc. Calif. Acad. Sci.* 6: 63. Type genus: *Trichodectes* Nitzsch, 1818.

### Genus *Bovicola* Ewing, 1929

#### Subgenus *Bovicola* Ewing, 1929

*Bovicola* Ewing, 1929. *Manual External Parasites*: 121, 123, 193. Type species: *Trichodectes caprae* Gurlt, 1843 = *Bovicola (Bovicola) caprae* (Gurlt, 1843) (by original designation).

*Rhabdopedilon* Kéler, 1938a. *Nova Acat Leop.-Carol. (N.F.)* 5(32): 453. Type species: *Trichodectes longicornis* Nitzsch, 1818 = *Bovicola (Bovicola) longicornis* (Nitzsch, 1818) (by original designation).

#### *Bovicola (Bovicola) bovis* (Linnaeus, 1758)

*Pediculus bovis* Linnaeus, 1758: 611.

*Trichodectes scalaris* Nitzsch, 1818: 296. Unnecessary *nomen novum* for *Pediculus bovis* Linnaeus, 1758.

*Trichodectes bovis* Linnaeus, 1758 [sic]; Harrison 1916: 69.

*Trichodectes scalaris* Nitzsch; Thomson 1922: 269.

*Trichodectes bovis* L. [sic]; Tillyard 1926: 134.

*Bovicola bovis* (Linnaeus, 1758); Kéler 1938a: 450, fig. 34.

*Damalinia (Bovicola) bovis* (Linnaeus, 1758); Hopkins 1949b: 527.

*Damalinia bovis* (Linnaeus, 1758); Hopkins & Clay 1952: 103.

*Damalinia bovis* (L.); Helson 1956: 13, 16.

*Damalinia bovis* (Linnaeus, 1758); Wise 1977: 57.

*Bovicola (Bovicola) bovis* (Linnaeus, 1758); Lyal 1985a: 251, figs 3, 48, 51.

*Bovicola (Damalinia) bovis* [sic]; Watson *et al.* 1996: 401.

*Bovicola (Bovicola) bovis* (Linnaeus, 1758); Tenquist & Charleston 2001: 486.

*Bovicola (Bovicola) bovis* (Linnaeus, 1758); Palma 2010: 409.

Neotype ♀ in NHML (Clay & Hopkins 1950: 227).

Type host: *Bos taurus* Linnaeus, 1758.

New Zealand host: *Bos taurus* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: HB, WN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Thomson (1922); Tillyard (1926); Helson (1956); Whitten (1970: 146); Helson (1970: 81); Whitten (1971: 161); Buchanan & Coles (1971: 197); Heath (1973: 330); Kettle (1974a); Kettle & Lukies (1974); Heath (1976: 57); Wise (1977); Tenquist (1977: 286); Chalmers & Charleston (1980a,b,c); Tenquist & Charleston (1981: 262); Watson *et al.* (1996); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Werneck (1936: 550, figs 183–185); Kéler (1938); Séguy (1944: 388, figs 574–578); Webb (1946: 55); Hopkins (1949b); Werneck (1950: 63); Clay & Hopkins (1950: 227); Hopkins (1960: 82); Emerson & Price (1975: 68, figs 197–198); Rudolph (1983: 16); Lyal (1985a); Price (1987: 220, figs 22.26–22.28); Butler & O'Connor (1994: 451); Barker (1996: 231); Price *et al.* (2003: 253); Palma & Jensen (2005: 59, 69); Martín-Mateo (2009: 258, figs 52–53); Bartlow *et al.* (2016: 222).

Remarks: *Bovicola (Bovicola) bovis*, also known as “cattle biting louse”, was introduced to New Zealand with its host by human agency (King 2005: 347).

***Bovicola (Bovicola) caprae* (Gurtl, 1843)**

- Trichodectes caprae* Gurtl, 1843: 3, pl. 1: fig. 2.  
*Trichodectes climax* Nitzsch [in Giebel], 1861b: 81, pl. 1: figs 1–2.  
*Trichodectes climax* Nitzsch, in Giebel, 1874 [sic]; Johnston & Harrison 1912: 373.  
*Trichodectes caprae* Gurtl, 1843; Harrison 1916: 69.  
*Trichodectes climax* Nitzsch; Thomson 1922: 269.  
*Trichodectes climax* Nitzsch; Tillyard 1926: 134.  
*Bovicola caprae* (Gurtl, 1843); Hopkins 1942a: 452.  
*Damalinea (Bovicola) caprae* (Gurtl, 1843); Hopkins 1949b: 533.  
*Damalinea caprae* (Gurtl, 1843); Hopkins & Clay 1952: 103.  
*Bovicola caprae* (Gurtl); Neuffer 1954: 452, figs 1, 6–7, 11, 16, 29, 37, 40a.  
*Damalinea caprae* (Gurtl); Helson 1956: 13, 16.  
*Damalinea caprae* (Gurtl, 1843); Watt 1971: 233, 244, fig. 4.  
*Damalinea caprae* (Gurtl, 1843); Wise 1977: 57.  
*Bovicola (Bovicola) caprae* (Gurtl, 1843); Lyal 1985a: 251, figs 3, 45–46, 52.  
*Bovicola (Bovicola) caprae* (Gurtl, 1843); Tenquist & Charleston 2001: 486.  
*Bovicola caprae*; King 2005: 387.  
*Bovicola (B.) caprae* (Gurtl, 1843); Palma 2010: 409.

Syntypes ♂♀, probably lost (Janet Weigner pers. comm. July 2012).

Type host: *Capra hircus* Linnaeus, 1758.

New Zealand host: *Capra hircus* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: TK, HB, WN, SD, MB, NC, MC, SC, WD, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Johnston & Harrison (1912); Thomson (1922); Tillyard (1926); Helson (1956); Watt (1971); Miller (1971: 131); Whitten (1971: 161); Andrews (1973: 326); Heath (1973: 330); Wise (1977); Tenquist & Charleston (1981: 262); King (1990: 420); Tenquist & Charleston (2001); King (2005); Palma (2010).

Other significant references: Werneck (1936: 540, figs 175–178); Hopkins (1942a); Séguy (1944: 391, figs 584–586, as *T. climax*); Hopkins (1949b); Werneck (1950: 60); Neuffer (1954); Clay & Hopkins (1955: 70); Emerson & Price (1975: 59, figs 193–196); Rudolph (1983: 16); Lyal (1985a); Barker (1996: 232); Price *et al.* (2003: 253); Martín-Mateo (2009: 258); Palma & Peck (2013: 65).

Remarks: *Bovicola (Bovicola) caprae*, also known as “goat biting louse”, was introduced to New Zealand with its host by human agency (King 2005: 377).

***Bovicola (Bovicola) limbatus* (Gervais, 1844)**

- Trichodectes limbatus* Gervais, 1844: 313, pl. 48: fig. 4.  
*Trichodectes limbatus* Gervais, 1847 [sic]; Harrison 1916: 70.  
*Bovicola limbatus* (Gervais, 1844); Hopkins 1942a: 452.  
*Damalinea (Bovicola) limbata* (Gervais, 1844); Hopkins 1949b: 533.  
*Damalinea limbata* (Gervais, 1844); Hopkins & Clay 1952: 105.  
*Damalinea limbata*; Andrews 1973: 328.  
*Damalinea limbata* (Gervais, 1844); Wise 1977: 58.  
*Bovicola (Bovicola) limbatus* (Gervais, 1844); Lyal 1985a: 251.  
*Bovicola (Bovicola) limbatus* (Gervais, 1844); Tenquist & Charleston 2001: 487.  
*Bovicola limbatus*; King 2005: 387.  
*Bovicola (B.) limbatus* (Gervais, 1844); Palma 2010: 409.

Neotype ♂ in SAIM (Hopkins 1942a: 448).

Type host: *Capra hircus* Linnaeus, 1758.

New Zealand host: *Capra hircus* Linnaeus, 1758.

Other host: *Capra hircus angorensis* (Shaw, 1800).

New Zealand localities: HB, WN, BR, NC, MC, SC, WD, CO, DN, KE, AU.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Andrews (1973); Wise (1977); Tenquist & Charleston (1981: 263); King (1990: 420); Tenquist & Charleston (2001); King (2005: 387); Palma (2010).

Other significant references: Werneck (1936: 545, figs 179–182); Hopkins (1942a); Séguy (1944: 394, figs 593–594); Hopkins (1949b); Werneck (1950: 62); Lyal (1985a); Butler & O'Connor (1994: 451); Barker (1996: 232); Price *et al.* (2003: 253); Martín-Mateo (2009: 257).

Remarks: *Bovicola (Bovicola) limbatus* was introduced to New Zealand with goats by human agency (King 2005: 377).

### ***Bovicola (Bovicola) longicornis* (Nitzsch, 1818)**

Figs 176–177

*Trichodectes longicornis* Nitzsch, 1818: 296.

*Bovicola americana* Jellison, 1935: 410, fig.

*Rhabdopedilon longicornis* (Nitzsch, 1818); Kéler 1938a: 456, figs 39–40.

*Trichodectes longicornis* Nitzsch, 1818; Séguy 1944: 395, figs 595–597.

*Damalinea (Rhabdopedilon) longicornis* (Nitzsch, 1818); Hopkins 1949b: 524.

*Damalinea (Rhabdopedilon) americanus* Jellison, 1935; Hopkins 1949b: 524.

*Damalinea americana* (Jellison, 1935); Hopkins & Clay 1952: 102.

*Damalinea longicornis* (Nitzsch, 1818); Hopkins & Clay 1952: 105.

*Damalinea (Bovicola) longicornis* (Nitzsch, 1818); Hopkins 1960: 86.

*Damalinea (Bovicola) concavifrons* Hopkins 1960: 87, figs 7–8.

*Damalinea longicornis* (Nitzsch, 1818); Andrews 1964: 104, figs 2, 3a.

*Damalinea (Bovicola) longicornis* (Nitzsch, 1818); Andrews 1972: 153, fig. 1.

*Damalinea longicornis* (Nitzsch, 1818); Wise 1977: 58.

*Bovicola (Bovicola) longicornis* (Nitzsch, 1818); Lyal 1985a: 251.

*Damalinea longicornis*; King 1990: 389, 451, 465.

*Bovicola (Bovicola) longicornis* (Nitzsch, 1818); Tenquist & Charleston 2001: 487.

*Bovicola longicornis*; King 2005: 358.

*Damalinea longicornis*; King 2005: 415, 426.

*Bovicola (B.) longicornis* (Nitzsch, 1818); Palma 2010: 409.

Neotype ♀ in the MLUH (Clay & Hopkins 1960: 45).

Type host: *Cervus elaphus* Linnaeus, 1758.

New Zealand hosts: *Cervus elaphus scoticus* Lönnberg, 1906; *Cervus elaphus nelsoni* (Bailey, 1935); *Rupicapra rupicapra rupicapra* Couturier, 1938.

Other hosts: None.

New Zealand localities: BP, WN, NC, MC, SC, BR, CO, DN, SL, FD, SI.

Geographic distribution: Eurasia; North America; South America; Australasia.

New Zealand references: Andrews (1964; 1972); Andrews (1973: 326, 328); Wise (1977); Charleston (1980: 150); Tenquist & Charleston (1981: 263); King (1990); Tenquist & Charleston (2001); King (2005); Palma (2010).

Other significant references: Kéler (1938); Séguy (1944); Hopkins (1949b); Clay & Hopkins (1960: 45); Hopkins (1960); Lyal (1985a); Butler & O'Connor (1994: 452); Price *et al.* (2003: 253).

Remarks: *Bovicola (Bovicola) longicornis* was introduced to New Zealand with red deer and wapiti by human agency (King 2005: 404, 422). New Zealand records of this louse species from *Rupicapra rupicapra* (e.g. Tenquist & Charleston 1981: 263; King 1990: 389) are likely due to a natural host-switch from *Cervus elaphus*, because *R. rupicapra* is parasitised by a different species —*Bovicola (Bovicola) alpinus* Kéler, 1942— in its native range (Price *et al.* 2003: 388).

The male of *Bovicola (Bovicola) longicornis* is extremely rare (Lyal 1985a: 251). However, two males (held in MONZ) have so far been collected from New Zealand deer, with the first recorded and described by Andrews (1972: 153, fig. 1).

Lyal (1985a: 251) resurrected *Bovicola (B.) concavifrons* from synonymy under *B. (B.) longicornis*, but I follow Andrews (1972: 153; 1973: 328) who, after a morphological assessment of the status of *B. (B.) concavifrons* [as *Damalinea concavifrons*] in New Zealand, did not recognise it as a species different from *B. (B.) longicornis*.

***Bovicola (Bovicola) ovis* (Schrank, 1781)**

*Pediculus ovis* Schrank, 1781: 502, pl. 1: figs 8–9.

*Trichodectes sphaerocephalus* Nitzsch, 1818: 296. Unnecessary *nomen novum* for *Pediculus ovis* Schrank, 1781.

*Trichodectes ovis* Schrank, 1781 [sic]; Harrison 1916: 71.

*Trichodectes sphaerocephalus* Nitzsch; Thomson 1922: 269.

*Trichodectes ovis* L. [sic]; Tillyard 1926: 134.

*Bovicola ovis* (Schrank, 1781); Kéler 1938a: 448, fig. 33.

*Damalinia (Bovicola) ovis* (Schrank, 1781); Hopkins 1949b: 533.

*Damalinia ovis* (Schrank, 1781); Hopkins & Clay 1952: 106.

*Damalinia ovis* (L.) [sic]; Helson 1956: 14, 16.

*Damalinia ovis* (Schrank, 1781); Clay 1964a: 233.

*Damalinia ovis*; Miller 1971: 131, fig. 345.

*Damalinia ovis* (Schrank, 1781); Wise 1977: 58.

*Bovicola (Bovicola) ovis* (Schrank, 1781); Lyal 1985a: 251.

*Bovicola (Bovicola) ovis* (Schrank, 1781); Tenquist & Charleston 2001: 487.

*Bovicola ovis*; King 2005: 396.

*Bovicola (B.) ovis* (Schrank, 1781); Palma 2010: 409.

Neotype ♂ in NHML (Clay & Hopkins 1954: 255).

Type host: *Ovis aries* Linnaeus, 1758.

New Zealand host: *Ovis aries* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: WN, NC, MC, SC, CH, CA, Macquarie Island.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Thomson (1922); Tillyard (1926); Helson (1956); Thomas (1958: 217); Clay (1964a); Gressitt (1964: 539); Watson (1967: 71); Heath & Millar (1970: 211); Miller (1971: 131); Whitten (1971: 161); Andrews (1972: 155, fig. 2); Kettle & Pearce (1974); Wise (1977); Tenquist (1977: 285); Heath (1978: 364); Heath (1979: 141); Tenquist & Charleston (1981: 263); Kettle & Lukies (1982a,b); Heath (1983: 367); Kettle *et al.* (1983); Kettle & Lukies (1984); Heath & Bishop (1988); Heath *et al.* (1992a,b); Heath (1994: 29); Bany *et al.* (1995); Heath *et al.* (1995a,b; 1996a,b); Pfeffer *et al.* (1996); Wilson *et al.* (1997); Tenquist & Charleston (2001); Heath (2001: 9); Palma & Horning (2002: 14, 18); Heath (2002: 48); Heath *et al.* (2004); King (2005: 396); Heath (2005: 138); Palma (2010); Heath & Levot (2015: 202).

Other significant references: Werneck (1936: 535, figs 169–174); Kéler (1938); Séguy (1944: 399, figs 604–608); Webb (1946: 58); Hopkins (1949b); Werneck (1950: 73, figs 54–56); Clay & Hopkins (1954: 254); Murray (1955b; 1957a: 13; 1957b,c; 1960b; 1962; 1963c); Hopkins (1960: 84, fig. 4, pl. 1: fig. 5); Murray & Gordon (1969: 179); Emerson & Price (1975: 68, figs 199–202); Lyal (1985a); Barker (1996: 232); Price *et al.* (2003: 253); Palma & Jensen (2005: 59, 69); Martín-Mateo (2009: 255).

Remarks: *Bovicola (Bovicola) ovis*, also known as “sheep body louse”, was introduced to New Zealand with its host by human agency (King 2005: 393).

**Subgenus *Lepikentron* Kéler, 1938**

*Lepikentron* Kéler, 1938a. *Nova Acta Leop.-Carol. (N.F.)* 5: 452. Type species: *Trichodectes breviceps* Rudow, 1866a = *Bovicola (Lepikentron) breviceps* (Rudow, 1866) (by original designation).

***Bovicola (Lepikentron) breviceps* (Rudow, 1866)**

Fig. 178

*Trichodectes breviceps* Rudow, 1866a: 110, pl. 5: fig. 2.

*Trichodectes breviceps* Rudow, 1866; Harrison 1916: 69.

*Trichodectes breviceps* Rudow, 1866; Werneck 1936: 531, figs 161–168.

*Lepikentron breviceps* (Rudow, 1866); Kéler 1938a: 453, figs 37–38.

*Bovicola breviceps* (Rudow, 1866); Werneck 1950: 73.

*Damalinia breviceps* (Rudow, 1866); Hopkins & Clay 1952: 103.

*Damalinia (Lepikentron) breviceps* (Rudow, 1866); Weidner 1966: 261.

*Bovicola (Lepikentron) breviceps* (Rudow, 1866); Lyal 1985a: 253, figs 43, 55.

*Bovicola (Lepikentron) breviceps* (Rudow, 1866); Price *et al.* 2003: 253.

*Bovicola (Lepikentron) breviceps* (Rudow, 1866); Palma *et al.* 2006: 253.

Lectotype ♀ in ZMHG (Clay & Hopkins 1955: 69; Weidner 1966: 261).

Type host: *Lama glama* (Linnaeus, 1758).

New Zealand host: *Lama pacos* (Linnaeus, 1758).

Other hosts: *Lama guanicoe* (Müller, 1776).

New Zealand localities: WI, SC.

Geographic distribution: South America; Australasia.

New Zealand references: McKenna (2001: 15; 2003: 12); Palma *et al.* (2006).

Other significant references: Werneck (1936); Kéler (1938); Hopkins (1949b: 522); Werneck (1950); Clay & Hopkins (1955); Weidner (1966); Lyal (1985a); Price *et al.* (2003); Mey & González-Acuña (2007: 71, figs 1–12).

Remarks: *Bovicola (Lepikentron) breviceps* was introduced to New Zealand with alpacas by human agency (Corrin & Burnett 1989). The male of *B. (L.) breviceps* is extremely rare (Lyal 1985a: 253, fig. 55); records from New Zealand include females and nymphs only (Palma *et al.* 2006: 253).

### Subgenus *Spinibovicola* Lyal, 1985

*Spinibovicola* Lyal, 1985. *Bull. Brit. Mus. (Nat. Hist.)* 51(3): 253. Type species: *Trichodectes hemitragi* Cummings, 1916a = *Bovicola (Spinibovicola) hemitragi* (Cummings, 1916a) (by original designation).

### ***Bovicola (Spinibovicola) hemitragi* (Cummings, 1916)**

Figs 179–180

*Trichodectes hemitragi* Cummings, 1916a: 273, figs 11–12.

*Bovicola hemitragi* (Cummings, 1916); Werneck 1950: 71, figs 52–53.

*Damalinea hemitragi* (Cummings, 1916); Hopkins & Clay 1952: 104.

*Damalinea hemitragi* (Cummings, 1916); Christie & Andrews 1964: 76.

*Damalinea hemitragi* (Cummings, 1916); Andrews 1970: 846.

*Damalinea (Bovicola) hemitragi* (Cummings, 1916); Andrews 1972: 156, fig. 3.

*Damalinea hemitragi* (Cummings, 1916); Wise 1977: 58.

*Bovicola (Spinibovicola) hemitragi* (Cummings, 1916); Lyal 1985a: 253, figs 12, 41, 50, 56.

*Damalinea hemitragi*; King 1990: 405

*Bovicola (Spinibovicola) hemitragi* (Cummings, 1916); Tenquist & Charleston 2001: 487.

*Bovicola hemitragi*; King 2005: 371.

*Bovicola (Spinibovicola) hemitragi* (Cummings, 1916); Palma 2010: 409.

Syntypes ♀♀ in NHML.

Type host: *Hemitragus jemlahicus* (Smith, 1826).

New Zealand host: *Hemitragus jemlahicus* (Smith, 1826).

Other hosts: None.

New Zealand localities: MC, WD.

Geographic distribution: Asia; New Zealand; South Africa; Argentina.

New Zealand references: Christie & Andrews (1964); Andrews (1970; 1972); Andrews (1973: 326, 328); Wise (1977); Tenquist & Charleston (1981: 262); King (1990); Tenquist & Charleston (2001); King (2005); Palma (2010).

Other significant references: Hopkins (1949b: 533); Werneck (1950); Lyal (1985a); Price *et al.* (2003: 253).

Remarks: *Bovicola (Spinibovicola) hemitragi* was introduced to New Zealand with Himalayan tahrs by human agency (King 2005: 362).

### Genus *Felicola* Ewing, 1929

#### Subgenus *Felicola* Ewing, 1929

*Felicola* Ewing, 1929. *Manual External Parasites*: 121, 122, 192. Type species: *Trichodectes subrostratus* “Nitzsch” = *Felicola (Felicola) subrostratus* (Burmeister, 1838a) (by original designation).

***Felicola (Felicola) subrostratus* (Burmeister, 1838)**

Figs 181–182

- Trichodectes subrostratus* Burmeister, 1838a: 438.  
*Trichodectes subrostratus* Nitzsch in Burmeister, 1838; Harrison 1916: 73.  
*Trichodectes subrostratus* Nitzsch [sic]; Tillyard 1926: 134.  
*Felicola subrostrata* [sic] Nitzsch, 1818 [sic]; Kéler 1938a: 443, figs 30–31.  
*Trichodectes subrostratus* Nitzsch, 1818 [sic]; Séguy 1944: 403, figs 618–620.  
*Felicola (Felicola) subrostratus* (Burmeister, 1838); Hopkins 1949b: 507.  
*Felicola subrostratus* (Burmeister, 1838); Hopkins & Clay 1952: 140.  
*Felicola subrostrata* (Nitz.) [sic]; Helson 1956: 13, 16.  
*Felicola subrostratus*; Miller 1971: 131.  
*Felicola subrostrata* [sic]; Whitten 1971: 161.  
*Felicola subrostratus* (Burmeister, 1838); Wise 1977: 58.  
*Felicola (Felicola) subrostratus* (Burmeister, 1838); Lyal 1985a: 312, figs 5, 204.  
*Felicola (Felicola) subrostratus* (Burmeister, 1838); Tenquist & Charleston 2001: 500.  
*Felicola (Felicola) subrostratus* (Burmeister, 1838); Palma 2010: 409.

Status, sex and repository of types unknown, presumed lost. See Clay (1949a: 1) and Palma & Pilgrim (1984: 150).

Type host: *Felis catus* Linnaeus, 1758.

New Zealand host: *Felis catus* Linnaeus, 1758.

Other hosts: *Felis silvestris* Schreber, 1777; *Ichneumia albicauda* (Cuvier, 1829); *Lynx rufus* (Schreber, 1777); *Salanoia concolor* (Geoffroy Saint-Hilaire, 1837); *Civettictis civetta* (Schreber, 1776).

New Zealand localities: HB, WN, NN, NC, MC, SC, CO, DN.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Tillyard (1926); Helson (1956); Pilgrim (1970: 76); Miller (1971: 131); Whitten (1971); Pilgrim (1974: 1034, fig. 1); Wise (1977); Tenquist & Charleston (1981: 265); King (1990: 344); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Werneck (1936: 526, figs 157–160); Kéler (1938); Séguy (1944); Hopkins (1949b); Hopkins (1960: 80); Emerson & Price (1975: 52, figs 147–150); Lyal (1985a); Price (1987: 220, fig. 22.30); Barker (1996: 232); Price *et al.* (2003: 258); Martín-Mateo (2009: 247).

Remarks: *Felicola (Felicola) subrostratus* was introduced to New Zealand with cats by human agency (King 2005: 310).

**Genus *Trichodectes* Nitzsch, 1818****Subgenus *Trichodectes* Nitzsch, 1818**

*Trichodectes* Nitzsch, 1818. *Germar's Mag. Entomol.* 3: 294. Type species: *Trichodectes canis* (De Geer, 1778b) (by subsequent designation).

***Trichodectes (Trichodectes) canis* (De Geer, 1778)**

Figs 183–184

- Ricinus canis* De Geer, 1778b: 81, pl. 4: fig. 16.  
*Pediculus canis* O. Fabricius, 1780: 215.  
*Trichodectes latus* Nitzsch, 1818: 296. Unnecessary *nomen novum* for *Ricinus canis* De Geer, 1778.  
*Trichodectes canis* De Geer, 1778 [sic]; Harrison 1916: 69.  
*Trichodectes latus* Nitzsch; Thomson 1922: 269.  
*Trichodectes canis* (De Geer, 1778); Hopkins & Clay 1952: 350.  
*Trichodectes canis* (Degeer) [sic]; Helson 1956: 17.  
*Trichodectes (Trichodectes) canis* (De Geer, 1778); Hopkins 1960: 77.  
*Trichodectes canis* (De Geer, 1778); Wise 1977: 58.  
*Trichodectes (Trichodectes) canis* (De Geer, 1778); Lyal 1985a: 300, figs 150, 158, 169.  
*Trichodectes (Trichodectes) canis* (De Geer, 1778); Tenquist & Charleston 2001: 522.  
*Trichodectes (Trichodectes) canis* (De Geer, 1778); Palma 2010: 409.

Neotype ♂ in NHML (Clay & Hopkins 1954: 244).

Type host: *Canis familiaris* Linnaeus, 1758.



New Zealand host: *Canis familiaris* Linnaeus, 1758.

Other hosts: *Canis aureus* Linnaeus, 1758; *Canis latrans* Say, 1823; *Canis lupus* Linnaeus, 1758; *Cerdocyon thous* (Linnaeus, 1766); *Pseudalopex culpaeus* (Molina, 1782); *Civettictis civetta* (Schreber, 1776); *Vulpes bengalensis* (Shaw, 1800).

New Zealand localities: WO, WN, NC, MC, SC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Thomson (1922); Helson (1956: 17); Pilgrim (1970: 76); Whitten (1971: 161); Wise (1977); Tenquist & Charleston (1981: 273); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Werneck (1936: 502, figs 130–135); Séguy (1944: 389, figs 579–583); Webb (1946: 55); Hopkins (1949b: 496); Symmons (1952: 396, figs 38–39); Clay & Hopkins (1954: 243); Kéler (1957c: figs 15, 17); Hopkins (1960); Emerson & Price (1975: 45, figs 119–122); Lyal (1985a); Price (1987: 220, fig. 22.29); Butler & O'Connor (1994: 452); Barker (1996: 233); Price *et al.* (2003: 274); Palma & Jensen (2005: 59, 69).

Remarks: *Trichodectes (Trichodectes) canis* was introduced to New Zealand with dogs by human agency (King 2005: 9, 258).

### Subgenus *Stachiella* Kéler, 1938

*Stachiella* Kéler, 1938a. *Nova Acta Leop.-Carol. (N.F.)* 5: 428. Type species: *Trichodectes pusillus* “Nitzsch, 1861” = *Trichodectes mustelae* (Schrank, 1803) (by original designation).

### *Trichodectes (Stachiella) ermineae* (Hopkins, 1941)

Figs 185–186

*Stachiella ermineae* Hopkins, 1941: 38.

*Trichodectes (Stachiella) ermineae* (Hopkins, 1941); Hopkins 1949b: 499.

*Trichodectes ermineae* (Hopkins, 1941); Hopkins & Clay 1952: 351.

*Trichodectes (Stachiella) ermineae* (Hopkins, 1941); Hopkins 1960: 78.

*Trichodectes (Stachiella) erminiae* [sic] (Hopkins, 1941); Lyal 1985a: 301, figs 156, 174.

*Stachiella ermineae* Hopkins, 1941; Price *et al.* 2003: 271.

*Trichodectes (Stachiella) ermineae* (Hopkins, 1941); Tenquist & Charleston 2001: 522.

*Trichodectes ermineae*; King 2005: 282.

*Trichodectes (Stachiella) ermineae* (Hopkins, 1941); Palma 2010: 409.

Holotype ♀ in NHML.

Type host: *Mustela erminea stabilis* Barrett-Hamilton, 1904.

New Zealand hosts: *Mustela erminea* Linnaeus, 1758; *Mustela furo* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: WN, NN, NC, MC, SC, WD, CO, DN, FD.

Geographic distribution: Eurasia; Australasia.

New Zealand references: King (1990: 307); Tenquist & Charleston (2001); King (2005); Palma (2010).

Other significant references: Hopkins (1949b); Hopkins (1960: 78); Lyal (1985a); Butler & O'Connor (1994: 452); Price *et al.* (2003).

Remarks: *Trichodectes (Stachiella) ermineae* was introduced to New Zealand with stoats by human agency (King 2005: 264). In agreement with Lyal (1985a: 301) and contrary to Price *et al.* (2003: 271), I regard *Stachiella* as a subgenus.

New Zealand records of *Trichodectes (Stachiella) ermineae* from *Mustela furo* are likely due to a natural host-switch from *Mustela erminea*, because *M. furo* is parasitised by a different species—*Trichodectes (Stachiella) jacobi* Eichler, 1941b—in its native range (Price *et al.* 2003: 388).

### *Trichodectes (Stachiella) mustelae* (Schrank, 1803)

*Pediculus mustelae* Schrank, 1803: 186.

*Trichodectes mustelae* Schrank, 1803 [sic]; Harrison 1916: 71.

*Trichodectes (Stachiella) mustelae* (Schrank, 1803); Hopkins 1949b: 499.

*Trichodectes mustelae* (Schrank, 1803); Hopkins & Clay 1952: 353.

*Trichodectes (Stachiella) mustelae* (Schrank, 1903 [sic]); Lyal 1985a: 301.

*Trichodectes (Stachiella) mustelae* (Schrank, 1803); Tenquist & Charleston 2001: 522.

*Stachiella mustelae* (Schrank, 1803); Price *et al.* 2003: 271.

*Trichodectes mustelae*; King 2005: 292.

*Trichodectes (S.) mustelae* (Schrank, 1803); Palma 2010: 409.

Neotype ♀ in the MLUH (Clay & Hopkins 1960: 11).

Type host: *Mustela nivalis nivalis* Linnaeus, 1766.

New Zealand host: *Mustela nivalis vulgaris* Erxleben, 1777.

Other host: *Mustela sibirica* Pallas, 1773.

New Zealand localities: WA, NC, MC, SC.

Geographic distribution: Eurasia; North America; New Zealand.

New Zealand references: Tenquist & Charleston (2001); King (2005); Palma (2010).

Other significant references: Kéler (1938: 429, figs 22–23); Séguy (1944: 398, figs 600–603); Hopkins (1949b); Clay & Hopkins (1960: 11); Lyal (1985a); Price *et al.* (2003); Martín-Mateo (2009: 243).

Remarks: *Trichodectes (Stachiella) mustelae* was introduced to New Zealand with weasels by human agency (King 2005: 288). In agreement with Lyal (1985a: 301) and contrary to Price *et al.* (2003: 271), I regard *Stachiella* as a subgenus of *Trichodectes*.

### Genus *Tricholipeurus* Bedford, 1929

*Tricholipeurus* Bedford, 1929. *15th Annual Rep. Director Veterinary Services*: 514. Type species: *Tricholipeurus aepycerus* Bedford, 1929 = (by original designation).

### *Tricholipeurus lipeuroides* (Mégnin, 1884)

Figs 187–188

*Trichodectes lipeuroides* Mégnin, 1884: 494, figs a,b,c.

*Eutrichophilus lipeuroides* Mégnin, 1884 [sic]; Harrison 1916: 74.

*Damalinea (Cervicola) lipeuroides* (Mégnin, 1884); Hopkins 1949b: 524.

*Tricholipeurus lipeuroides* (Mégnin, 1884); Werneck 1950: 175, figs 259–265.

*Damalinea lipeuroides* (Mégnin, 1884); Hopkins & Clay 1952: 105.

*Damalinea (Tricholipeurus) lipeuroides* (Mégnin, 1884); Hopkins 1960: 88.

*Tricholipeurus lipeuroides* (Mégnin, 1884); Emerson & Price 1975: 68, figs 211–214.

*Damalinea lipeuroides* (Mégnin, 1884); Wise 1977: 58.

*Damalinea (Tricholipeurus) lipeuroides* (Mégnin, 1884); Lyal 1985a: 265.

*Damalinea (Tricholipeurus) lipeuroides* (Mégnin, 1884); Tenquist & Charleston 2001: 494.

*Tricholipeurus lipeuroides* (Mégnin, 1884); Price *et al.* 2003: 275.

*Damalinea lipeuroides*; King 2005: 464.

*Tricholipeurus lipeuroides* (Mégnin, 1884); Palma 2010: 409.

Syntypes ♂♀, repository unknown.

Type host: *Odocoileus virginianus mexicanus* (J.F. Gmelin, 1788).

New Zealand host: *Odocoileus virginianus borealis* (Miller, 1900).

Other hosts: *Odocoileus virginianus virginianus* (Zimmerman, 1780); *Odocoileus virginianus texanus* (Mearns, 1898); *Odocoileus hemionus* (Rafinesque, 1817).

New Zealand locality: SI.

Geographic distribution: Americas; Europe; New Zealand.

New Zealand references: Andrews (1973: 326, 328); Wise (1977); Tenquist & Charleston (1981: 263); King (1990: 513); Tenquist & Charleston (2001); King (2005: 464); Palma (2010).

Other significant references: Hopkins (1949b); Werneck (1950); Hopkins (1960); Emerson & Price (1975); Lyal (1985a); Price *et al.* (2003).

Remarks: *Tricholipeurus lipeuroides* was introduced to New Zealand with white-tailed deer by human agency (King 2005: 460).

***Tricholipeurus parallelus* (Osborn, 1896)**

- Trichodectes parallelus* Osborn, 1896: 240, fig. 148.  
*Trichodectes parallelus* Osborn, 1896; Harrison 1916: 72.  
*Damalinea (Cervicola) parallelus* (Osborn, 1896); Hopkins 1949b: 524.  
*Tricholipeurus parallelus* (Osborn, 1896); Werneck 1950: 184.  
*Damalinea parallela* (Osborn, 1896); Hopkins & Clay 1952: 106.  
*Damalinea (Tricholipeurus) parallela* (Osborn, 1896); Hopkins 1960: 89.  
*Damalinea parallela* (Osborn, 1896); Andrews 1973: 326, 328.  
*Tricholipeurus parallelus* (Osborn, 1896); Emerson & Price 1975: 68, figs 215–218.  
*Damalinea parallela* (Osborn, 1896); Wise 1977: 58.  
*Damalinea (Tricholipeurus) parallela* (Osborn, 1896); Lyal 1985a: 265.  
*Damalinea (Tricholipeurus) parallela* (Osborn, 1896); Tenquist & Charleston 2001: 494.  
*Tricholipeurus parallelus* (Osborn, 1896); Price *et al.* 2003: 275.  
*Damalinea parallela*; King 2005: 464.  
*Tricholipeurus parallelus* (Osborn, 1896); Palma 2010: 409.

Syntypes ♀♀, but only one still extant in CUIC [<http://cuic.entomology.cornell.edu/insects/view/1014871>].

Type host: *Odocoileus virginianus* (Zimmermann, 1780).

New Zealand host: *Odocoileus virginianus borealis* (Miller, 1900).

Other host: *Odocoileus hemionus* (Rafinesque, 1817).

New Zealand localities: FD, SI.

Geographic distribution: Americas; Europe; New Zealand.

New Zealand references: Andrews (1973); Wise (1977); Tenquist & Charleston (1981: 263); King (1990: 513); Tenquist & Charleston (2001); King (2005); Palma (2010).

Other significant references: Hopkins (1949b); Werneck (1950); Hopkins (1960); Samuel & Trainer (1971: 507); Emerson & Price (1975); Lyal (1985a); Price *et al.* (2003).

Remarks: *Tricholipeurus parallelus* was introduced to New Zealand with white-tailed deer by human agency (King 2005: 460).

**Genus *Werneckiella* Eichler, 1940**

*Werneckiella* Eichler, 1940a. *Zool. Anz.* 129: 160. Type species: *Trichodectes equi* Denny, 1842 = *Werneckiella equi* (Denny, 1842) (by original designation).

***Werneckiella equi* (Denny, 1842)**

- Trichodectes equi* Denny, 1842: 61, 191, pl. 17: fig. 7.  
*Trichodectes pilosus* Giebel, 1874: 59.  
*Trichodectes parumpilosus* Piaget, 1880: 397.  
*Trichodectes equi* Linnaeus, 1758 [sic]; Harrison 1916: 70.  
*Trichodectes pilosus* Giebel, 1874; Tillyard 1926: 134.  
*Trichodectes equi* (Linnaeus, 1758) [sic]; Werneck 1936: 554, figs 186–188.  
*Bovicola equi* (Denny, 1842); Kéler 1938a: 450, figs 35–36.  
*Werneckiella equi* (Denny, 1842); Eichler 1940: 160.  
*Damalinea (Werneckiella) equi* (Denny, 1842); Hopkins 1949b: 520.  
*Damalinea equi* (Denny, 1842); Hopkins & Clay 1952: 104.  
*Bovicola equi* (L.) [sic]; Helson 1956: 13, 16.  
*Damalinea (Bovicola) equi* (Denny, 1842); Hopkins 1960: 82.  
*Damalinea equi*; Miller 1971: 131.  
*Bovicola equi* (Linnaeus, 1758) [sic]; Emerson & Price 1975: 68, figs 203–206.  
*Damalinea equi* (Denny, 1842); Wise 1977: 58.  
*Werneckiella equi* (Denny, 1842); Moreby 1978: 399, figs 2–5, 7, 11, 15, 20, 25, 36.  
*Werneckiella equi* (Denny, 1842); Tenquist & Charleston 1981: 274.  
*Werneckiella equi* (Denny, 1842); Lyal 1985a: 255, figs 61–62.  
*Werneckiella equi* (Denny, 1842); Tenquist & Charleston 2001: 523.  
*Bovicola (Werneckiella) equi* (Denny, 1842); Price *et al.* 2003: 253.

*Bovicola (Werneckiella) equi* (Denny, 1842); Martín-Mateo 2009: 253, 329.

*Werneckiella equi* (Denny, 1842); Palma 2010: 409.

Lectotype ♀ in NHML (Moreby 1978: 400).

Type host: *Equus caballus* Linnaeus, 1758.

New Zealand host: *Equus caballus* Linnaeus, 1758.

Other hosts: *Equus przewalskii* (Poliakov, 1881); *Equus hemionus kulan* Groves & Mazak, 1967.

New Zealand localities: WN, NC, MC, SL.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Tillyard (1926); Helson (1956); Miller (1971); Whitten (1971: 161); Wise (1977); Tenquist & Charleston (1981); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Werneck (1936); Kéler (1938); Eichler (1940); Séguy (1944: 392, figs 587–588); Hopkins (1949b); Werneck (1950: 79, figs 66–67); Murray (1957d: 183; 1963d); Hopkins (1960); Emerson & Price (1975); Moreby (1978); Lyal (1985a); Barker (1996: 233); Price *et al.* (2003); Palma & Jensen (2005: 59, 69); Martín-Mateo (2009).

Remarks: *Werneckiella equi*, also known as “horse biting louse”, was introduced to New Zealand with its host by human agency (King 2005: 329). In agreement with Lyal (1985a: 255) and contrary to Price *et al.* (2003: 253), I regard *Werneckiella* as a full genus. Although a few males of *W. equi* are known (Lyal 1985a: 255), no male has been found on New Zealand horses yet.

### ***Werneckiella ocellata* (Piaget, 1880)**

Fig. 189

*Trichodectes parumpilosus* var. *ocellata* Piaget, 1880: 398.

*Trichodectes ocellatus* Piaget, 1880; Harrison 1916: 71.

*Werneckiella ocellata* (Piaget, 1880); Eichler 1940: 161.

*Damalinia (Werneckiella) ocellata* (Piaget, 1880); Hopkins 1949b: 520.

*Bovicola ocellatus* (Piaget, 1880); Werneck 1950: 84, figs 75–81.

*Damalinia ocellata* (Piaget, 1880); Hopkins & Clay 1952: 106.

*Werneckiella equi asini* Eichler, 1953: 445, fig.

*Damalinia asini* (Eichler, 1953); Hopkins & Clay 1955: 180.

*Werneckiella ocellata* (Piaget, 1880); Moreby 1978: 402, figs 1, 12, 26.

*Werneckiella ocellata* (Piaget, 1880); Lyal 1985a: 255.

*Werneckiella ocellata* (Piaget, 1880); Tenquist & Charleston 2001: 523.

*Bovicola (Werneckiella) ocellatus* (Piaget, 1880); Price *et al.* 2003: 253.

*Werneckiella ocellata* (Piaget, 1880); Palma 2010: 409.

Lectotype ♀ in NHML (Moreby 1978: 403).

Type host: “*Equus burchelli*”, in error (see Moreby 1978: 402).

New Zealand host: *Equus asinus* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: WN, NC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Tenquist & Charleston (2001); Palma (2010).

Other significant references: Eichler (1940); Hopkins (1949b); Werneck (1950); Moreby (1978); Lyal (1985a); Butler & O'Connor (1994: 452); Price *et al.* (2003).

Remarks: *Werneckiella ocellata* was introduced to New Zealand with donkeys by human agency (Barclay 2002: 26). In agreement with Lyal (1985a: 255) and contrary to Price *et al.* (2003: 253), I regard *Werneckiella* as a full genus. The male of *W. ocellata* is still unknown, therefore, this species is believed to be parthenogenetic (Lyal 1985a: 255).

### **Suborder ANOPLURA Leach, 1815**

Anoplura Leach, 1815a: *Edinburgh Encyclopaedia* 9: 77.

**Family ECHINOPHTHIRIIDAE Enderlein, 1904**

Echinophthiriidae Enderlein, 1904b. *Zool. Anz.* 28: 136. Type genus: *Echinophthirus* Giebel, 1874.

**Genus *Antarctophthirus* Enderlein, 1906**

*Antarctophthirus* Enderlein, 1906. *Zool. Anz.* 29: 661. Type species: *Antarctophthirus ogmorhini* Enderlein, 1906 (by original designation).

*Arctophthirus* Mjöberg, 1910a. *Arkiv Zool.* 6(13): 177. Type species: *Arctophthirus trichechi* (Bohemann, 1865) = *Antarctophthirus trichechi* (Bohemann, 1865) (by monotypy).

***Antarctophthirus carlinii* Leonardi et al., 2014**

*Antarctophthirus* sp.; Harrison 1937: 13.

*Antarctophthirus* sp. ?; Clay 1940a: 296.

“*Antarctophthirus ? ogmorhini*” Hopkins, 1949b: 510 (not *Antarctophthirus ogmorhini* Enderlein, 1906).

“*Antarctophthirus ogmorhini*” Murray, 1964: 243 (not *Antarctophthirus ogmorhini* Enderlein, 1906).

“*Antarctophthirus ogmorhini*” Murray et al., 1965: 761, fig. 1, pls 1–2 (not *Antarctophthirus ogmorhini* Enderlein, 1906).

“*Antarctophthirus ogmorhini*” Clay & Moreby, 1967: 166, 169, fig. 179 (not *Antarctophthirus ogmorhini* Enderlein, 1906).

*Antarctophthirus ogmorhini* Enderlein, 1906; Gressitt 1970: 329. In part.

*Antarctophthirus ogmorhini*; Kim et al. 1975: 547. In part.

*Antarctophthirus ogmorhini*; King 1990: 254. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Durden & Musser 1994a: 7. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Barker 1996: 237. In part.

“*Antarctophthirus ogmorhini*” Mehlhorn et al., 2002: 651, figs 1–17 (not *Antarctophthirus ogmorhini* Enderlein, 1906).

*Antarctophthirus ogmorhini*; King 2005: 234. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Palma 2010: 409. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Leonardi & Palma 2013: 451. In part.

*Antarctophthirus carlinii* Leonardi et al., 2014: 3948, figs 1–6.

Holotype ♂ in CENPAT.

Type host: *Leptonychotes weddellii* (Lesson, 1826).

New Zealand host: *Leptonychotes weddellii* (Lesson, 1826).

Other hosts: None.

New Zealand localities: Macquarie Island, RO.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Harrison (1937); Murray et al. (1965); Clay & Moreby (1967); Gressitt (1970); King (1990); Barker (1996); King (2005); Palma (2010).

Other significant references: Clay (1940a); Hopkins (1949b); Murray (1964); Kim et al. (1975); Durden & Musser (1994a); Durden & Musser (1994b: 141); Mehlhorn et al. (2002: 651, figs 1–17); Leonardi & Palma (2013); Leonardi et al. (2014).

Remarks: The population of *Antarctophthirus* from *Leptonychotes weddellii* was previously known as *A. ogmorhini*, but has been recognised as a separate species by Leonardi et al. (2014).

***Antarctophthirus lobodontis* Enderlein, 1909**

“*Antarctophthirus ogmorhini*” Neumann, 1907a: 13 (not *Antarctophthirus ogmorhini* Enderlein, 1906).

*Antarctophthirus lobodontis* Enderlein, 1909: 508, 510, figs KK–NN.

*Antarctophthirus lobodontis* Enderlein, 1909; Hopkins 1949b: 509.

*Antarctophthirus lobodontis* Enderlein, 1909; Clay & Moreby 1967: 166, 169, fig. 180.

*Antarctophthirus lobodontis* Enderlein, 1909; Pilgrim 1974: 1031, fig. 3.

*Antarctophthirus lobodontis* Enderlein, 1909; Palma 2010: 409.

Syntypes ♂♀, repository unknown, probably lost (Leonardi & Palma 2013: 452).

Type host: *Lobodon carcinophagus* Hombron & Jacquinot, 1842.

New Zealand host: *Lobodon carcinophagus* Hombron & Jacquinot, 1842.

Other hosts: None.

New Zealand localities: WN, RO.

Geographic distribution: Antarctica; Southern Oceans; New Zealand.

New Zealand references: Pilgrim (1974); King (1990: 254, 277); King (2005: 234); Palma (2010).

Other significant references: Neumann (1907a); Freund (1928: 20, figs 14–16); Ferris (1934: 488, fig. 284); Clay (1940a: 296); Hopkins (1949b); Barker (1996: 236); Clay & Moreby (1967); Clay & Moreby (1970: 220); Kim *et al.* (1975: 547); Murray (1976: 92, fig. 4.8); Kim (1988: 108); Durden & Musser (1994a: 7); Durden & Musser (1994b: 141); Leonardi & Palma (2013: 452); Leonardi *et al.* (2016: 672, figs 1–6).

Remarks: There are only two records of *Antarctophthirus lobodontis* from the New Zealand Subregion: one from Antarctica dated 1963, and a sample from a crabeater seal found dead in the southern end of the North Island in March 2015 (voucher specimens in MONZ).

### ***Antarctophthirus mawsoni* Harrison, 1937**

*Antarctophthirus mawsoni* Harrison, 1937: 11, pl. 1: fig. 1.

*Antarctophthirus mawsoni* Harrison, 1937; Hopkins 1949b: 509.

*Antarctophthirus mawsoni* Harrison, 1937; King 1990: 254, 278.

*Antarctophthirus mawsoni* Harrison, 1937; King 2005: 234.

Holotype ♀ in AMSA (Australian Museum 2012).

Type host: *Ommatophoca rossii* J.E. Gray, 1844.

New Zealand host: *Ommatophoca rossii* J.E. Gray, 1844.

Other hosts: None

New Zealand localities: None.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: King (1990; 2005).

Other significant references: Hopkins (1949b); Clay & Moreby (1967: 166, 168); Kim *et al.* (1975: 547); Durden & Musser (1994a: 7); Durden & Musser (1994b: 141); Barker (1996: 236); Leonardi & Palma (2013: 452).

Remarks: Although *Ommatophoca rossii* occurs in the New Zealand Subregion as defined by the Checklist Committee (2010: 2, 497), *Antarctophthirus mawsoni* has not yet been recorded from the Subregion. This species is included here because of King's (1990: 254; 2005: 234) citations.

### ***Antarctophthirus microchir* (Trouessart & Neumann, 1888)**

Figs 190–191

*Echinophthirus microchir* Trouessart & Neumann, 1888: 80, figs a,b,c.

*Echinophthirus microchir* Trouessart [sic]; Hutton 1904: 228.

*Antarctophthirus microchir* (Trouessart & Neumann, 1888); Enderlein 1906: 663, figs 3–4.

*Echinophthirus microchir* Trouessart [sic]; Myers 1922: 12.

*Antarctophthirus microchir* (Trouessart & Neumann, 1888); Tillyard 1926: 135.

*Antarctophthirus microchir californianus* Fahrenholz, 1939: 42.

*Antarctophthirus microchir microchir* (Trouessart & Neumann, 1888); Hopkins 1949b: 508.

*Antarctophthirus microchir* (Trouessart & Neumann, 1888); Clay 1964a: 233.

*Antarctophthirus microchir* (Trouessart & Neumann, 1888); Wise 1977: 67.

*Antarctophthirus microchir* (Trouessart & Neumann, 1888); Horning *et al.* 1980: 7, 13.

*Antarctophthirus* sp.; Horning *et al.* 1980: 7, 12.

*Antarctophthirus microchir* (Trouessart & Neumann, 1888); Palma 2010: 409.

Syntypes ♂♀, probably lost (Palma & Peck 2013: 65; Leonardi & Palma 2013: 450).

Type host: *Phocarctos hookeri* (J.E. Gray, 1844).

New Zealand hosts: *Phocarctos hookeri* (J.E. Gray, 1844); *Arctocephalus forsteri* (Lesson, 1828).

Other hosts: *Neophoca cinerea* (Péron, 1816); *Zalophus californianus* (Lesson, 1828); *Zalophus wollebaeki* Silvertsen, 1953; *Otaria flavescens* (Shaw, 1800); *Eumetopias jubatus* (Schreber, 1776).

New Zealand localities: KA, WD, SN, AU.

Geographic distribution: Palearctic, Nearctic, Neotropical and Australasian Regions.

New Zealand references: Trouessart & Neumann (1888); Hutton (1904); Enderlein (1906); Enderlein (1909: 504); Myers (1922); Tillyard (1926); Clay (1964a); Gressitt (1964: 539); Miller (1971: 131); Wise (1977); Horning *et al.*

(1980); King (1990: 254); Durden & Musser (1994a: 7); King (2005: 234); Palma (2010); Leonardi & Palma (2013: 450, 456); Leonardi & Lazzari (2014).

Other significant references: Enderlein (1909: 511, figs 176–177, 183–184); Freund (1928: 21, figs 17–19); Ferris (1934: 489, figs 285–286); Harrison (1937: 10); Webb (1946: 95); Hopkins (1949b); Ferris (1951: 75); Kim *et al.* (1975: 545); Kim *et al.* (1986: 46, pl. 2); Kim (1987: 230); Durden & Musser (1994b: 140); Barker (1996: 236); McIntosh & Murray (2007: 103); Leonardi *et al.* (2009: 1086, figs 1–7). Aznar *et al.* (2009: 293, figs 1–4); Leonardi *et al.* (2011: 62, figs 2–3, 6); Leonardi *et al.* (2012a: 929, figs 1–12); Leonardi *et al.* (2012b: 2, figs 1–3); Palma & Peck (2013: 65).

Remarks: Although some populations of *Antarctophthirus microchir* from different hosts show no morphological differences, they probably represent a complex of cryptic species (Leonardi & Palma 2013: 450). This is the first record of *Antarctophthirus microchir* from the New Zealand fur seal, *Arctocephalus forsteri*.

### ***Antarctophthirus ogmorhini* Enderlein, 1906**

*Antarctophthirus ogmorhini* Enderlein, 1906: 662, figs 1–2.

*Antarctophthirus ogmorhini* Enderlein; Tillyard 1926: 135.

*Antarctophthirus ogmorhini* Enderlein, 1906; Harrison 1937: 11.

*Antarctophthirus ogmorhini* Enderlein, 1906; Watson 1967: 74.

*Antarctophthirus ogmorhini* Enderlein, 1906; Gressitt 1970: 329. In part.

*Antarctophthirus ogmorhini*; Kim *et al.* 1975: 547. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Wise 1977: 67.

*Antarctophthirus ogmorhini*; King 1990: 254. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Durden & Musser 1994a: 7. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Barker 1996: 237. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Palma & Horning 2002: 14, 18.

*Antarctophthirus ogmorhini*; King 2005: 234. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Palma 2010: 409. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Leonardi & Palma 2013: 451, 456. In part.

*Antarctophthirus ogmorhini* Enderlein, 1906; Leonardi *et al.* 2014: 3950.

Syntypes ♂♀ in NHML (Leonardi & Palma 2013: 451).

Type host: *Hydrurga leptonyx* (Blainville, 1820).

New Zealand host: *Hydrurga leptonyx* (Blainville, 1820).

Other hosts: None.

New Zealand localities: “New Zealand waters”; HB, Macquarie Island.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Tillyard (1926); Harrison (1937); Watson (1967); Gressitt (1970); Wise (1977); King (1990); Palma & Horning (2002); King (2005); Palma (2010).

Other significant references: Neumann (1907a); Enderlein (1909: 509, figs 174–175, 181–182); Cummings (1916c: 172); Freund (1928: 19, figs 12–13); Ferris (1934: 486, figs 282–283); Hopkins (1949b: 509); Ferris (1951: 73, fig. 31); Séguy (1953: 593, figs 49–50); Kim *et al.* (1975); Durden & Musser (1994a); Durden & Musser (1994b: 141); Barker (1996); Leonardi & Palma (2013); Leonardi *et al.* (2014).

Remarks: The population of *Antarctophthirus* from *Leptonychotes weddellii*—previously known as *A. ogmorhini*—has been described as a separate species: *A. carlinii* (see above).

### **Genus *Lepidophthirus* Enderlein, 1904**

*Lepidophthirus* Enderlein, 1904a. *Zool. Anz.* 28: 44. Type species: *Lepidophthirus macrorhini* Enderlein, 1904 (by original designation).

#### ***Lepidophthirus macrorhini* Enderlein, 1904**

Figs 192–193

*Lepidophthirus macrorhini* Enderlein, 1904a: 46, figs 1–5.

*Lepidophthirus macrorhini* Enderlein, 1904; Enderlein 1909: 515, figs OO, PP, QQ, pl. 59, pl. 60: fig. 180.

- Lepidophthirus macrorhini* Enderlein, 1904; Tillyard 1926: 135.  
*Lepidophthirus macrorhini* Enderlein, 1904; Clay & Moreby 1967: 166, 169, fig. 181.  
*Lepidophthirus macrorhini* Enderlein, 1904; Watson 1967: 74.  
*Lepidophthirus macrorhini* Enderlein, 1904; Wise 1977: 68.  
*Lepidophthirus macrorhini* Enderlein, 1904; Horning *et al.* 1980: 8, 13.  
*Lepidophthirus* [sic] *macrorhini*; King 2005: 234.  
*Lepidophthirus macrorhini* Enderlein, 1904; Palma & Horning 2002: 14, 18.  
*Lepidophthirus macrorhini* Enderlein, 1904; Palma 2010: 409.

Syntypes ♂♀, repository unknown, probably lost (Barker 1996: 237).

Type host: *Mirounga leonina* (Linnaeus, 1758).

New Zealand host: *Mirounga leonina* (Linnaeus, 1758).

Other hosts: None.

New Zealand localities: KA, SN, CA, Macquarie Island.

Geographic distribution: Antarctica; Southern Oceans.

New Zealand references: Tillyard (1926); Harrison (1937: 13); Murray (1958: 404); Gressitt (1964: 539); Murray & Nicholls (1965: 437); Watson (1967); Gressitt (1970: 329); Clay & Moreby (1970: 220); Wise (1977); Lowry *et al.* (1978: 137); Horning *et al.* (1980); King (1990: 254, 267); Durden & Musser (1994a: 8); Barker (1996: 237); Palma & Horning (2002); King (2005); Palma (2010).

Other significant references: Enderlein (1909); Freund (1928: 32, figs 30–35); Ferris (1934: 499, figs 291–292); Hopkins (1949b: 510); Ferris (1951: 78, fig. 34); Murray (1964: 242); Clay & Moreby (1967); Kim *et al.* (1975: 547); Durden & Musser (1994b: 141); Green & Turner (2004: 74); Leonardi & Palma (2013: 453, 456).

Remarks: *Lepidophthirus macrorhini* is a large species frequently found on southern elephant seals, and with an unusual life cycle (Murray & Nicholls 1965).

#### Family HAEMATOPINIDAE Enderlein, 1904

Haematopinidae Enderlein, 1904b. *Zool. Anz.* 28: 136. Type genus: *Haematopinus* Leach, 1815b.

#### Genus *Haematopinus* Leach, 1815

*Haematopinus* Leach, 1815b. *Encyclopaedia Britannica, Suppl. 1*: 24. Type species: *Haematopinus suis* (Linnaeus, 1758) (by monotypy).

#### *Haematopinus asini* (Linnaeus, 1758)

Figs 194–195

- Pediculus asini* Linnaeus, 1758: 612.  
*Pediculus macrocephalus* Burmeister, 1838b: Species 18.  
*Haematopinus asini* (Linnaeus); Denny 1842: 32, pl. 25: fig. 1.  
*Haematopinus macrocephalus* (Burmeister, 1838); Giebel 1874: 44, pl. 2: fig. 5.  
 “*Haematopinus eurystermus* [sic]” Kirk, 1900: 303, fig. (not *Haematopinus eurysternus* Denny, 1842).  
*Haematopinus asini* Linnaeus [sic]; Myers 1922: 12.  
*Haematopinus macrocephalus* Burmeister [sic]; Thomson 1922: 339.  
*Haematopinus asini* (Linnaeus, 1758); Ferris 1933: 464, figs 273–274.  
*Haematopinus asini asini* (Linnaeus, 1758); Webb 1948: 578.  
*Haematopinus asini macrocephalus* (Burmeister, 1838); Webb 1948: 578.  
*Haematopinus asini asini* (Linnaeus, 1758); Hopkins 1949b: 520.  
*Haematopinus asini macrocephalus* [sic] (Burmeister, 1838); Hopkins 1949b: 520.  
*Haematopinus asini* (Linnaeus, 1758); Ferris 1951: 85, figs 37–38.  
*Haematopinus asini* (L.); Helson 1956: 13, 16.  
*Haematopinus asini*; Miller 1971: 131.  
*Haematopinus asini* (Linnaeus, 1758); Wise 1977: 67.  
*Haematopinus asini* (Linnaeus, 1758); Tenquist & Charleston 2001: 502.  
*Haematopinus asini* (Linnaeus, 1758); Palma 2010: 409.

Status, sex and repository of types unknown.



Type host: *Equus asinus* Linnaeus, 1758.

New Zealand host: *Equus caballus* Linnaeus, 1758.

Other host: *Equus burchelli* (J.E. Gray, 1824).

New Zealand localities: WI, MC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Kirk (1900); Myers (1922); Thomson (1922); Tillyard (1926: 135); Helson (1956); Miller (1971: 131); Whitten (1971: 161); Wise (1977); Tenquist & Charleston (1981: 266); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Cummings (1916c: 172); Ferris (1933); Séguy (1944: 448, figs 722–723); Webb (1948); Hopkins (1949b); Webb (1949: 149); Ferris (1951); Murray (1957d: 183); Kim *et al.* (1986: 80, pl. 17); Durden & Musser (1994a: 15); Durden & Musser (1994b: 141); Barker (1996: 238); Durden (2001: 8).

Remarks: *Haematopinus asini*, also known as “horse sucking louse”, was introduced to New Zealand with horses by human agency (King 2005: 329).

### ***Haematopinus eurysternus* Denny, 1842**

*Pediculus eurysternus* Nitzsch, 1818: 305. *Nomen nudum*.

*Pediculus eurysternus* Burmeister, 1838b: Species 14. Suppressed by Opinion 1050 (I.C.Z.N. 1976).

*Haematopinus eurysternus* Denny, 1842: 29, pl. 25: fig. 5. Preserved by Opinion 1050 (I.C.Z.N. 1976).

*Haematopinus eurysternus* Nitzsch [sic]; Hutton 1904: 353.

*Haematopinus eurysternus* Nitzsch [sic]; Myers 1922: 12.

*Haematopinus eurysternus* Nitzsch [sic]; Thomson 1922: 339.

*Haematopinus eurysternus* (Nitzsch, 1818) [sic]; Ferris 1933: 448, figs 263–264.

*Haematopinus eurysternus* (Nitzsch, 1818) [sic]; Ferris 1951: 88, figs 39–40.

*Haematopinus eurysternus* (Nitzsch) [sic]; Helson 1956: 13, 16.

*Haematopinus eurysternus*; Miller 1971: 131.

*Haematopinus eurysternus* Denny, 1842; Kim & Weisser 1973: 45.

*Haematopinus eurysternus* Denny, 1842; Meleney & Kim 1974: 511, figs 1–3, 18, 22, 24, 27, 30, 33.

*Haematopinus eurysternus* (Nitzsch, 1818) [sic]; Wise 1977: 67.

*Haematopinus eurysternus* Denny, 1842; Kim *et al.* 1986: 82, pl. 18.

*Haematopinus eurysternus* (Nitzsch, 1818) [sic]; Durden & Musser 1994a: 15.

*Haematopinus eurysternus* (Nitzsch, 1818) [sic]; Tenquist & Charleston 2001: 502.

*Haematopinus eurysternus* (Nitzsch, 1818) [sic]; Palma 2010: 409.

Lectotype ♀ in NHML (Kim *et al.* 1986: 82).

Type host: *Bos taurus* Linnaeus, 1758.

New Zealand host: *Bos taurus* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: Not specified.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Hutton (1904); Myers (1922); Thomson (1922); Tillyard (1926: 135); Helson (1956); Whitten (1970: 146); Helson (1970: 81); Miller (1971); Whitten (1971: 161); Buchanan & Coles (1971: 197); Wise (1977); Tenquist (1977: 286); Chalmers & Charleston (1980a,c); Tenquist & Charleston (1981: 266); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Ferris (1933); Séguy (1944: 449, figs 724–725); Webb (1946: 91); Webb (1949: 150); Hopkins (1949b: 527); Ferris (1951); Murray (1957a: 17); Kim & Weisser (1973); Meleney & Kim (1974); International Commission on Zoological Nomenclature (1976); Kim *et al.* (1986); Green & Palma (1991: 21, 23); Durden & Musser (1994a); Durden & Musser (1994b: 144); Barker (1996: 238); Durden (2001: 8); Bartlow *et al.* (2016: 222).

Remarks: *Haematopinus eurysternus*, also known as “short-nosed cattle louse”, was introduced to New Zealand with its host by human agency (King 2005: 347). Notwithstanding the I.C.Z.N. (1976) Opinion designating “Denny, 1842” as the author and date of this species, several post-1976 authors incorrectly ascribed it to “Nitzsch, 1818”.

***Haematopinus suis* (Linnaeus, 1758)**

- Pediculus suis* Linnaeus, 1758: 611.  
*Pediculus urius* Nitzsch, 1818: 305.  
*Haematopinus suis* (Linnaeus); Denny 1842: 34, pl. 25: fig. 2.  
*Haematopinus urius* (Nitzsch, 1818); Giebel, 1874: 45, pl. 2: fig. 6.  
*Haematopinus suis* Linnaeus [sic]; Myers 1922: 12.  
*Haematopinus urius* Nitzsch [sic]; Thomson 1922: 339.  
*Haematopinus suis* (Linnaeus, 1758); Ferris 1933: 425, figs 252–256.  
*Haematopinus suis suis* (Linnaeus, 1758); Hopkins 1949b: 521.  
*Haematopinus suis* (Linnaeus, 1758); Ferris 1951: 91, figs 41–42.  
*Haematopinus suis* Linnaeus [sic]; Ineson 1954: 603.  
*Haematopinus suis* (L.); Helson 1956: 14, 16.  
*Haematopinus suis*; Miller 1971: 131.  
*Haematopinus suis* (Linnaeus, 1758); Wise 1977: 67.  
*Haematopinus suis* (Linnaeus, 1758); Tenquist & Charleston 2001: 503.  
*Haematopinus suis* (Linnaeus, 1758); Palma 2010: 409.

Syntypes presumed lost. See Ferris (1951: 94) and Kim *et al.* (1986: 87).

Type host: *Sus scrofa* Linnaeus, 1758.

New Zealand host: *Sus scrofa* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: ND, CL, TO, WA, WN, SD, MB, NC, MC, SC, Norfolk Island.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Myers (1922); Thomson (1922); Tillyard (1926: 135); Symmons (1952: 398, fig. 41); Ineson (1954); Helson (1956); Miller (1971); Whitten (1971: 161); Wise (1977); Tenquist & Charleston (1981: 266); King (1990: 369); Tenquist & Charleston (2001); Heath (2002: 48); King (2005: 343); Palma (2010).

Other significant references: Ferris (1933); Eichler (1941a: 369, fig. 35); Séguy (1944: 450, figs 721, 726); Webb (1949: 152); Hopkins (1949b); Ferris (1951: 91, figs 41–42); Ramcke (1965: 547, figs 1–31, 8 pls); Rudolph (1983: 16); Kim *et al.* (1986: 86, pl. 20); Green & Palma (1991: 21, 23); Durden & Musser (1994a: 17); Durden & Musser (1994b: 142); Tombesi & Castro (1995: 653); Barker (1996: 239); Durden (2001: 4, 8, fig. 1.1F).

Remarks: *Haematopinus suis*, also known as the “pig louse”, was introduced to New Zealand with its host by human agency (King 2005: 336).

**Family HOPLOPLEURIDAE Ewing, 1929**

Hoplopleuridae Ewing, 1929. *Manual External Parasites*: 133. Type genus: *Hoplopleura* Enderlein, 1904 (as Hoplopleurinae).

**Genus *Hoplopleura* Enderlein, 1904**

*Hoplopleura* Enderlein, 1904c. *Zool. Anz.* 28: 221. Type species: *Hoplopleura acanthopus* (Burmeister, 1838b) (by original designation).

***Hoplopleura pacifica* Ewing, 1924**

Figs 196–197

- Hoplopleura pacifica* Ewing, 1924: 9, figs 1b,c, 2.  
*Hoplopleura pacifica* Ewing, 1924; Hopkins 1949b: 481.  
*Hoplopleura oenomydis* Ferris, 1921; Ferris 1951: 139. In part.  
*Hoplopleura pacifica* Ewing, 1924; Ford-Robertson & Bull 1966: 223.  
*Hoplopleura pacifica* Ewing, 1924; Wise 1977: 67.  
 “*Hoplopleura oenomydis*” Tenquist & Charleston, 1981: 267 (not *Hoplopleura oenomydis* Ferris, 1921).  
*Hoplopleura pacifica* Ewing, 1924; Kim *et al.* 1986: 112, pl. 32.  
*Hoplopleura pacifica* Ewing, 1924; Tenquist & Charleston 2001: 504.  
*Hoplopleura pacifica* Ewing, 1924; Palma 2010: 409.

Lectotype ♀ in BPBM (Voss 1966: 30; Kim *et al.* 1986: 112).

Type host: *Rattus exulans hawaiiensis* Stone, 1917.

New Zealand host: *Rattus exulans* (Peale, 1848).

Other hosts: *Rattus argentiventer* (Robinson & Kloss, 1916); *Rattus norvegicus* (Berkenhout, 1769); *Rattus rattus* (Linnaeus, 1758); *Rattus tiomanicus* (Miller, 1900).

New Zealand localities: ND, AK, CL, NC, MC, SC, SI, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Ford-Robertson & Bull (1966); Pilgrim (1970: 77); Wise (1977); Tenquist & Charleston (1981); Gibson & Pilgrim (1986: 95); King (1990: 189); Roberts (1991: 354); Tenquist & Charleston (2001); King (2005: 171); Palma (2010).

Other significant references: Hopkins (1949b); Ferris (1951); Voss (1966: 29, figs 1–6); Johnson (1964: 71, figs 1, 5, 9–11, 15); Johnson (1972: 220, figs 13, 15–17, 19, 26); Kim *et al.* (1986); Green & Palma (1991: 21, 24); Durden & Musser (1994a: 31); Durden & Musser (1994b: 160); Barker (1996: 241); Durden (2001: 8).

Remarks: *Hoplopleura pacifica* was introduced to New Zealand and other Pacific Ocean islands with kiore by human agency (King 2005: 161).

### Family LINOGNATHIDAE Webb, 1946

Linognathidae Webb, 1946. *Proc. Zool. Soc. London*: 116: 107. Type genus: *Linognathus* Enderlein, 1905.

### Genus *Linognathus* Enderlein, 1905

*Trichaulus* Enderlein, 1904b. *Zool. Anz.* 28: 139, 141. Type species: *Trichaulus piliferus* (Burmeister, 1838b) = *Linognathus setosus* (von Olfers, 1816) (by original designation). Preoccupied by *Trichaulus* Mayr, 1885.

*Linognathus* Enderlein, 1905. *Zool. Anz.* 29: 194. *Nomen novum* for *Trichaulus* Enderlein, 1904.

### *Linognathus ovillus* (Neumann, 1907)

*Haematopinus ovillus* Neumann, 1907b: 522, figs 1–3.

*Haematopinus ovillus* Neumann, 1907; Evans 1907: 225, fig.

*Haematopinus ovillus* Neumann, 1907; Gilruth 1908: 194, 3 figs.

*Haematopinus ovillus* NeumaNN, Myers 1922: 12.

*Haematopinus ovillus* NeumaNN, Thomson 1922: 339.

*Linognathus ovillus* (Neumann, 1907); Tillyard 1926: 135.

*Linognathus ovillus* (Neumann, 1907); Ferris 1932b: 346, figs 209–210.

*Linognathus ovillus* (Neumann); Helson 1956: 14, 17.

*Linognathus ovillus*; Miller 1971: 131.

*Linognathus ovillus* (Neumann, 1907); Wise 1977: 66.

*Linognathus ovillus* (Neumann, 1907); Tenquist & Charleston 2001: 507.

*Linognathus ovillus* (Neumann, 1907); Palma 2010: 409.

Syntypes ♂♀, repository not confirmed, probably in the Ecole Vétérinaire de Toulouse, France (Kim *et al.* 1986: 126).

Type host: *Ovis aries* Linnaeus, 1758.

New Zealand host: *Ovis aries* Linnaeus, 1758.

Other hosts: None.

New Zealand locality: WN.

Geographic distribution: Australasia; North America; Asia; Falkland Islands.

New Zealand references: Neumann (1907b); Evans (1907); Gilruth (1908); Myers (1922); Thomson (1922); Tillyard (1926); Ferris (1932b); Murray (1955a: 22); Helson (1956); Murray (1963a,b); Miller (1971); Whitten (1971: 161); Wise (1977); Tenquist & Charleston (1981: 267); Durden & Musser (1994a: 41); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Webb (1949: 143); Hopkins (1949b: 534); Ferris (1951: 231, figs 99–100); Kim *et al.* (1986: 126, pl. 38); Durden & Musser (1994a: 41); Durden & Musser (1994b: 144); Barker (1996: 242); Durden (2001: 8).

Remarks: *Linognathus ovillus*, also known as “sheep face louse”, was introduced to New Zealand with its host by human agency (King 2005: 393), and originally described from specimens collected in New Zealand and Scotland.

***Linognathus pedalis* (Osborn, 1896)**

- Haematopinus pedalis* Osborn, 1896: 170.  
*Trichaulus pedalis* (Osborn, 1896); Enderlein, 1904b: 142.  
*Haematopinus pedalis* Osborn; Myers 1922: 12.  
*Haematopinus pedalis* Osborn; Thomson 1922: 339.  
*Linognathus pedalis* (Osborn, 1896); Tillyard 1926: 135, fig. O3.  
*Linognathus pedalis* (Osborn, 1896); Ferris 1932b: 344, figs 207–208.  
*Linognathus pedalis* (Osborn); Helson 1956: 14, 17.  
*Linognathus pedalis*; Miller 1971: 131, fig. 344.  
*Linognathus pedalis* (Osborn, 1896); Wise 1977: 66.  
*Linognathus pedalis* (Osborn, 1896); Tenquist & Charleston 2001: 508.  
*Linognathus pedalis* (Osborn, 1896); Palma 2010: 409.

Neotype ♀ in NHML (Kim *et al.* 1986: 128).

Type host: *Ovis aries* Linnaeus, 1758.

New Zealand host: *Ovis aries* Linnaeus, 1758.

Other host: *Oreamnos americanus* (Blainville, 1816).

New Zealand localities: WN, KA, NC, MC, SC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Myers (1922); Thomson (1922); Tillyard (1926); Helson (1956); Miller (1971: 131); Whitten (1971: 161); Wise (1977); Tenquist & Charleston (1981: 268); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Ferris (1932b); Webb (1949: 134); Hopkins (1949b: 534); Ferris (1951: 231, figs 101–102); Murray (1960a; 1963a); Kim *et al.* (1986: 128, pl. 39); Durden & Musser (1994a: 42); Durden & Musser (1994b: 144); Barker (1996: 242); Durden (2001: 4, 8).

Remarks: *Linognathus pedalis*, also known as “sheep foot louse”, was introduced to New Zealand with its host by human agency (King 2005: 393).

***Linognathus setosus* (von Olfers, 1816)**

Figs 198–199

- Pediculus setosus* von Olfers, 1816: 80.  
*Pediculus piliferus* Burmeister, 1838b: Species 13.  
*Haematopinus piliferous* [sic] (Burmeister); Denny 1842: 28, pl. 25: fig. 4.  
*Trichaulus piliferus* (Burmeister, 1838); Enderlein, 1904b: 142.  
*Linognathus piliferus* (Burmeister, 1838); Enderlein 1905: 194.  
*Haematopinus piliferus* Burmeister; Myers 1922: 12.  
*Haematopinus piliferus* [sic] Burm.; Thomson 1922: 339.  
*Linognathus piliferus* Burmeister; Tillyard 1926: 135.  
*Linognathus setosus* (von Olfers, 1816); Ferris 1932b: 340, figs 206–207, 216E.  
*Linognathus setosus* (Olfers); Helson 1956: 13, 17.  
*Linognathus setosus*; Miller 1971: 131.  
*Linognathus setosus* (von Olfers, 1816); Wise 1977: 66.  
*Linognathus setosus* (von Olfers, 1816); Tenquist & Charleston 2001: 508.  
*Linognathus setosus* (von Olfers, 1816); Palma 2010: 409.

Neotype ♀ in University of Heidelberg, Germany (Kim *et al.* 1986: 130).

Type host: *Canis familiaris* Linnaeus, 1758.

New Zealand host: *Canis familiaris* Linnaeus, 1758.

Other hosts: *Canis lupus* Linnaeus, 1758; *Canis latrans* Say, 1823; *Vulpes lagopus* (Linnaeus, 1758).

New Zealand localities: WI, WN, MC.

Geographic distribution: All continents, except Antarctica

New Zealand references: Myers (1922); Thomson (1922); Tillyard (1926); Helson (1956); Miller (1971: 131); Whitten (1971: 161); Pilgrim (1974: 1031, fig. 2); Wise (1977); Tenquist & Charleston (1981: 268); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Ferris (1932b); Séguy (1944: 439, figs 698–702); Webb (1946: 85); Webb (1949: 142); Hopkins (1949b: 496); Ferris (1951: 235, figs 103–104); Kim *et al.* (1986: 130, pl. 40); Green & Palma (1991: 21, 23); Durden & Musser (1994a: 42); Durden & Musser (1994b: 140); Barker (1996: 243); Palma & Jensen (2005: 59, 69).

Remarks: *Linognathus setosus*, also known as “dog sucking louse”, was introduced to New Zealand with its host by human agency (King 2005: 9, 258).

### ***Linognathus stenopsis* (Burmeister, 1838)**

*Pediculus stenopsis* Burmeister, 1838b: Species 16, fig. 3.

*Haematopinus stenopsis*; Denny 1842: 36.

*Trichaulus stenopsis* (Burmeister, 1838); Enderlein, 1904b: 142, fig. 13.

*Linognathus stenopsis* (Burmeister, 1838); Ferris 1932b: 349, figs 210–212.

*Linognathus stenopsis* (Burmeister); Helson 1956: 13, 17.

*Linognathus stenopsis* (Burmeister, 1838); Wise 1977: 66.

*Linognathus stenopsis* (Burmeister, 1838); Tenquist & Charleston 2001: 508.

*Linognathus stenopsis* (Burmeister, 1838); Palma 2010: 409.

Neotype ♀ in University of Heidelberg, Germany (Kim *et al.* 1986: 132).

Type host: *Capra hircus* Linnaeus, 1758.

New Zealand host: *Capra hircus* Linnaeus, 1758.

Other hosts: *Capra ibex* Linnaeus, 1758; *Rupicapra rupicapra* Couturier, 1938.

New Zealand localities: HB, TK, WN, SD, MB, BR, NC, MC, SC, WD.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Helson (1956); Whitten (1971: 161); Pilgrim (1974: 1031, fig. 1); Wise (1977); Tenquist & Charleston (1981: 268); King (1990: 420); Tenquist & Charleston (2001); King (2005: 387); Palma (2010).

Other significant references: Ferris (1932b); Séguy (1944: 440, figs 703–704); Webb (1949: 136); Hopkins (1949b: 533); Ferris (1951: 238, figs 105–106); Murray (1957a: 17); Ramchurn (1980: 6); Kim *et al.* (1986: 132, pl. 41); Durden & Musser (1994a: 42); Durden & Musser (1994b: 144); Barker (1996: 243); Durden (2001: 8).

Remarks: *Linognathus stenopsis*, also known as “goat sucking louse”, was introduced to New Zealand with its host by human agency (King 2005: 377).

### ***Linognathus vituli* (Linnaeus, 1758)**

*Pediculus vituli* Linnaeus, 1758: 611.

*Haematopinus vituli* (Linnaeus); Denny 1842: 31, pl. 25: fig. 3.

*Trichaulus vituli* (L.); Enderlein, 1904b: 142.

*Linognathus vituli* Linnaeus [sic]; Myers 1922: 12.

*Haematopinus vituli* Linnaeus; Thomson 1922: 339.

*Linognathus vituli* L. [sic]; Tillyard 1926: 135.

*Linognathus vituli* (Linnaeus, 1758); Ferris 1932b: 356, figs 214–216.

*Linognathus vituli* (L.); Helson 1956: 13, 17.

*Linognathus vituli*; Miller 1971: 131, fig. 343.

*Linognathus vituli* (Linnaeus, 1758); Wise 1977: 66.

*Linognathus vituli* (Linnaeus, 1758); Tenquist & Charleston 2001: 508.

*Linognathus vituli* (Linnaeus, 1758); Palma 2010: 409.

Neotype ♂ in University of Heidelberg, Germany (Kim *et al.* 1986: 134).

Type host: *Bos taurus* Linnaeus, 1758.

New Zealand host: *Bos taurus* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: HB, WN, NN, NC, MC, SC, SL, KE.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Myers (1922); Thomson (1922); Tillyard (1926); Helson (1956); Whitten (1970: 146); Helson (1970: 81); Miller (1971: 131); Whitten (1971: 161); Buchanan & Coles (1971: 197); Kettle (1972; 1974a); Kettle

& Lukies (1974; 1979); Wise (1977); Tenquist (1977: 286); Chalmers & Charleston (1980a,b,c); Kettle & Watson (1981); Tenquist & Charleston (1981: 268); Watson *et al.* (1996); Tenquist & Charleston (2001); Heath (2002: 48); Palma (2010); Heath (2013: 14, fig. centre).

Other significant references: Bedford (1929: 502); Ferris (1932b); Séguy (1944: 441, figs 705–707); Webb (1946: 83); Webb (1949: 143); Hopkins (1949b: 527); Ferris (1951: 241, figs 107–108); Callinan (1980: 484); Rudolph (1983: 16); Kim *et al.* (1986: 134, pl. 42); Green & Palma (1991: 21, 23); Durden & Musser (1994a: 44); Durden & Musser (1994b: 144); Barker (1996: 243); Durden (2001: 8); Bartlow *et al.* (2016: 222).

Remarks: *Linognathus vituli*, also known as “long-nosed cattle louse”, was introduced to New Zealand with its host by human agency (King 2005: 347).

### Genus *Solenopotes* Enderlein, 1904

*Solenopotes* Enderlein, 1904b. *Zool. Anz.* 28: 139, 143. Type species: *Solenopotes capillatus*, Enderlein, 1904 (by original designation).

#### *Solenopotes burmeisteri* (Fahrenheit, 1919)

Figs 200–201

*Linognathus burmeisteri* Fahrenheit, 1919: 23.

*Solenopotes burmeisteri* (Fahrenheit, 1919); Ferris 1932b: 406, figs 249a,b,d.

*Solenopotes burmeisteri* (Fahrenheit, 1919); Ferris 1951: 253.

*Solenopotes burmeisteri* (Fahrenheit, 1919); Andrews 1964: 106, fig. 3b.

*Solenopotes burmeisteri* (Fahrenheit, 1919); Kim & Weisser 1974: 117, figs 3–4, 16, 25, 33, 40, 48.

*Solenopotes burmeisteri* (Fahrenheit, 1919); Wise 1977: 67.

*Solenopotes burmeisteri* (Fahrenheit, 1919); Tenquist & Charleston 2001: 521.

*Solenopotes burmeisteri* (Fahrenheit, 1919); Palma 2010: 409.

Lectotype ♂ in the MLUH (Kim & Weisser 1974: 117).

Type host: *Cervus elaphus* Linnaeus, 1758.

New Zealand hosts: *Cervus elaphus scoticus* Lönningberg, 1906; *Odocoileus virginianus borealis* (Miller, 1900).

Other host: *Cervus nippon* Temminck, 1838.

New Zealand localities: WA, CO, DN, SI.

Geographic distribution: Eurasia; North Africa; North America; Southern South America; Australasia.

New Zealand references: Andrews (1964); Pilgrim (1970: 78); Wise (1977); Charleston (1980: 150); Tenquist & Charleston (1981: 273); King (1990: 451); Tenquist & Charleston (2001); King (2005: 415); Palma (2010).

Other significant references: Ferris (1932b); Webb (1949: 147); Hopkins (1949b: 524); Ferris (1951); Kim & Weisser (1974); Durden & Musser (1994a: 45); Durden & Musser (1994b: 142).

Remarks: *Solenopotes burmeisteri* was introduced to New Zealand with red deer by human agency (King 2005: 404). *Odocoileus virginianus borealis* is a new host record for *S. burmeisteri* (voucher specimens in MONZ), and it may be the result of a host switch from *Cervus elaphus*.

#### *Solenopotes capillatus* Enderlein, 1904

*Solenopotes capillatus* Enderlein, 1904b: 144, figs 14–15. Preserved by Opinion 1050 (I.C.Z.N. 1976).

*Solenopotes capillatus* Enderlein, 1904; Ewing 1929: 139, fig. 76.

*Solenopotes capillatus* Enderlein, 1904; Ferris 1932b: 397, figs 243–244.

*Solenopotes capillatus* Enderlein, 1904; Ferris 1951: 253, figs 113–114.

*Solenopotes capillatus*; Helson 1970: 81.

*Solenopotes capillatus*; Whitten 1971: 161.

*Solenopotes capillatus* Enderlein, 1904; Kim & Weisser 1974: 118, figs 5–6, 17, 27, 34, 43, 46, 49.

*Solenopotes capillatus* Enderlein, 1904; Tenquist & Charleston 2001: 521.

*Solenopotes capillatus* Enderlein, 1904; Palma 2010: 409.

Holotype ♂ in ZMHU, but subsequently lost (Kim & Weisser 1974: 120).

Type host: *Bos taurus* Linnaeus, 1758.

New Zealand host: *Bos taurus* Linnaeus, 1758.

Other hosts: None.

New Zealand locality: WI.

Geographic distribution: Eurasia; North America; Australasia; South Africa.

New Zealand references: Helson (1970); Whitten (1970: 146); Whitten (1971); Chalmers & Charleston (1980a: 198); Tenquist & Charleston (1981: 273); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Ferris (1932b); Séguy (1944: 446, figs 715–718); Webb (1946: 88); Webb (1949: 147); Hopkins (1949b: 527); Kim & Weisser (1973); Kim & Weisser (1974); International Commission on Zoological Nomenclature (1976); Kim *et al.* (1986: 138, pl. 44); Durden & Musser (1994a: 45); Durden & Musser (1994b: 144); Barker (1996: 244); Durden (2001: 8).

Remarks: *Solenopotes capillatus*, also known as “little blue cattle louse”, was introduced to New Zealand with cattle by human agency (King 2005: 347).

### Family PEDICULIDAE Leach, 1817

Pediculidae Leach, 1817. *Zool. Miscellany* 3: 64. Type genus: *Pediculus* Linnaeus, 1758.

### Genus *Pediculus* Linnaeus, 1758

*Pediculus* Linnaeus, 1758. *Systema Naturae* 10: 610. Type species: *Pediculus humanus* Linnaeus, 1758 (by subsequent designation).

### *Pediculus humanus capitis* De Geer, 1778

Figs 202–203

*Pediculus humanus capitis* De Geer, 1778a: 67, pl. 1: fig. 6. Preserved by Opinion 2333 (I.C.Z.N. 2014).

*pediculus* [sic] *humanus*; Polack 1838: 320.

*Pediculus capitis* Nitzsch [sic]; Hutton 1904: 353.

*Pediculus capitis* Nitzsch [sic]; Myers 1922: 12.

*Pediculus capitis* Nitzsch [sic]; Thomson 1922: 338.

“*Pediculus (Pediculus) humanus humanus*” Ewing, 1926: 22, text figs 1a, 2, 3a, 7, pl. 3: fig. 8 (not *Pediculus humanus* Linnaeus, 1758).

*Pediculus capitis* L. [sic]; Tillyard 1926: 135.

*Pediculus humanus* Linnaeus, 1758; Ferris 1935: 543, figs 307, 312B, 321B, 322G, 323B–324B, pls I, III. In part.

*Pediculus humanus capitis*; Buxton 1939: 10, figs 3, 7B.

*Pediculus humanus capitis* De Geer, 1778; Hopkins 1949b: 452.

*Pediculus humanus capitis* Deg. [sic]; Helson 1956: 14, 17.

*Pediculus capitis*; Miller 1971: 131, fig. 341.

*Pediculus humanus capitis*; Andrews 1976a: 61.

*Pediculus humanus capitis* De Geer, 1778; Wise 1977: 66.

*Pediculus capitis* De Geer, 1778; Busvine 1978: 7.

*Pediculus humanus capitis* De Geer, 1778; Savill 1990: 69.

*Pediculus humanus capitis* De Geer, 1778; Palma 2010: 295, 409.

Status, sex and repository of types unknown.

Type host: *Homo sapiens* Linnaeus, 1758.

New Zealand host: *Homo sapiens* Linnaeus, 1758.

Other hosts: None.

New Zealand localities: AK, HB, TK, WN, NC, MC, SC.

Geographic distribution: Cosmopolitan.

New Zealand references: Banks (1769: 186); Anonymous (1837: 3); Polack (1838); White & Doubleday (1843: 283); Hutton (1904); Myers (1922); Thomson (1922); Tillyard (1926); Helson (1956); Miller (1971); Andrews (1976a, 1976b: 62); Wise (1977); Andrews (1980); Andrews & Tonkin (1989: 199); Savill (1990); Palma (2010).

Other significant references: Ewing (1926); Ferris (1935); Buxton (1939); Busvine (1948: 1); Hopkins (1949b); Schaefer (1978: 669); Busvine (1978: 6); Maunder (1983: 8, figs 7–11); Bresciani *et al.* (1983: 24); Mumcuoglu & Zias (1988: 545); Palma (1991b: 194, pl. 21); Durden & Musser (1994a: 50); Durden & Musser (1994b: 140); Barker

(1996: 245); Palma & Jensen (2005: 60, 69); Leo & Barker (2005); Leo *et al.* (2005); Hänel & Palma (2007: 118, 128, 131); Reed *et al.* (2007); Light *et al.* (2008a,b); Arriaza *et al.* (2012); Rózsa & Apari (2012); Clayton *et al.* (2015: 169, 173); Reed *et al.* (2015: 203).

Remarks: The human head louse was introduced to New Zealand with humans. Both Polynesians and Europeans brought head lice with them (Anonymous 1837; Andrews 1976a: 59; Andrews 1976b: 64).

### ***Pediculus humanus humanus* Linnaeus, 1758**

Figs 204–205

*Pediculus humanus* Linnaeus, 1758: 610.

*Pediculus humanus corporis* De Geer, 1778a: 67, pl. 1: fig. 7.

*Pediculus humanus corporis* Retzius, 1783: 201. Suppressed by Opinion 2333 (I.C.Z.N. 2014).

*Pediculus vestimenti* Nitzsch, 1818: 305.

*Pediculus vestimenti* Nitzsch; Hutton 1904: 353.

*Pediculus humanus* Linnaeus; Myers 1922: 12.

*Pediculus corporis* De Geer; Thomson 1922: 338.

*Pediculus (Pediculus) humanus corporis* [sic] De Geer, 1778; Ewing 1926: 18.

*Pediculus vestimenti* Nitzsch; Tillyard 1926: 135.

*Pediculus humanus* Linnaeus, 1758; Ferris 1935: 543, figs 306, 308, 312A, 321A–324A. In part.

*Pediculus humanus corporis*; Buxton 1939: 10, figs 1, 4, 7A.

*Pediculus humanus humanus* Linnaeus, 1758; Hopkins 1949b: 452.

*Pediculus humanus humanus* L.; Helson 1956: 14, 17.

*Pediculus corporis*; Miller 1971: 131.

*Pediculus humanus humanus*; Andrews 1976a: 61.

*Pediculus humanus humanus* Linnaeus, 1758; Wise 1977: 66.

*Pediculus humanus* Linnaeus, 1758; Busvine 1978: 7.

*Pediculus humanus* Linnaeus, 1758; Tenquist & Charleston 2001: 515.

*Pediculus h. humanus* Linnaeus, 1758; Palma 2010: 296, 409.

Status, sex and repository of types unknown.

Type host: *Homo sapiens* Linnaeus, 1758.

New Zealand host: *Homo sapiens* Linnaeus, 1758.

Other hosts: None.

New Zealand locality: WN.

Geographic distribution: Cosmopolitan.

New Zealand references: Anonymous (1837: 3); Hutton (1904); Myers (1922); Thomson (1922); Tillyard (1926); Helson (1956); Miller (1971); Pilgrim (1974: 1033, figs 2, 3); Andrews (1976a, 1976b: 62); Wise (1977); Andrews (1980); Tenquist & Charleston (1981: 271); Tenquist & Charleston (2001); Leo & Barker (2005); Leo *et al.* (2005); Palma (2010).

Other significant references: Nuttall (1917: 80); Nuttall (1919: 335); Ewing (1926); Ferris (1935); Buxton (1939); Webb (1946: 77); Busvine (1948: 1); Hopkins (1949b); Ferris (1951: 261, figs 116–119); Schaefer (1978: 669); Busvine (1978: 6); Maunder (1983: 8, 18, fig. 12); Bresciani *et al.* (1983: 24); Rudolph (1983: 16); Kim *et al.* (1986: 150, pl. 48); Durden & Musser (1994a: 50); Durden & Musser (1994b: 140); Barker (1996: 245); Reed *et al.* (2007); Light *et al.* (2008b); Clayton *et al.* (2015: 169, 173); Reed *et al.* (2015: 203).

Remarks: The human body louse was introduced to New Zealand with humans. Both Polynesians and Europeans brought body lice with them. (Anonymous 1837; Thomson 1922; Andrews 1976a: 59; Andrews 1976b: 64).

### **Family PTHIRIDAE Ewing, 1929**

Pthiridae Ewing, 1929. *Manual External Parasites*: 132. Type genus: *Pthirus* Leach, 1815a (as “Phthiridae”).

### **Genus *Pthirus* Leach, 1815**

*Pthirus* Leach, 1815a. Brewster’s *Edinburgh Encyclopaedia* 9(1): 77. Type species: *Pediculus pubis* Linnaeus, 1758 (by subsequent designation) (see Hemming 1958: 54).



*Phthirus* Leach, 1817. *Zool. Miscellany* 3: 65. Invalid emendation (Kim *et al.* 1986: 209).

*Phthirius* Burmeister, 1835. *Handbuch Entomologie* 1: 58. Invalid emendation (Kim *et al.* 1986: 209).

*Phtirius* Mjöberg, 1910a. *Arkiv Zool.* 6(13): 171. Misspelling.

### ***Pthirus pubis* (Linnaeus, 1758)**

Figs 206–207

*Pediculus pubis* Linnaeus, 1758: 611.

*Pthirus inguinalis* Leach, 1815a: 77.

*Phthirus inguinalis* Leach, 1815; Leach, 1817: 65.

*Phthirius inguinalis* Leach, 1815; Burmeister 1835: 58.

*Phtirius inguinalis* Leach; Mjöberg 1910: 171.

*Phthirius pubis* Linnaeus [sic]; Myers 1922: 12.

*Phthirius inguinalis*; Thomson 1922: 338.

*Pthirus pubis* L. [sic]; Tillyard 1926: 135.

*Pthirus pubis* (Linnaeus, 1758); Ferris 1935: 603, figs 335–337.

*Pthirus pubis* (Linnaeus, 1758); Helson 1956: 14, 17.

*Pthirus pubis*; Miller 1971: 131, fig. 342.

*Pthirus pubis*; Andrews 1976a: 61.

*Pthirus pubis* (Linnaeus, 1758); Wise 1977: 66.

*Pthirus pubis* (Linnaeus, 1758); Kim *et al.* 1986: 210, pl. 76.

*Pthirus pubis* (Linnaeus, 1758); Tenquist & Charleston 2001: 518.

*Pthirus pubis* (Linnaeus, 1758); Palma 2010: 296.

*Pthirus pubis* (Linnaeus, 1758); Palma 2010: 409.

Syntypes probably lost (Kim *et al.* 1986: 210).

Type host: *Homo sapiens* Linnaeus, 1758.

New Zealand host: *Homo sapiens* Linnaeus, 1758.

Other hosts: Occasionally recorded on dogs and other animals (Lupidio 1980: 30; Kim *et al.* 1986: 210).

New Zealand localities: HB, TK, WN, NN, NC, MC, SC.

Geographic distribution: Cosmopolitan.

New Zealand references: Myers (1922); Thomson (1922); Tillyard (1926); Helson (1956); Miller (1971); Pilgrim (1974: 1033, fig. 4); Andrews (1976a, 1976b: 62); Wise (1977); Tenquist & Charleston (1981: 272); Tenquist & Charleston (2001); Palma (2010).

Other significant references: Nuttall (1918: 383, figs 1, 3–5, 7–9); Nuttall (1919: 345); Ferris (1935); Buxton (1939: 93, fig. 25); Séguy (1944: 458, figs 735–736); Webb (1946: 82); Hopkins (1949b: 452); Ferris (1951: 281, figs 122–124); Kraus & Glassman (1976); Lupidio (1980: 30); Maunder (1983: 7, figs 6, 13); Girling (1984: 207); Kim *et al.* (1986); Burns (1987: 741); Durden & Musser (1994a: 76); Durden & Musser (1994b: 140); Barker (1996: 247); Kenward (1999); Palma & Jensen (2005: 60, 69); Reed *et al.* (2007); Clayton *et al.* (2015: 169); Reed *et al.* (2015: 203).

Remarks: The pubic louse was introduced to New Zealand with humans. Both Polynesians and Europeans brought pubic lice with them (Andrews 1976a: 59; Andrews 1976b: 64).

### **Family POLYPLACIDAE Fahrenholz, 1912**

Polyplacidae Fahrenholz, 1912. *Jahrb. Niedersächs. Zool. Ver. Hannover* 2–4: 58. Type genus: *Polyplax* Enderlein, 1904 (as Polyplacinae).

### **Genus *Haemodipsus* Enderlein, 1904**

*Haemodipsus* Enderlein, 1904b. *Zool. Anz.* 28: 139, 143. Type species: *Haemodipsus lyriocephalus* (Burmeister, 1838b) (by original designation).

***Haemodipsus lyriocephalus* (Burmeister, 1838)**

Figs 208–209

- Pediculus lyriocephalus* Burmeister, 1838b: Species 11, fig. 7.  
*Haematopinus lyriocephalus* (Burmeister); Denny 1842: 27, pl. 24: fig. 4.  
*Haemodipsus lyriocephalus* (Burmeister, 1838); Enderlein 1904b: 143.  
*Haemodipsus lyriocephalus* (Burmeister, 1839) [sic]; Ferris 1932b: 330, figs 202–203.  
*Haemodipsus lyriocephalus* (Burmeister, 1839) [sic]; Wise 1977: 67.  
*Haemodipsus lyriocephalus* (Burmeister, 1839) [sic]; Tenquist & Charleston 2001: 503.  
*Haemodipsus lyriocephalus* (Burmeister, 1839) [sic]; Palma 2010: 409.

Status, sex and repository of types unknown.

Type host: *Lepus timidus* Linnaeus, 1758.New Zealand host: *Lepus europaeus occidentalis* de Winton, 1898.Other hosts: *Lepus europaeus* Pallas, 1778; *Lepus sinensis* J.E. Gray, 1832; *Lepus tolai* Pallas, 1778; *Lepus arcticus* Ross, 1819.

New Zealand locality: HB.

Geographic distribution: Eurasia; Australasia; South America.

New Zealand references: Pilgrim (1970: 78); Wise (1977); Tenquist &amp; Charleston (1981: 267); King (1990: 170); Tenquist &amp; Charleston (2001); King (2005: 156); Palma (2010).

Other significant references: Ferris (1932b); Séguy (1944: 443, figs 708–709); Webb (1946: 72); Hopkins (1949b: 453); Ferris (1951: 179, fig. 80); Broekhuizen (1971: 158, figs 1, 7); Durden &amp; Musser (1994a: 56); Durden &amp; Musser (1994b: 166).

Remarks: *Haemodipsus lyriocephalus* was introduced to New Zealand with hares by human agency (King 2005: 152).

As it can be seen in the above synonymy, there is a discrepancy in the literature regarding the original date of publication of *Pediculus lyriocephalus* Burmeister. This also applies to other species described in the same book (see Remarks under *Polyplax serrata* and *P. spinulosa* below). A comprehensive literature research has failed to show any evidence that the date “1839” for these species is correct. Therefore, I have changed it to “1838”, as stated on the title page of Burmeister (1838b). See also Durden *et al.* (2014: 258).

***Haemodipsus ventricosus* (Denny, 1842)**

- Haematopinus ventricosus* Denny, 1842: 30, pl. 25: fig. 6.  
*Haematopinus ventricosus* Denny, 1842; Enderlein 1904b: 143.  
*Haematopinus ventricosus* Denny; Myers 1922: 12.  
*Haematopinus ventricosus* Denny; Thomson 1922: 339.  
*Haemodipsus ventricosus* Denny [sic]; Tillyard 1926: 135.  
*Haemodipsus ventricosus* (Denny, 1842); Ferris 1932b: 332, figs 204–205.  
*Haemodipsus ventricosus* (Denny); Helson 1956: 14, 17.  
*Haemodipsus ventricosus*; Miller 1971: 131.  
*Haemodipsus ventricosus* (Denny, 1842); Wise 1977: 67.  
*Haemodipsus ventricosus* (Denny, 1842); Tenquist & Charleston 2001: 503.  
*Haemodipsus ventricosus* (Denny, 1842); Palma 2010: 409.

Lectotype ♀ in NHML (Kim *et al.* 1986: 170).Type host: *Oryctolagus cuniculus* (Linnaeus, 1758).New Zealand host: *Oryctolagus cuniculus* (Linnaeus, 1758).

Other hosts: None. See Durden &amp; Musser (1994a: 57) for erroneous host records.

New Zealand localities: BP, HB, WA, MC, AU, Macquarie Island.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Myers (1922); Thomson (1922); Tillyard (1926); Helson (1956); Bull (1960: 268); Miller (1971: 131); Wise (1977); Tenquist &amp; Charleston (1981: 267); King (1990: 156); Tenquist &amp; Charleston (2001); King (2005: 145); Palma (2010).

Other significant references: Ferris (1916b: 102, fig. 3); Ferris (1932b); Séguy (1944: 443, figs 710–711); Hopkins (1949b: 454); Ferris (1951: 179, figs 81–82); Kim *et al.* (1986: 170, pl. 57); Durden & Musser (1994a: 57); Durden & Musser (1994b: 166); Barker (1996: 246); Durden (2001: 8).

Remarks: *Haemodipsus ventricosus* was introduced to the New Zealand mainland and some Subantarctic islands with rabbits by human agency (Bull 1960; King 2005: 133).

### Genus *Polyplax* Enderlein, 1904

*Polyplax* Enderlein, 1904b. *Zool. Anz.* 28: 139, 142. Type species: *Polyplax spinulosa* (Burmeister, 1838b) (by original designation).

#### *Polyplax serrata* (Burmeister, 1838)

Figs 210–211

*Pediculus serratus* Burmeister, 1838b: Species 6.

*Haematopinus serratus*; Denny 1842: 36.

*Haematopinus serratus* (Burmeister, 1838); Enderlein 1904b: 142.

*Polyplax serrata* (Burmeister, 1839) [sic]; Ferris 1923: 191, figs 120B, E.

*Polyplax serrata* (Burmeister, 1839) [sic]; Gibson & Pilgrim 1986: 95.

*Polyplax serrata* (Burmeister, 1839) [sic]; Tenquist & Charleston 2001: 516.

*Polyplax serrata* (Burmeister, 1839) [sic]; Palma: 409.

Status, sex and repository of types uncertain (Kim *et al.* 1986: 204).

Type host: *Mus musculus* Linnaeus, 1758.

New Zealand host: *Mus musculus* Linnaeus, 1758.

Other hosts: *Crocidura leucodon* (Hermann, 1780); *Clethrionomys glareolus* (Schreber, 1780); *Microtus arvalis* (Pallas, 1778); and nine species of *Apodemus* (see Durden & Musser 1994a: 73).

New Zealand localities: WN, MC.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Tenquist & Charleston (1981: 271); Gibson & Pilgrim (1986); King (1990: 241); Tenquist & Charleston (2001); King (2005: 171, 219); Palma (2010).

Other significant references: Ferris (1923); Hopkins (1949b: 482); Ferris (1951: 210); Murray (1961: 1); Wegner (1974: 203); Kim *et al.* (1986: 204, pl. 74); Durden & Musser (1994a: 73); Durden & Musser (1994b: 159); Barker (1996: 246); Durden (2001: 8).

Remarks: *Polyplax serrata* was introduced to New Zealand with mice by human agency (King 2005: 205). King (1990: 189) cites *P. serrata* as a parasite of *Rattus exulans*, but that record is regarded here as accidental, either from natural straggling or from contamination by human agency.

As it can be seen in the above synonymy, there is a discrepancy in the literature regarding the original date of publication of *Pediculus serratus* Burmeister. This also applies to other species described in the same book (see Remarks under *Haemodipsus lyriocephalus* and *Polyplax spinulosa*). A comprehensive literature research has failed to show any evidence that the date “1839” for these species is correct. Therefore, I have changed it to “1838”, as stated on the title page of Burmeister (1838b). See also Durden *et al.* (2014: 258).

#### *Polyplax spinulosa* (Burmeister, 1838)

*Pediculus spinulosus* Burmeister, 1838b: Species 8.

*Haematopinus spinulosus* (Burmeister); Denny 1842: 26, pl. 24: fig. 5.

*Polyplax spinulosa* (Burmeister, 1838); Enderlein 1904b: 142.

*Polyplax spinulosa* (Burmeister, 1839) [sic]; Ferris 1923: 187, figs 119, 120A, D, F, H.

*Polyplax spinulosa* (Burmeister, 1839) [sic]; Clay 1964a: 233.

*Polyplax spinulosa* (Burmeister, 1839) [sic]; Watt, 1971: 238, 244.

*Polyplax spinulosa* (Burmeister, 1839) [sic]; Wise 1977: 67.

*Polyplax spinulosa* (Burmeister, 1839) [sic]; Gibson & Pilgrim 1986: 95.

*Polyplax spinulosa* (Burmeister, 1839) [sic]; Tenquist & Charleston 2001: 517.

*Polyplax spinulosa* (Burmeister, 1839) [sic]; Palma 2010: 409.

Status, sex and repository of types uncertain (Kim *et al.* 1986: 206).

Type host: *Rattus norvegicus* (Berkenhout, 1769).

New Zealand hosts: *Rattus exulans* (Peale, 1848); *Rattus norvegicus* (Berkenhout, 1769); *Rattus rattus* (Linnaeus, 1758).

Other hosts: *Rattus argentiventer* (Robinson & Kloss, 1916); *Rattus nitidus* (Hodgson, 1845); *Rattus tanezumi* Temminck, 1844; *Rattus turkestanicus* (Satunin, 1903); *Bandicota bengalensis* (J.E. Gray & Hardwicke, 1833).

New Zealand localities: ND, AK, CL, BP, TO, HB, WA, WN, BR, NC, MC, SC, KE, SI, Macquarie Island.

Geographic distribution: All continents, except Antarctica.

New Zealand references: Clay (1964a); Gressitt (1964: 539); Watson (1967: 74); Watt (1971); Wise (1977); Tenquist & Charleston (1981: 271); Gibson & Pilgrim (1986); King (1990: 204, 221); Tenquist & Charleston (2001); Palma & Horning (2002: 14, 18); King (2005: 184, 199); Palma (2010).

Other significant references: Enderlein (1905: 192); Fahrenheit (1912: 30, figs 8–10, pl. 2: figs 8–13); Cummings (1915: 256, 268, figs 7, 15–16); Ferris (1923); Séguy (1944: 432, figs 677–682); Hopkins (1949b: 481); Kim *et al.* (1986: 206, pl. 75); Green & Palma (1991: 21, 24); Durden & Musser (1994a: 73); Durden & Musser (1994b: 160); Barker (1996: 246); Durden (2001: 8); Palma & Peck (2013: 67).

Remarks: *Polyplax spinulosa* was introduced to New Zealand with rats by human agency (King 2005: 176, 189). As it can be seen in the above synonymy, there is a discrepancy in the literature regarding the original date of publication of *Pediculus spinulosus* Burmeister. This also applies to other species described in the same book (see Remarks under *Haemodipsus lyriocephalus* and *Polyplax serrata* above). A comprehensive literature research has failed to show any evidence that the date “1839” for these species is correct. Therefore, I have changed it to “1838”, as stated on the title page of Burmeister (1838b). See also Durden *et al.* (2014: 258).

## SPECIES & SUBSPECIES DELETED FROM THE NEW ZEALAND LOUSE FAUNA

### ***Goniodes gigas* (Taschenberg, 1879)**

Wise (1977: 60) recorded this species following its inclusion by Helson (1956: 13, as *Goniocotes gigas*) in a list of arthropods affecting livestock in New Zealand. However, no specimens of this species have been collected, examined or recorded since that report. Instead, another similar species, *Goniodes dissimilis* (Denny, 1842) has been found parasitising chickens in New Zealand (Thomson 1922; Whitten 1971; Palma 1999: 384). Helson’s (1956) record is most likely the result of a misidentification. Therefore, I delete *Goniodes gigas* from the New Zealand louse fauna.

### ***Holomenopon clypeilargum* Eichler, 1943**

Price (1971: 643) identified one female from *Aythya novaeseelandiae* as *Holomenopon clypeilargum*, and Price *et al.* (2003: 112, 279) repeated that host-lice association. However, examination of more specimens of both sexes showed that they belong to a yet unnamed and undescribed species. Therefore, Pilgrim & Palma (1982: 17) listed this record under *Aythya novaeseelandiae* as “*Holomenopon* sp.” and included an explanatory note (Pilgrim & Palma 1982: 30, note 19). See also entry “*Holomenopon* species” above.

### ***Lunaceps numenii numenii* (Denny, 1842)**

Pilgrim & Palma (1982: 20) recorded *Lunaceps numenii numenii* (Denny, 1842) from *Numenius madagascariensis*. However, that population is now known as *Lunaceps numenii madagascariensis* Gustafsson & Olsson, 2012. Therefore, I delete the nominate subspecies from the New Zealand louse fauna.

### ***Lunaceps incoenis* (Kellogg & Chapman, 1899)**

Pilgrim & Palma (1982: 21) recorded *Lunaceps incoenis* (Kellogg & Chapman, 1899) *s. l.* from *Calidris ferruginea*. However, that population is now known as *Lunaceps falcinellus* Timmermann, 1954, as determined by Gustafsson & Olsson (2012b: 24). Therefore, I delete *Lunaceps incoenis* from the New Zealand louse fauna.

***Perineus concinnus* (Kellogg & Chapman, 1899)**

Wise (1977: 63) listed this species following a report by Harrison (1937: 29). However, Pilgrim & Palma (1979: 177; 1982: 29) showed that it was a misidentification of *Perineus circumfasciatus* Kéler, 1957. Therefore, I delete *Perineus concinnus* from the New Zealand louse fauna.

***Sturnidoecus* sp.**

Watt (1971: 238, 244) listed 4 nymphs of “*Sturnidoecus* sp.” from *Prosthemadera novaeseelandiae novaeseelandiae*. Re-examination of those nymphs shows that it is a misidentification of a *Saemundsonia* sp. and most likely the result of a contamination (see also Pilgrim & Palma 1982: 31, note 25).

## HOST-PARASITE LIST

Louse taxa are listed under each host taxon in alphabetical order according to genera.

- ◆ Denotes louse species or subspecies which are **endemic** to the New Zealand Subregion.
- ♣ Denotes louse species or subspecies which are **new records** from the New Zealand Subregion.
- ε Denotes host species or subspecies which breed in the New Zealand Subregion **only**, even if their geographical ranges extend outside the Subregion.
- \* Denotes host species or subspecies which have been **introduced** to the New Zealand Subregion by human agency.

**AVES****CASUARIIFORMES****APTERYGIDAE**

- |  |                                  |
|--|----------------------------------|
| ° <i>Apteryx mantelli</i> Bartlett, 1852<br>◆ <i>Apterygon mirum</i> Clay, 1961<br>◆ <i>Rallicola</i> ( <i>Aptericola</i> ) <i>rodericki</i> Palma, 1991   | <b>North Island brown kiwi</b>   |
| ° <i>Apteryx rowi</i> Tennyson, Palma, Robertson, Worthy & Gill, 2003<br>◆ <i>Apterygon okarito</i> Palma & Price, 2004<br>◆ <i>Rallicola</i> ( <i>Aptericola</i> ) <i>gadawi</i> Harrison, 1915 <i>sensu lato</i>                               | <b>Okarito brown kiwi</b>        |
| ° <i>Apteryx australis australis</i> Shaw, 1813<br>◆ <i>Apterygon dumosum</i> Tandan, 1972<br>◆ <i>Rallicola</i> ( <i>Aptericola</i> ) <i>gadawi</i> Harrison, 1915  | <b>South Island brown kiwi</b>   |
| ° <i>Apteryx australis lawryi</i> Rothschild, 1893<br>◆ <i>Apterygon dumosum</i> Tandan, 1972<br>◆ <i>Rallicola</i> ( <i>Aptericola</i> ) <i>gadawi</i> Harrison, 1915 <i>sensu lato</i>   | <b>Stewart Island brown kiwi</b> |
| ° <i>Apteryx owenii</i> Gould, 1847<br>◆ <i>Apterygon dumosum</i> Tandan, 1972<br>◆ <i>Rallicola</i> ( <i>Aptericola</i> ) <i>gadawi</i> Harrison, 1915 <i>sensu lato</i><br>◆ <i>Rallicola</i> ( <i>Aptericola</i> ) <i>pilgrimi</i> Clay, 1972 | <b>Little spotted kiwi</b>       |
| ° <i>Apteryx haastii</i> Potts, 1872<br>◆ <i>Apterygon hintoni</i> Clay, 1966<br>◆ <i>Rallicola</i> ( <i>Aptericola</i> ) <i>gracilentus</i> Clay, 1953  | <b>Great spotted kiwi</b>        |

**GALLIFORMES****PHASIANIDAE**

- |   |                         |
|---|-------------------------|
| * <i>Callipepla californica brunnescens</i> (Ridgway, 1884)<br><i>Goniodes stefani</i> Clay & Hopkins 1955<br><i>Lagopoecus docophoroides</i> (Piaget, 1880)<br><i>Oxylipeurus ellipticus</i> (Kéler, 1958) | <b>California quail</b> |
|---|-------------------------|

- \* ***Numida meleagris* (Linnaeus, 1758) [captive]** **Helmeted Guineafowl**  
*Menopon gallinae* (Linnaeus, 1758)  
*Lipeurus caponis* (Linnaeus, 1758)
- \* ***Alectoris chukar* (J.E. Gray, 1830)** **Chukor**  
*Cuclotogaster heterographus* (Nitzsch [*in* Giebel], 1866)  
*Goniocotes pusillus* (Nitzsch [*in* Giebel], 1866)  
*Goniodes dispar* Burmeister, 1838
- \* ***Perdix perdix perdix* (Linnaeus, 1758)** **Grey partridge**  
*Amyrsidea (Argimenopon) perdicis* (Denny, 1842)  
*Lipeurus maculosus maculosus* Clay, 1938
- \* ***Coturnix ypsilophora australis* (Latham, 1802)** **Australian brown quail**  
*Cuclotogaster synoicus* (Clay, 1938)  
*Goniodes retractus* Le Souëf, 1902
- \* ***Colinus virginianus taylori* Lincoln, 1915** **Midwestern bobwhite quail**  
*Goniodes ortygis* (Denny, 1842)  
*Oxylpeurus clavatus* (McGregor, 1917)
- \* ***Gallus gallus gallus* (Linnaeus, 1758)** **Feral chicken**  
*Cuclotogaster heterographus* (Nitzsch [*in* Giebel], 1866)  
*Goniocotes gallinae* (De Geer, 1778)  
*Goniodes dissimilis* (Denny, 1842)  
*Lipeurus caponis* (Linnaeus, 1758)  
*Menacanthus pallidulus* (Neumann, 1912)  
*Menacanthus stramineus* (Nitzsch, 1818)  
*Menopon gallinae* (Linnaeus, 1758)
- \* ***Phasianus colchicus* Linnaeus, 1758** **Common pheasant**  
*Amyrsidea (Argimenopon) perdicis* (Denny, 1842)  
*Goniocotes chrysocephalus* Giebel, 1874  
*Goniodes colchici* (Denny, 1842)  
*Lipeurus maculosus maculosus* Clay, 1938  
*Menacanthus pallidulus* (Neumann, 1912)  
*Oxylpeurus mesopelios colchicus* Clay, 1938
- \* ***Pavo cristatus* Linnaeus, 1758** **Peafowl**  
*Amyrsidea (Argimenopon) minuta* Emerson, 1961  
*Colpocephalum tausi* (Ansari, 1951)  
*Goniodes pavonis* (Linnaeus, 1758)
- \* ***Meleagris gallopavo gallopavo* Linnaeus, 1758** **Wild turkey**  
*Chelopistes meleagridis* (Linnaeus, 1758)  
*Lipeurus caponis* (Linnaeus, 1758)  
*Menacanthus stramineus* (Nitzsch, 1818)  
*Oxylpeurus polytrapezius polytrapezius* (Burmeister, 1838)

## ANSERIFORMES

## ANATIDAE

- Dendrocygna eytoni* (Eyton, 1838)** **Plumed whistling duck**  
*Holomenopon leucoxanthum* (Burmeister, 1838)  
♣ *Acidoproctus emersoni* Timmermann, 1962
- \* ***Cygnus olor* (Gmelin, 1789)** **Mute swan**  
*Anatoecus icterodes olor* Złotorzycka, 1970  
*Ornithobius bucephalus* (Giebel, 1874)
- Cygnus atratus* (Latham, 1790)** **Black swan**  
*Anatoecus dentatus magnicornutus* Złotorzycka, 1970  
*Anatoecus icterodes olor* Złotorzycka, 1970

- Anatoecus singhi* Palma, 2015  
*Ciconiphilus pectiniventris* (Harrison, 1916)  
*Holomenopon leucoxanthum* (Burmeister, 1838)  
*Ornithobius fuscus* Le Souëf, 1902  
*Trinoton nigrum* Le Souëf, 1902
- \* ***Anser anser* (Linnaeus, 1758)** Greylag goose  
*Anaticola anseris* (Linnaeus, 1758)  
*Ciconiphilus pectiniventris* (Harrison, 1916)
- \* ***Branta canadensis maxima* Delacour, 1951** Canada goose  
*Anaticola anseris* (Linnaeus, 1758)  
*Ciconiphilus pectiniventris* (Harrison, 1916)  
*Holomenopon leucoxanthum* (Burmeister, 1838)  
*Ornithobius goniopleurus* (Denny, 1842)  
*Trinoton querquedulae* (Linnaeus, 1758)
- ° ***Tadorna variegata* (Gmelin, 1789)** Paradise shelduck  
*Anaticola magnificus* Ansari, 1955  
*Anatoecus dentatus* (Scopoli, 1763) *sensu lato*  
*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*  
*Holomenopon tadornae* (Gervais, 1844)  
*Trinoton querquedulae* (Linnaeus, 1758)
- Tadorna tadornoides* (Jardine & Selby, 1828)** Chestnut-breasted shelduck  
*Anaticola magnificus* Ansari, 1955  
*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*  
*Holomenopon tadornae* (Gervais, 1844)
- Chenonetta jubata* (Latham, 1802)** Australian wood duck  
*Anaticola* species  
*Anatoecus dentatus* (Scopoli, 1763) *sensu lato*
- ° ***Hymenolaimus malacorhynchos* (Gmelin, 1789)** Blue duck  
♦ *Acidoproctus gottwaldhirschi* Eichler, 1958  
*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*
- Anas gracilis* Buller, 1869** Grey teal  
*Anaticola crassicornis* (Scopoli, 1763)  
*Anatoecus dentatus* (Scopoli, 1763) *sensu lato*  
*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*  
*Holomenopon leucoxanthum* (Burmeister, 1838)  
*Trinoton querquedulae* (Linnaeus, 1758)
- ° ***Anas chlorotis* G.R. Gray, 1845** Brown teal  
*Anaticola* species  
*Anatoecus dentatus* (Scopoli, 1763) *sensu lato*  
*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*  
*Trinoton querquedulae* (Linnaeus, 1758)
- ° ***Anas aucklandica* (G.R. Gray, 1849)** Auckland Island teal  
*Holomenopon leucoxanthum* (Burmeister, 1838)
- \* ***Anas platyrhynchos platyrhynchos* Linnaeus, 1758** Mallard  
*Anaticola crassicornis* (Scopoli, 1763)  
*Anatoecus dentatus* (Scopoli, 1763) *sensu lato*  
*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*  
*Holomenopon leucoxanthum* (Burmeister, 1838)  
*Trinoton querquedulae* (Linnaeus, 1758)
- Anas superciliosa* Gmelin, 1789** Grey duck  
*Anaticola crassicornis* (Scopoli, 1763)  
*Anatoecus dentatus* (Scopoli, 1763) *sensu lato*

*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*  
*Trinoton querquedulae* (Linnaeus, 1758)

***Anas rhynchotis* Latham, 1802**

*Anaticola crassicornis* (Scopoli, 1763)  
*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*  
*Holomenopon* species 2  
*Trinoton querquedulae* (Linnaeus, 1758)

**Australasian shoveler**

**\**Aythya novaeseelandiae* (Gmelin, 1789)**

*Anaticola crassicornis* (Scopoli, 1763)  
*Anatoecus dentatus* (Scopoli, 1763) *sensu lato*  
*Anatoecus icterodes* (Nitzsch, 1818) *sensu lato*  
 ♦ *Holomenopon* species 1  
*Trinoton querquedulae* (Linnaeus, 1758)

**New Zealand scaup**

**PODICIPEDIFORMES**

**PODICIPEDIDAE**

***Podiceps cristatus australis* Gould, 1844**

♣ *Aquanirmus podiceps* (Denny, 1842)

**Australasian crested grebe**

**\**Poliiocephalus rufopectus* (G.R. Gray, 1843)**

♦ *Aquanirmus australis* Kettle, 1974

**New Zealand dabchick**

**SPHENISCIFORMES**

**SPHENISCIDAE**

***Aptenodytes forsteri* G.R. Gray, 1844**

*Austrogoniodes mawsoni* Harrison, 1937

**Emperor penguin**

***Aptenodytes patagonicus* J.F. Miller, 1778**

*Nesiotinus demersus* Kellogg, 1903

**King penguin**

***Pygoscelis adeliae* (Hombron & Jacquinot, 1841)**

*Austrogoniodes antarcticus* Harrison, 1937

**Adelie penguin**

***Eudyptes filholi* Hutton, 1879**

*Austrogoniodes cristati* Kéler, 1952  
*Austrogoniodes hamiltoni* Harrison, 1937  
*Austrogoniodes macquariensis* Harrison, 1937

**Eastern rockhopper penguin**

**\**Eudyptes pachyrhynchus* G.R. Gray, 1845**

*Austrogoniodes concii* (Kéler, 1952)  
*Austrogoniodes cristati* Kéler, 1952

**Fiordland crested penguin**

**\**Eudyptes robustus* Oliver, 1953**

*Austrogoniodes concii* (Kéler, 1952)  
*Austrogoniodes cristati* Kéler, 1952

**Snares crested penguin**

**\**Eudyptes sclateri* Buller, 1888**

*Austrogoniodes concii* (Kéler, 1952)  
*Austrogoniodes cristati* Kéler, 1952  
*Austrogoniodes strutheus* Harrison, 1915 *nomen dubium*

**Erect-crested penguin**

***Eudyptes chrysolophus* (Brandt, 1837)**

*Austrogoniodes cristati* Kéler, 1952  
*Austrogoniodes macquariensis* Harrison, 1937 *sensu lato*

**Macaroni penguin**

**\**Eudyptes schlegeli* Finsch, 1876**

*Austrogoniodes cristati* Kéler, 1952  
*Austrogoniodes hamiltoni* Harrison, 1937

**Royal penguin**



*Austrogoniodes macquariensis* Harrison, 1937

° *Megadyptes antipodes* (Hombron & Jacquinot, 1841)

*Austrogoniodes concii* (Kéler, 1952)

◆ *Austrogoniodes vanalphenae* Banks & Palma, 2003

Yellow-eyed penguin

*Eudyptula minor* (J.R. Forster, 1781)

*Austrogoniodes waterstoni* (Cummings, 1914)

Little penguin

PROCELLARIIFORMES

DIOMEDEIDAE

*Diomedea exulans* Linnaeus, 1758

*Austromenopon affine* (Piaget, 1890)

*Docophoroides brevis* (Dufour, 1835)

*Episbates pederiformis* (Dufour, 1835)

*Harrisoniella hopkinsi* Eichler, 1952

*Paraclisis hyalina* (Neumann, 1911)

*Perineus concinnoides* Kéler, 1957

Wandering albatross

° *Diomedea antipodensis antipodensis* Robertson & Warham, 1992

*Austromenopon affine* (Piaget, 1890)

*Docophoroides brevis* (Dufour, 1835)

*Episbates pederiformis* (Dufour, 1835)

*Harrisoniella hopkinsi* Eichler, 1952

*Paraclisis hyalina* (Neumann, 1911)

Antipodean albatross

° *Diomedea antipodensis gibsoni* Robertson & Warham, 1992

*Docophoroides brevis* (Dufour, 1835)

*Episbates pederiformis* (Dufour, 1835)

*Paraclisis hyalina* (Neumann, 1911)

Gibson's albatross

° *Diomedea epomophora* Lesson, 1825

*Austromenopon affine* (Piaget, 1890)

*Docophoroides brevis* (Dufour, 1835)

*Episbates pederiformis* (Dufour, 1835)

*Harrisoniella hopkinsi* Eichler, 1952

*Paraclisis hyalina* (Neumann, 1911)

*Perineus concinnoides* Kéler, 1957

Southern royal albatross

° *Diomedea sanfordi* Murphy, 1917

*Austromenopon affine* (Piaget, 1890)

*Docophoroides brevis* (Dufour, 1835)

*Episbates pederiformis* (Dufour, 1835)

*Harrisoniella hopkinsi* Eichler, 1952

*Paraclisis hyalina* (Neumann, 1911)

*Perineus concinnoides* Kéler, 1957

Northern royal albatross

*Thalassarche chlororhynchus* (Gmelin, 1789)

*Paraclisis diomedae* (J.C. Fabricius, 1775)

*Perineus circumfasciatus* Kéler, 1957

Atlantic yellow-nosed-albatross

*Thalassarche chrysostoma* (J.R. Forster, 1785)

*Austromenopon navigans* (Kellogg, 1896)

*Austromenopon pinguis* (Kellogg, 1896)

*Docophoroides simplex* (Waterston, 1914)

*Paraclisis diomedae* (J.C. Fabricius, 1775)

*Perineus circumfasciatus* Kéler, 1957

*Saemundssonina* (*Saemundssonina*) *albatrossa* Palma, 2012

Grey-headed albatross

*Thalassarche melanophris* (Temminck, 1828)

*Austromenopon navigans* (Kellogg, 1896)

*Austromenopon pinguis* (Kellogg, 1896)

Black-browed albatross

*Docophoroides harrisoni* Waterston, 1917  
*Docophoroides simplex* (Waterston, 1914)  
*Harrisoniella ferox* (Giebel, 1867)  
*Paraclisis diomedea* (J.C. Fabricius, 1775)  
*Perineus circumfasciatus* Kéler, 1957

° ***Thalassarche impavida* Mathews, 1912**

*Austromenopon pinguis* (Kellogg, 1896)  
*Docophoroides simplex* (Waterston, 1914)  
*Harrisoniella ferox* (Giebel, 1867)  
*Paraclisis diomedea* (J.C. Fabricius, 1775)  
*Perineus circumfasciatus* Kéler, 1957  
*Saemundssonina (Saemundssonina) albatrossa* Palma, 2012

**Campbell black-browed albatross**

° ***Thalassarche bulleri bulleri* (Rothschild, 1888)**

*Austromenopon navigans* (Kellogg, 1896)  
*Austromenopon pinguis* (Kellogg, 1896)  
*Docophoroides harrisoni* Waterston, 1917  
*Harrisoniella ferox* (Giebel, 1867)  
*Paraclisis diomedea* (J.C. Fabricius, 1775)  
*Perineus circumfasciatus* Kéler, 1957

**Southern Buller's albatross**

° ***Thalassarche bulleri platei* (Reichenow, 1898)**

*Austromenopon navigans* (Kellogg, 1896)  
*Austromenopon pinguis* (Kellogg, 1896)  
*Docophoroides harrisoni* Waterston, 1917  
*Harrisoniella ferox* (Giebel, 1867)  
*Paraclisis diomedea* (J.C. Fabricius, 1775)  
*Perineus circumfasciatus* Kéler, 1957

**Northern Buller's albatross**

° ***Thalassarche cauta stadi* Falla, 1933**

*Austromenopon navigans* (Kellogg, 1896)  
*Austromenopon pinguis* (Kellogg, 1896)  
*Docophoroides harrisoni* Waterston, 1917  
*Harrisoniella ferox* (Giebel, 1867)  
*Paraclisis diomedea* (J.C. Fabricius, 1775)  
*Perineus circumfasciatus* Kéler, 1957

**New Zealand white-capped albatross**

° ***Thalassarche eremita* Murphy, 1930**

*Austromenopon navigans* (Kellogg, 1896)  
*Docophoroides harrisoni* Waterston, 1917  
*Harrisoniella ferox* (Giebel, 1867)  
*Paraclisis diomedea* (J.C. Fabricius, 1775)  
*Perineus circumfasciatus* Kéler, 1957

**Chatham Island albatross**

° ***Thalassarche salvini* (Rothschild, 1878)**

*Austromenopon navigans* (Kellogg, 1896)  
*Austromenopon pinguis* (Kellogg, 1896)  
*Docophoroides harrisoni* Waterston, 1917  
*Harrisoniella ferox* (Giebel, 1867)  
*Paraclisis diomedea* (J.C. Fabricius, 1775)  
*Perineus circumfasciatus* Kéler, 1957

**Salvin's albatross**

***Phoebetria fusca* (Hilsenberg, 1822)**

*Perineus circumfasciatus* Kéler, 1957

**Sooty albatross**

***Phoebetria palpebrata* (J.R. Forster, 1785)**

*Austromenopon pinguis* (Kellogg, 1896)  
*Docophoroides simplex* (Waterston, 1914)  
*Paraclisis diomedea* (J.C. Fabricius, 1775)  
*Perineus circumfasciatus* Kéler, 1957  
*Saemundssonina (Saemundssonina) albatrossa* Palma, 2012

**Light-mantled sooty albatross**

## PROCELLARIIDAE

- Macronectes giganteus* (Gmelin, 1789)** Southern giant petrel  
*Austromenopon ossifragae* (Eichler, 1949)  
*Docophoroides murphyi* (Kellogg, 1914)  
*Paraclisis miriceps* (Kellogg & Kuwana, 1902)  
*Paraclisis obscura* (Rudow, 1869)  
*Perineus macronekti* Palma & Pilgrim, 1988  
*Saemundssonina* (*Saemundssonina*) *gaini* (Neumann, 1913)
- Macronectes halli* Mathews, 1912** Northern giant petrel  
*Austromenopon ossifragae* (Eichler, 1949)  
*Docophoroides murphyi* (Kellogg, 1914)  
*Paraclisis obscura* (Rudow, 1869)  
*Perineus macronekti* Palma & Pilgrim, 1988  
*Saemundssonina* (*Saemundssonina*) *gaini* (Neumann, 1913)
- Fulmarus glacialisoides* (A. Smith, 1826)** Antarctic fulmar  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon brevifimbriatum* (Piaget, 1880)  
*Perineus nigrolimbatus* (Giebel, 1874)  
*Saemundssonina* (*Saemundssonina*) *bicolor* (Rudow, 1870)
- Thalassoica antarctica* (Gmelin, 1789)** Antarctic petrel  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon brevifimbriatum* (Piaget, 1880)  
*Pseudonirmus lugubris* (Taschenberg, 1882)  
*Saemundssonina* (*Saemundssonina*) *antarctica* (Wood, 1937)
- Daption capense capense* (Linnaeus, 1758)** Cape petrel  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon brevifimbriatum* (Piaget, 1880)  
*Pseudonirmus gurlti* (Taschenberg, 1882)  
*Saemundssonina* (*Saemundssonina*) *stammeri* Timmermann, 1959
- Daption capense australe* Mathews, 1913** Snares Cape petrel  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Pseudonirmus gurlti* (Taschenberg, 1882)  
*Saemundssonina* (*Saemundssonina*) *stammeri* Timmermann, 1959
- Pagodroma nivea nivea* (G. Forster, 1777)** Lesser snow petrel  
*Pseudonirmus charcoti* (Neumann, 1907)
- Pagodroma nivea major* (Schlegel, 1863)** Greater snow petrel  
*Pseudonirmus charcoti* (Neumann, 1907)  
*Saemundssonina* (*Saemundssonina*) *antarctica* (Wood, 1937)
- Lugensa brevirostris* (Lesson, 1833)** Kerguelen petrel  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Bedfordiella unica* Thompson, 1937  
*Longimenopon* species  
*Saemundssonina* (*Saemundssonina*) *pterodromae* Timmermann, 1959
- \**Pterodroma macroptera gouldi* (Hutton, 1869)** Grey-faced petrel  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Halipeurus procellariae* (J.C. Fabricius, 1775)  
*Naubates* (*Guenterion*) *heteroproctus* Harrison, 1937  
*Saemundssonina* (*Puffinoecus*) *enderleini* (Eichler, 1949)  
*Trabeculus schillingi* Rudow, 1866 *sensu lato*

- Pterodroma lessonii* (Garnot, 1826)**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Halipeurus procellariae* (J.C. Fabricius, 1775)  
*Naubates (Guenterion) lessonii* Palma & Pilgrim, 2002  
*Saemundssonina (Puffinoecus) enderleini* (Eichler, 1949)  
*Trabeculus schillingi* Rudow, 1866 *sensu lato*
- White-headed petrel**
- Pterodroma solandri* (Gould, 1844)**  
*Naubates (Naubates) thieli* Timmermann, 1965  
*Trabeculus schillingi* Rudow, 1866 *sensu lato*
- Providence petrel**
- \**Pterodroma magentae* (Giglioli & Salvadori, 1869)**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Halipeurus procellariae* (J.C. Fabricius, 1775)  
*Halipeurus theresae* Timmermann, 1969  
*Longimenopon* species  
*Saemundssonina (Puffinoecus)* species
- Chatham Island taiko**
- Pterodroma neglecta neglecta* (Schlegel, 1863)**  
*Austromenopon popellus* (Piaget, 1890)  
*Halipeurus kermadecensis* (Johnston & Harrison, 1912)  
*Longimenopon* species  
*Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912)
- Kermadec petrel**
- Pterodroma mollis* (Gould, 1844)**  
*Austromenopon popellus* (Piaget, 1890)  
*Halipeurus procellariae* (J.C. Fabricius, 1775)  
*Naubates (Guenterion) pterodromi* Bedford, 1930  
*Saemundssonina (Puffinoecus) enderleini* (Eichler, 1949)  
*Trabeculus schillingi* Rudow, 1866
- Soft-plumaged petrel**
- \**Pterodroma inexpectata* (J.R. Forster, 1844)**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon* species 2  
*Halipeurus consimilis* Timmermann, 1960  
*Longimenopon* species  
*Naubates (Guenterion) pterodromi* Bedford, 1930  
*Trabeculus schillingi* Rudow, 1866 *sensu lato*
- Mottled petrel**
- Pterodroma externa* (Salvin, 1875)**  
*Austromenopon popellus* (Piaget, 1890)  
*Halipeurus kermadecensis* (Johnston & Harrison, 1912)  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*
- Juan Fernández petrel**
- \**Pterodroma cervicalis* (Salvin, 1891)**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Halipeurus noctivagus* Timmermann, 1960  
*Naubates (Guenterion) damma* Timmermann, 1961  
*Trabeculus* species 1
- White-naped petrel**
- Pterodroma nigripennis* (Rothschild, 1893)**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon* species 1  
*Halipeurus confusus* Palma, 2011  
*Longimenopon* species  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*
- Black-winged petrel**
- \**Pterodroma axillaris* (Salvin, 1893)**  
*Halipeurus theresae* Timmermann, 1969  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*
- Chatham petrel**

- ° ***Pterodroma cookii* (G.R. Gray, 1843)** **Cook's petrel**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon* species 1  
*Halipeurus turtur* Edwards, 1961  
*Longimenopon* species  
*Naubates (Guenterion) damma* Timmermann, 1961  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*
- Pterodroma longirostris* (Stejneger, 1893)** **Stejneger's petrel**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon* species 1  
*Halipeurus leucophryna* Timmermann, 1960  
*Longimenopon* species  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*
- ° ***Pterodroma pycrofti* Falla, 1933** **Pycroft's petrel**  
*Austromenopon* species 1  
*Halipeurus leucophryna* Timmermann, 1960  
*Longimenopon* species  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*
- Pterodroma leucoptera caledonica* Imber & Jenkins, 1981** **New Caledonian petrel**  
*Austromenopon* species 1  
*Halipeurus pricei* Palma, 2011  
*Naubates (Guenterion) damma* Timmermann, 1961  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*
- Halobaena caerulea* (Gmelin, 1789)** **Blue petrel**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon stammeri* Timmermann, 1963  
*Longimenopon* species  
*Naubates (Guenterion) clypeatus* (Giebel, 1874)  
*Saemundssonina (Saemundssonina) pterodromae* Timmermann, 1959
- Pachyptila vittata* (G. Forster, 1777)** **Broad-billed prion**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon stammeri* Timmermann, 1963  
*Longimenopon* species  
*Naubates (Guenterion) prioni* (Enderlein, 1908)  
*Saemundssonina (Saemundssonina) desolata* Timmermann, 1959
- Pachyptila salvini salvini* (Mathews, 1912)** **Salvin's prion**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon stammeri* Timmermann, 1963  
*Longimenopon* species  
*Naubates (Guenterion) prioni* (Enderlein, 1908)  
*Saemundssonina (Saemundssonina) desolata* Timmermann, 1959
- Pachyptila desolata* (Gmelin, 1789)** **Antarctic prion**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon stammeri* Timmermann, 1963  
*Longimenopon galeatum* Timmermann, 1957  
*Naubates (Guenterion) prioni* (Enderlein, 1908)  
*Saemundssonina (Saemundssonina) desolata* Timmermann, 1959
- Pachyptila belcheri* (Mathews, 1912)** **Thin-billed prion**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon stammeri* Timmermann, 1963  
*Longimenopon* species  
*Naubates (Guenterion) prioni* (Enderlein, 1908)  
*Saemundssonina (Saemundssonina) desolata* Timmermann, 1959

- Pachyptila turtur* (Kuhl, 1820)**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon stammeri* Timmermann, 1963  
*Longimenopon* species  
*Naubates* (*Guenterion*) *prioni* (Enderlein, 1908)  
*Saemundssonina* (*Saemundssonina*) *desolata* Timmermann, 1959  
**Fairy prion**
- \**Pachyptila crassirostris crassirostris* (Mathews, 1912)**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon stammeri* Timmermann, 1963  
*Naubates* (*Guenterion*) *prioni* (Enderlein, 1908)  
*Saemundssonina* (*Saemundssonina*) *desolata* Timmermann, 1959  
**Fulmar prion**
- \**Pachyptila crassirostris pyramidalis* Fleming, 1939**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon stammeri* Timmermann, 1963  
*Longimenopon* species  
*Naubates* (*Guenterion*) *prioni* (Enderlein, 1908)  
*Saemundssonina* (*Saemundssonina*) *desolata* Timmermann, 1959  
**Chatham fulmar prion**
- Pachyptila crassirostris flemingi* Tennyson & Bartle, 2005**  
*Naubates* (*Guenterion*) *prioni* (Enderlein, 1908)  
**Lesser fulmar prion**
- Bulweria bulwerii* (Jardine & Selby, 1828)**  
*Austromenopon bulweriae* Timmermann, 1963  
*Halipeurus bulweriae* Timmermann, 1960  
**Bulwer's petrel**
- Procellaria aequinoctialis* Linnaeus, 1758**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Naubates* (*Naubates*) *fuliginosus* (Taschenberg, 1882)  
*Trabeculus hexakon* (Waterston, 1914)  
**White-chinned petrel**
- \**Procellaria westlandica* Falla, 1946**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Naubates* (*Naubates*) *fuliginosus* (Taschenberg, 1882)  
*Trabeculus hexakon* (Waterston, 1914)  
**Westland petrel**
- \**Procellaria parkinsoni* G.R. Gray, 1862**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Naubates* (*Naubates*) *fuliginosus* (Taschenberg, 1882)  
*Trabeculus hexakon* (Waterston, 1914)  
**Black petrel**
- Procellaria cinerea* Gmelin, 1789**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon popellus* (Piaget, 1890)  
*Naubates* (*Naubates*) *fuliginosus* (Taschenberg, 1882)  
*Trabeculus hexakon* (Waterston, 1914)  
**Grey petrel**
- Pseudobulweria rostrata* (Peale, 1848)**  
*Halipeurus marquesanus* (Ferris, 1932)  
♣ *Saemundssonina* (*Puffinoecus*) *jamaicensis* Timmermann, 1962  
**Tahiti petrel**
- Calonectris leucomelas* (Temminck, 1836)**  
*Austromenopon longithoracicum* (Piaget, 1880)  
*Halipeurus angusticeps* (Piaget, 1880)  
*Saemundssonina* (*Puffinoecus*) *orientalis* (Uchida, 1949)  
**Streaked shearwater**
- Puffinus pacificus pacificus* (Gmelin, 1789)**  
*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon paululum* (Kellogg & Chapman, 1899)  
**Wedge-tailed shearwater**

*Halipeurus (Halipeurus) mirabilis* Thompson, 1940  
*Naubates (Naubates) harrisoni* Bedford, 1930  
*Saemundssonina (Puffinoecus) puellula* Timmermann, 1965  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*

***Puffinus pacificus chlororhynchus* Lesson, 1831**

*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus (Halipeurus) mirabilis* Thompson, 1940  
*Naubates (Naubates) harrisoni* Bedford, 1930  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*

**Wedge-tailed shearwater**

**\* *Puffinus bulleri* Salvin, 1888**

*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus (Halipeurus) thompsoni* Edwards, 1961  
*Naubates (Naubates) harrisoni* Bedford, 1930

**Buller's shearwater**

***Puffinus carneipes* Gould, 1844**

*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus gravis priapululus* Timmermann, 1961  
*Naubates (Naubates) harrisoni* Bedford, 1930  
*Saemundssonina (Puffinoecus) species*  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*

**Flesh-footed shearwater**

***Puffinus griseus* (Gmelin, 1789)**

*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus diversus* (Kellogg, 1896)  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*

**Sooty shearwater**

***Puffinus tenuirostris* (Temminck, 1836)**

*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus diversus* (Kellogg, 1896)  
*Trabeculus hexakon* (Waterston, 1914) *sensu lato*

**Short-tailed shearwater**

***Puffinus nativitatis* Streets, 1877**

*Trabeculus species 2*

**Christmas Island shearwater**

***Puffinus newelli* Henshaw, 1900**

*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus placodus* Edwards, 1961  
*Trabeculus mirabilis* (Kellogg, 1896)

**Newell's shearwater**

***Puffinus puffinus* (Brünnich, 1764)**

*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus diversus* (Kellogg, 1896)  
*Trabeculus aviator* (Evans, 1912)

**Manx shearwater**

**\* *Puffinus gavia* (J.R. Forster, 1844)**

*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus spadix subclavus* Timmermann, 1961  
*Saemundssonina (Puffinoecus) valida* (Kellogg & Chapman, 1899)

**Fluttering shearwater**

**\* *Puffinus huttoni* Mathews, 1912**

*Ancistrona vagelli* (J.C. Fabricius, 1787)  
*Austromenopon paululum* (Kellogg & Chapman, 1899)  
*Halipeurus spadix spadix* Timmermann, 1961  
*Naubates (Naubates) harrisoni* Bedford, 1930  
 ♦ *Trabeculus flemingi* Timmermann, 1959

**Hutton's shearwater**

<p>° <i>Puffinus assimilis kermadecensis</i> <b>Murphy, 1927</b>  <i>Ancistrona vagelli</i> (J.C. Fabricius, 1787)  <i>Austromenopon paululum</i> (Kellogg &amp; Chapman, 1899)  <i>Halipeurus placodus</i> Edwards, 1961  <i>Trabeculus mirabilis</i> (Kellogg, 1896)</p>	<b>Kermadec little shearwater</b>
<p>° <i>Puffinus assimilis haurakiensis</i> <b>Fleming &amp; Serventy, 1943</b>  <i>Ancistrona vagelli</i> (J.C. Fabricius, 1787)  <i>Austromenopon paululum</i> (Kellogg &amp; Chapman, 1899)  <i>Halipeurus mundae</i> Edwards, 1961  <i>Trabeculus mirabilis</i> (Kellogg, 1896)</p>	<b>North Island little shearwater</b>
<p><i>Puffinus elegans</i> <b>Giglioli &amp; Salvadori, 1869</b>  <i>Austromenopon paululum</i> (Kellogg &amp; Chapman, 1899)  <i>Halipeurus mundae</i> Edwards, 1961  <i>Trabeculus mirabilis</i> (Kellogg, 1896)</p>	<b>Subantarctic little shearwater</b>
<b>HYDROBATIDAE</b>	
<p><i>Oceanites oceanicus exasperatus</i> <b>Mathews, 1912</b>  <i>Philoceanus robertsi</i> (Clay, 1940)</p>	<b>Wilson's storm petrel</b>
<p><i>Garrodia nereis</i> (<b>Gould, 1841</b>)  <i>Philoceanus garrodiae</i> (Clay, 1940)  <i>Saemundssonina (Saemundssonina) nereis</i> Timmermann, 1956</p>	<b>Grey-backed storm petrel</b>
<p><i>Pelagodroma marina dulciae</i> <b>Mathews, 1912</b>  <i>Austromenopon enigki</i> Timmermann, 1963  <i>Halipeurus pelagodromae</i> Palma, 2011  <i>Saemundssonina (Saemundssonina) marina</i> Timmermann, 1956</p>	<b>Australian white-faced storm petrel</b>
<p>° <i>Pelagodroma marina maoriana</i> <b>Mathews, 1912</b>  <i>Austromenopon enigki</i> Timmermann, 1963  <i>Halipeurus pelagodromae</i> Palma, 2011  <i>Longimenopon</i> species  <i>Saemundssonina (Saemundssonina) marina</i> Timmermann, 1956</p>	<b>New Zealand white-faced storm petrel</b>
<p>° <i>Pelagodroma albiclunis</i> <b>Murphy &amp; Irving, 1951</b>  <i>Ancistrona vagelli</i> (J.C. Fabricius, 1787)  <i>Halipeurus pelagodromae</i> Palma, 2011</p>	<b>Kermadec storm petrel</b>
<p>° <i>Pealeornis maoriana</i> <b>Mathews, 1932</b>  <i>Austromenopon enigki</i> Timmermann, 1963  <i>Philoceanus fasciatus</i> (Carriker, 1958)  ♣ <i>Saemundssonina (Saemundssonina)</i> species</p>	<b>New Zealand storm petrel</b>
<p><i>Fregetta tropica</i> (<b>Gould, 1844</b>)  <i>Austromenopon enigki</i> Timmermann, 1963  <i>Halipeurus pelagicus</i> (Denny, 1842)  <i>Philoceanus fasciatus</i> (Carriker, 1958)  ♣ <i>Saemundssonina (Saemundssonina) platycephalus</i> (Kellogg &amp; Kuwana, 1902)</p>	<b>Black-billed storm petrel</b>
<p><i>Fregetta grallaria grallaria</i> (<b>Vieillot, 1818</b>)  <i>Halipeurus pelagicus</i> (Denny, 1842)  <i>Philoceanus fasciatus</i> (Carriker, 1958)</p>	<b>White-bellied storm petrel</b>
<p><i>Oceanodroma leucorhoa leucorhoa</i> (<b>Vieillot, 1818</b>)  <i>Halipeurus pelagicus</i> (Denny, 1842)  <i>Saemundssonina (Saemundssonina) thalassidromae incisa</i> Timmermann, 1950</p>	<b>Leach's storm petrel</b>
<b>PELECANOIDIDAE</b>	
<p><i>Pelecanoides urinatrix urinatrix</i> (<b>Gmelin, 1789</b>)</p>	<b>Northern diving petrel</b>



*Austromenopon elliotti* Timmermann, 1954  
*Halipeurus falsus pacificus* Edwards, 1961  
*Pelmatocerandra setosa* (Giebel, 1876)

\**Pelecanoides urinatrix chathamensis* Murphy & Harper, 1916

Southern diving petrel

*Austromenopon elliotti* Timmermann, 1954  
*Halipeurus falsus pacificus* Edwards, 1961  
*Pelmatocerandra setosa* (Giebel, 1876)

*Pelecanoides urinatrix exsul* Salvin, 1896

Subantarctic diving petrel

*Austromenopon elliotti* Timmermann, 1954  
*Halipeurus falsus pacificus* Edwards, 1961  
*Pelmatocerandra setosa* (Giebel, 1876)

*Pelecanoides georgicus* Murphy & Harper, 1916

South Georgian diving petrel

*Austromenopon elliotti* Timmermann, 1954  
*Pelmatocerandra setosa* (Giebel, 1876)

PHAETHONTIFORMES

PHAETHONTIDAE

*Phaethon rubricauda* Boddaert, 1783

Red-tailed tropicbird

*Austromenopon beckii* (Kellogg, 1906)  
*Saemundssonina* (*Saemundssonina*) *hexagona* (Giebel, 1874)

*Phaethon lepturus dorotheae* Mathews, 1913

White-tailed tropicbird

*Austromenopon beckii* (Kellogg, 1906)  
*Saemundssonina* (*Saemundssonina*) *uppalensis* (Rudow, 1870)

PELECANIFORMES

PELECANIDAE

*Pelecanus conspicillatus* Temminck, 1824

Australian pelican

*Colpocephalum eucarenum* Burmeister, 1838  
*Pectinopygus australis* Thompson, 1948

SULIDAE

*Morus serrator* (G.R. Gray, 1843)

Australasian gannet

*Eidmanniella pustulosa* (Nitzsch [*in* Giebel], 1866)  
*Pectinopygus bassani* (O. Fabricius, 1780)

*Sula leucogaster plotus* (Forster, 1844)

Brown booby

*Eidmanniella albescens* (Piaget, 1880)  
*Pectinopygus annulatus* (Piaget, 1880)  
*Pectinopygus garbei* (Pessôa & Guimarães, 1935)

*Sula dactylatra tasmani* van Tets, Meredith, Fullagar & Davidson, 1988

Masked booby

*Eidmanniella albescens* (Piaget, 1880)  
*Pectinopygus annulatus* (Piaget, 1880)

PHALACROCORACIDAE

\**Phalacrocorax melanoleucos brevirostris* Gould, 1837

Little shag

*Eidmanniella eurygaster* (Nitzsch [*in* Giebel], 1866)  
*Pectinopygus dispar* (Piaget, 1880)

*Phalacrocorax carbo novaehollandiae* Stephens, 1826

Black shag

*Eidmanniella pellucida* (Rudow, 1869)  
*Pectinopygus gyricornis* (Denny, 1842)

\**Phalacrocorax varius varius* (Gmelin, 1789)

Pied shag

*Eidmanniella pellucida* (Rudow, 1869)  
*Pectinopygus varius* Timmermann, 1964

<b><i>Phalacrocorax sulcirostris</i> (Brandt, 1837)</b> <i>Eidmanniella eurygaster</i> (Nitzsch [ <i>in</i> Giebel], 1866) <i>Pectinopygus setosus</i> (Piaget, 1880)	<b>Little black shag</b>
<sup>o</sup> <b><i>Leucocarbo carunculatus</i> (Gmelin, 1789)</b> <i>Eidmanniella pellucida</i> (Rudow, 1869) ◆ <i>Pectinopygus carunculatus</i> Timmermann, 1964 <i>sensu lato</i>	<b>New Zealand king shag</b>
<sup>o</sup> <b><i>Leucocarbo chalconotus</i> (G.R. Gray, 1845)</b> <i>Eidmanniella pellucida</i> (Rudow, 1869) ◆ <i>Pectinopygus carunculatus</i> Timmermann, 1964 <i>sensu lato</i>	<b>Stewart Island shag</b>
<sup>o</sup> <b><i>Leucocarbo onslowi</i> (Forbes, 1893)</b> ◆ <i>Pectinopygus carunculatus</i> Timmermann, 1964 <i>sensu lato</i>	<b>Chatham Island shag</b>
<sup>o</sup> <b><i>Leucocarbo ranfurlyi</i> (Ogilvie-Grant, 1901)</b> ◆ <i>Pectinopygus carunculatus</i> Timmermann, 1964 <i>sensu lato</i>	<b>Bounty Island shag</b>
<sup>o</sup> <b><i>Leucocarbo colensoi</i> (Buller, 1888)</b> ◆ <i>Pectinopygus carunculatus</i> Timmermann, 1964	<b>Auckland Island shag</b>
<sup>o</sup> <b><i>Leucocarbo campbelli</i> (Filhol, 1878)</b> ◆ <i>Pectinopygus carunculatus</i> Timmermann, 1964 <i>sensu lato</i>	<b>Campbell Island shag</b>
<sup>o</sup> <b><i>Leucocarbo purpurascens</i> (Brandt, 1837)</b> <i>Pectinopygus turbinatus</i> (Piaget, 1890)	<b>Macquarie Island shag</b>
<sup>o</sup> <b><i>Stictocarbo punctatus punctatus</i> (Sparman, 1786)</b> <i>Eidmanniella pellucida</i> (Rudow, 1869) ◆ <i>Pectinopygus punctatus</i> Timmermann, 1964	<b>Spotted shag</b>
<sup>o</sup> <b><i>Stictocarbo punctatus oliveri</i> Mathews, 1931</b> <i>Eidmanniella pellucida</i> (Rudow, 1869) ◆ <i>Pectinopygus punctatus</i> Timmermann, 1964	<b>Blue shag</b>
<sup>o</sup> <b><i>Stictocarbo featherstoni</i> (Buller, 1873)</b> <i>Eidmanniella pellucida</i> (Rudow, 1869) <i>Pectinopygus</i> species	<b>Pitt Island shag</b>
<b>CICONIIFORMES</b>	
<b>ARDEIDAE</b>	
<b><i>Ardea modesta</i> J.E. Gray, 1831</b> <i>Ardeicola expallidus</i> Blagoveshtchensky, 1940 <i>Ciconiphilus decimfasciatus</i> (Boisduval & Lacordaire, 1835)	<b>White heron</b>
<b><i>Ardea ibis coromanda</i> (Boddaert, 1783)</b> <i>Ardeicola expallidus</i> Blagoveshtchensky, 1940 <i>Ciconiphilus decimfasciatus</i> (Boisduval & Lacordaire, 1835)	<b>Eastern cattle egret</b>
<b><i>Egretta novaehollandiae novaehollandiae</i> (Latham, 1790)</b> <i>Ardeicola pilgrimi</i> Tandan, 1972 <i>Ciconiphilus decimfasciatus</i> (Boisduval & Lacordaire, 1835)	<b>White-faced heron</b>
<b><i>Egretta garzetta immaculata</i> (Gould, 1846)</b> <i>Ardeicola expallidus</i> Blagoveshtchensky, 1940 <i>Ciconiphilus decimfasciatus</i> (Boisduval & Lacordaire, 1835)	<b>Little egret</b>
<b><i>Egretta sacra sacra</i> (Gmelin, 1789)</b> <i>Ardeicola neopallidus</i> Price, Hellenthal & Palma, 2003 <i>Ciconiphilus decimfasciatus</i> (Boisduval & Lacordaire, 1835)	<b>Reef heron</b>

- Botaurus poiciloptilus* (Wagler, 1827) Australasian bittern  
*Ardeicola stellaris* (Denny, 1842)  
*Ciconiphilus decimfasciatus* (Boisduval & Lacordaire, 1835)

## THRESKIORNITHIDAE

- Plegadis falcinellus* (Linnaeus, 1766) Glossy ibis  
*Ardeicola raphidius* (Nitzsch [in Giebel], 1866)  
*Colpocephalum leptopygos* Nitzsch [in Giebel], 1874  
*Ibidoecus bisignatus* (Nitzsch [in Giebel], 1866)  
*Plegadiphilus plegadis* (Dubinin, 1938)
- Threskiornis molucca strictipennis* (Gould, 1838) Australian white ibis  
*Ibidoecus diana* Tandan, 1958  
*Plegadiphilus threskiornis* Bedford, 1939
- Platalea regia* Gould, 1838 Royal spoonbill  
*Ardeicola plataleae* (Linnaeus, 1758)  
*Eucolpocephalum femorale* (Piaget, 1880)  
*Ibidoecus plataleae* (Denny, 1842)

## ACCIPITRIFORMES

## ACCIPITRIDAE

- Circus approximans* Peale, 1848 Swamp harrier  
*Colpocephalum turbinatum* Denny, 1842  
*Degeeriella fusca* (Denny, 1842)  
*Nosopon lucidum* (Rudow, 1869)

## FALCONIFORMES

## FALCONIDAE

- Falco cenchroides cenchroides* Vigors & Horsfield, 1827 Nankeen kestrel  
*Colpocephalum subzerafae* Tendeiro, 1988  
*Laemobothrion tinnunculi* (Linnaeus, 1758)
- <sup>°</sup>*Falco novaeseelandiae* Gmelin, 1788 New Zealand falcon  
*Degeeriella rufa rufa* (Burmeister, 1838)  
*Nosopon lucidum* (Rudow, 1869)

## GRUIFORMES

## RALLIDAE

- <sup>°</sup>*Gallirallus philippensis assimilis* (G.R. Gray, 1843) Banded rail  
*Pseudomenopon scopulacorne* (Denny, 1842)  
*Rallicola (Rallicola) ortygometae philippensis* Emerson, 1966
- <sup>°</sup>*Gallirallus australis greyi* (Buller, 1888) North Island weka  
 ◆ *Pseudomenopon pilgrimi* Price, 1974  
 ◆ *Rallicola (Rallicola) harrisoni* Emerson, 1955
- <sup>°</sup>*Gallirallus australis australis* (Sparman, 1786) Western weka  
 ◆ *Rallicola (Rallicola) harrisoni* Emerson, 1955
- <sup>°</sup>*Gallirallus australis hectori* (Hutton, 1873) Buff weka  
 ◆ *Pseudomenopon pilgrimi* Price, 1974  
 ◆ *Rallicola (Rallicola) harrisoni* Emerson, 1955
- <sup>°</sup>*Gallirallus australis scotti* (Ogilvie-Grant, 1905) Stewart Island weka  
 ◆ *Rallicola (Rallicola) harrisoni* Emerson, 1955
- Porzana tabuensis tabuensis* (Gmelin, 1789) Spotless crane  
 ♣ *Fulicoffula stammeri* Eichler, 1958  
*Rallicola (Rallicola) tabuensis* Emerson, 1966

<p>° <i>Porzana pusilla affinis</i> (J.E. Gray, 1845)  ♣ <i>Fulicoffula stammeri</i> Eichler, 1958  ♣ <i>Incidifrons porzanae</i> Blagoveshtchensky, 1951  <i>Rallicola (Rallicola) tabuensis</i> Emerson, 1966</p>	Marsh crake
<p><i>Porphyrio melanotus melanotus</i> Temminck, 1820  <i>Pseudomenopon concretum</i> (Piaget, 1880)  <i>Rallicola (Rallicola) lugens</i> (Giebel, 1874)</p>	Pukeko
<p>° <i>Porphyrio hochstetteri</i> (A.B. Meyer, 1883)  ♦ <i>Rallicola (Rallicola) takahe</i> Holloway, 1956</p>	South Island takahe
<p><i>Fulica atra australis</i> Gould, 1845  <i>Fulicoffula lurida</i> (Nitzsch, 1818)  <i>Incidifrons fulicae</i> (Linnaeus, 1758)  <i>Pseudomenopon pilosum</i> (Scopoli, 1763)  <i>Rallicola (Rallicola) fulicae</i> (Denny, 1842)</p>	Australian coot
<b>CHARADRIIFORMES</b>	
<b>SCOLOPACIDAE</b>	
<p>° <i>Coenocorypha pusilla</i> (Buller, 1869)  ♦ <i>Quadriceps coenocoryphae</i> Timmermann, 1955</p>	Chatham Island snipe
<p>° <i>Coenocorypha huegeli</i> (Tristram, 1893)  ♦ <i>Quadriceps coenocoryphae</i> Timmermann, 1955</p>	Snares Island snipe
<p>° <i>Coenocorypha aucklandica aucklandica</i> (G.R. Gray, 1845)  <i>Austromenopon</i> species 3  ♦ <i>Quadriceps coenocoryphae</i> Timmermann, 1955</p>	Auckland Island snipe
<p>° <i>Coenocorypha aucklandica meinertzhagenae</i> Rothschild, 1927  <i>Austromenopon</i> species 3</p>	Antipodes Island snipe
<p>° <i>Coenocorypha aucklandica perseverance</i> Miskelly &amp; Baker, 2010  ♦ <i>Quadriceps coenocoryphae</i> Timmermann, 1955</p>	Campbell Island snipe
<p><i>Calidris canutus rogersi</i> (Mathews, 1913)  <i>Actornithophilus umbrinus</i> (Burmeister, 1838)  <i>Austromenopon lutescens</i> (Burmeister, 1838)  <i>Carduiceps zonarius</i> (Nitzsch [in Giebel], 1866)  <i>Luniceps drosti</i> Timmermann, 1954  <i>Saemundssonina (Saemundssonina) tringae</i> (O. Fabricius, 1780) <i>sensu lato</i></p>	Lesser knot
<p><i>Calidris alba</i> (Pallas, 1776)  <i>Luniceps actophilus</i> (Kellogg &amp; Chapman, 1899)</p>	Sanderling
<p><i>Calidris ferruginea</i> (Pontoppidan, 1763)  <i>Luniceps falcinellus</i> Timmermann, 1954</p>	Curlew sandpiper
<p><i>Calidris acuminata</i> (Horsfield, 1821)  <i>Actornithophilus umbrinus</i> (Burmeister, 1838)  <i>Carduiceps zonarius</i> (Nitzsch [in Giebel], 1866)  <i>Luniceps superciliosus</i> Gustafsson &amp; Olsson, 2012</p>	Sharp-tailed sandpiper
<p><i>Calidris ruficollis</i> (Pallas, 1776)  <i>Luniceps falcinellus</i> Timmermann, 1954</p>	Red-necked stint
<p><i>Numenius madagascariensis</i> (Linnaeus, 1766)  <i>Luniceps numenii madagascariensis</i> Gustafsson &amp; Olsson, 2012</p>	Eastern curlew

- Numenius phaeopus variegatus* (Scopoli, 1786)** Asiatic whimbrel  
*Austromenopon phaeopodis* (Schränk, 1802)  
*Lunaceps numenii oliveri* (Johnston & Harrison, 1912)  
*Saemundssonina* (*Saemundssonina*) *scolopacisphaeopodis scolopacisphaeopodis* (Schränk, 1803)
- Limosa lapponica baueri* Naumann, 1836** Eastern bar-tailed godwit  
*Actornithophilus limosae* (Kellogg, 1908)  
*Austromenopon meyeri* (Giebel, 1874)  
*Carduiceps cingulatus lapponicus* Emerson, 1953  
*Lunaceps limosella* Timmermann, 1954  
*Saemundssonina* (*Saemundssonina*) *limosae* (Denny, 1842)
- Limosa limosa melanuroides* Gould, 1846** Asiatic black-tailed godwit  
*Actornithophilus spinulosus* (Piaget, 1880)  
*Austromenopon limosae* Timmermann, 1954  
*Carduiceps cingulatus cingulatus* (Denny, 1842)  
*Lunaceps limosae* Bechet, 1968  
*Saemundssonina* (*Saemundssonina*) *thompsoni* Timmermann, 1951
- Tringa incana* (Gmelin, 1789)** Wandering tattler  
*Saemundssonina* (*Saemundssonina*) *platygaster* (Denny, 1842) *sensu lato*
- Arenaria interpres* (Linnaeus, 1758)** Ruddy turnstone  
*Actornithophilus bicolor* (Piaget, 1880)  
*Actornithophilus pediculoides* (Mjöberg, 1910)  
*Austromenopon lutescens* (Burmeister, 1838)  
*Quadriceps strepsilaris* (Denny, 1842)
- HAEMATOPODIDAE**
- <sup>o</sup>*Haematopus unicolor* J.R. Forster, 1844** Variable oystercatcher  
*Actornithophilus grandiceps* (Piaget, 1880)  
*Quadriceps auratus* (Haan, 1829)  
*Quadriceps ridgwayi* (Kellogg, 1906)  
*Saemundssonina* (*Saemundssonina*) *haematopi* (Linnaeus, 1758)
- <sup>o</sup>*Haematopus finschi* Martens, 1897** South Island pied oystercatcher  
*Actornithophilus grandiceps* (Piaget, 1880)  
*Austromenopon haematopi* Timmermann, 1954  
*Quadriceps auratus* (Haan, 1829)  
*Saemundssonina* (*Saemundssonina*) *haematopi* (Linnaeus, 1758)
- <sup>o</sup>*Haematopus chathamensis* Hartert, 1927** Chatham Island oystercatcher  
*Actornithophilus grandiceps* (Piaget, 1880)  
*Quadriceps auratus* (Haan, 1829)
- RECURVIROSTRIDAE**
- Himantopus himantopus leucocephalus* Gould, 1837** Pied stilt  
*Austromenopon himantopi* Timmermann, 1954  
*Quadriceps hemichrous* (Nitzsch *in* Giebel, 1866)  
*Quadriceps semifissus* (Nitzsch [*in* Giebel], 1866)  
*Saemundssonina* (*Saemundssonina*) *platygaster* (Denny, 1842) *sensu lato*
- <sup>o</sup>*Himantopus novaezelandiae* Gould, 1841** Black stilt  
*Quadriceps hemichrous* (Nitzsch *in* Giebel, 1866)  
*Quadriceps semifissus* (Nitzsch [*in* Giebel], 1866)  
*Saemundssonina* (*Saemundssonina*) *platygaster* (Denny, 1842) *sensu lato*
- CHARADRIIDAE**
- Pluvialis fulva* (Gmelin, 1789)** Pacific golden plover  
*Actornithophilus ochraceus* (Nitzsch, 1818) *sensu lato*  
*Quadriceps charadrii orarius* (Kellogg, 1896)  
*Saemundssonina* (*Saemundssonina*) *conica conica* (Denny, 1842)

- ° ***Charadrius obscurus aquilonius* Dowding, 1994** Northern New Zealand dotterel  
*Actornithophilus ochraceus* (Nitzsch, 1818) *sensu lato*  
 ♦ *Quadriceps dominella* Timmermann, 1953  
*Saemundssonina (Saemundssonina) platygaster* (Denny, 1842) *sensu lato*
- ° ***Charadrius obscurus obscurus* Gmelin, 1789** Southern New Zealand dotterel  
*Actornithophilus ochraceus* (Nitzsch, 1818) *sensu lato*  
 ♦ *Quadriceps dominella* Timmermann, 1953  
*Saemundssonina (Saemundssonina) platygaster* (Denny, 1842) *sensu lato*
- ° ***Charadrius bicinctus bicinctus* Jardine & Selby, 1827** Banded dotterel  
*Quadriceps cedemajori* Timmermann, 1969  
*Saemundssonina (Saemundssonina) platygaster balati* Timmermann, 1969
- ° ***Charadrius bicinctus exilis* Falla, 1978** Auckland Island banded dotterel  
*Austromenopon aegialitidis* (Durrant, 1906) *sensu lato*  
*Quadriceps cedemajori* Timmermann, 1969  
 ♦ *Quadriceps novaeseelandiae* Timmermann, 1953
- Charadrius veredus* Gould, 1848** Oriental dotterel  
*Quadriceps assimilis* (Piaget, 1890)
- ° ***Anarhynchus frontalis* (Quoy & Gaimard, 1830)** Wrybill  
*Quadriceps cedemajori* Timmermann, 1969
- ° ***Thinornis novaeseelandiae* (Gmelin, 1789)** Shore plover  
 ♦ *Quadriceps novaeseelandiae* Timmermann, 1953  
 ♦ *Saemundssonina (Saemundssonina) chathamensis* Timmermann, 1977
- Vanellus miles novaehollandiae* Stephens, 1819** Spur-winged plover  
*Actornithophilus hoplopteri* (Mjöberg, 1910) *sensu lato*  
*Austromenopon aegialitidis* (Durrant, 1906) *sensu lato*  
*Quadriceps renschi* Timmermann, 1954
- GLAREOLIDAE**
- Glareola maldivarum* J.R. Forster, 1795** Oriental pratincole  
 ♣ *Actornithophilus sedes* Eichler, 1944  
 ♣ *Austromenopon cursorium* (Giebel, 1874)  
*Quadriceps ellipticus* (Nitzsch [in Giebel], 1866) *sensu lato*
- STERCORARIIDAE**
- Catharacta antarctica lonnbergi* Mathews, 1912** Subantarctic skua  
*Austromenopon fuscofasciatum* (Piaget, 1880)  
*Haffneria grandis* (Piaget, 1880)  
*Saemundssonina (Saemundssonina) euryrhynga* (Giebel, 1874)
- Catharacta maccormicki* (Saunders, 1893)** South polar skua  
*Austromenopon fuscofasciatum* (Piaget, 1880)  
*Haffneria grandis* (Piaget, 1880)  
*Quadriceps normifer alpha* (Kellogg, 1914)  
*Saemundssonina (Saemundssonina) euryrhynga* (Giebel, 1874)
- Stercorarius parasiticus* (Linnaeus, 1758)** Arctic skua  
*Austromenopon fuscofasciatum* (Piaget, 1880)  
*Haffneria grandis* (Piaget, 1880)  
*Quadriceps normifer normifer* (Grube, 1851)  
*Saemundssonina (Saemundssonina) cephalus* (Denny, 1842)
- Stercorarius longicaudus* Vieillot, 1819** Long-tailed skua  
*Austromenopon fuscofasciatum* (Piaget, 1880)  
*Quadriceps normifer parvopallidus* (Eichler, 1951)

*Saemundssonina* (*Saemundssonina*) *inexpectata* Timmermann, 1951

## LARIDAE

- Larus dominicanus dominicanus* Lichtenstein, 1823** Southern black-backed gull  
*Actornithophilus piceus lari* (Packard, 1870)  
*Austromenopon transversum* (Denny, 1842)  
*Quadriceps ornatus fuscolaminulatus* (Enderlein, 1908)  
♣ *Quadriceps punctatus sublingulatus* Timmermann, 1952  
*Saemundssonina* (*Saemundssonina*) *lari* (O. Fabricius, 1780) *sensu lato*
- ° ***Larus novaehollandiae scopulinus* J.R. Forster, 1844** Red-billed gull  
*Actornithophilus piceus lari* (Packard, 1870)  
*Austromenopon transversum* (Denny, 1842)  
*Quadriceps punctatus lingulatus* (Waterston, 1914)  
*Saemundssonina* (*Saemundssonina*) *lari* (O. Fabricius, 1780) *sensu lato*
- ° ***Larus bulleri* Hutton, 1871** Black-billed gull  
*Austromenopon transversum* (Denny, 1842)  
*Quadriceps punctatus lingulatus* (Waterston, 1914)  
*Saemundssonina* (*Saemundssonina*) *lari* (O. Fabricius, 1780) *sensu lato*
- ## STERNIDAE
- Anous stolidus pileatus* (Scopoli, 1786)** Brown noddy  
*Quadriceps separatus* (Kellogg & Kuwana, 1902)
- Anous minutus minutus* Boie, 1844** Black noddy  
*Actornithophilus ceruleus* (Timmermann, 1954)  
*Austromenopon atrofultum* (Piaget, 1880)  
*Quadriceps hopkinsi hopkinsi* Timmermann, 1952  
*Saemundssonina* (*Saemundssonina*) *remota* Timmermann, 1951
- Procelsterna cerulea albivitta* Bonaparte, 1856** Grey noddy  
*Actornithophilus ceruleus* (Timmermann, 1954)  
*Austromenopon atrofultum* (Piaget, 1880)  
*Quadriceps hopkinsi apophoretus* Timmermann, 1969
- Gygis alba candida* (Gmelin, 1789)** White tern  
*Saemundssonina* (*Saemundssonina*) *gygisa* Palma, 2012
- Onychoprion fuscatus serratus* (J.R. Forster, 1830)** Sooty tern  
*Austromenopon atrofultum* (Piaget, 1880)  
*Quadriceps birostris* (Giebel, 1874)  
*Saemundssonina* (*Saemundssonina*) *albemarlensis* (Kellogg & Kuwana, 1902)
- Sternula albifrons sinensis* (Gmelin, 1789)** Eastern little tern  
*Quadriceps nycthemerus* (Burmeister, 1838)  
*Saemundssonina* (*Saemundssonina*) *melanocephalus* (Burmeister, 1838)
- Hydroprogne caspia* (Pallas, 1770)** Caspian tern  
*Austromenopon atrofultum* (Piaget, 1880)  
*Quadriceps caspius* (Giebel, 1874)
- ° ***Chlidonias albostratus* (G.R. Gray, 1845)** Black-fronted tern  
*Quadriceps* species  
*Saemundssonina* (*Saemundssonina*) *lockleyi* Clay, 1949
- ° ***Sterna striata* Gmelin, 1789** White-fronted tern  
*Austromenopon atrofultum* (Piaget, 1880)  
*Quadriceps sellatus sellatus* (Burmeister, 1838)  
*Saemundssonina* (*Saemundssonina*) *sternae* (Linnaeus, 1758)

- \* *Sterna vittata bethunei* Travers, 1896 New Zealand Antarctic tern  
*Austromenopon atrofulvum* (Piaget, 1880)  
*Quadriceps sellatus houri* Hopkins, 1949  
*Quadriceps sellatus sellatus* (Burmeister, 1838)  
*Saemundssonina (Saemundssonina) lockleyi* Clay, 1949
- Sterna paradisaea* Pontoppidan, 1763 Arctic tern  
*Austromenopon atrofulvum* (Piaget, 1880)  
*Quadriceps sellatus houri* Hopkins, 1949  
*Saemundssonina (Saemundssonina) lockleyi* Clay, 1949
- COLUMBIFORMES**  
**COLUMBIDAE**
- \* *Columba livia* Gmelin, 1789 Rock pigeon  
*Bonomiella columbae* Emerson, 1957  
*Campanulotes bidentatus compar* (Burmeister, 1838)  
*Colpocephalum turbinatum* Denny, 1842  
*Hohorstiella lata* (Piaget, 1880)  
*Columbicola columbae columbae* (Linnaeus, 1758)
- \* *Streptopelia chinensis tigrina* (Temminck, 1810) Spotted dove  
♣ *Coloceras chinense* (Kellogg & Chapman, 1902)  
*Columbicola columbae columbae* (Linnaeus, 1758)
- \* *Hemiphaga novaeseelandiae* (Gmelin, 1789) New Zealand pigeon  
♣ *Hohorstiella timorensis* Tendeiro, 1980  
♦ *Coloceras novaeseelandiae* (Tendeiro, 1972)
- \* *Hemiphaga chathamensis* (Rothschild, 1891) Chatham Island pigeon  
♦ *Coloceras harrisoni* (Tendeiro, 1972)
- \* *Hemiphaga spadicea* (Latham, 1802) Norfolk Island pigeon  
♦ *Coloceras hemiphagae* (Tendeiro, 1972)  
♦ *Coloceras restinctum* (Tendeiro, 1972)
- PSITTACIFORMES**  
**STRIGOPIDAE**
- \* *Strigops habroptilus* G.R. Gray, 1845 Kakapo  
♦ *Heteromenopon (Keamenopon) kea* (Kellogg, 1907)
- \* *Nestor meridionalis meridionalis* (Gmelin, 1788) South Island kaka  
♦ *Heteromenopon (Keamenopon) kea* (Kellogg, 1907)  
♦ *Forficuloecus meinertzhageni* Guimarães, 1974  
♦ *Neopsittaconirmus kea* (Kellogg, 1907)
- \* *Nestor meridionalis septentrionalis* Lorenz, 1896 North Island kaka  
♦ *Heteromenopon (Keamenopon) kea* (Kellogg, 1907)  
♦ *Forficuloecus meinertzhageni* Guimarães, 1974  
♦ *Neopsittaconirmus kea* (Kellogg, 1907)
- \* *Nestor notabilis* Gould, 1856 Kea  
♦ *Colpocephalum pilgrimi* Price, 1967  
♦ *Heteromenopon (Keamenopon) kea* (Kellogg, 1907)  
♦ *Forficuloecus meinertzhageni* Guimarães, 1974  
♦ *Neopsittaconirmus kea* (Kellogg, 1907)
- CACATUIDAE**
- Cacatua galerita* (Latham, 1790) Sulphur-crested cockatoo  
*Franciscoloa (Franciscoloa) pallida* (Piaget, 1880)



*Neopsittaconirmus albus* (Le Souëf & Bullen, 1902)

*Psittoeacus vanzolinii* Guimarães, 1974

## PSITTACIDAE

- \* *Platycercus eximius* (Shaw, 1792)  
♣ *Forficuloecus* species Eastern rosella
- ° *Cyanoramphus novaezelandiae novaezelandiae* (Sparrman, 1787)  
♦ *Forficuloecus pilgrimi* Guimarães, 1985 Red-crowned parakeet
- ° *Cyanoramphus novaezelandiae chathamensis* Oliver, 1930  
♦ *Forficuloecus pilgrimi* Guimarães, 1985 Chatham Island red-crowned parakeet
- ° *Cyanoramphus auriceps* (Kuhl, 1820)  
♦ *Forficuloecus pilgrimi* Guimarães, 1985 Yellow-crowned parakeet
- ° *Cyanoramphus malherbi* Souancé, 1857  
♦ *Forficuloecus pilgrimi* Guimarães, 1985 Orange-fronted parakeet
- ° *Cyanoramphus forbesi* Rothschild, 1893  
♦ *Forficuloecus pilgrimi* Guimarães, 1985 Forbes' parakeet
- ° *Cyanoramphus unicolor* (Lear, 1831)  
♦ *Forficuloecus pilgrimi* Guimarães, 1985  
*Heteromenopon* (*Keamenopon*) species Antipodes Island parakeet
- ° *Cyanoramphus hochstetteri* (Reischek, 1889)  
♦ *Forficuloecus pilgrimi* Guimarães, 1985 Reischek's parakeet

## CUCULIFORMES

### CUCULIDAE

- Cuculus optatus* Gould, 1845  
*Cuculicola latirostris* (Burmeister, 1838) Oriental cuckoo
- ° *Chrysococcyx lucidus lucidus* (Gmelin, 1788)  
♦ *Cuculicola kui* Kettle, 1980 Shining cuckoo
- ° *Eudynamys taitensis* (Sparrman, 1787)  
*Cuculiphilus* (*Cuculiphilus*) *fasciatiiventris* Carriker, 1955 Long-tailed cuckoo
- Scythrops novaehollandiae* Latham, 1790  
*Cuculiphilus* (*Cuculiphilus*) *platygaster* (Giebel, 1874) Channel-billed cuckoo

## STRIGIFORMES

### STRIGIDAE

- ° *Ninox novaeseelandiae novaeseelandiae* (Gmelin, 1788)  
*Kurodaia* (*Conciella*) *cryptostigmatia* (Nitzsch [*in* Giebel], 1861)  
*Strigiphilus vapidus* Clay, 1977 Morepork
- ° *Sceloglaux albifacies albifacies* (G.R. Gray, 1844)  
*Strigiphilus* species South Island laughing owl
- \* *Athene noctua* (Scopoli, 1769)  
*Strigiphilus cursitans* (Nitzsch [*in* Giebel], 1861) Little owl

## TYTONIDAE

- Tyto alba delicatula* (Gould, 1837)  
*Strigiphilus aitkeni* Clay, 1966 Australian barn owl

## APODIFORMES

## APODIDAE

- Hirundapus caudacutus caudacutus* (Latham, 1802) White-throated needletail  
 ♣ *Dennyus (Takamatsua)* species

## CORACIIFORMES

## HALCYONIDAE

- Dacelo novaeguineae novaeguineae* (Hermann, 1783) Laughing kookaburra  
 ♣ *Alcedoecus delphax* (Nitzsch [*in* Giebel], 1866)  
 ♣ *Emersoniella bracteata* (Nitzsch [*in* Giebel], 1866)
- ° *Todiramphus sanctus vagans* (Lesson, 1828) New Zealand kingfisher  
*Alcedoecus alatoctypeatus* (Piaget, 1885)

## CORACIIDAE

- Eurystomus orientalis pacificus* (Latham, 1802) Dollarbird  
*Capraiella* species

## PASSERIFORMES

## ACANTHISITTIDAE

- ° *Acanthisitta chloris chloris* (Sparrman, 1787) South Island rifleman  
*Menacanthus* species  
 ◆ *Philopteroides novaezealandiae* Mey, 2004
- ° *Xenicus longipes longipes* (Gmelin, 1789) South Island bush wren  
 ◆ *Philopteroides xenicus* Mey, 2004

## CALLAEIDAE

- ° *Callaeas wilsoni* (Bonaparte, 1851) North Island kokako  
 ◆ *Brueelia callaeincola* Valim & Palma, 2015  
 ◆ *Philopterus novaezealandiae* Palma & Price, 2000
- ° *Callaeas cinerea* (Gmelin, 1788) South Island kokako  
 ◆ *Brueelia callaeincola* Valim & Palma, 2015  
 ◆ *Philopterus novaezealandiae* Palma & Price, 2000
- ° *Philesturnus rufusater* (Lesson, 1828) North Island saddleback  
 ◆ *Brueelia callaeincola* Valim & Palma, 2015
- ° *Philesturnus carunculatus* (Gmelin, 1789) South Island saddleback  
 ◆ *Brueelia callaeincola* Valim & Palma, 2015
- ° *Heteralocha acutirostris* (Gould, 1837) Huia  
 ◆ *Rallicola (Huiacola) extinctus* (Mey, 1990)

## NOTIOMYSTIDAE

- ° *Notiomystis cincta* (du Bus de Gisignies, 1839) Stitchbird  
 ◆ *Myrsidea hihi* Sychra, Kolencik & Palma, 2016

## TURNAGRIDAE

- ° *Turnagra capensis capensis* (Sparrman, 1787) South Island piopio  
*Brueelia* species

**ACANTHIZIDAE**

- \* *Gerygone igata* (Quoy & Gaimard, 1830) **Grey warbler**  
*Myrsidea* species  
 ◆ *Philopteroides pilgrimi* Valim & Palma, 2013  
*Ricinus* species

**MELIPHAGIDAE**

- \* *Anthornis melanura obscura* Falla, 1948 **Three Kings bellbird**  
 ◆ *Melibrueelia novaeseelandiae* Valim & Palma, 2015  
 ◆ *Myrsidea novaeseelandiae* Sychra, Kolencik & Palma, 2016
- \* *Anthornis melanura oneho* Bartle & Sagar, 1987 **Poor Knights bellbird**  
 ◆ *Melibrueelia novaeseelandiae* Valim & Palma, 2015  
 ◆ *Myrsidea novaeseelandiae* Sychra, Kolencik & Palma, 2016
- \* *Anthornis melanura melanura* (Sparrman, 1786) **Bellbird**  
 ◆ *Melibrueelia novaeseelandiae* Valim & Palma, 2015  
 ◆ *Myrsidea novaeseelandiae* Sychra, Kolencik & Palma, 2016
- \* *Prosthemadera novaeseelandiae novaeseelandiae* (Gmelin, 1788) **Tui**  
 ◆ *Melibrueelia novaeseelandiae* Valim & Palma, 2015  
 ◆ *Myrsidea novaeseelandiae* Sychra, Kolencik & Palma, 2016

**PACHYCEPHALIDAE**

- \* *Mohoua albicilla* (Lesson, 1830) **Whitehead**  
*Brueelia* species  
*Myrsidea* species  
*Philopterus* species 2

**ARTAMIDAE**

- \* *Gymnorhina tibicen* (Latham, 1802) **Australian magpie**  
*Brueelia semiannulata* (Piaget, 1883)  
*Myrsidea ivanliteraki* Sychra, Kolencik & Palma, 2016  
*Philopterus* species 1

**RHIPIDURIDAE**

- \* *Rhipidura fuliginosa placabilis* Bangs, 1921 **North Island fantail**  
 ◆ *Philopteroides fuliginosus* Valim & Palma, 2013
- \* *Rhipidura fuliginosa fuliginosa* (Sparrman, 1787) **South Island fantail**  
 ◆ *Menacanthus rhipidurae* Palma & Price, 2005  
 ◆ *Philopteroides fuliginosus* Valim & Palma, 2013

**CORVIDAE**

- \* *Corvus frugilegus* Linnaeus, 1758 **Rook**  
*Colpocephalum fregili* Denny, 1842

**MONARCHIDAE**

- Monarcha melanopsis* (Vieillot, 1818) **Black-faced monarch**  
*Brueelia* species

**PETROICIDAE**

- \* *Petroica (Petroica) macrocephala toitoi* (Lesson, 1828) **North Island tomtit**  
 ◆ *Philopteroides macrocephalus* Valim & Palma, 2013

° <i>Petroica (Petroica) macrocephala macrocephala</i> (Gmelin, 1789) ◆ <i>Philopteroides macrocephalus</i> Valim & Palma, 2013	South Island tomtit
° <i>Petroica (Petroica) macrocephala dannefaerdi</i> (Rothschild, 1894) ◆ <i>Philopteroides macrocephalus</i> Valim & Palma, 2013	Snares Island tomtit
° <i>Petroica (Miro) australis australis</i> (Sparrman, 1788) <i>Menacanthus eurysternus</i> (Burmeister, 1838)	South Island robin
<b>ALAUDIDAE</b>	
* <i>Alauda arvensis</i> Linnaeus, 1758 ♣ <i>Brueelia parviguttata</i> (Blagoveshtchensky, 1940) ♣ <i>Philopterus stadleri</i> (Eichler, 1959)	Eurasian skylark
<b>MEGALURIDAE</b>	
° <i>Bowdleria punctata wilsoni</i> Stead, 1936 <i>Penenirmus</i> species	Codfish Island Fernbird
<b>ZOSTEROPIDAE</b>	
<i>Zosterops lateralis lateralis</i> (Latham, 1802) <i>Menacanthus eurysternus</i> (Burmeister, 1838)	Silvereye
<b>TURDIDAE</b>	
* <i>Turdus merula merula</i> Linnaeus, 1758 <i>Brueelia amsel</i> (Eichler, 1951) <i>Brueelia merulensis</i> (Denny, 1842) <i>Menacanthus eurysternus</i> (Burmeister, 1838) <i>Myrsidea thoracica</i> (Giebel, 1874) <i>Philopterus turdi</i> Denny, 1842	Eurasian blackbird
* <i>Turdus philomelos</i> Brehm, 1831 <i>Brueelia turdinulae</i> Ansari, 1956 <i>Menacanthus eurysternus</i> (Burmeister, 1838) <i>Philopterus turdi</i> Denny, 1842	Song thrush
<b>STURNIDAE</b>	
* <i>Sturnus vulgaris vulgaris</i> Linnaeus, 1758 <i>Brueelia nebulosa</i> (Burmeister, 1838) <i>Menacanthus eurysternus</i> (Burmeister, 1838) <i>Sturnidoecus sturni</i> (Schrank, 1776)	Common starling
* <i>Acridothères tristis</i> (Linnaeus, 1766) <i>Menacanthus eurysternus</i> (Burmeister, 1838) <i>Sturnidoecus</i> species	Common myna
<b>PASSERIDAE</b>	
* <i>Passer domesticus domesticus</i> (Linnaeus, 1758) <i>Brueelia cyclothorax</i> (Burmeister, 1838) <i>Menacanthus eurysternus</i> (Burmeister, 1838)	House sparrow
<b>MOTACILLIDAE</b>	
° <i>Anthus novaeseelandiae novaeseelandiae</i> (Gmelin, 1789) <i>Menacanthus eurysternus</i> (Burmeister, 1838) <i>Myrsidea</i> species ♣ <i>Philopterus irkutensis</i> Fedorenko, 1985	New Zealand pipit

- \* *Anthus novaeseelandiae steindachneri* Reischek, 1889  
 ♣ *Philopterus irkutensis* Fedorenko, 1985

Antipodes Island pipit

## PRUNELLIDAE

- \* *Prunella modularis* (Linnaeus, 1758)  
 ♣ *Philopterus modularis* Denny, 1842

Dunnock

## FRINGILLIDAE

- \* *Carduelis chloris* (Linnaeus, 1758)  
 ♣ *Brueelia breueri* Balát, 1955  
*Menacanthus eurysternus* (Burmeister, 1838)  
*Myrsidea serini* (Séguy, 1944)

European greenfinch

- \* *Carduelis carduelis britannica* (Hartert, 1871)  
 ♣ *Brueelia densilimba* (Nitzsch [in Giebel], 1866)  
*Myrsidea serini* (Séguy, 1944)

European goldfinch

- \* *Serinus canaria* (Linnaeus, 1758) [captive]  
*Menacanthus eurysternus* (Burmeister, 1838)  
*Myrsidea serini* (Séguy, 1944)

Island canary

## EMBERIZIDAE

- \* *Emberiza citrinella* Linnaeus, 1758  
*Brueelia delicata* (Nitzsch [in Giebel], 1866)  
*Myrsidea serini* (Séguy, 1944)

Yellowhammer

## MAMMALIA

## MARSUPIALIA

## MACROPODIDAE

- \* *Petrogale penicillata* (J.E. Gray, 1825)  
*Boopia notafusca* Le Souëf, 1902  
*Heterodoxus ampullatus* Kéler, 1971

Brush-tailed rock wallaby

## LAGOMORPHA

## LEPORIDAE

- \* *Oryctolagus cuniculus cuniculus* (Linnaeus, 1758)  
*Haemodipsus ventricosus* (Denny, 1842)
- \* *Lepus europaeus occidentalis* de Winton, 1898  
*Haemodipsus lyriocephalus* (Burmeister, 1838b)

European rabbit

Brown hare

## RODENTIA

## CAVIIDAE

- \* *Cavia porcellus* (Linnaeus, 1758)  
*Gliricola (Gliricola) porcelli* (Schrank, 1781)  
*Gyropus ovalis* Burmeister, 1838a

Guinea pig

## MURIDAE

- \* *Rattus exulans* (Peale, 1848)  
*Hoplopleura pacifica* Ewing, 1924
- \* *Rattus norvegicus* (Berkenhout, 1769)  
*Polyplax spinulosa* (Burmeister, 1838b)

Kiore

Norway rat

- 
- \* ***Rattus rattus* (Linnaeus, 1758)** Ship rat  
*Polyplax spinulosa* (Burmeister, 1838b)
- \* ***Mus musculus* Linnaeus, 1758** House mouse  
*Polyplax serrata* (Burmeister, 1838b)
- CARNIVORA**  
**CANIFORMIA**
- OTARIIDAE**
- Arctocephalus forsteri* (Lesson, 1828)** New Zealand fur seal  
*Antarctophthirus microchir* (Trouessart & Neumann, 1888)
- Phocarctos hookeri* (J.E. Gray, 1844)** New Zealand sea lion  
*Antarctophthirus microchir* (Trouessart & Neumann, 1888)
- PHOCIDAE**
- Mirounga leonina* (Linnaeus, 1758)** Southern elephant seal  
*Lepidophthirus macrorhini* Enderlein, 1904
- Leptonychotes weddellii* (Lesson, 1826)** Weddell seal  
*Antarctophthirus carlinii* Leonardi *et al.*, 2014
- Hydrurga leptonyx* (Blainville, 1820)** Leopard seal  
*Antarctophthirus ogmorhini* Enderlein, 1906
- Lobodon carcinophagus* (Hombron & Jacquinot, 1842)** Crabeater seal  
*Antarctophthirus lobodontis* Enderlein, 1909
- Ommatophoca rossii* J.E. Gray, 1844** Ross seal  
*Antarctophthirus mawsoni* Harrison, 1937
- CANIDAE**
- \* ***Canis familiaris* Linnaeus, 1758** Dog  
*Linognathus setosus* (von Olfers, 1816)  
*Trichodectes (Trichodectes) canis* (De Geer, 1778)
- MUSTELIDAE**
- \* ***Mustela erminea* Linnaeus, 1758** Stoat  
*Trichodectes (Stachiella) ermineae* (Hopkins, 1941)
- \* ***Mustela nivalis vulgaris* Erxleben, 1777** Weasel  
*Trichodectes (Stachiella) mustelae* (Schrank, 1803)
- \* ***Mustela furo* Linnaeus, 1758** Feral ferret  
*Trichodectes (Stachiella) ermineae* (Hopkins, 1941)
- FELIFORMIA**
- FELIDAE**
- \* ***Felis catus* Linnaeus, 1758** Feral Cat  
*Felicola (Felicola) subrostratus* (Burmeister, 1838a)

**PERISSODACTYLA**  
**EQUIDAE**

\* *Equus caballus* **Linnaeus, 1758** **Horse**  
*Haematopinus asini* (Linnaeus, 1758)  
*Werneckiella equi* (Denny, 1842)

\* *Equus asinus* **Linnaeus, 1758** **Donkey**  
*Haematopinus asini* (Linnaeus, 1758)  
*Werneckiella ocellata* (Piaget, 1880)

**ARCTIODACTYLA**  
**SUIDAE**

\* *Sus scrofa* **Linnaeus, 1758** **Feral pig**  
*Haematopinus suis* (Linnaeus, 1758)

**BOVIDAE**

\* *Bos taurus* **Linnaeus, 1758** **Feral cattle**  
*Bovicola (Bovicola) bovis* (Linnaeus, 1758)  
*Haematopinus eurysternus* Denny, 1842  
*Linognathus vituli* (Linnaeus, 1758)  
*Solenopotes capillatus* Enderlein, 1904

\* *Rupicapra rupicapra rupicapra* **Couturier, 1938** **Chamois**  
*Bovicola (Bovicola) longicornis* (Nitzsch, 1818)

\* *Hemitragus jemlahicus* **(Smith, 1826)** **Himalayan tahr**  
*Bovicola (Spinibovicola) hemitrangi* (Cummings, 1916)

\* *Capra hircus* **Linnaeus, 1758** **Feral goat**  
*Bovicola (Bovicola) caprae* (Gurlt, 1843)  
*Bovicola (Bovicola) limbatus* (Gervais, 1844)  
*Linognathus stenopsis* (Burmeister, 1838b)

\* *Ovis aries* **Linnaeus, 1758** **Feral sheep**  
*Bovicola (Bovicola) ovis* (Schrank, 1781)  
*Linognathus ovillus* (Neumann, 1907)  
*Linognathus pedalis* (Osborn, 1896)

**CAMELIDAE**

\* *Lama pacos* **(Linnaeus, 1758)** **Alpaca**  
*Bovicola (Lepikentron) breviceps* (Rudow, 1866)

**CERVIDAE**

\* *Cervus elaphus scoticus* **Lönnberg, 1906** **Red deer**  
*Bovicola (Bovicola) longicornis* (Nitzsch, 1818)  
*Solenopotes burmeisteri* (Fahrenheit, 1919)

\* *Cervus elaphus nelsoni* **(Bailey, 1935)** **Wapiti**  
*Bovicola (Bovicola) longicornis* (Nitzsch, 1818)

\* *Odocoileus virginianus borealis* **(Miller, 1900)** **White-tailed deer**  
*Solenopotes burmeisteri* (Fahrenheit, 1919)  
*Tricholipeurus lipeuroides* (Mégnin, 1884)  
*Tricholipeurus parallelus* (Osborn, 1896)

## PRIMATES HOMINIDAE

### \* *Homo sapiens* Linnaeus, 1758

Human

*Pediculus humanus capitis* De Geer, 1778

*Pediculus humanus humanus* Linnaeus, 1758

*Pthirus pubis* (Linnaeus, 1758)

## LIST OF BIRD SPECIES WHICH BREED IN THE NEW ZEALAND SUBREGION BUT WITH NO LICE COLLECTED FROM THEM IN THE SUBREGION

There are 18 bird species which breed in the New Zealand Subregion, but with no lice recorded from them. Among them, six species are rare endemics seldom searched for lice, six other species have been introduced by humans either purposely or by accident, and the remaining six are natives that also live in other parts of the world. Most of the native and introduced species have had lice recorded from them elsewhere, but in New Zealand they are rare or have restricted distributions. Despite some opportunities to search for lice from a few of these 18 species, no specimens have been found.

<i>Tachybaptus novaehollandiae novaehollandiae</i> (Stephens, 1826)	Australasian little grebe
<i>Pygoscelis papua papua</i> (J.R. Forster, 1781)	Northern gentoo penguin
<i>Pygoscelis antarctica</i> (J.R. Forster, 1781)	Chinstrap penguin
* <i>Cereopsis novaehollandiae</i> Latham, 1802	Cape Barren goose
<sup>ε</sup> <i>Lewinia muelleri</i> (Rothschild, 1893)	Auckland Island rail
<i>Elseyaornis melanops</i> (Vieillot, 1818)	Black-fronted dotterel
<i>Sternula nereis davisae</i> Mathews & Iredale, 1913	New Zealand fairy tern
* <i>Streptopelia risoria</i> (Linnaeus, 1758)	Barbary dove
* <i>Platycercus elegans</i> (Gmelin, 1788)	Crimson rosella
<sup>ε</sup> <i>Xenicus gilviventris</i> Pelzeln, 1867	Rock wren
<sup>ε</sup> <i>Gerygone albofrontata</i> G.R. Gray, 1845	Chatham Island warbler
<sup>ε</sup> <i>Mohoua ochrocephala</i> (Gmelin, 1789)	Yellowhead
<sup>ε</sup> <i>Mohoua novaeseelandiae</i> (Gmelin, 1789)	Brown creeper
<sup>ε</sup> <i>Petroica (Miro) traversi</i> (Buller, 1872)	Black robin
<i>Hirundo neoxena neoxena</i> Gould, 1842	Welcome swallow
* <i>Fringilla coelebs</i> Linnaeus, 1758	Chaffinch
* <i>Carduelis flammea</i> (Linnaeus, 1758)	Common redpoll
* <i>Emberiza cirrus</i> Linnaeus, 1766	Cirl bunting

ε Denotes species or subspecies which are **endemic** to the New Zealand Subregion.

\* Denotes species or subspecies which have been **introduced** to the New Zealand Subregion by human agency.

## ACKNOWLEDGEMENTS

I am extremely grateful to Jean-Claude Stahl (MONZ) for his outstanding photographic work, shooting and preparing all the photographs included in this book. I am indebted to Julia Kasper (Wellington, New Zealand) for her translations of German papers and for her assistance in locating types held in German institutions, and to Stephen Marder (Washington, D.C., United States of America) for his translation of Russian papers. I greatly acknowledge the assistance of Terry D. Galloway (Winnipeg, Canada) for supplying useful historical references, and that of Brian Michael Fitzgerald (Stokes Valley, Wellington, New Zealand) for reviewing the historical section of the text. For providing information about the locations of primary type specimens, I thank: Costica Adam (Grigore Antipa National Museum of Natural History, Bucharest, Romania); Jürgen Deckert (Museum für Naturkunde, Berlin, Germany); Daniel Gustafsson (University of Utah, Salt Lake City, Utah, U.S.A.); Takuya



Kiyoshi (National Science Museum, Tokyo, Japan); Robert McGowan (National Museum of Scotland); Karen Needham (University of British Columbia, Vancouver, Canada); Peter T. Oboyski (Essig Museum of Entomology, University of California, Berkeley, California, U.S.A.); Graham Rotheray (National Museum of Scotland); Jürgen Schmidl (University of Erlangen-Nuremberg, Germany); Karla Schneider (Institut für Zoologie und Zoologische Sammlungen, Martin-Luther-Universität Halle-Wittenberg, Germany); Vincent S. Smith (Natural History Museum, London, England); Oldřich Sychra (University of Veterinary and Pharmaceutical Sciences, Brno, Czech Republic); Gert Tröster (Zoologisches Museum, Georg-August-Universität Göttingen, Germany); Michel P. Valim (Museu de Zoologia da Universidade de São Paulo, São Paulo, Brasil) and Janet Weigner (Freien Universität, Berlin, Germany).

## REFERENCES

- Adam, C. 2007: Data on the chewing louse fauna (Phthiraptera: Amblycera, Ischnocera) from some Romanian autochthonous and exotic birds. *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"* 50: 145–210.
- Adams, R.J.; Price, R.D. & Clayton, D.H. 2005: Taxonomic revision of Old World members of the feather louse genus *Columbicola* (Phthiraptera: Ischnocera), including descriptions of eight new species. *Journal of Natural History* 39: 3545–6318.
- Ahmad, A.; Gupta, N. & Saxena, A.K. 2013: Reproductive system of an ischnoceran species, *Ardeicola expallidus* infesting cattle egret (*Bubulcus ibis*). *The Bioscan* 8(2): 443–446.
- Ahmad, A.; Gupta, N.; Saxena, A.K. & Gupta, D.K. 2015: Population levels of Phthiraptera on domestic ducks (*Anas platyrhynchos*) (Anseriformes: Anatidae). *Journal of Parasitic Diseases* 39(3): 567–571.
- Amerson, A.B. & Emerson, K.C. 1971: Records of Mallophaga from Pacific birds. *Atoll Research Bulletin* 146: 1–30.
- Andrews, J.R.H. 1964: The arthropod and helminth parasites of red deer (*Cervus elaphus* L.) in New Zealand. *Transactions of the Royal Society of New Zealand, Zoology* 5(9): 97–121.
- Andrews, J.R.H. 1970: Notes on infestations of *Damalinia hemitragi* (Cummings, 1916) (Mallophaga) on tahr in New Zealand. *The Journal of Parasitology* 56(4): 846–848.
- Andrews, J.R.H. 1972: Descriptions of the hitherto unknown males of *Damalinia longicornis* (Nitzsch, 1818) and *Damalinia hemitragi* (Cummings, 1916)—Trichodectidae; Mallophaga. *Journal of Natural History* 6: 153–157.
- Andrews, J.R.H. 1973: Records of Mallophaga of the genus *Damalinia* from wild ruminants in New Zealand. *The New Zealand Entomologist* 5(3–4): 324–330.
- Andrews, J.R.H. 1976a: The parasites of man in New Zealand: a review. *New Zealand Journal of Zoology* 3: 59–67.
- Andrews, J.R.H. 1976b: The parasitology of the Maori in pre-European times. *New Zealand Medical Journal* 84(568): 62–65.
- Andrews, J.R.H. 1980: Pediculosis in New Zealand. *The International Journal of Dermatology* 19(1): 32–34.
- Andrews, J.R.H. & Tonkin, S.L. 1989: Scabies and pediculosis in Tokelau Island children in New Zealand. *Journal of Research in Social Health* 109(6): 199–203.
- Anonymous (1837: New Zealanders. *The Morning Chronicle*. London. Issue 21193: 3 (17 October).
- Ansari, M.A.R. 1947: Mallophaga (Ischnocera) infesting birds in the Punjab (India). *Proceedings of the National Institute of Sciences of India* 13(6): 253–303.
- Ansari, M.A.R. 1951: Mallophaga (Amblycera) infesting birds in the Panjab (India). *Proceedings of the National Institute of Sciences of India* 17(2): 127–203.
- Ansari, M.A.R. 1955a: 11. The genera *Alcedoecus* Clay and Meinertzhagen, 1939; *Alcedoffula* Clay and Meinertzhagen, and *Capraiella* Conci, 1941 *Proceedings of the Seventh Pakistan Science Conference. Section: Biology*. Bahawalpur, Pakistan. Pp. 45–47.
- Ansari, M.A.R. 1955b: 15. The genera *Anaticola* Clay, 1936; *Anatoecus* Cummings, 1916 and *Incidifrons* Ewing, 1929. *Proceedings of the Seventh Pakistan Science Conference. Section: Biology*. Bahawalpur, Pakistan. Pp. 51–52.
- Ansari, M.A.R. 1956: Studies on *Bruelia* [sic] species (Mallophaga) occurring on true thrushes. *Biologia (Lahore)* 2: 102–143.
- Ansari, M.A.R. 1957: *Bruelia elegans* sp. nov., a new louse parasitic on the Australian crow, *Gymnohorina* [sic] *tibicen*, with a supplementary note on the genus *Bruelia* Kéler (Mallophaga: Ischnocera). *Biologia (Lahore)* 3: 122–131.
- Arnold, D.C. 2005: Review of the genus *Ornithobius* (Phthiraptera: Ischnocera: Philopteridae), with descriptions of two new species. *Journal of the Kansas Entomological Society* 78(2): 158–166.
- Arnold, D.C. 2006: Review of the genus *Acidoproctus* (Phthiraptera: Ischnocera: Philopteridae), with description of a new species. *Journal of the Kansas Entomological Society* 79(3): 272–282.
- Arriaza, B.; Orellana, N.C.; Barbosa, H.S.; Menna-Barreto, R.F.S.; Araújo, A. & Standen, V. 2012: Severe head lice infestation in an Andean mummy of Arica, Chile. *Journal of Parasitology* 98(2): 433–436.

- Australian Museum 2012: Australian species of Family Echinophthiriidae. *Antarctophthirus mawsoni* Harrison, 1937. Accessed via the *Atlas of Living Australia* (www.ala.org.au). <http://biocache.ala.org.au/occurrence/2a1b9922-072a-4cfe-ac49-8272b2c3cfc2>
- Aznar, F.J.; Leonardi, M.S.; Berón-Vera, B.; Vales, D.G.; Ameghino, S.; Raga, J.A. & Crespo, E.A. 2009: Population dynamics of *Antarctophthirus microchir* (Anoplura: Echinophthiriidae) in pups from South American sea lion, *Otaria flavescens*, in Northern Patagonia. *Parasitology* 136(3): 293–303.
- Baker, A.J. 1974: Ecological and behavioural evidence for the systematic status of New Zealand oystercatchers (Charadriiformes: Haematopodidae). *Life Sciences Contributions, Royal Ontario Museum* 96: 1–34.
- Baker, A.J.; Daugherty, C.H.; Colbourne, R. & McLennan, J.L. 1995: Flightless brown kiwis of New Zealand possess extremely subdivided population structure and cryptic species like small mammals. *Proceedings of the National Academy of Sciences of the United States of America* 92: 8254–8258.
- Balát, F. 1955: Příspěvek k poznání všenek rodu *Brüelia* I. *Práce Brněnské Základny Československé Akademie Věd* 27: 499–524, 2 pls.
- Balát, F. 1958: Beitrag zur Kenntnis der Mallophagenfauna der bulgarischen Vögel. *Práce Brněnské Základny Československé Akademie Věd* 30: 397–421.
- Balát, F. 1974: *Ornithobius matthewsi*—Eine neue Mallophagenart der Graugans, *Anser anser* L. *Annotationes Zoologicae et Botanicae* 94: 1–6.
- Banks, J. 1769: In: Hooker, J.D. (Ed.) 1896: *Journal of the Right Hon. Sir Joseph Banks during Captain Cook's first voyage in H.M.S. Endeavour in 1768–71 to Terra del Fuego, Otahite, New Zealand, Australia, the Dutch East Indies, etc.* London: Macmillan & Co Ltd. xlviii + 466 pp.
- Banks, J.C. & Palma, R.L. 2003: A new species and new host records of *Austrogoniodes* (Insecta: Phthiraptera: Philopteridae) from penguins (Aves: Sphenisciformes). *New Zealand Journal of Zoology* 30: 69–75.
- Banks, J.C.; Palma, R.L. & Paterson, A.M. 2006: Cophylogenetic relationships between penguins and their chewing lice. *Journal of Evolutionary Biology* 19: 156–166.
- Banks, J.C. & Paterson, A.M. 2004: A penguin-chewing louse (Insecta: Phthiraptera) phylogeny derived from morphology. *Invertebrate Systematics* 18(1): 89–100.
- Banks, J.C. & Paterson, A.M. 2005: Multi-host parasite species in cophylogenetic studies. *International Journal for Parasitology* 35: 741–746.
- Bany, J.; Pfeffer, A.; Phegan, M. & Heath, A.C.G. 1995: Proliferative responses of lymphocytes in *Bovicola ovis*-infested lambs. *International Journal for Parasitology* 25(6): 765–768.
- Barclay, L.E. 2002: Donkeys in New Zealand. *Donkey Dispatch* 30(3): 26.
- Barker, S.C. 1996: Boopiidae; Gyropidae; Trichodectidae; Anoplura. Pp. 91–105, 231–247. In: Wells, A. (Ed.) *Zoological Catalogue of Australia*. Volume 26. *Psocoptera, Phthiraptera, Thysanoptera*. Melbourne: CSIRO Publishing, Australia.
- Barker, S.C. & Close, R.L. 1990: Zoogeography and host associations of the *Heterodoxus octoseriatus* group and *H. ampullatus* (Phthiraptera: Boopiidae) from rock-wallabies (Marsupialia: *Petrogale*). *International Journal for Parasitology* 20(8): 1081–1087.
- Bartlow, A.W.; Villa, S.M.; Thompson, M.W. & Bush, S.E. 2016: Walk or ride? Phoretic behaviour of amblyceran and ischnoceran lice. *International Journal for Parasitology* 46(4): 221–227.
- Baum, H. 1968: Biologie und Ökologie der Amselfederläuse. *Angewandte Parasitologie* 9: 129–175.
- Bechet, I. 1968: Specii de *Luniceps* (Insecta, Mallophaga) din fauna României. *Comunicări de Zoologie. Societatea Științe Biologice din Republica Socialistă România* 6: 125–129.
- Bedford, G.A.H. 1929: Anoplura (Siphunculata and Mallophaga) from South African hosts. *15th Annual report of the Director of Veterinary Services, Union of South Africa*: 501–549.
- Bedford, G.A.H. 1930: New genera and species of Mallophaga from South African hosts. *16th Report of the Director of Veterinary Services and Animal Industry, Union of South Africa*: 153–173.
- Bedford, G.A.H. 1939: Notes on Menoponidae (Mallophaga) with descriptions of new genera and species. *Onderstepoort Journal of Veterinary Science and Animal Industry* 12(1): 121–152.
- Benoit, P.L.G. 1976: 17. Mallophaga. Pp. 233–235. In: La faune terrestre de l'Île de Sainte-Hélène. Troisième Partie. *Annales du Musée Royal de l'Afrique Centrale, Tervuren (Sciences Zoologiques)* 215: 1–533.
- Blagoveshtchensky, D.I. 1940: Mallophaga from birds of the Talysh lowlands. *Parazitologiceskij Sbornik (Moscow & Leningrad)* [= *Magasin de Parasitologie de l'institut Zoologique de l'Académie des Sciences de l'URSS* ] 8: 25–90 [in Russian, with English summary].
- Blagoveshtchensky, D.I. 1951: Mallophaga Tadzykistana. *Parazitologiceskij Sbornik (Moscow & Leningrad)* [= *Magasin de Parasitologie de l'institut Zoologique de l'Académie des Sciences de l'URSS* ] 13: 272–327 [in Russian].

- Bohemann, C.H. 1865: Spetsbergens insekt-fauna. *Öfversigt af Kungliga Svenska Vetenskaps-Akademiens Förhandlingar, Kobnhaven* 22(8): 563–580, 1 pl.
- Boeiro, M.; Rego, C. & Palma, R. 2008: Phthiraptera. Pp. 267–269, 273, 297. In: Borges, P.A.V.; Abreu, C.; Aguiar, A.M.F.; Carvalho, P.; Jardim, R.; Melo, I.; Oliveira, P.; Serrano, A.R.M. & Vieira, P. (Eds) *A list of the terrestrial fungi, flora and fauna of Madeira and Selvagens archipelagos*. Funchal and Angra do Heroísmo: Direcção Regional do Ambiente da Madeira and Universidade dos Açores. 438 pp.
- Boisduval, J.-B.A.D. de & Lacordaire, J.T. 1835: Second ordre. Anoploures. Pp. 117–125. In: *Faune entomologique des environs de Paris; ou species général des insectes qui se trouvent dans un rayon de quinze a vingt lieues aux alentours de Paris*. Paris: Méquignon-Marvis, Père et Fils, Libraires. Volume 1. 696 pp., 3 pls.
- Bourgeois, C.E. & Threlfall, W. 1979: Parasites of the greater shearwater (*Puffinus gravis*) from Newfoundland, Canada. *Canadian Journal of Zoology* 57(7): 1355–1357.
- Breljih, S. & Tovornik, D. 1961: Prispevek k poznavanju tekutov (Mallophaga) Jugoslavije I. *Biološki Vestnik* 9: 93–107.
- Breljih, S. & Tovornik, D. 1962: Prispevek k poznavanju tekutov (Mallophaga) Jugoslavije II. *Biološki Vestnik* 10: 85–100.
- Bresciani, J.; Haarløv, N.; Nansen, P. & Møller, G. 1983: Head louse (*Pediculus humanus* subsp. *capitis* de Geer) from mummified corpses of Greenlanders, A.D. 1460 ( $\pm$  50). *Acta Entomologica Fennica* 42: 24–27.
- Broekhuizen, S. 1971: On the occurrence of hare lice, *Haemodipsus* spp. (Anoplura: Hoplopleuridae) on hares, *Lepus europaeus*, in The Netherlands. *Zeitschrift für Parasitenkunde* 36: 158–168.
- Brooke, M. de L. & Nakamura, H. 1998: The acquisition of host-specific feather lice by common cuckoos (*Cuculus canorus*). *Journal of Zoology (London)* 244: 167–173.
- Browne, R.A.; Anderson, D.J.; Houser, J.N.; Cruz, F.; Glasgow, K.J.; Hodges, C.N. & Massey, G. 1997: Genetic diversity and divergence of endangered Galápagos and Hawaiian petrel populations. *The Condor* 99: 812–815.
- Buchanan, R.D. & Coles, P.G. 1971: Control of lice on cattle with Dursban. *New Zealand Veterinary Journal* 19(9): 197–202.
- Buckley, T.R.; Palma, R.L.; Johns, P.M.; Gleeson, D.M.; Heath, A.C.G.; Hitchmough, R.A. & Stringer, I.A.N. 2012: The conservation status of small or less well known groups of New Zealand terrestrial invertebrates. *The New Zealand Entomologist* 35(2): 137–143, 2 Appendices.
- Bull, P.C. 1960: Parasites of the European rabbit, *Oryctolagus cuniculus* (L.), on some Subantarctic Islands. *New Zealand Journal of Science* 3(2): 258–273.
- Burbidge, M.L.; Colbourne, R.M.; Robertson, H.A. & Baker, A.J. 2003: Molecular and other biological evidence supports the recognition of at least three species of brown kiwi. *Conservation Genetics* 4: 167–177.
- Burmeister, H. 1835: Pediculina. In: *Handbuch der Entomologie*, Berlin: Enslin, 1: 56–60.
- Burmeister, H. 1838a: Mallophaga. In: *Handbuch der Entomologie*, Berlin: Enslin, 2(1): 418–443.
- Burmeister, H. 1838b: Pediculina. 8 pp., 2 pls. In: Burmeister, H. (Ed.) *Genera quaedam Insectorum. Iconibus illustravit et descripsit*. Volume 1. *Rhynchotha*. Berolini: Burmeister et Stange. 132 pp. (unnumbered).
- Burns, D.A. 1987: The treatment of *Pthirus pubis* infestation of the eyelashes. *British Journal of Dermatology* 117(6): 741–743.
- Bush, S.E.; Weckstein, J.D.; Gustafsson, D.R.; Allen, J.; DiBlasi, E.; Shreve, S.C.; Boldt, R.; Skeen, H.R. & Johnson, K.P. 2015: Data supporting a molecular phylogeny of the hyper-diverse genus *Brueelia*. *Data in Brief* 5: 1078–1091.
- Bush, S.E.; Weckstein, J.D.; Gustafsson, D.R.; Allen, J.; DiBlasi, E.; Shreve, S.C.; Boldt, R.; Skeen, H.R. & Johnson, K.P. 2016: Unlocking the black box of feather louse diversity: A molecular phylogeny of the hyper-diverse genus *Brueelia*. *Molecular Phylogenetics and Evolution* 94: 737–751.
- Busvine, J.R. 1948: The ‘head’ and ‘body’ races of *Pediculus humanus* L. *Parasitology* 39(1, 2): 1–16.
- Busvine, J.R. 1978: Evidence from double infestations for the specific status of human head lice and body lice. *Systematic Entomology* 3(1): 1–8.
- Butler, F.T. & O’Connor, J.P. 1994: A review of the Irish Ischnocera and Amblycera (Phthiraptera). *The Irish Naturalist’s Journal* 24(11): 449–457.
- Buxton, P.A. 1939: *The louse. An account of the lice which infest man, their medical importance and control*. London: Edward Arnold & Co. x + 115 pp.
- Cabot, D. 1975: Feather lice (Mallophaga) from charadriiform birds collected at Galway Bay, Co. Galway. *The Irish Naturalist Journal* 18(5): 145–150.
- Callinan, A.P. 1980: Effects of artificially induced infestations of the cattle louse, *Linognathus vituli*. *Australian Veterinary Journal* 56: 484–486.
- Carriker, M.A., Jr. 1936: Studies in Neotropical Mallophaga, Part I.— Lice of the Tinamous. *Proceedings of the Academy of Natural Sciences of Philadelphia* 88: 45–218, 32 pls.
- Carriker, M.A., Jr. 1944: Studies in Neotropical Mallophaga— N° IV. New genera and species. *Boletín de Entomología Venezolana* 3(2): 65–104, 6 pls.

- Carriker, M.A., Jr. 1945a: Studies in Neotropical Mallophaga (V). *The lipeuroid forms of the New World "Galliformes"*. Part 2. *Revista Brasileira de Biologia* 5(1): 91–112.
- Carriker, M.A., Jr. 1945b: Studies in Neotropical Mallophaga (VII). *Goniodes* and allied genera from gallinaceous hosts. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 6: 355–399.
- Carriker, M.A., Jr. 1947: Neotropical Mallophaga miscellany N° 2. The genus *Ibidoecus* Cummings. *Boletín de Entomología Venezolana* 6(2–4): 111–136.
- Carriker, M.A., Jr. 1949: On a collection of Mallophaga from Guam, Marianas Islands. *Proceedings of the United States National Museum* 100(3254): 1–24.
- Carriker, M.A., Jr. 1954: Studies in Neotropical Mallophaga (XIII)—The Menoponidae of the Neotropical Psittacidae. *Revista Brasileira de Entomologia* 2: 145–173.
- Carriker, M.A., Jr. 1955: A corrected list of the Venezuelan Mallophaga published by E.W. Stafford, 1943. *Boletín de Entomología Venezolana* 11(1–2): 31–54.
- Carriker, M.A., Jr. 1957: Notes on some of the Vernon L. Kellogg types of Mallophaga. *Microentomology* 22(5): 95–110.
- Carriker, M.A., Jr. 1958: Neotropical Mallophaga miscellany N°. 7. New Mallophaga from the Antarctic. *Acta Zoológica Lilloana* 15: 183–188.
- Carriker, M.A., Jr. 1964: Descriptions of new and little known species of Mallophaga (Insecta) from maritime hosts of Chile, South America. *Publicaciones del Centro de Estudios Entomológicos* 6: 1–26.
- Carriker, M.R. 2001: *Vista Nieve: the remarkable, true adventures of an early twentieth-century naturalist and his family in Colombia, South America*. Rio Hondo, Texas: Blue Mantle Press. xiii + 312 pp.
- Castro, D.C. & Cicchino, A.C. 1978: Contribución al conocimiento de los malófagos argentinos III. Sobre algunos Menoponidae de la avifauna bonaerense: *Menacanthus eurysternus* (Burmeister) y *Menacanthus pici* (Denny) (Insecta: Mallophaga). *Revista de la Sociedad Entomológica Argentina* 37(1–4): 77–83.
- Castro, D.C. & Cicchino, A.C. 1983: Contribución al conocimiento de los malófagos argentinos. XIII. Dos nuevas especies del género *Incidifrons* Ewing, 1929 (Mallophaga – Philopteridae) parásitas del género *Fulica* (Aves – Gruiformes) de Argentina. *Revista de la Sociedad Entomológica Argentina* 42(1–4): 273–281.
- Castro, D.C. & Cicchino, A.C. 1996: Algunas Menoponidae (Insecta – Phthiraptera) de aves bonaerenses. *Revista de la Asociación de Ciencias Naturales del Litoral* 27(2): 137–140.
- Catanach, T.A. & Johnson, K.P. 2015: Independent origins of the feather lice (Insecta: *Degeeriella*) of raptors. *Biological Journal of the Linnean Society* 114(4): 837–847.
- Chalmers, K. & Charleston, W.A.G. 1980a: Cattle lice in New Zealand: observations on the prevalence, distribution and seasonal; patterns of infestation. *New Zealand Veterinary Journal* 28: 198–200.
- Chalmers, K. & Charleston, W.A.G. 1980b: Cattle lice in New Zealand: observations on the biology and ecology of *Damalinea bovis* and *Linognathus vituli*. *New Zealand Veterinary Journal* 28: 214–216.
- Chalmers, K. & Charleston, W.A.G. 1980c: Cattle lice in New Zealand: effects on host liveweight gain and haematocrit levels. *New Zealand Veterinary Journal* 28: 235–237.
- Charleston, W.A.G. 1980: Lungworm and lice of the red deer (*Cervus elaphus*) and the fallow deer (*Dama dama*)—a review. *New Zealand Veterinary Journal* 28(8): 150–152.
- Checklist Committee (OSNZ): Gill, B.J. (Convener); Bell, B.D.; Chambers, G.K.; Medway, D.G.; Palma, R.L.; Scofield, R.P.; Tennyson, A.J.D. & Worthy, T.H. 2010: *Checklist of the birds of New Zealand, Norfolk and Macquarie Islands, and the Ross Dependency, Antarctica*. Fourth Edition. Wellington: Te Papa Press in association with the Ornithological Society of New Zealand. x + 501 pp.
- Choe, J.C. & Kim, K.C. 1987: Community structure of arthropod ectoparasites on Alaskan seabirds. *Canadian Journal of Zoology* 65(12): 2998–3005.
- Choe, J.C. & Kim, K.C. 1988: Microhabitat preference and coexistence of ectoparasitic arthropods on Alaskan seabirds. *Canadian Journal of Zoology* 66(4): 987–997.
- Christie, A.H.C. & Andrews, J.R.H. 1964: Introduced ungulates in New Zealand (a) Himalayan tahr *Tuataria* 12(2): 69–77.
- Cicchino, A.C. & González-Acuña, D.A. 2012: Species in the genus *Bonomiella* Conci, 1942 (Phthiraptera: Menoponidae) from Argentina and Chile. *Zootaxa* 3427: 47–56.
- Cicchino, A.C. & Valim, M.P. 2015: Notes on taxonomy and distribution of *Myrsidea serini* (Séguy, 1944) (Phthiraptera: Amblycera: Menoponidae) on southern South American passerine birds (Aves: Passeriformes). *Papéis Avulsos de Zoologia* 55(16): 231–243.
- Clay, T. 1936: Two new genera of Mallophaga. *Proceedings of the Zoological Society of London*. [no volume numbering]: 615–618.
- Clay, T. 1938: A revision of the genera and species of Mallophaga occurring on gallinaceous hosts. Part I. *Lipeurus* and related genera. *Proceedings of the Zoological Society of London*. [no volume number]: 109–204, 14 pls.

- Clay, T. 1940a: Anoplura. *British Graham Land Expedition, 1934–37, Scientific Reports 1*(5): 295–318, 1 pl.
- Clay, T. 1940b: A revision of the genera and species of Mallophaga occurring on Gallinaceous hosts. Part II. *Goniodes*. *Proceedings of the Zoological Society of London* (Series B) 110: 1–120.
- Clay, T. 1941: A new genus and species of Mallophaga. *Parasitology* 33(1): 119–129.
- Clay, T. 1949a: Species of the genus *Saemundssonina* (Mallophaga) from the Sterninae. *American Museum Novitates* 1409: 1–25.
- Clay, T. 1949b: Systematic notes on the Piaget collections of Mallophaga.— Pt I. *Annals and Magazine of Natural History* (Series 12) 2: 811–838, 895–921.
- Clay, T. 1951a: Systematic notes on the Piaget collections of Mallophaga.— Pt II. *Annals and Magazine of Natural History* (Series 12) 4(38): 173–182.
- Clay, T. 1951b: An introduction to a classification of the avian Ischnocera (Mallophaga): Part I. *The Transactions of the Royal Entomological Society of London* 102(2): 171–194, 1 pl.
- Clay, T. 1953a: Revisions of the genera of Mallophaga. I. The *Rallicola*-complex. *Proceedings of the Zoological Society of London* 123(3): 563–587, pl. 1.
- Clay, T. 1953b: Systematic notes on the Piaget collections of Mallophaga.— Pt IV. *Annals and Magazine of Natural History* (Series 12) 6: 641–657.
- Clay, T. 1958a: A note on some Antarctic Mallophaga. *Annals and Magazine of Natural History* (Series 13) 1: 250–256, 1 pl.
- Clay, T. 1958b: Revisions of Mallophaga genera. *Degeeriella* from the Falconiformes. *Bulletin of the British Museum (Natural History) Entomology* 7(4): 121–207, 9 pls.
- Clay, T. 1958c: Mallophaga (Insecta). Part I. *The Natural History of Rennell Island, British Solomon Islands* (Copenhagen) 2(15): 143–149.
- Clay, T. 1959: Key to the species of *Austromenopon* Bedford (Mallophaga) parasitic on the Charadriiformes. *Proceedings of the Royal Entomological Society of London* (Series B) 28(11–12): 157–168.
- Clay, T. 1961: A new genus and species of Menoponidae (Mallophaga, Insecta) from *Apteryx*. *Annals and Magazine of Natural History* (Series 13) 3(33): 571–576.
- Clay, T. 1962: A key to the species of *Actornithophilus* Ferris with notes and descriptions of new species. *Bulletin of the British Museum (Natural History) Entomology* 11(5): 189–244, 8 pls.
- Clay, T. 1964a: Insects of Campbell Island. Phthiraptera. *Pacific Insects Monograph* 7: 230–234.
- Clay, T. 1964b: Geographical distribution of the Mallophaga (Insecta). *Bulletin of the British Ornithologist's Club* 84: 14–16.
- Clay, T. 1966a: A new species of *Apterygon* (Mallophaga: Menoponidae). *The Entomologist* 99: 292–293, 2 pls.
- Clay, T. 1966b: The species of *Strigiphilus* (Mallophaga: Philopteridae) parasitic on the barn owls, *Tyto* (Tytonidae). *Journal of the Entomological Society of Queensland* 5: 10–17.
- Clay, T. 1966c: Contributions towards a revision of *Myrsidea* Waterston. I. (Menoponidae: Mallophaga). *Bulletin of the British Museum (Natural History) Entomology* 17(8): 327–395, 2 pls.
- Clay, T. 1967: Mallophaga (biting lice) and Anoplura (sucking lice). Part I: *Austrogoniodes* (Mallophaga) parasitic on penguins (Sphenisciformes). Pp. 149–155, 170–176. In: Gressitt, J.L. (Ed.) *Antarctic Research Series*, volume 10, *Entomology of Antarctica*. Washington D.C.: American Geophysical Union. xii + 395 pp.
- Clay, T. 1970: The Amblycera (Phthiraptera: Insecta). *Bulletin of the British Museum (Natural History) Entomology* 25(3): 75–98, 5 pls.
- Clay, T. 1971: The identity of *Nirmus bracteatus* Nitzsch (Mallophaga: Insecta). *The Western Australian Naturalist* 12(2): 44–45.
- Clay, T. 1972: The species of *Rallicola* (Insecta: Phthiraptera: Ischnocera) parasitic on Kiwis (*Apteryx*). *New Zealand Journal of Science* 15(1): 70–76.
- Clay, T. 1973: The species groups of *Pectinopygus* (Phthiraptera: Philopteridae). *Bulletin of the British Museum (Natural History) Entomology* 29(4): 200–224, 3 pls.
- Clay, T. 1976a: The species of *Ibidoecus* (Phthiraptera) on Threskiornis (Aves). *Systematic Entomology* 1: 1–7.
- Clay, T. 1976b: Geographical distribution of the avian lice (Phthiraptera): A review. *Journal of the Bombay Natural History Society* 71(3): 536–547.
- Clay, T. 1977: The *Strigiphilus cursitans* group (Phthiraptera: Insecta). *Records of the Queen Victoria Museum* 56: 1–4.
- Clay, T. 1981a: A report on a collection of lice (Boopidae [sic]: Phthiraptera) on *Petrogale* (Rock wallabies). *Proceedings of the Linnean Society of New South Wales* 105(1): 65–78.
- Clay, T. 1981b: The ischnoceran lice (Phthiraptera) of the oystercatchers (Aves: Haematopodidae). *Canadian Journal of Zoology* 59(6): 933–938.
- Clay, T. & Hopkins, G.H.E. 1950: The early literature on Mallophaga. Part I. 1758–62. *Bulletin of the British Museum (Natural History) Entomology* 1(3): 221–272, 2 pls.

- Clay, T. & Hopkins, G.H.E. 1951: The early literature on Mallophaga. Part II, 1763–1775. *Bulletin of the British Museum (Natural History) Entomology* 2(1): 1–37, 3 pls.
- Clay, T. & Hopkins, G.H.E. 1954: The early literature on Mallophaga. (Part III, 1776–1786). *Bulletin of the British Museum (Natural History) Entomology* 3(6): 221–266, 3 pls.
- Clay, T. & Hopkins, G.H.E. 1955: Notes on the Rudow Collection of Mallophaga at Hamburg. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 53: 49–73.
- Clay, T. & Hopkins, G.H.E. 1960: The early literature on Mallophaga. (Part IV, 1787–1818). *Bulletin of the British Museum (Natural History) Entomology* 9(1): 1–61, 6 pls.
- Clay, T. & Hopkins, G.H.E. 1961: *Harrisoniella* Bedford, 1928 [sic]: proposed designation of a type-species under the plenary powers (Insecta, Mallophaga). *Bulletin of Zoological Nomenclature* 18(3): 195–198.
- Clay, T. & Meinertzhagen, R. 1938a: Two new genera of Mallophaga. *The Entomologist* 71: 73–76.
- Clay, T. & Meinertzhagen, R. 1938b: New genera and species of Mallophaga. *The Entomologist* 71: 275–279.
- Clay, T. & Meinertzhagen, R. 1939a: New genera and species of Mallophaga. *The Entomologist* 72: 161–168.
- Clay, T. & Meinertzhagen, R. 1939b: Three new genera of Mallophaga from Charadriiformes. *Annals and Magazine of Natural History (Series 11)* 4: 450–454.
- Clay, T. & Moreby, C. 1967: Mallophaga (biting lice) and Anoplura (sucking lice). Part II: Keys and locality lists of Mallophaga and Anoplura. Pp. 157–169, 177–196. In: Gressitt, J.L. (Ed.) *Antarctic Research Series*, volume 10, *Entomology of Antarctica*. Washington D.C.: American Geophysical Union. xii + 395 pp.
- Clay, T. & Moreby, C. 1970: Mallophaga and Anoplura of Subantarctic islands. *Pacific Insects Monographs* 23: 216–220.
- Clayton, D.H. 1990: Host specificity of *Strigiphilus* owl lice (Ischnocera: Philopteridae), with the description of new species and host associations. *Journal of Medical Entomology* 27(3): 257–265.
- Clayton, D.H.; Bush, S.E. & Johnson, K.P. 2015: *Coevolution of life on hosts. Integrating ecology and history*. Chicago & London: The University of Chicago Press. xiv + 294 pp., 16 pls.
- Clayton, D.H. & Price, R.D. 1984: Taxonomy of the *Strigiphilus cursitans* group (Ischnocera: Philopteridae), parasites of owls (Strigiformes). *Annals of the Entomological Society of America* 77(4): 340–363.
- Cocker, M. 1989: *Richard Meinertzhagen: Soldier, scientist, and spy*. London: Martin Secker & Warburg Ltd. 256 pp.
- Cohen, B.L.; Baker, A.J.; Blechschmidt, K.; Dittmann, D.L.; Furness, R.W.; Gerwin, J.A.; Helbig, A.J.; De Korte, J.; Marshall, H.D.; Palma, R.L.; Peter, H.-U.; Ramli, R.; Siebold, I.; Willcox, M.S.; Wilson, R.H. & Zink, R.M. 1997: Enigmatic phylogeny of skuas (Aves: Stercorariidae). *Proceedings of the Royal Society of London (B)* 264: 181–190.
- Conci, C. 1941: Nuovi generi di mallofagi. *Bolletino della Società Entomologica Italiana (Genova)* 73(8) 126–127.
- Conci, C. 1942a: Un nuovo genere di Somaphantidae, ascrivibile ad una nuova sottofamiglia (Mallophaga–Liotheida). *Studi Trentini di Scienze Naturali* 23(2): 123–126.
- Conci, C. 1942b: Quattro nuovi generi ed una nuova sottofamiglia di Menoponidae dei passeracei (Mallophaga). *Bolletino della Società Entomologica Italiana (Genova)* 74: 30–31.
- Conci, C. 1942c: I mallofagi degli Psittaciformes. *Bolletino della Società Entomologica Italiana (Genova)* 74: 33–41.
- Conci, C. 1952: L'allevamento in condizioni sperimentali dei Mallofagi. I.— *Cuclotogaster heterographus* Nitzsch. *Bolletino dei Musei e degli Istituti Biologici della Università di Genova* 24(150): 17–40.
- Conci, C. 1956: L'allevamento in condizioni sperimentali dei Mallofagi. III.— *Columbicola c. columbae* (Linnaeus, 1758). *Bolletino dei Musei e degli Istituti Biologici della Università di Genova* 26(162): 47–70.
- Conti, J.A. & Forrester, D.J. 1981: Interrelationships of parasites of white-winged doves and mourning doves in Florida. *Journal of Wildlife Diseases* 17(4): 529–536.
- Corrin, K. & Burnett, K. 1989: Alpacas and llamas at sea. *Surveillance* 16(2): 21–23.
- Crosby, T.K; Dugdale, J.S. & Watt, J.C. 1976: Recording specimen localities in New Zealand: an arbitrary system of areas and codes defined. *New Zealand Journal of Zoology* 3: 69 + map.
- Crosby, T.K; Dugdale, J.S. & Watt, J.C. 1998: Area codes for recording specimen localities in the New Zealand subregion. *New Zealand Journal of Zoology* 25: 175–183.
- Crossland, A.C. 1993: Do Arctic skuas sometimes roost on land at night?. *Notornis* 40(4): 305–306.
- Cummings, B.F. 1914: Descriptions of five new species of Anoplura and Mallophaga. *Bulletin of Entomological Research* 5(2): 155–177.
- Cummings, B.F. 1915: On two new species of *Polyplax* (Anoplura) from Egypt. *Proceedings of the Zoological Society of London 1915*: 245–272.
- Cummings, B.F. 1916a: Studies on the Anoplura and Mallophaga, being a report upon a collection from the mammals and birds in the Society's Gardens.— Part I., with a preface. *Proceedings of the Zoological Society of London 1916*: 253–295.
- Cummings, B.F. 1916b: Studies on the Anoplura and Mallophaga, being a report upon a collection from the mammals and birds in the Society's Gardens.— Part II. *Proceedings of the Zoological Society of London 1916*: 643–693.

- Cummings, B.F. 1916c: Note on the thorax in Anoplura and in the genus *Nesiotinus* of the Mallophaga. *Annals and Magazine of Natural History* (Series 8) 17: 171–174.
- Dalgleish, R.C. 1972: The *Penenirmus* (Mallophaga: Ischnocera) of the Picidae (Aves: Piciformes). *Journal of the New York Entomological Society* 80: 83–104.
- Davies, S. 1995: Feather lice (Insecta: Phthiraptera) on wrybill. *The Stilt* 27: 46–47.
- De Geer, C. 1778a: Des poux. Pp. 62–69, pl. 1. In: *Mémoires pour servir à l'histoire des insectes*. Volume 7. Stockholm: Pierre Hesselberg. xii + 950 pp., 49 pls.
- De Geer, C. 1778b: Des Ricins. Pp. 69–82, pl. 4. In: *Mémoires pour servir à l'histoire des insectes*. Volume 7. Stockholm: Pierre Hesselberg. xii + 950 pp., 49 pls.
- Denny, H. 1842: *Monographia Anoplurorum Britanniae or, an essay on the British species of parasitic insects belonging to the order of Anoplura of Leach, with the modern divisions of the genera according to the views of Leach, Nitzsch, and Burmeister, with highly magnified figures of each species*. London: Henry G. Bohn. xxiv + 262 pp., 26 pls.
- Derylo, A. 1970: Mallophaga as a reservoir of *Pasteurella multocida*. *Acta Parasitologica Polonica* 17(35): 301–313.
- Derylo, A. 1977: Badania nad rola wszołow *Eomenacanthus stramineus* (Nitzsch) w przenoszeniu toksoplazmozy u kur. *Wiadomości Parazytologiczne* 23: 131–134.
- Dickinson, E.C. (Ed.) 2003: *The Howard & Moore complete checklist of the birds of the world* (Third edition). London: Christopher Helm. Pp. 1–1040.
- Dik, B.; Per, E.; Erciyas Yavuz, K. & Yamaç, E. 2015: Chewing lice (Phthiraptera: Amblycera, Ischnocera) species found on birds in Turkey, with new records and a new host association. *Turkish Journal of Zoology* 39: 790–798.
- Dubinín, V.B. 1938: Ismeneniya parasitofauny karavajki (*Plegadis falcinellus*) vysyvajemyje vosrostorn i migraczej chosiaina [Changements de la parasitofaune *Plegadis falcinellus* L., provoqués par l'âge et la migration de l'hôte]. *Trudy Astrachanskogo Gosudarstvennogo Zapovednika* 2: 114–213 [in Russian, with French summary].
- Dufour, L. 1835: Description et iconographe de trois espèces du genre *Philopterus*, parasites de l'albatros. *Annales de la Société Entomologique de France* 4: 669–681, 1 pl.
- Dunn, R.R.; Harris, N.C.; Colwell, R.K.; Koh, L.P. & Sodhi, N.S. 2009: The sixth mass coextinction: are most endangered species parasites and mutualists? *Proceedings of the Royal Society B, Biological Sciences* 276: 3037–3045.
- Durden, L.A. 2001: Lice (Phthiraptera). Pp. 3–17. In: Samuel, W.M.; Pybus, M.J. & Kocan, A.A. (Eds) *Parasitic diseases of wild mammals*. Second edition. Ames: Iowa State University Press. viii + 568 pp.
- Durden, L.A. & Musser, G.G. 1994a: The sucking lice (Insecta, Anoplura) of the world: a taxonomic checklist with records of mammalian hosts and geographical distributions. *Bulletin of the American Museum of Natural History* 218: 1–90.
- Durden, L.A. & Musser, G.G. 1994b: The mammalian hosts of the sucking lice (Anoplura) of the world: A host-parasite list. *Bulletin of the Society for Vector Ecology* 19(2): 130–168.
- Durden, L.A.; Willis, E.E.; Garretson, A.M. & Ereemeeva, M.E. 2014: The sucking lice (Phthiraptera: Anoplura) of Georgia, USA: hosts, geographical distributions, and medical/veterinary importance. *Journal of Entomological Science* 49(3): 246–271.
- Durrant, E.P. 1906: Descriptions of new Mallophaga. *The Ohio Naturalist* 6(7): 528–530.
- Edwards, R.L. 1961: Studies of the Philopteridae (Mallophaga) from the birds of the order Procellariiformes [sic]. 1. The genus *Halipeurus*. *The Journal of Parasitology* 47(1): 125–157.
- Edwards, R.L. 1965: Revision of the genus *Aquanirmus* (Mallophaga: Philopteridae), parasitic on grebes (Podicipidae). *The Canadian Entomologist* 97(9): 920–935.
- Eichler, W. 1940a: Notulae Mallophagologicae.—I. Neue Gattungen und Subfamilien von Haarlingen. *Zoologischer Anzeiger* 129(5/6): 158–162.
- Eichler, W. 1940b: Notulae Mallophagologicae. —IV. Neue Gattungen und höhere Einheiten von Kletterfederlingen. *Zoologischer Anzeiger* 130(5/6): 97–103.
- Eichler, W. 1940c: Notulae Mallophagologicae. III. Die Unterfamilie Menacanthinae nov. subfam. *Zentralblatt für Bakteriologie, Parasitenkunde und Infektionskrankheiten* 145: 361–365.
- Eichler, W. 1941a: Zur Klassifikation der Lauskerfe (Phthiraptera Haeckel: Rhynchophthirina, Mallophaga und Anoplura). *Archiv für Naturgeschichte. Neue Folge* 10: 345–398.
- Eichler, W. 1941b: Mallophagen-Synopsis. II. Genus *Stachiella*. *Zoologischer Anzeiger* 136: 187–189.
- Eichler, W. 1941c: Notulae Mallophagologicae. II. Neue Gattungen bei Haufffußfederlingen. *Stettiner Entomologische Zeitung* 102(1): 125–128.
- Eichler, W. 1943: Mallophagen-Synopsis. IX. Genus. *Mitteilungen der Münchener Entomologischen Gesellschaft* 33(1): 236–239.
- Eichler, W. 1944a: Notulae Mallophagologicae. X. *Anseriphilus* nov. gen. und andere Neuerungen bei amblyceren Federlingen. *Deutsche Entomologische Zeitschrift* 1943(1/2): 56–64.

- Eichler, W. 1944b: Notulae Mallophagologicae. XI. Acht neue Gattungen der Nirmi und Docophori. *Stettiner Entomologische Zeitung* 105: 80–82.
- Eichler, W. 1947: Dr. E. Mjöberg's zoological collections from Sumatra. 15. Mallophaga. *Arkiv för Zoologi* 39A(2): 1–21.
- Eichler, W. 1948: Schutzfärbung bei Federlingen. *Vögel der Heimat* 6: 103–108.
- Eichler, W. 1949a: Phthirapterorum nova genera. *Bollettino della Società Entomologica Italiana* 79: 11–13.
- Eichler, W. 1949b: Notulae Mallophagologicae. XV. Sturmvogel-Federlinge. *Revista Brasileira de Biologia* 9(3): 337–347.
- Eichler, W. 1951a: Die Federlinge der Drosseln. Pp. 29–47. *Bedeutung der Vogelwelt in Forschung und Praxis – Zusammenstellung der Vorträge gehalten auf der Ersten Ornithologen-Tagung in der Deutschen Demokratischen Republik am 21 und 22 Oktober 1950 in Leipzig*. 63 pp.
- Eichler, W. 1951b: Mallophagen-Synopsis XVII. Genus *Koeniginirmus*. *Bonner zoologische Beiträge* 2(1–2): 125–134.
- Eichler, W. 1952a: Mallophagen-Synopsis XXV. Genus *Harrisoniella*. *Beiträge zur Vogelkunde* 2: 40–43.
- Eichler, W. 1952b: Notulae Mallophagologicae. XXVI. *Rhombiceps* n. g. und andere neue Federlingsgattungen. *Zoologischer Anzeiger* 149: 74–78.
- Eichler, W. 1953: Neue oder wenig bekanntehaustierparasiten. I. *Werneckiella equi asini* nov. subsp. vom Esel. *Monatsheften für Veterinärmedizin* 8(18): 445–447.
- Eichler, W. 1958: Notulae Mallophagologicae. XX. Neue Federlingsformen in meiner Federlingsbearbeitung der "Neuen Brehm-Bucherei". *Naturwissenschaftliches Museum Nachrichten, Aschaffenburg* 58: 59–64, pl. 3.
- Eichler, W. 1959: Notizen zur angewandten Parasitenkunde (III). 24. Ein neuer Lerchenfederling. *Das Deutsche Gesundheitswesen* 14(25): 1173.
- Eichler, W. & Vasjukova, T.T. 1980: Die Mallophagengattung *Anaticola* (Phthiraptera, Mallophaga). *Deutsche Entomologische Zeitschrift (Neue Folge)*. 27(4–5): 335–375, 15 pls.
- Eichler, W. & Vasjukova, T.T. 1981: Die Mallophagengattung *Trinoton*. *Mitteilungen aus dem Zoologischen Museum in Berlin* 57(1): 23–62, 9 pls.
- Emerson, K.C. 1949: A species of Mallophaga from a partridge. *Entomological News* 60: 116–117.
- Emerson, K.C. 1953: A new species of *Carduceps*. *Proceedings of the Entomological Society of Washington* 55(5): 209–211.
- Emerson, K.C. 1954: A review of the genus *Menopon* Nitzsch, 1818 (Mallophaga). *Annals and Magazine of Natural History (Series 12)* 7: 225–232.
- Emerson, K.C. 1955: A review of the genus *Rallicola* (Phloptoridae, Mallophaga) found on Aramidae, Psophiidae and Rallidae. *Annals of the Entomological Society of America* 48(4): 284–299.
- Emerson, K.C. 1956a: Mallophaga (chewing lice) occurring on the domestic chicken. *Journal of the Kansas Entomological Society* 29(2): 63–79.
- Emerson, K.C. 1956b: A note on the identity of *Longimenopon pediculoides* (Mjöberg). *Proceedings of the Entomological Society of Washington* 58(5): 295–296.
- Emerson, K.C. 1957: A new species of Mallophaga from the pigeon. *The Florida Entomologist* 40(2): 63–64.
- Emerson, K.C. 1961a: A new species of Mallophaga from the peafowl. *The Florida Entomologist* 44(3): 117–118.
- Emerson, K.C. 1961b: The Vernon L. Kellogg Mallophaga type material in the United States National Museum. *Proceedings of the Entomological Society of Washington* 63(4): 247–254.
- Emerson, K.C. 1962: Mallophaga (chewing lice) occurring on the turkey. *Journal of the Kansas Entomological Society* 35(1): 196–201.
- Emerson, K.C. 1965: The Vernon L. Kellogg Mallophaga type material in the Cornell University Collection. *Proceedings of the Entomological Society of Washington* 67(1): 46–50.
- Emerson, K.C. 1966: New species and records of *Rallicola* (Mallophaga: Phloptoridae) from New Guinea. *Journal of Medical Entomology* 3(3–4): 336–337.
- Emerson, K.C. (Ed.) 1967: Carriker on Mallophaga. Posthumous papers, catalog of forms described as new, and bibliography. Melbourne A. Carriker, Jr. 1879–1965. *Bulletin of the United States National Museum* 248: x + 150 pp.
- Emerson, K.C. 1971: Mallophaga determinations. Chapter 28: 360. In: van Zinderen Bakker Sr, E.M.; Winterbottom J.M. & Dyer, R.A. (Eds) *Marion and Prince Edward Islands*. Cape Town: A.A. Balkema. 427 pp., 25 pls, 89 photos.
- Emerson, K.C. 1972a: *Checklist of the Mallophaga of North America (North of Mexico)* Part I. Suborder Ischnocera. Dugway (Utah): Deseret Test Center. 200 pp.
- Emerson, K.C. 1972b: *Checklist of the Mallophaga of North America (North of Mexico)* Part II. Suborder Amblycera. Dugway (Utah): Deseret Test Center. 118 pp.
- Emerson, K.C. 1979: *Lice in my life*. Arlington, Virginia: published by the author. 106 pp.
- Emerson, K.C. & Johnson, J.C. 1961: The genus *Penenirmus* (Mallophaga) found on North American woodpeckers. *Journal of the Kansas Entomological Society* 34(1): 34–43.



- Emerson, K.C. & Price, R.D. 1968: A new species of *Dennyus* (Mallophaga: Menoponidae) from the Malaysian spine-tailed swift. *Proceedings of the Biological Society of Washington* 81: 87–90.
- Emerson, K.C. & Price, R.D. 1975: Mallophaga of Venezuelan mammals. *Brigham Young University Science Bulletin. Biological Series* 20(3): 1–77.
- Emerson, K.C. & Ward, R.A. 1958: Notes on Philippine Mallophaga, I. Species from Ciconiiformes, Anseriformes, Falconiformes, Galliformes, Gruiformes and Charadriiformes. *Fieldiana – Zoology* 42(4): 49–61.
- Enderlein, G. 1904a: *Lepidophthirus* nov. gen., eine Laus der Elefantenrobbe von der Kerguelen-Insel. *Zoologischer Anzeiger* 28: 43–47.
- Enderlein, G. 1904b: Läusestudien. I. Über die Morphologie, Klassifikation und systematische Stellung der Anopluren nebst Bemerkungen zur Systematik der Insektenordnungen. *Zoologischer Anzeiger* 28: 121–147.
- Enderlein, G. 1904c: Läuse-Studien. *Zoologischer Anzeiger* 28: 220–223.
- Enderlein, G. 1905: Läusestudien. IV. Über einen auffälligen Sexualdimorphismus bei *Polyplax spinulosa* (Burm.). *Zoologischer Anzeiger* 29: 192–194.
- Enderlein, G. 1906: Läusestudien V. Schuppen als sekundäre Atmungsorgane, sowie über eine neue antarktische Echinophthiriiden-Gattung. 12. Beitrag zur Kenntnis der antarktischen Fauna. *Zoologischer Anzeiger* 29: 659–665.
- Enderlein, G. 1908–1909: Die Insekten des antarktischen Gebietes. *Deutsche Südpolar-Expedition 1901–1903* 10, *Zoologie* 2: 361–528, 24 pls.
- Enderlein, G. 1917: Über einige subantarktische Mallophagen. *Zoologischer Anzeiger* 49: 240–245.
- Escalante, G.C.; Sweet, A.D.; McCracken, K.G.; Gustafsson, D.R.; Wilson, R.E. & Johnson, K.P. 2016: Patterns of cryptic host specificity in duck lice based on molecular data. *Medical and Veterinary Entomology* 30: 200–208.
- Evans, W. 1907: A new louse (*Haematopinus ovillus*, Neum.) from the sheep. *The Annals of Scottish Natural History* 1907: 225–226.
- Evans, W. 1912: Note on Mallophaga from the little auk or rotchie (*Alle alle*); with list of species taken on birds and mammals in the Forth Area. *Proceedings of the Royal Physical Society of Edinburgh* 18(4): 265–276.
- Ewing, H.E. 1924: Ectoparasites of some Polynesian and Malaysian rats of the genus *Rattus*. *Bulletin of the Bernice P. Bishop Museum* 14: 7–11.
- Ewing, H.E. 1926: A revision of the American lice of the genus *Pediculus*, together with a consideration of the significance of their geographical and host distribution. *Proceedings of the United States National Museum* 68(19): 1–30, pls. 1–3.
- Ewing, H.E. 1927: Descriptions of new genera and species of Mallophaga, together with keys to some related genera of Menoponidae and Philopteridae. *Journal of the Washington Academy of Sciences* 17(4): 86–96.
- Ewing, H.E. 1929: *A manual of external parasites*. London: Baillière, Tindall & Cox. xvi + 225 pp.
- Ewing, H.E. 1933: *Neocolpocephalum*, a new name for the mallophagan genus *Ferrisia* Uchida. *The Journal of Parasitology* 20: 65–66.
- Fabricius, J.C. 1775: *Systema Entomologiae, sistens insectorum, classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus*. Flensburgi et Lipsiae: Officina Libraria Kortii. xxxii + 832 pp.
- Fabricius, J.C. 1787: *Mantissa insectorum sistens species nuper detectas adiectis synonymis, observationibus, descriptionibus, emendationibus*. Hafniae: Impensis Christ. gottl. Proft. Volume 2. 382 pp.
- Fabricius, O. 1780: *Fauna Groenlandica systematice sistens animalia Groenlandiae occidentalis hactenus indagata*. Hafniae et Lipsiae: J.G. Rothe. xvi + 452 pp., pl. 1.
- Fahrenholz, H. 1912: Beiträge zur Kenntnis der Anopluren. *Jahresbericht des Niedersächsischen Zoologischen Vereins* 2–4: 1–60, 3 pls.
- Fahrenholz, H. 1919: Zur Nomenklatur einiger Anopluren-Arten. II. *Jahresbericht des Niedersächsischen Zoologischen Vereins zu Hannover* 5–10: 22–27.
- Fahrenholz, H. 1939: Beiträge zur Kenntnis der Anopluren. IV. *Mitteilungen aus dem entomologischen Verein Bremen* 26: 32–47.
- Fedorenko, I.A. 1978: Materials on the Mallophaga fauna in Passeriformes of the Ukraine. Communication IV. Philopterinae (Mallophaga, Ischnocera). Part 2. *Vestnik Zoologii*, Kiev. 1978(2): 55–60 [in Russian with English summary].
- Fedorenko, I.A. 1985: New species of the genus *Philopterus* (Mallophaga) from passeriform birds of the USSR fauna. *Vestnik Zoologii* 6: 11–16 [in Russian, with English summary].
- Ferris, G.F. 1916a: Some generic groups in the mallophagan family Menoponidae. *The Canadian Entomologist* 48: 301–311.
- Ferris, G.F. 1916b: Notes on Anoplura and Mallophaga, from mammals, with descriptions of four new species and a new variety of Anoplura. *Psyche* 23(4): 97–120.
- Ferris, G.F. 1921: Contributions toward a monograph of the sucking lice. Part II. *Stanford University Publications, Biological Sciences* 2(2): 53–133. Stanford University Press: California.

- Ferris, G.F. 1923: Contributions toward a monograph of the sucking lice. Part IV. *Stanford University Publications, Biological Sciences* 2(4): 179–270. Stanford University Press: California.
- Ferris, G.F. 1924: The mallophagan family Menoponidae. *Parasitology* 16(1): 55–66.
- Ferris, G.F. 1932a: New species and other records of Mallophaga from the Marquesas. *Bulletin of the Bernice P. Bishop Museum* 98: 53–72.
- Ferris, G.F. 1932b: Contributions toward a monograph of the sucking lice. Part V. *Stanford University Publications, Biological Sciences* 2(5): 273–413. Stanford University Press: California.
- Ferris, G.F. 1933: Contributions toward a monograph of the sucking lice. Part VI. *Stanford University Publications, Biological Sciences* 2(6): 415–470. Stanford University Press: California.
- Ferris, G.F. 1934: Contributions toward a monograph of the sucking lice. Part VII. *Stanford University Publications, Biological Sciences* 2(7): 473–526. Stanford University Press, California.
- Ferris, G.F. 1935: Contributions toward a monograph of the sucking lice. Part VIII. *Stanford University Publications, Biological Sciences* 2(8): 527–634. Stanford University Press: California.
- Ferris, G.F. 1951: The sucking lice. *Memoirs of the Pacific Coast Entomological Society* 1: 1–320.
- Ford-Robertson, J. de C. & Bull, P.C. 1966: Some parasites of the kiore, *Rattus exulans*, on Little Barrier and Hen Islands, New Zealand. *New Zealand Journal Science* 9: 221–224.
- Forrester, D.J.; Kale II, H.W.; Price, R.D.; Emerson, K.C. & Foster, G.W. 1995: Chewing lice (Mallophaga) from birds in Florida: A listing by host. *Bulletin of the Florida Museum of Natural History* 39(1): 1–43.
- Foster, G.W.; Kinsella, J.M.; Price, R.D.; Mertins, J.W. & Forrester, D.J. 1996: Parasitic helminths and arthropods of greater shearwaters (*Puffinus gravis*) from Florida. *Journal of the Helminthological Society of Washington* 63(1): 83–88.
- Fowler, J.A. & Hodson, D. 1988: The Mallophaga of Leach's petrels *Oceanodroma leucorhoa* from North Rona, Scotland. *Seabird* 11: 47–49.
- Fowler, J.A. & Miller, C.J. 1984: Non-haematophagous ectoparasite populations of procellariiform birds in Shetland, Scotland. *Seabird* 7: 23–30.
- Fowler, J.A.; Miller, C.J. & Cohen, S. 1984: Ectoparasite populations from breeding and wandering storm petrels. *Bird Study* 31(2): 126–130.
- Fowler, J.A. & Price, R.A. 1987: A comparative study of the Ischnoceran Mallophaga of Wilson's Petrel *Oceanites oceanicus* and British Storm Petrel *Hydrobates pelagicus*. *Seabird* 10: 43–49.
- Fowler, J.A. & Shaw, G.J. 1990: The Mallophaga of Manx Shearwaters *Puffinus p. puffinus* from Ynys Enlli, Wales. *Seabird* 12: 14–19.
- Freund, L. 1928: Anoplura Pinnipedium (Robbenläuse). In: Grimpe G. & Wagler, E. (Eds) *Tierwelt der Nord- und Ostsee. Geest und Portig, Leipzig* 4(11): 1–36.
- Furness, R.W. & Palma, R.L. 1992: Phthiraptera of petrels and skuas from Gough Island, South Atlantic Ocean. *Seabird* 14: 33–42.
- Galloway, T.D. 2005: Ectoparasites from native and introduced birds from Christchurch and surrounding areas, New Zealand. *Tuhinga – Records of the Museum of New Zealand Te Papa Tongarewa* 16: 13–20.
- Galloway, T.D. & Lamb, R.J. 2014: Abundance and stability are species traits for four chewing lice (Phthiraptera: Menoponidae, Philopteridae) on feral pigeons, *Columba livia* (Aves: Columbiformes: Columbidae). *The Canadian Entomologist* 146: 444–456.
- Galloway, T.D. & Lamb, R.J. 2015: Seasonal population dynamics of four species of chewing lice (Phthiraptera: Menoponidae, Philopteridae) on feral pigeons (Aves: Columbiformes: Columbidae). *The Canadian Entomologist* 147: 712–722.
- Garfield, B. 2007: *The Meinertzhagen mystery. The life and legend of a colossal fraud*. Washington, D.C.: Potomac Books. xiv + 353 pp.
- George, R.S. 1981: Obituary – Gordon Burnett Thompson, 1919–1979. *Entomologist's Monthly Magazine* 117: 87–88.
- Gervais, F.L.P. 1844: Dicères épizoïques. Pp. 290–361, pls 48–49. In: Walckenaer, C.A. (Ed.) *Histoire Naturelle des Insectes. Aptères*. Tome 3. Paris: Librairie Encyclopédique de Roret. viii + 476 pp.
- Gibson, R.N. & Pilgrim, R.L.C. 1986: Some ectoparasites on rodents in New Zealand. II. Sucking lice (Insecta: Anoplura). *Mauri Ora* 13: 93–102.
- Giebel, C.G.A. 1861a: Die Federlinge der Raubvögel aus Chr. L. Nitzsch's handschriftlichem Nachlass zusammengestellt. *Zeitschrift für die gesammten Naturwissenschaften (Halle)* 17(6): 515–529.
- Giebel, C.G.A. 1861b: Die Haarlinge der Gattungen *Trichodectes* und *Gyropus* nach Chr. L. Nitzsch's Untersuchungen. *Zeitschrift für die gesammten Naturwissenschaften (Halle)* 18(8/9): 81–93, 2 pls.
- Giebel, C.G.A. 1866: Die im zoologischen Museum der Universität Halle aufgestellten Epizoen nebst Beobachtungen über dieselben. *Zeitschrift für die gesammten Naturwissenschaften (Halle)* 28(11/12): 353–397.

- Giebel, C.G.A. 1867: "Über *Lipeurus ferox*", n. sp. Correspondenzblatt. *Zeitschrift für die gesammten Naturwissenschaften* (Halle) 29: 195–196.
- Giebel, C.G.A. 1874: *Insecta epizoa. Die auf Säugetieren und Vögeln schmarotzenden Insecten nach Chr. L. Nitzsch's Nachlass bearbeitet*. Leipzig: Otto Wigand. xvi + 308 pp., 20 pls.
- Giebel, C.G.A. 1876: Diagnoses of some species of Mallophaga collected by the Rev. A. E. Eaton during the late Transit-of-Venus Expedition to Kerguelen's Island. *Annals and Magazine of Natural History* (Series 4) 17(101): 388–389.
- Giglioli, H. 1864: On some parasitical insects from China. *The Quarterly Journal of Microscopical Science, London (N.S.)* 4: 18–26, pl. 1.
- Gill, B.J. 1980: Abundance, feeding, and morphology of passerine birds at Kowhai Bush, Kaikoura, New Zealand. Appendix 1. Ectoparasites collected or noted while handling birds at Kowhai Bush. *New Zealand Journal of Zoology* 7(2): 246.
- Gilruth, J.A. 1908: *Haematopinus* (blood sucking louse) of sheep. *New Zealand Department of Agriculture Annual Report for 1908*. Pp. 194–196, 1 pl.
- Girling, M.A. 1984: Eighteenth century records of human lice (Phthiraptera, Anoplura) and fleas (Siphonaptera, Pulicidae) in the City of London. *Entomologist's Monthly Magazine* 120: 207–210.
- Göllner-Scheiding, U. 1973: Katalog der im Zoologischen Museum Berlin vorhandenen Mallophagentypen. *Lounais-Hämeen Luonto* 46: 29–46.
- González-Acuña, D.; Corvalan, F.; Barrientos, C.; Doussang, D.; Mathieu, C.; Nilsson, L.; Casanueva, M.E. & Palma, R.L. 2011: Community structure of lice (Insecta: Phthiraptera) from two sympatric gull species: kelp gull (*Larus dominicanus*) and Franklin's gull (*Larus pipixcan*) in Talcahuano, Chile. *Neotropical Entomology* 40(3): 300–304.
- Gordon, D.P. (Ed.) 2010: *New Zealand Inventory of Biodiversity*. Volume Two. Kingdom Animalia – Chaetognatha, Ecdysozoa, Ichnofossils. Christchurch: Canterbury University Press. 528 pp.
- Green, E.D. & Turner, M.L. 2003a: Micromorphology of the kiwi feather louse (*Rallicola gracilentus*). *Journal of the South African Veterinary Association* 74(3): 97–98.
- Green, E.D. & Turner, M.L. 2003b: Functional micromorphology of the louse *Docophoroides brevis* from the wandering albatross. *Microscopy Society of Southern Africa – Proceedings* 33: 77.
- Green, E.D. & Turner, M.L. 2004: Functional micromorphology of the elephant seal louse *Lepidophthirus macrorhini*. *Microscopy Society of Southern Africa – Proceedings* 34: 74.
- Green, R.H. & Palma, R.L. 1991: A list of lice (Insecta: Phthiraptera) recorded from Tasmania. *Records of the Queen Victoria Museum, Launceston* 100: 1–43.
- Greenslade, P. 2006: *The invertebrates of Macquarie Island*. Kingston: Australian Antarctic Division. xvi + 326 pp.
- Gressitt, J.L. 1964: Insects of Campbell Island. Summary. *Pacific Insects Monograph* 7: 531–600.
- Gressitt, J.L. 1970: Subantarctic entomology and biogeography. *Pacific Insects Monograph* 23: 295–374.
- Grossi, A.A.; Sharanowski, B.J. & Galloway, T.D. 2014: *Anatoecus* species (Phthiraptera: Philopteridae) from Anseriformes in North America and taxonomic status of *Anatoecus dentatus* and *Anatoecus icterodes*. *The Canadian Entomologist* 146: 598–608.
- Grube, A.W. 1851: Klasse Insecta — Parasitae. Fam. Mallophaga. Pp. 467–497, 516, pls 31–32. In: Middendorff, A.T. von (Ed.) *Reise in den äussersten Norden und Osten Sibiriens während der Jahre 1843–1844*. St Petersburg. *Zoologie* 2(1): 1–516, 32 pls.
- Guimarães, L.R. 1938: Nota sobre um mallophago (*Austrogoniodes bifasciatus* (Piaget) parasita do pinguim. *Revista de Biologia e Higiene* 9(1): 39–46.
- Guimarães, L.R. 1942: Dois novos gêneros de malófagos de psitacídeos exóticos. *Papéis Avulsos do Departamento de Zoologia (São Paulo)* 2(4): 79–95.
- Guimarães, L.R. 1943: Contribuição para o conhecimento dos malofagos das aves da Argentina. *Revista de la Sociedad Entomológica Argentina* 11: 423–439.
- Guimarães, L.R. 1945: Sôbre alguns ectoparasitos de aves e mamíferos do litoral paranaense. *Arquivos do Museu Paranaense* 4(7): 179–190, 2 pls.
- Guimarães, L.R. 1974a: Ischnocera (Mallophaga) infesting parrots (Psittaciformes) I. Genera *Neopsittaconirmus* Conci, 1942, and *Psittaconirmus* Harrison, 1915. *Arquivos de Zoologia* 25(3): 121–201.
- Guimarães, L.R. 1974b: Ischnocera (Mallophaga) infesting parrots (Psittaciformes). II. Genera *Psittoecus* Conci, 1942, and *Forficuloecus* Conci, 1941. *Papéis Avulsos de Zoologia* 28(9): 163–180.
- Guimarães, L.R. 1985: Ischnocera (Mallophaga) infesting parrots (Psittaciformes). V. Four new species of *Forficuloecus* Conci, 1941 (Philopteridae) from the South Pacific. *Papéis Avulsos de Zoologia* 36(5): 41–49.
- Gurlt, E.F. 1843: Über die auf den Haus-Säugetieren und Haus-Vögeln lebenden Schmarotzer – Insekten und Arachniden. *Magazin für die gesammte Thierheilkunde* 9: 1–24, 1 pl.

- Gustafsson, D.R. & Bush, S.E. 2014: Three new species of chewing lice of the genus *Emersoniella* Tendeiro, 1965 (Insecta: Phthiraptera: Ischnocera: Philopteridae) from Papua New Guinean kingfishers and kookaburras (Aves: Coraciiformes: Alcedinidae). *Zootaxa* 3796: 528–544.
- Gustafsson, D.R. & Olsson, U. 2012a: Flyway homogenisation or differentiation? Insights from the phylogeny of the sandpiper (Charadriiformes: Scolopacidae: Calidrinae) wing louse genus *Lunaceps* (Phthiraptera: Ischnocera). *International Journal for Parasitology* 42: 93–102.
- Gustafsson, D.R. & Olsson, U. 2012b: The “Very thankless task”: Revision of *Lunaceps* Clay and Meinertzhagen, 1939 (Insecta: Phthiraptera: Ischnocera: Philopteridae), with descriptions of six new species and one new subspecies. *Zootaxa* 3377: 1–85.
- Haan, M.W. de 1829: Explication des Planches. Pp. 304–312, pls 1–7. In: Lyonet, P. (Ed.) Anatomie de différentes espèces d’insectes. *Mémoires du Muséum d’Histoire Naturelle* 18: 233–312, 7 pls.
- Hackman, W. & Nyholm, E.S. 1968: Notes on the arthropod fauna of Spitsbergen II. Mallophaga from Spitsbergen and Bear Island. *Annales Entomologici Fennici* 34(2): 75–82.
- Haeckel, E. 1896: *Systematische Phylogenie*. 2. Theil. *Systematische Phylogenie der wirbellosen Thiere (Invertebrata)*. Berlin: Verlag von Georg Reimer. 720 pp.
- Hammer, S.; Brown, R.M.; Bugoni, L.; Palma, R.L. & Hughes, J. 2010: On the origin of *Halipeurus heraldicus* on Round Island petrels: cophylogenetic relationships between petrels and their chewing lice. *Molecular Phylogenetics and Evolution* 55: 1111–1120.
- Hänel, C. & Palma, R.L. 2007: The lice of the Tristan da Cunha Archipelago (Insecta: Phthiraptera). *Beiträge zur Entomologie* 57(1): 105–133.
- Harrison, L. 1915a: On a new family and five new genera of Mallophaga. *Parasitology* 7: 383–407, 2 pls.
- Harrison, L. 1915b: Mallophaga from *Apteryx*, and their significance; with a note on the genus *Rallicola*. *Parasitology* 8(1): 88–100.
- Harrison, L. 1916: The genera and species of Mallophaga. *Parasitology* 9(1): 1–156.
- Harrison, L. 1937: Mallophaga and Siphunculata. 2(1): 1–47, pls 1–3. In: Johnston, T.H. (Ed.) *Scientific reports, Australasian Antarctic Expedition 1911–14, series C, Zoology and Botany*. Sydney: Government Printer.
- Harrison, P.; Sallaberry, M.; Gaskin, C.P.; Baird, K.A.; Jaramillo, A.; Metz, S.M.; Pearman, M.; O’Keeffe, M.; Dowdall, J.; Enright, S.; Fahy, K.; Gilligan, J. & Lillie, G. 2013: A new storm-petrel species from Chile. *The Auk* 130(1): 180–191.
- Heath, A.C.G. 1973: The biology and survival of starved cattle and goat biting lice (Mallophaga) at different temperatures and relative humidities. *The New Zealand Entomologist* 5(3–4): 330–334.
- Heath, A.C.G. 1976: Lice infestations. *New Zealand Journal of Agriculture* 132(5): 57.
- Heath, A.C.G. 1978: Seasonality in ectoparasites. *The New Zealand Entomologist* 6(4): 364–365.
- Heath, A.C.G. 1979: Ectoparasites of feral sheep on Campbell Island. *New Zealand Journal of Zoology* 6(1): 141–144.
- Heath, A.C.G. 1983: Parasites of the feral sheep of Pitt Island, Chatham group. *New Zealand Journal of Zoology* 10(4): 365–370.
- Heath, A.C.G. 1994: Ectoparasites of livestock in New Zealand. *New Zealand Journal of Zoology* 21(1): 23–38.
- Heath, A.C.G. 2001: Lice. Pp. 9–11, 3 figs. In: Edwards, S.; Marshall, A.; Cole, D. & Heath, A. (Eds) *Fly and lice: Numbering their days*. Wellington: WoolPro – Operation Clean Fleeces. 52 pp.
- Heath, A.C.G. 2002: Ectoparasites of livestock and companion animals in New Zealand. *New Zealand Veterinary Journal* 50(3) Supplement: 48.
- Heath, A.C.G. 2005: Sheep biting-lice resistance management. Pp.138–142. In: Martin, N.A.; Beresford, R.M. & Harrington, K.C. (Eds) *Pesticide resistance: Prevention and management strategies 2005*. Hastings: The New Zealand Plant Protection Society Inc. vi + 166.
- Heath, A.C.G. 2010: A review of ectoparasites of *Apteryx* spp. (kiwi) in New Zealand, with new host records, and the biology of *Ixodes anatis* (Acari: Ixodidae). *Tuhinga – Records of the Museum of New Zealand Te Papa Tongarewa* 21: 147–159.
- Heath, A.C.G. 2013: The role of ticks, biting flies and lice in the transmission of theileriosis. *VetScript* 26(6): 13–14.
- Heath, A.C.G. & Bishop, D.M. 1988: Evaluation of two ‘pour-on’ insecticides against the sheep-biting louse, *Bovicola ovis* and the sheep ked, *Melophagus ovinus*. *New Zealand Journal of Agricultural Research* 31(1): 9–12.
- Heath, A.C.G.; Bishop, D.M.; Cole, D.J.W. & Pfeffer, A. 1996a: The development of cockle, a sheep pelt defect, in relation to size of infestation and time of exposure to *Bovicola ovis*, the sheep-biting louse. *Veterinary Parasitology* 67: 259–267.
- Heath, A.C.G.; Cole, D.J.W. & Bishop, D.M. 1992a: Some currently available insecticides and their comparative efficacy on louse-infested, long-woolled sheep. *New Zealand Veterinary Journal* 40(3): 101–103.
- Heath, A.C.G.; Cole, D.J.W.; Bishop, D.M. & Cooper, S.M. 1996b: The comparative ability of some lousicides to reduce cockle in sheep pelts. *New Zealand Veterinary Journal* 44(4): 135–137.

- Heath, A.C.G.; Cooper, S.M.; Cole, D.J.W. & Bishop, D.M. 1995a: Evidence for the role of the sheep biting-louse *Bovicola ovis* in producing cockle, a sheep pelt defect. *Veterinary Parasitology* 59: 53–58.
- Heath, A.C.G.; Lampkin, N. & Jowett, J.H. 1995b: Evaluation of non-conventional treatments for control of the biting louse (*Bovicola ovis*) on sheep. *Medical and Veterinary Entomology* 9(4): 407–412.
- Heath, A.C.G. & Levot, G.W. 2015: Parasiticide resistance in flies, lice and ticks in New Zealand and Australia: mechanisms, prevalence and prevention. *New Zealand Veterinary Journal* 63(4): 199–210.
- Heath, A.C.G. & Millar, E.S. 1970: Recent insecticides: their efficacy as plunge dips against the biting louse, *Damalinia ovis*, and the ked, *Melophagus ovinus*, on sheep. *New Zealand Veterinary Journal* 18(10): 211–213.
- Heath, A.C.G.; Millthorpe, A.P. & Eves, N. 1971: Pigeon mites and human infestation. *The New Zealand Entomologist* 5(1): 90–92.
- Heath, A.C.G.; Nottingham, R.M.; Bishop, D.M. & Cole, D.J.W. 1992b: An evaluation of two cypermethrin-based pour-on formulations on sheep infected with the biting louse, *Bovicola ovis*. *New Zealand Veterinary Journal* 40(3): 104–106.
- Heath, A.C.G.; Pfeffer, A.T. & Morrison, L. 2004: Association of sheep body louse (*Bovicola ovis*) infestation with weaning weight, birth rank and cockle in lambs. *Wool Technology and Sheep Breeding* 52(1): 8–18.
- Helson, G.A.H. 1956: Some arthropods affecting man and livestock in New Zealand. *The New Zealand Veterinary Journal* 4: 11–18.
- Helson, G.A.H. 1970: Insect pests. Cattle lice. *New Zealand Journal of Agriculture* 120(5): 81.
- Hemming, F. 1958: Opinion 104. Pp. 49–55. In: *Official list of generic names in zoology. First installment: Names 1–1274*. London: International Commission on Zoological Nomenclature. xxxviii + 200 pp.
- Hill, W.W. & Tuff, D.W. 1978: A review of the Mallophaga parasitizing the Columbiformes of North America north of Mexico. *Journal of the Kansas Entomological Society* 51(2): 307–327.
- Hinojos, J.G. & Canaris, A.G. 1988: Metazoan parasites of *Himantopus mexicanus* Müller (Aves) from southwestern Texas, with a checklist of helminth parasites from North America. *The Journal of Parasitology* 74(2): 326–331.
- Holdaway, R.N.; Worthy, T.H. & Tennyson, A.J.D. 2001: A working list of breeding bird species of the New Zealand region at first human contact. *New Zealand Journal of Zoology* 28: 119–187.
- Holloway, B.A. 1956: A new species of *Rallicola* (Insecta: Mallophaga) from *Notornis mantelli* Owen. *Records of the Dominion Museum* 2(3): 113–119.
- Hopkins, G.H.E. 1940: Stray notes on Mallophaga.— II. 7. Notes on some Mallophaga of the coot. *Annals and Magazine of Natural History* (Series 11) 5: 421–426.
- Hopkins, G.H.E. 1941: Stray notes on Mallophaga.— III. 11. The identity of two species of *Trichodectes*, s. l. *Annals and Magazine of Natural History* (Series 11) 7: 36–40.
- Hopkins, G.H.E. 1942a: Notes on Trichodectidae (Mallophaga). *Revista Brasileira de Biologia* 2(4), 439–453.
- Hopkins, G.H.E. 1942b: The Mallophaga as an aid to the classification of birds. *Ibis* 84: 94–106.
- Hopkins, G.H.E. 1949a: Stray notes on Mallophaga— IX. 56. The various identities of *Docophorus auratus*. *Annals and Magazine of Natural History* (Series 12) 2: 29–32, 1 pl.
- Hopkins, G.H.E. 1949b: The host-associations of the lice of mammals. *Proceedings of the Zoological Society of London* 119(2): 387–604.
- Hopkins, G.H.E. 1949c: Stray notes on Mallophaga— IX. 72. A new name for *Nirmus punctulatus* Giebel, 1874. *Annals and Magazine of Natural History* (Series 12) 2: 50–51.
- Hopkins, G.H.E. 1949d: Stray notes on Mallophaga— IX. 74. A new species of *Quadriceps* from a tern. *Annals and Magazine of Natural History* (Series 12) 2: 51–52, 1 pl.
- Hopkins, G.H.E. 1950: Stray notes on Mallophaga— X. 80. The type species of *Nosopios* Eichler. *Annals and Magazine of Natural History* (Series 12) 3: 230–242.
- Hopkins, G.H.E. 1951a: Stray notes on Mallophaga— XI. 91. The host-distribution of *Quadriceps houri* Hopkins. *Annals and Magazine of Natural History* (Series 12) 4: 373.
- Hopkins, G.H.E. 1951b: Stray notes on Mallophaga— XI. 92. The host of *Saemundssonina uppalensis* (Rudow), 1870. *Annals and Magazine of Natural History* (Series 12) 4: 374.
- Hopkins, G.H.E. 1960: Notes on some Mallophaga from mammals. *Bulletin of the British Museum (Natural History). Entomology* 10(2): 77–95, 2 pls.
- Hopkins, G.H.E. & Clay, T. 1952: *A check list of the genera & species of Mallophaga*. London: Trustees of the British Museum. 362 pp.
- Hopkins, G.H.E. & Clay, T. 1953: Additions and corrections to the check list of Mallophaga. *Annals and Magazine of Natural History* (Series. 12) 6: 434–448.
- Hopkins, G.H.E. & Clay, T. 1955: Additions and corrections to the check list of Mallophaga. *Annals and Magazine of Natural History* (Series. 12) 8: 177–190.

- Horning, D.S.; Palma, R.L. & Pilgrim, R.L.C. 1980: The lice (Insecta, Phthiraptera) from the Snares Islands, New Zealand. *National Museum of New Zealand Miscellaneous Series* 3: 1–17.
- Hughes, J.M. 1984a: Distribution of Mallophaga on the Australian magpie (*Gymnorhina tibicen* Latham) (Family: Cracticidae). *Australian Journal of Zoology* 21(5): 459–466.
- Hughes, J.M. 1984b: Morphometric variation in the Mallophaga of the Australian magpie (*Gymnorhina tibicen* Latham). *Australian Journal of Zoology* 21(5): 467–477.
- Hughes, J.; Kennedy, M.; Johnson, K.P.; Palma, R.L. & Page, R.D.M. 2007: Multiple cophylogenetic analyses reveal frequent cospeciation between peleciform birds and *Pectinopygus* lice. *Systematic Biology* 56(2): 232–251.
- Humphreys, P.N. 1975: Wet-feather associated with *Holomenopon leucoanthum* in a duck. *The Veterinary Record* 97: 96–97.
- Hunter, J.E. & Colwell, M.A. 1994: Phthiraptera infestation of five shorebird species. *Wilson Bulletin* 106(2): 400–403.
- Hutton, F.W. 1904: *Index Faunae Novae Zealandiae*. London: Philosophical Institute of Canterbury, New Zealand. viii + 372 pp.
- Imber, M.J. & Tennyson, A.J.D. 2001: A new petrel species (Procellariidae) from the south-west Pacific. *Emu* 101: 123–127.
- Ineson, M.J. 1954: A comparison of the parasites of wild and domestic pigs in New Zealand. *Transactions of the Royal Society of New Zealand* 82(2): 579–609.
- International Commission on Zoological Nomenclature [I.C.Z.N.] 1963: Opinion no. 656 *Harrisoniella* Bedford, 1929 (Insecta: Mallophaga): designation of a type species under the plenary powers. *Bulletin of Zoological Nomenclature* 20(3): 178–180.
- International Commission on Zoological Nomenclature [I.C.Z.N.] 1976: Opinion 1050. *Pediculus eurysternus* Burmeister, 1838 (Insecta, Anoplura): suppressed under the plenary powers. *Bulletin of Zoological Nomenclature* 32(4): 238–239.
- International Commission on Zoological Nomenclature [I.C.Z.N.] 2014: Opinion 2333. *Mémoires pour servir à l'histoire des insectes* by De Geer (1752–1778) and the additional volume by Retzius (1783): ruled to be binominal and available. *Bulletin of Zoological Nomenclature* 71(1): 53–59.
- Jałoszyński, P.; Gustafsson, D.R.; Wanat, A. & Wanat, M. 2014: Type specimens of Phthiraptera in the collection of Jadwiga Złotorzycka preserved in the Museum of Natural History, University of Wrocław. *Genus* 25(4): 645–661.
- Jellison, W.L. 1935: A new species of *Bovicola* (Mallophaga). *The Journal of Parasitology* 21: 410–411.
- Jensen, K. & Palma, R.L. 2005: Insecta (insects). Pp. 226–230, 491–492. In: Rohde, K. (Ed.) *Marine Parasitology*. Collingwood, Victoria: CSIRO Publishing. 592 pp.
- Jiménez-González, A.; Rodríguez-Caabeiro, F. & Martín-Mateo, M.P. 1980: Estudio taxonómico de malófagos parásitos de *Turdus philomelos* Brehm. *Nouvelle Revue d'Entomologie* 10(2): 205–218.
- Johnson, K.P. & Clayton, D.H. 2003: *The biology, ecology and evolution of chewing lice*. Pp. 449–476. In: Price, R.D.; Hellenthal, R.A.; Palma, R.L.; Johnson, K.P. & Clayton, D.H. *The chewing lice: world checklist and biological overview*. Illinois Natural History Survey Special Publication 24. x + 501 pp.
- Johnson, K.P. & Clayton, D.H. 2004: Untangling coevolutionary history. *Systematic Biology* 53(1): 92–94.
- Johnson, P.T. 1964: The hoplopleurid lice of the Indo-Malayan Subregion (Anoplura: Hoplopleuridae). *Miscellaneous Publications of the Entomological Society of America* 4(3): 67–102.
- Johnson, P.T. 1972: Some Anoplura of the Oriental Region. A study of *Hoplopleura pacifica* Ewing and allies. *Journal of Medical Entomology* 9(3): 219–227.
- Johnston, T.H. & Harrison, L. 1911: Notes on some Mallophagan generic names. *Proceedings of the Linnean Society of New South Wales* 36(2): 321–328.
- Johnston, T.H. & Harrison, L. 1912: A collection of Mallophaga from the Kermadecs. *Transactions of the New Zealand Institute* 44: 363–373.
- Kéler, S. von 1936: Über einige Mallophagen aus Rossitten. *Arbeiten über Morphologische und Taxonomische Entomologie aus Berlin-Dahlem* 3(4): 256–264.
- Kéler, S. von 1937: Ein Beitrag zur Kenntnis der Mallophagen des Rebhuhnes (*Perdix perdix* Linn.). *Arbeiten über Morphologische und Taxonomische Entomologie aus Berlin-Dahlem* 4(2): 126–135.
- Kéler, S. von 1938a: Baustoffe zu einer Monographie der Mallophagen. I. Teil: Überfamilie Trichodectoidea. *Nova Acta Leopoldina Abhandlungen der Kaiserlich Leopoldinisch-Carolinisch Deutschen Akademie der Naturforscher (Neue Folge)* 5(32): 393–467.
- Kéler, S. von 1938b: Über eine neue Gattung von Mallophagen, *Eidmanniella* n. gen. *Annales Musei Zoologici Polonici* 13: 81–87.
- Kéler, S. von 1939: Baustoffe zu einer Monographie der Mallophagen. II. Teil: Überfamilie der Nirmoidea (1). *Nova Acta Leopoldina Abhandlungen der Kaiserlich Leopoldinisch-Carolinisch Deutschen Akademie der Naturforscher (Neue Folge)* 8(51): 1–254, 4 pls.
- Kéler, S. von 1942: Ein Beitrag zur Kenntnis der Mallophagen. *Arbeiten über Morphologische und Taxonomische Entomologie aus Berlin-Dahlem* 9(2): 69–85.

- Kéler, S. von 1952: On some Mallophaga of sea-birds from the Tristan da Cunha Group and the Dyer Island. *Journal of the Entomological Society of Southern Africa* 15(2): 204–238.
- Kéler, S. von 1954: The Mallophaga from *Eudypetes chrysolophus* (Brand) and *E. cristatus* (Miller). *Proceedings of the Royal Entomological Society of London* (B) 23(3–4): 49–59.
- Kéler, S. von 1956: Die Mallophagen von Sturmvögeln und Ruderfüßern. I. *Harrisoniella* Bedford und *Perineus* Thompson (Mallophaga). *Beiträge zur Entomologie* 6(5/6): 521–534.
- Kéler, S. von 1957a: Die Mallophagen von Sturmvögeln und Ruderfüßern. I. *Harrisoniella* Bedford und *Perineus* Thompson (Mallophaga) (1. Fortsetzung). *Beiträge zur Entomologie* 7(3/4): 281–297.
- Kéler, S. von 1957b: Die Mallophagen von Sturmvögeln und Ruderfüßern. I. *Harrisoniella* Bedford und *Perineus* Thompson (Mallophaga) (2. Fortsetzung). *Beiträge zur Entomologie* 7(5/6): 493–527.
- Kéler, S. von 1957c. Über die Deszendenz und die differenzierung der Mallophagen. *Zeitschrift für Parasitenkunde* 18: 55–160.
- Kéler, S. von 1958a: Die Mallophagen von Sturmvögeln und Ruderfüßern. I. *Harrisoniella* Bedford und *Perineus* Thompson (Mallophaga) (Nachtrag). *Beiträge zur Entomologie* 8(3/4): 378–384.
- Kéler, S. von 1958b: The genera *Oxylipeurus* Mjöberg and *Splendoroffula* Clay and Meinertzhagen (Mallophaga). *Deutsche Entomologische Zeitschrift, Neue Folge* 5(3/4): 299–362.
- Kéler, S. von 1960a: Über die dualistische differenzierung der Gattung *Anatoecus* Cummings (Mallophaga). *Zeitschrift für Parasitenkunde* 20: 207–316.
- Kéler, S. von 1960b: Über die Nymphe von *Nesiotinus demersus* (Kellogg) (Mallophaga). *Deutsche Entomologische Zeitschrift, Neue Folge* 7(1/2): 177–184.
- Kéler, S. von 1971: A revision of the Australasian Boopiidae (Insecta: Phthiraptera), with notes on the Trimenoponidae. *Australian Journal of Zoology, Supplement* 6: 1–126.
- Kellogg, V.L. 1896a: New Mallophaga, I,— with special reference to a collection made from maritime birds of the Bay of Monterey, California. *Proceedings of the California Academy of Sciences* (Series 2) 6: 31–168, 14 pls.
- Kellogg, V.L. 1896b: New Mallophaga, II,— from land birds; together with an account of the Mallophagous mouth-parts. *Proceedings of the California Academy of Sciences* (Series 2) 6: 431–548, 14 pls.
- Kellogg, V.L. 1903: Two new genera of Mallophaga. *Biological Bulletin Wood's Hole, Massachusetts* 5(2): 85–91.
- Kellogg, V.L. 1906: A second collection of Mallophaga from birds of the Galapagos and Revillagigedo Islands and neighboring waters. *Transactions of the American Entomological Society* 32: 315–324.
- Kellogg, V.L. 1907: The mallophagan parasites of the kea. *Psyche* 14(6): 122–123.
- Kellogg, V.L. 1908: Mallophaga. In: Wytzman, P. (Ed.) *Genera Insectorum*. Brussels. 66: 1–87, 3 pls.
- Kellogg, V.L. 1914: Mallophaga from birds of the South Atlantic. *Science Bulletin, Brooklyn Institute Museum* 2(4): 80–89, 1 pl.
- Kellogg, V.L. & Chapman, B.L. 1899: New Mallophaga, III. Mallophaga from birds of California. *Occasional Papers of the California Academy of Sciences* 6: 53–143, 5 pls.
- Kellogg, V.L. & Chapman, B.L. 1902: Mallophaga from birds of the Hawaiian Islands. *Journal of the New York Entomological Society* 10: 155–169, 3 pls.
- Kellogg, V.L. & Kuwana, S.I. 1901: Mallophaga from Alaskan birds. *Proceedings of the Academy of Natural Sciences of Philadelphia 1900*: 151–159, 1 pl.
- Kellogg, V.L. & Kuwana, S.I. 1902: Papers from the Hopkins Stanford Galapagos Expedition, 1898–1899. X. Entomological results (8). Mallophaga from birds. *Proceedings of the Washington. Academy of Sciences* 4: 457–499, 4 pls.
- Kenward, H. 1999: Pubic lice (*Phthirus pubis* L.) were present in Roman and medieval Britain. *Antiquity* 73(282): 911–915.
- Kettle, P.R. 1972: Pour-on insecticides for the control of *Linognathus vituli*. *New Zealand Veterinary Journal* 20(9): 167.
- Kettle, P.R. 1974a: The influence of cattle lice (*Damalinea bovis* and *Linognathus vituli*) on weight gain in beef animals. *New Zealand Veterinary Journal* 22(1): 10–11.
- Kettle, P.R. 1974b: *Aquanirmus australis* n. sp. (Mallophaga: Philopteridae) from the New Zealand dabchick, *Podiceps rufopectus*. *New Zealand Journal of Zoology* 1(3): 337–340.
- Kettle, P.R. 1980: *Cuculicola kui* n. sp. (Phthiraptera: Philopteridae) from the New Zealand shining cuckoo. *New Zealand Journal of Zoology* 7(1): 89–90.
- Kettle, P.R. 1983: The seasonal incidence of parasitism by Phthiraptera on starlings (*Sturnus vulgaris*) in England. *New Zealand Entomologist* 7(4): 403–408.
- Kettle, P.R. & Lukies, J.M. 1974: Pour-on insecticides for the control of cattle lice (*Linognathus vituli* and *Damalinea bovis*). *New Zealand Veterinary Journal* 22(5): 76–77.
- Kettle, P.R. 1979: The efficacy of some pour-on insecticides for the control of long-nosed sucking lice (*Linognathus vituli*) on cattle. *New Zealand Veterinary Journal* 27(4): 78–79.

- Kettle, P.R. 1982a: Effect of sheep lice (*Damalinia ovis*) on wool colour. *New Zealand Journal of Experimental Agriculture* 10(1): 15–17.
- Kettle, P.R. 1982b: Long-term effect of sheep body lice (*Damalinia ovis*) on body weight and wool production. *New Zealand Journal of Agriculture Research* 25(4): 531–534.
- Kettle, P.R. 1984: Recovery of sheep lice (*Damalinia ovis*) from baled wool: a technique enabling nation-wide surveillance of louse ridden flocks. *New Zealand Journal of Experimental Agriculture* 12(1): 39–42.
- Kettle, P.R. & Pearce, D.M. 1974: Effect of the sheep body louse (*Damalinia ovis*) on host weight gain and fleece value. *New Zealand Journal of Experimental Agriculture* 2: 219–221.
- Kettle, P.R. & Watson, A.J. 1981: The efficacy of phosmet as a "pour-on" for the control of long-nosed sucking lice (*Linognathus vituli*) on cattle. *New Zealand Veterinary Journal* 29(10): 187–188.
- Kettle, P.R.; Watson, A.J. & White, D.A. 1983: Evaluation of a deltamethrin formulation as a back line treatment of sheep for the control of the sheep body louse (*Damalinia ovis*). *New Zealand Journal of Experimental Agriculture* 11: 321–324.
- Kim, K.C. 1987: Chapter 23. Order Anoplura. Pp. 224–245. In: Stehr, F.W. (Ed.) *Immature insects*. Volume 1. Dubuque, Iowa: Kendall / Hunt Publishing Company. xiv + 754 pp.
- Kim, K.C. 1988: Chapter 7. Evolutionary parallelism in Anoplura and eutherian mammals. Pp. 91–114. In: Service, M.W. (Ed.) *Biosystematics of haematophagous insects*. Oxford: Systematics Association, Clarendon Press. xii + 363 pp.
- Kim, K.C.; Pratt, H.D. & Stojanovich, C.J. 1986: *The sucking lice of North America — An illustrated manual for identification*. University Park & London: The Pennsylvania State University Press. xii + 241 pp.
- Kim, K.C.; Repenning, C.A. & Morejohn, G.V. 1975: Specific antiquity of the sucking lice and evolution of otariid seals. *Rapport et Procès verbaux des Réunions du conseil permanent International pour l'Exploration de la Mer* 169: 544–549.
- Kim, K.C. & Weisser, C.F. 1973: *Haematopinus eurysternus* (Denny, 1842) (Haematopinidae, Anoplura, Insecta): proposed validation under the Plenary Powers. *Bulletin of Zoological Nomenclature* 30(1): 42–46.
- Kim, K.C. & Weisser, C.F. 1974: Taxonomy of *Solenopotes* Enderlein, 1904, with redescription of *Linognathus panamensis* Ewing (Linognathidae: Anoplura). *Parasitology* 69: 107–135.
- King, C.M. (Ed.) 1990: *The handbook of New Zealand mammals*. Auckland: Oxford University Press. xii + 600 pp.
- King, C.M. 2005: *The handbook of New Zealand mammals*. Second Edition. South Melbourne: Oxford University Press. xx + 610 pp.
- Kirk, T.W. 1900: Appendix VII. *Eighth Report of the New Zealand Department of Agriculture*. – Division of biology and pomology. Pp. 276–326.
- Klockenhoff, H.F. 1984: A redescription of *Myrsidea serini* (Mallophaga: Menoponidae), a parasite from passerine birds. *New Zealand Journal of Zoology* 11(1): 17–22.
- Koh, L.P.; Dunn, R.R.; Sodhi, N.S.; Colwell, R.K.; Proctor, H.C. & Smith, V.S. 2004: Species coextinctions and the biodiversity crisis. *Science* 305: 1632–1634.
- Kopociński, B.; Lonc, E. & Modrzejewska, M. 1998: Fitting a bivariate negative binomial model to the distribution of bird lice (Phthiraptera, Mallophaga) parasitizing the pheasant (*Phasianus colchicus* L.). *Acta Parasitologica* 43(2): 81–85.
- Kraus, S.J. & Glassman, L.H. 1976: The crab louse – Review of physiology and study of anatomy as seen by the scanning electron microscope. *Journal of the American Venereal Disease Association* 2(4): 12–18.
- Lakshminarayana, K.V. 1970: Mallophaga Indica IV. Trends in evolution in the *Laemobothrion*-complex (Phthiraptera: Mallophaga), with description of a new genus. *Oriental Insects* 4(2): 131–142.
- Lakshminarayana, K.V. 1977: Notes on the genus *Pseudomenopon* with remarks on host relationships. *Angewandte Parasitologie* 18(3): 152–162.
- Lakshminarayana, K.V. 1979: A synoptic list of Mallophaga sens. lat. (Phthiraptera: Insecta) from India and adjacent countries together with host and regional indices. *Records of the Zoological Survey of India* 75: 39–201.
- Latreille, P.A. 1804: Ordre Second. Parasites; parasita. Pp. 83–112, pl. 68. In: *Histoire naturelle, générale et particulière, des Crustacés et des Insectes*. Paris: F. Dufart. Volume 8. 411 pp., 7 pls.
- Leach, W.E. 1815a: Entomology. Pp. 57–172. In: Brewster, D. (Ed.) *The Edinburgh Encyclopaedia*. Volume 9. Edinburgh: William Blackwood & Sons.
- Leach, W.E. 1815b: Entomology. *Encyclopaedia Britannica, Supplement 1*: 24.
- Leach, W.E. 1817: On the families, stirpes, and genera of the Order Anoplura. *The Zoological Miscellany* 3: 64–67, 1 pl.
- Ledger, J.A. 1980: The arthropod parasites of vertebrates in Africa south of the Sahara. IV. Phthiraptera (Insecta). *Publications of the South African Institute for Medical Research* 56: 1–327.
- Leo, N.P. & Barker, S.C. 2005: Unravelling the evolution of the head lice and body lice of humans. *Parasitology Research* 98(1): 44–47.
- Leo, N.P.; Hughes, J.M.; Yang, X.; Poudel, S.K.; Brogdon, W.G. & Barker, S.C. 2005: The head and body lice of humans are genetically distinct (Insecta: Phthiraptera, Pediculidae): evidence from double infestations. *Heredity* 95(1): 34–40.



- Leonardi, M.S.; Bobinac, M. & Negrete, J. 2016: Redescription of *Antarctophthirus lobodontis* (Anoplura: Echinophthiriidae) from the crabeater seal and identification key for Antarctic lice. *Polar Biology* 39: 671–676.
- Leonardi, M.S.; Crespo, E.A.; Raga, J.A. & Aznar, F. 2011: Insectos oceánicos: los piojos buceadores. *Ciencia Hoy* 21(123): 58–64.
- Leonardi, M.S.; Crespo, E.A.; Raga, J.A. & Fernández, M. 2012a: Scanning electron microscopy of *Antarctophthirus microchir* (Phthiraptera: Anoplura: Echinophthiriidae): Studying morphological adaptations to aquatic life. *Micron* 43: 929–936.
- Leonardi, M.S.; Crespo, E.A.; Raga, J.A. & Fernández, M. 2009: Redescription of *Antarctophthirus microchir* (Anoplura: Echinophthiriidae) from South American sea lion, *Otaria flavescens*, from Patagonia, Argentina. *The Journal of Parasitology* 95: 1086–1092.
- Leonardi, M.S.; Crespo, E.A.; Vales, D.G.; Feijoo, M.; Raga, J.A. & Aznar, F.J. 2012b: Life begins when the sea lion is ashore: habitat use by a louse of a diving mammal host. *Bulletin of Entomological Research* 102: 444–452.
- Leonardi, M.S. & Lazzari, C.R. 2014: Uncovering deep mysteries: The underwater life of an amphibious louse. *Journal of Insect Physiology* 71: 164–169.
- Leonardi, M.S. & Palma, R.L. 2013: Review of the systematics, biology and ecology of lice from pinnipeds and river otters (Insecta: Phthiraptera: Anoplura: Echinophthiriidae). *Zootaxa* 3630(3): 445–466.
- Leonardi, M.S.; Poljak, S.; Carlini, P.; Galliari, J.; Bobinac, M.; Santos, M.; Márquez, M.E. & Negrete, J. 2014: *Antarctophthirus carlinii* (Anoplura: Echinophthiriidae), a new species from the Weddell seal *Leptonychotes weddelli*. *Parasitology Research* 113(11): 3947–3951.
- Le Souëf, S.A. 1902a: Descriptions of some new Mallophaga from marsupials. *The Victorian Naturalist* 19: 50–51, 1 pl.
- Le Souëf, S.A. 1902b: Descriptions of some new species of Mallophaga from Australian birds.— Part IV. *The Victorian Naturalist* 19: 90–91.
- Le Souëf, S.A. & Bullen, H. 1902a: Descriptions of some Mallophaga from Australian birds. *The Victorian Naturalist* 18: 155–158, 1 pl.
- Le Souëf, S.A. & Bullen, H. 1902b: Description of a mallophagous parasite from the kangaroo. *The Victorian Naturalist* 18: 159.
- Light, J.E.; Allen, J.M.; Long, L.M.; Carter, T.E.; Barrow, L.; Suren, G.; Raoult, D. & Reed D.L. 2008a: Geographic distributions and origins of human head lice (*Pediculus humanus capitis*) based on mitochondrial data. *The Journal of Parasitology* 94(6): 1275–1281.
- Light, J.E.; Touns, M.A. & Reed, D.L. 2008b: What's in a name: the taxonomic status of human head and body lice. *Molecular Phylogenetics and Evolution*. 47(3): 1203–1216.
- Linnaeus, C. von 1758: *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Tenth Edition. Volume I. Holmiae: Laurentii Salvii. iv + 824 pp.
- Lowry, J.K.; Horning, D.S.; Poore, G.C.B. & Ricker, R.W. 1978: *The Australian Museum Macquarie Island Expedition, Summer 1977–1978*. Sydney: The Australian Museum Trust. 152 pp.
- Lupidio, M.C. 1980: *Phthirus pubis* en animales en Tandil, Provincia de Buenos Aires, Argentina. *Revista AVEPA – Asociación Veterinaria Española de Especialistas en Pequeños Animales* 2(9): 30.
- Lyal, C.H.C. 1985a: A cladistic analysis and classification of trichodectid mammal lice (Phthiraptera: Ischnocera). *Bulletin of the British Museum (Natural History)* 51(3): 187–346.
- Lyal, C.H.C. 1985b: Phylogeny and classification of the Psocodea, with particular reference to the lice (Psocodea: Phthiraptera). *Systematic Entomology* 10: 145–165.
- MacLeod, C.J.; Paterson, A.M.; Tompkins, D.M. & Duncan, R.P. 2010: Parasites lost – do invaders miss the boat or drown on arrival? *Ecology Letters* 13(4): 516–527.
- Marconcini, A. & Macchioni, G. 1975 [1974]: Contributo alla conoscenza des mallofagi del fagiano (*Phasianus colchicus* L.) in Toscana. *Annali della Facoltà di Medicina Veterinaria dell'Università di Pisa* 27: 101–112.
- Marples, B.J. 1942: A study of the little owl, *Athene noctua*, in New Zealand. *Transactions and Proceedings of the Royal Society of New Zealand* 72: 237–252.
- Marris, J.W.M. 2000: The beetle (Coleoptera) fauna of the Antipodes Islands, with comments on the impact of mice; and an annotated checklist of the insect and arachnid fauna. *Journal of the Royal Society of New Zealand* 30(2): 169–195.
- Marshall, A.G. 1981: *The ecology of ectoparasitic insects*. London: Academic Press. xvi + 459 pp.
- Martens, J.M. 1974: Zur Taxonomie der Gattung *Saemundssonina* Timmermann (Mallophaga: Ischnocera) auf Schnepfen (Scolopacinae) und Strandläufern (Eroliinae). *Mitteilungen aus den Hamburgischen Zoologischen Museum und Institut* 70: 119–163.
- Martens, J.M. 1980: Die Regenpfeifer-Parasiten *Quadriceps dominella* und *Quadriceps novaeseelandiae* von Neuseeland. *Senckenbergiana Biologica* 60(5/6): 349–354.

- Martens, J.M. & Palma, R.L. 1981: Species distribution of genus *Quadriceps* (Mallophaga: Philopteridae) on New Zealand endemic plovers. *New Zealand Journal of Zoology* 8: 83–85.
- Martin, M. 1934: Life history of the pigeon louse (*Columbicola columbae* (Linnaeus)). *The Canadian Entomologist* 66(1): 6–16.
- Martín-Mateo, M.P. 1990: Contribución al conocimiento de los malófagos parásitos de aves en la Isla de Tenerife (Mallophaga: Insecta). *Vieraeva* 19: 175–184.
- Martín-Mateo, M.P. 1992a: Malófagos de aves marinas. Especies parásitas de aves Pelecaniformes. (Insecta: Mallophaga). *García de Orta (Série Zoologia)* 17(1–2): 37–51.
- Martín-Mateo, M.P. 1992b: Estudio de las especies de malófagos (Phthiraptera) parásitas de Glareolidae (Aves) y consideraciones sobre el “status” taxonómico del género *Glareolites* Eichler. *Annales de la Société Entomologique de France (Nouvelle Série)* 28(4): 409–420.
- Martín-Mateo, M.P. 1994: Phthiraptera from *Platalea leucorodia* L. (Aves: Ciconiiformes: Threskiornithidae) in Spain. *Research and Reviews in Parasitology* 54(2): 109–115.
- Martín-Mateo, M.P. 1996: Species of the genus *Puffinoecus* Eichler, 1949 (Mallophaga: Philopteridae) parasites on shearwaters (Aves: Procellariidae). *Research and Reviews in Parasitology* 56(1): 49–62.
- Martín-Mateo, M.P. 2002: *Fauna Ibérica*. Volume 20. *Mallophaga, Amblycera*. Madrid: Museo Nacional de Ciencias Naturales & Consejo Superior de Investigaciones Científicas. 187 pp.
- Martín-Mateo, M.P. 2009: *Fauna Ibérica*. Volume 32. *Phthiraptera, Ischnocera*. Madrid: Museo Nacional de Ciencias Naturales & Consejo Superior de Investigaciones Científicas. 363 pp.
- Martinů, J.; Sychra, O.; Literák, I.; Čapek, M.; Gustafsson, D.L. & Štefka, J. 2015: Host generalists and specialists emerging side by side: an analysis of evolutionary patterns in the cosmopolitan chewing louse genus *Menacanthus*. *International Journal of Parasitology* 45(1): 63–73.
- Mason, G.E. 1921: Observations on certain external parasites found upon the New Zealand huia (*Neomorpha acutirostris* Gould) and not previously recorded. *Transactions of the New Zealand Institute* 53: 357–359.
- Maturano, R. & Daemon, E. 2014: Reproduction, development and habits of the large turkey louse *Chelopistes meleagridis* (Phthiraptera: Ischnocera) under laboratory conditions. *Brazilian Journal of Biology* 74(3): 712–719.
- Maunder, J.W. 1983: The appreciation of lice. *Proceedings of the Royal Institution of Great Britain* 55: 1–31.
- McGregor, E.A. 1912: A new mallophagan. *Entomological News* 23(7): 305–306.
- McGregor, E.A. 1917: Eight new Mallophaga of the genus *Lipeurus* from North American birds. *Psyche* 24(4): 105–117, 3 pls.
- McIntosh, R. & Murray, M.D. 2007: Louse infestations of the Australian sea lion *Neophoca cinerea*. *Australian Mammalogy* 29: 103–106.
- McKenna, P.B. 2001: Register of new host-parasite records. Louse infestation on a llama. *Surveillance* 28(4): 15–16.
- McKenna, P.B. 2003: Register of new host-parasite records. Louse infestation on an alpaca. *Surveillance* 30(1): 12–13.
- Mégnin, P. 1884: Le *Trichodectes lipeuroides* n. sp. (du *Cervus mexicanus*). *Le Naturaliste, Paris* 6: 494–495.
- Mehlhorn, B.; Mehlhorn, H. & Plötz, J. 2002: Light and scanning electron microscopical study on *Antarctophthirus ogmorhini* lice from the Antarctic seal *Leptonychotes weddellii*. *Parasitology Research* 88(7): 651–660.
- Meleney, W.P. & Kim, K.C. 1974: A comparative study of cattle-infesting *Haematopinus*, with redescription of *H. quadripertusus* Fahrenholz, 1916 (Anoplura: Haematopinidae). *The Journal of Parasitology* 60(3): 507–522.
- Melville, D.S. 1985: Long-tailed skuas *Stercorarius longicaudus* in New Zealand. *Notornis* 32: 51–73.
- Mester, H. 1971: Federlinge auf Limikolen-Gelegen. *Journal für Ornithologie* 112(2): 109–130.
- Mey, E. 1982: Mongolische Mallophagen I. Ergebnisse der mongolischen Gemeinschaftsreise von Ornithologen aus der DDR 1979. IX, zugleich Ergebnisse der Mongolisch-Deutschen Biologischen Expedition seit 1962, Nr. 107. *Mitteilungen aus dem Zoologischen Museum in Berlin* 58(1): 155–195, 5 pls.
- Mey, E. 1986) [1985]: Mallophagen aus Thüringen II. *Veröffentlichungen des Naturkundemuseum Erfurt* 4: 32–43.
- Mey, E. 1989: Ein neuer Sturmvogelfederling (Insecta, Phthiraptera) aus dem Südatlantik. *Rudolstädter Naturhistorische Schriften* 2: 51–56.
- Mey, E. 1990: Eine neue ausgestorbene Vogel-Ischnozere von Neuseeland, *Huiacola extinctus* (Insecta, Phthiraptera). *Zoologischer Anzeiger* 224(1/2): 49–73.
- Mey, E. 1994: Beziehungen zwischen Larvenmorphologie und Systematik der Adulti bei den Vogel-Ischnozeren (Insecta, Phthiraptera, Ischnocera). *Mitteilungen aus dem Zoologischen Museum in Berlin* 70(1): 3–84.
- Mey, E. 2003: On the development of animal louse systematics (Insecta, Phthiraptera) up to the present day. *Rudolstädter Naturhistorische Schriften* 11: 115–134.
- Mey, E. 2004: Zur Taxonomie, Verbreitung und parasitophyletischer Evidenz des *Philopterus*-Komplexes (Insecta, Phthiraptera, Ischnocera). *Ornithologischer Anzeiger* 43(2): 149–203.

- Mey, E. 2005: *Psittacobrosus bechsteini*: ein neuer ausgestorbener Federling (Insecta, Phthiraptera, Amblycera) vom Dreifarbenara *Ara tricolor* (Psittaciiformes), nebst einer annotierten Übersicht über fossile und rezent ausgestorbene Tierläuse. *Anzeiger des Vereins Thüringer Ornithologen* 5: 201–217.
- Mey, E. 2011 [2010]: *Nesiotinus kerguelensis* n. sp. – ein Federling (Phthiraptera, Ischnocera, Philopteridae s. l.) vom Lummensturmvogel *Pelecanoides urinatrix exsul* Salvin von den subantarktischen Kerguelen, nebst Anmerkungen zur Gattung *Nesiotinus* Kellogg. *Rudolstädter Naturhistorische Schriften* 17: 77–94.
- Mey, E. 2014: Naturhistorisches Museum im Thüringer Landesmuseum Heidecksburg Rudolstadt. Pp. 90–111. In: Werneburg, R. & Mey, E. (Eds) *Thüringer Natur-Schätze Naturkundliche Museen und Sammlungen im Freistadt Thüringen*. Regensburg: Verlag Schnell & Steiner. 280 pp.
- Mey, E.; Dalgleish, R.C. & Rékási, J. 2007: Berlin 1975: The forgotten symposium. *Rudolstädter Naturhistorische Schriften* 14: 61–69.
- Mey, E. & González-Acuña, D. 2007: Über einen Massenbefall von *Bovicola (Lepikentron) breviceps* (Rudow) (Insecta, Phthiraptera, Ischnocera, Bovicolidae) auf einem Alpaka *Vicugna vicugna* forma *pacos* in Thüringen (Deutschland), mit Anmerkungen zur Parthenogenese bei Tierläusen. *Rudolstädter Naturhistorische Schriften* 14: 71–82.
- Millán, J.; Gortazar, C.; Martín-Mateo, M.P. & Villafuerte, R. 2004: Comparative survey of the ectoparasite fauna of wild and farm-reared red-legged partridges (*Alectoris rufa*), with an ecological study in wild populations. *Parasitology Research* 93(1): 79–85.
- Miller, D. 1971: *Common insects in New Zealand*. Wellington: A.H. & A.W. Reed Ltd. xx + 178 pp., 9 pls.
- Miller, P. & Miller, K. 1986: A beach-wrecked red-tailed tropicbird. *Notornis* 33(1): 50–51.
- Mjöberg, E. 1910a: Studien über Mallophagen und Anopluren. *Arkiv för Zoologi* 6(13): 1–296, 5 pls.
- Mjöberg, E. 1910b: Studien über Pediculiden und Mallophagen. *Zoologischer Anzeiger* 35(9/10): 287–293.
- Modrzejewska, M. & Zlotorzyska, J. 1987: Studies on morphology of nymphs of selected Amblycera and Ischnocera (Mallophaga). *Polskie Pismo Entomologiczne* 57: 657–672.
- Monteiro, L.R. & Furness, R.W. 1995: Fea's petrel *Pterodroma feae* in the Azores. *Bulletin of the British Ornithologist's Club* 115(1): 9–14, pl. 1.
- Moreby, C. 1976: The Phthiraptera of birds from Rennell Island. *The Natural History of Rennell Island, British Solomon Islands* (Copenhagen) 7(15): 91–96.
- Moreby, C. 1978: The biting louse genus *Werneckiella* (Phthiraptera: Trichodectidae) ectoparasitic on the horse family Equidae (Mammalia: Perissodactyla). *Journal of Natural History* 12: 395–412.
- Mourik, S.C. van & Norman, F.I. 1985: Ectoparasites of some waterfowl (Anatidae) from Victoria. *Occasional Papers from the Museum of Victoria* 2: 1–3.
- Moyer, B.R. & Wagenbach, G.E. 1995: Sunning by black noddies (*Anous minutus*) may kill chewing lice (*Quadraceps hopkinsi*). *The Auk* 112(4): 1073–1077.
- Mumcuoglu, K.Y. & J. Zias, J. 1988: Head lice, *Pediculus humanus capitis* (Anoplura: Pediculidae) from hair combs excavated in Israel and dated from the first century B.C. to the eighth century A.D. *Journal of Medical Entomology* 25(6): 545–547.
- Murray, M.D. 1955a: Infestation of sheep with the face louse (*Linognathus ovillus*) *Australian Veterinary Journal* 31(2): 22–26.
- Murray, M.D. 1955b: Oviposition in lice with reference to *Damalinia ovis*. *Australian Veterinary Journal* 31: 320–321.
- Murray, M.D. 1957a: The distribution of the eggs of mammalian lice on their hosts I. Description of the oviposition behaviour. *Australian Journal of Zoology* 5: 13–18.
- Murray, M.D. 1957b: The distribution of the eggs of mammalian lice on their hosts II. Analysis of the oviposition behaviour of *Damalinia ovis* (L.) on the sheep. *Australian Journal of Zoology* 5: 19–29.
- Murray, M.D. 1957c: The distribution of the eggs of mammalian lice on their hosts III. The distribution of the eggs of *Damalinia ovis* (L.) on the sheep. *Australian Journal of Zoology* 5: 173–182.
- Murray, M.D. 1957d: The distribution of the eggs of mammalian lice on their hosts IV. The distribution of the eggs of *Damalinia equi* (Denny) and *Haematopinus asini* (L.) on the horse. *Australian Journal of Zoology* 5: 183–187.
- Murray, M.D. 1958: Ecology of the louse *Lepidophthirus macrorhini* Enderlein 1904 on the elephant seal *Mirounga leonina* (L.). *Nature* 182: 404–405.
- Murray, M.D. 1960a: The ecology of lice from sheep. I. The influence of skin temperature on populations of *Linognathus pedalis* (Osborne). *Australian Journal of Zoology* 8(3): 349–356.
- Murray, M.D. 1960b: The ecology of lice on sheep. II. The influence of temperature and humidity on the development and hatching of the eggs of *Damalinia ovis* (L.). *Australian Journal of Zoology* 8: 357–362.
- Murray, M.D. 1961: The ecology of the louse *Polyplax serrata* (Burm.) on the mouse *Mus musculus* L. *Australian Journal of Zoology* 9(1): 1–13.

- Murray, M.D. 1962: Efficiency of insecticides against the sheep body louse (*Damalinia ovis*). *Australian Veterinary Journal* 38: 308.
- Murray, M.D. 1963a: The ecology of lice from sheep. III. Differences between the biology of *Linognathus pedalis* (Osborne) and *L. ovillus* (Neumann). *Australian Journal of Zoology* 11: 153–156.
- Murray, M.D. 1963b: The ecology of lice from sheep. IV. The establishment and maintenance of populations of *Linognathus ovillus* (Neumann). *Australian Journal of Zoology* 11: 157–172, pl. 1.
- Murray, M.D. 1963c: The ecology of lice on sheep. V. Influence of heavy rain on populations of *Damalinia ovis* (L.). *Australian Journal of Zoology* 11: 173–182.
- Murray, M.D. 1963d: Influence of temperature on the reproduction of *Damalina equi* (Denny). *Australian Journal of Zoology* 11: 183–189.
- Murray, M.D. 1964: Ecology of the ectoparasites of seals and penguins. Pp. 241–245. In: Carrick, R.; Holdgate, M.W. & Prevost, J. (Eds) *Biologie Antarctique/Antarctic Biology*. Proceedings of 1st SCAR/TUBS Symposium on Antarctic Biology, Paris, 2–8 September 1962. Paris: Hermann. 651 pp.
- Murray, M.D. 1976: Insect parasites of marine birds and mammals. Chapter 4: 79–96. In: Cheng, L. (Ed.) *Marine Insects*. Amsterdam: North Holland Publishing Company. xii + 581 pp.
- Murray, M.D. & Calaby, J.H. 1971: The host relations of the Boopiidae. Appendix II: 81–84. In: Kéler, S. von 1971: A revision of the Australasian Boopiidae (Insecta: Phthiraptera), with notes on the Trimenoponidae. *Australian Journal of Zoology, Supplement* 6: 1–126.
- Murray, M.D. & Gordon, G. 1969: Ecology of lice on sheep. VII. Population dynamics of *Damalinia ovis* (Schrank). *Australian Journal of Zoology* 17: 179–186.
- Murray, M.D. & Nicholls, D.G. 1965: Studies on the ectoparasites of seals and penguins. I. The ecology of the louse *Lepidophthirus macrorhini* Enderlein on the southern elephant seal, *Mirounga leonina* (L.). *Australian Journal of Zoology* 13: 437–454, 4 pls.
- Murray, M.D.; Palma, R.L. & Pilgrim, R.L.C. 1990: Ectoparasites of Australian, New Zealand and Antarctic birds. Pp. 1365–1374. In: Marchant, S. & Higgins P.J. (Eds) *Handbook of Australian, New Zealand and Antarctic Birds*. Volume 1. *Ratites to Ducks*. Melbourne: Oxford University Press. 1400 pp.
- Murray, M.D.; Palma, R.L. & Pilgrim, R.L.C. 1993: Ectoparasites of Australian, New Zealand and Antarctic birds. Pp. 959–962. In: Marchant, S. & Higgins, P.J. (Eds) *Handbook of Australian, New Zealand and Antarctic Birds*. Volume 2. *Raptors to Lapwings*. Melbourne: Oxford University Press. 984 pp.
- Murray, M.D.; Palma, R.L. & Pilgrim, R.L.C. 1999: Ectoparasites of Australian, New Zealand and Antarctic birds. Pp. 1240–1242. In: Higgins, P.J. (Ed.) *Handbook of Australian, New Zealand and Antarctic Birds*. Volume 4. *Parrots to Dollarbird*. Melbourne: Oxford University Press. 1248 pp.
- Murray, M.D.; Palma, R.L. & Pilgrim, R.L.C. 2006a: Addendum: Appendix II— Volume 3. Ectoparasites of Australian, New Zealand and Antarctic birds. Pp. 1964–1966. In: Higgins, P.J.; Peter, J.M. & Cowling S.J. (Eds) *Handbook of Australian, New Zealand & Antarctic Birds*. Volume 7. *Boatbill to Starlings*. Melbourne: Oxford University Press. 1984 pp.
- Murray, M.D.; Palma, R.L.; Pilgrim, R.L.C. & Shaw, M.D. 2001: Ectoparasites of Australian, New Zealand and Antarctic birds. Pp. 1261–1263. In: Higgins, P.J.; Peter, J.M. & Steele, W.K. (Eds) *Handbook of Australian, New Zealand and Antarctic Birds*. Volume 5. *Tyrant-flycatchers to Chats*. Melbourne: Oxford University Press. 1269 pp.
- Murray, M.D.; Palma, R.L.; Pilgrim, R.L.C. & Shaw, M.D. 2002: Ectoparasites of Australian, New Zealand and Antarctic birds. Pp. 1215–1217. In: Higgins, P.J. & Peter, J.M. (Eds) *Handbook of Australian, New Zealand and Antarctic Birds*. Volume 6. *Pardalotes to shrike-thrushes*. Melbourne: Oxford University Press. 1225 pp.
- Murray, M.D.; Palma, R.L.; Pilgrim, R.L.C. & Shaw, M.D. 2006b: Appendix I: Ectoparasites of Australian, New Zealand and Antarctic birds. Pp. 1956–1960. In: Higgins, P.J.; Peter, J.M. & Cowling S.J. (Eds) *Handbook of Australian, New Zealand & Antarctic Birds*. Volume 7. *Boatbill to Starlings*. Melbourne: Oxford University Press. 1984 pp.
- Murray, M.D.; Smith, M.S.R. & Soucek, Z. 1965: Studies on the ectoparasites of seals and penguins II. The ecology of the louse *Antarctophthirus ogmorhini* Enderlein on the Weddell seal, *Leptonychotes weddelli* Lesson. *Australian Journal of Zoology* 13: 761–771, 2 pls.
- Myers, J.G. 1922: The order Hemiptera in New Zealand. With special reference to its biological and economic aspects. *The New Zealand Journal of Science and Technology* 5(1): 1–12.
- Najer, T.; Gustafsson, D.R. & Sychra, O. 2016: Two new species of *Philopteroides* (Phthiraptera: Ischnocera: Philopteridae) of the *beckeri* species-group, from New Guinean painted berrypeckers (Aves: Passeriformes: Paramythiidae). *Zootaxa* 4139(4): 527–541.
- Nakagawa, H. 1959a: New Mallophaga from the streaked shearwater, *Calonectris leucomelas* from Japanese waters. Part I. *Journal of Agricultural Sciences, Tokyo Nogyo Daigaku* 4: 384–389.

- Nakagawa, H. 1959b: A redescription of *Dennyus major* (Uchida) from the needle-tailed swift *Hirundapus caudacutus caudacutus* (Mallophaga). *Journal of Agricultural Sciences, Tokyo Nogyo Daigaku* 5: 23–27.
- Nelson, B.C. 1969: Bird ectoparasites from Norfolk Island. *Australian Zoologist* 15(2): 199–200.
- Nelson, B.C. 1972: A revision of the new world species of *Ricinus* (Mallophaga) occurring on Passeriformes (Aves). *University of California Publications in Entomology* 68: 1–130, 43 plates.
- Nelson, B.C. & Murray, M.D. 1971: The distribution of Mallophaga on the domestic pigeon (*Columba livia*). *International Journal for Parasitology* 1: 21–29, 2 pls.
- Nelson, R.C. & Price, R.D. 1965: The *Laemobothrion* (Mallophaga: Laemobothriidae) of the Falconiformes. *Journal of Medical Entomology* 2(3): 249–257.
- Neuffer, G. 1954: Die Mallophagenhaut und ihre Differenzierungen. *Zoologische Jahrbücher. Abteilung für Anatomie und Ontogenie der Tiere* 73(4): 449–519, 1 pl.
- Neumann, L.-G. 1890: Contribution a l'étude des Ricinidae parasites des oiseaux de la famille des Psittacidae. *Bulletin de la Société d'Histoire Naturelle*, Toulouse 24: 55–69.
- Neumann, L.-G. 1906: Notes sur les Mallophages. I.— Nomenclature. *Bulletin de la Société Zoologique de France* 31: 54–60.
- Neumann, L.-G. 1907a: Pédiculidés, Mallophages, Ixodidés. Pp. 13–17. In: Charcot, J. (Ed.) *Expédition Antarctique Française (1903–1905). Sciences naturelles: documents scientifiques. Arthropodes*. Paris: Mason et C<sup>ie</sup>, Editeurs. 9: 1–100.
- Neumann, L.-G. 1907b: Nouveau pou du mouton (*Haematopinus ovillus* n. sp.). *Revue Vétérinaire* 32(64): 520–524.
- Neumann, L.-G. 1911: Part III. Mallophaga. Pp. 19–22, pl. 3. In: *British Antarctic Expedition 1907–9 under the command of Sir E.H. Shackleton, c.v.o. Reports on the Scientific Investigations, Vol. 2, Biology*. 68 pp., 8 pls.
- Neumann, L.-G. 1912a: Notes sur les Mallophages.—II. I.— Sur le genre *Menopon* (sous-genre *Menacanthus* n. subgen.). *Archives de Parasitologie*, Paris 15(3): 353–368.
- Neumann, L.-G. 1912b: Notes sur les Mallophages. —II. II.— Espèces nouvelles (Gen. div.). *Archives de Parasitologie*, Paris 15(3): 368–384.
- Neumann, L.-G. 1913: Mallophaga. Deuxième Expédition Antarctique Française (1908–1910). *Sciences Naturelles: Documents Scientifiques*, Masson et Cie, Paris: 187–196.
- Nicholls, D.C.; Urquhart, E.M.; Ward, J.B.; Johns, P.M. 1998: A list of insect primary types held in Canterbury Museum, Christchurch, New Zealand. *Records of the Canterbury Museum* 12(2): 1–43.
- Nitzsch, C.L. 1818: Die Familien und Gattungen der Theierinsekten (Insecta epizoica); als ein Prodrömus einer Naturgeschichte derselben. *E.F. Germar's Magazin der Entomologie* 3: 261–318.
- Nitzsch, C.L. 1866: Die Federlinge der Sing-, Schrei-, Kletter- und Taubenvögel. *Zeitschrift für die gesammten Naturwissenschaften* (Halle) 27(2): 115–122.
- Nuttall, G.H.F. 1917: The biology of *Pediculus humanus*. *Parasitology* 10(1): 80–185.
- Nuttall, G.H.F. 1918: The biology of *Phthirus pubis*. *Parasitology* 10(3): 383–405.
- Nuttall, G.H.F. 1919: The systematic position, synonymy and iconography of *Pediculus humanus* and *Phthirus pubis*. *Parasitology* 11(3–4): 329–345.
- Olfers, I.F.J.M. von 1816: *De vegetativis et animatis corporibus in corporibus animatis reperiundis commentarius*. Berolini: Taberna Libraria Maureriana. vi + 113 pp., 1 pl.
- Osborn, H. 1890: Mallophaga. Pp. 188–189. In: Howard, L.O. (Ed.) *Scientific results of explorations by the U. S. fish commission steamer Albatross. No. V.— Annotated catalogue of the insects collected in 1887–88. Proceedings of the United States National Museum* 12(771): 185–216.
- Osborn, H. 1896: Insects affecting domestic animals: an account of the species of importance in North America, with mention of related forms occurring on other animals. *Bulletin U.S. Department of Agriculture Division Entomology (new series)* 5: 1–302, 2 pls.
- Overgaard, C. 1943: Mallophaga from gallinaceous birds. *Entomologiske Meddelelser uidegivet af Entomologiske Forening* 23: 1–17.
- Packard, A.S. 1870: Certain parasitic insects. *American Naturalist* 4: 83–99, pl. 1.
- Page, R.D.M.; Cruickshank, R.H.; Dickens, M.; Furness, R.W.; Kennedy, M.; Palma, R.L. & Smith, V.S. 2004: Phylogeny of “*Philoceanus* complex” seabird lice (Phthiraptera: Ischnocera) inferred from mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution* 30: 633–652.
- Palma, R.L. 1978: Slide-mounting of lice: a detailed description of the Canada balsam technique. *The New Zealand Entomologist* 6(4): 432–436.
- Palma, R.L. 1991a: A new species of *Rallicola* (Insecta: Phthiraptera: Philopteridae) from the North Island brown kiwi. *Journal of the Royal Society of New Zealand* 21(4): 313–322.
- Palma, R.L. 1991b: Ancient head lice on a wooden comb from Antinoë, Egypt. *The Journal of Egyptian Archaeology* 77: 194, pl. 21.

- Palma, R.L. 1991c: Two bird lice (Insecta: Phthiraptera) collected during Captain Cook's 2nd voyage around the world. *Archives of Natural History* 18(2): 237–247.
- Palma, R.L. 1994a: New synonymies in the lice (Insecta: Phthiraptera) infesting albatrosses and petrels (Procellariiformes). *The New Zealand Entomologist* 17: 64–69.
- Palma, R.L. 1994b: The identity of *Nirmus obtusus* and other *Quadriceps* species (Phthiraptera: Philopteridae) from Clipperton Island and the Galápagos Islands. *Journal of the Royal Society of New Zealand* 24(3): 267–276.
- Palma, R.L. 1995: A new synonymy and new records of *Quadriceps* (Insecta: Phthiraptera: Philopteridae) from the Galápagos Islands. *New Zealand Journal of Zoology* 22(2): 217–222.
- Palma, R.L. 1996a: First records of marsupial lice (Insecta: Phthiraptera: Boopiidae) on a brushtailed rock wallaby from New Zealand. *New Zealand Journal of Zoology* 23(2): 161–164.
- Palma, R.L. 1996b: Laemobothriidae; Menoponidae; Ricinidae; Philopteridae. Pp. 106–230. In: Wells, A. (Ed.) *Zoological Catalogue of Australia*. Volume 26. *Psocoptera, Phthiraptera, Thysanoptera*. Melbourne: CSIRO Publishing, Australia. xiv + 418 pp.
- Palma, R.L. 1999: Amendments and additions to the 1982 list of chewing lice (Insecta: Phthiraptera) from birds in New Zealand. *Notornis* 46: 373–387.
- Palma, R.L. 2000: The species of *Saemundssonina* (Insecta: Phthiraptera: Philopteridae) from skuas (Aves: Stercorariidae). *New Zealand Journal of Zoology* 27(2): 121–128.
- Palma, R.L. 2001: Museum marine file: Wingless fliers? Yes, they are feather lice! *Seafood New Zealand* 9(8): 65–68.
- Palma, R.L. 2004: Designation of a lectotype for *Docophorus atlanticus* Kellogg, 1914 (Insecta: Phthiraptera: Philopteridae). *Tuhinga – Records of the Museum of New Zealand Te Papa Tongarewa* 15: 13–16.
- Palma, R.L. 2010: Order Phthiraptera: Lice. Pp. 294–296, 407–409. In: Gordon, D.P. (Ed.) *New Zealand Inventory of Biodiversity*. Volume Two. *Kingdom Animalia – Chaetognatha, Ecdysozoa, Ichnofossils*. Christchurch: Canterbury University Press. 528 pp.
- Palma, R.L. 2011a: A tribute to Robert Louis Cecil Pilgrim. Research Associate, Museum of New Zealand Te Papa Tongarewa. *Tuhinga – Records of the Museum of New Zealand Te Papa Tongarewa* 22: 149–156.
- Palma, R.L. 2011b: New taxa, new synonymies and new host records in the louse genus *Halipeurus* (Insecta: Phthiraptera: Philopteridae) parasitic on petrels (Aves: Procellariiformes). *Zootaxa* 3017: 1–45.
- Palma, R.L. 2012: Three new species of the louse genus *Saemundssonina* (Insecta: Phthiraptera: Philopteridae). *Zootaxa* 3478: 38–48.
- Palma, R.L. 2015a: A new species of the genus *Anatoecus* (Phthiraptera: Philopteridae) from the black swan, *Cygnus atratus*. *The New Zealand Entomologist* 38(2): 142–148.
- Palma, R.L. 2015b: The correct type hosts and designation of a lectotype for the louse *Nirmus triangulatus alpha* (Insecta: Phthiraptera: Philopteridae). *The New Zealand Entomologist* 38(2): 149–155.
- Palma, R.L. & Horning, D.S. 2002: The lice (Insecta: Phthiraptera) from Macquarie Island. *ANARE Research Notes* 105: 1–27.
- Palma, R.L. & Imber, M.J. 2000: Coexistence of two species of *Halipeurus* (Phthiraptera) on Chatham Island Taiko (*Pterodroma magentae*) (Aves). *New Zealand Journal of Zoology* 27(3): 229–232.
- Palma, R.L. & Jensen, J.-K. 2005: Lice (Insecta: Phthiraptera) and their host associations in the Faroe Islands. *Steenstrupia* 29(1): 49–73.
- Palma, R.L.; Lovis, P.M. & Tither, C. 1989: An annotated list of primary types of the phyla Arthropoda (except Crustacea) and Tardigrada held in the National Museum of New Zealand. *National Museum of New Zealand Miscellaneous Series* 20: 1–49.
- Palma, R.L.; McKenna, P.B. & Aitken, P. 2006: Confirmation of the occurrence of the chewing louse *Bovicola (Lepikentron) breviceps* (Insecta: Phthiraptera: Trichodectidae) on alpacas (*Lama pacos*) in New Zealand. *New Zealand Veterinary Journal* 54(5): 253–254.
- Palma, R.L. & Peck, S.B. 2013: An annotated checklist of parasitic lice (Insecta: Phthiraptera) from the Galápagos Islands. *Zootaxa* 3627(1): 1–87.
- Palma, R.L. & Pilgrim, R.L.C. 1977: An abnormal male of *Halipeurus (Halipeurus) consimilis* (Mallophaga: Philopteridae) without genitalia. *The New Zealand Entomologist* 6(3): 290–292.
- Palma, R.L. & Pilgrim, R.L.C. 1983: The genus *Bedfordiella* (Mallophaga: Philopteridae) and a note on the lice from the Kerguelen petrel (*Pterodroma brevirostris*). *National Museum of New Zealand Records* 2(13): 145–150.
- Palma, R.L. & Pilgrim, R.L.C. 1984: A revision of the genus *Harrisoniella* (Mallophaga: Philopteridae). *New Zealand Journal of Zoology* 11(2): 145–166.
- Palma, R.L. & Pilgrim, R.L.C. 1988. [1987]: A revision of the genus *Perineus* (Phthiraptera: Philopteridae). *New Zealand Journal of Zoology* 14(4): 563–586.

- Palma, R.L. & Pilgrim, R.L.C. 2002: A revision of the genus *Naubates* (Insecta: Phthiraptera: Philopteridae). *Journal of the Royal Society of New Zealand* 32(1): 7–60.
- Palma, R.L. & Price, R.D. 2000: *Philopterus novaezealandiae*, a new species of chewing louse (Phthiraptera: Philopteridae) from the Kokako (Passeriformes: Callaeidae). *Journal of the Royal Society of New Zealand* 30(3): 293–297.
- Palma, R.L. & Price, R.D. 2004: *Apterygon okarito*, a new species of chewing louse (Insecta: Phthiraptera: Menoponidae) from the Okarito brown kiwi (Aves: Apterygiformes: Apterygidae). *New Zealand Journal of Zoology* 31(1): 67–73.
- Palma, R.L. & Price, R.D. 2005: *Menacanthus rhipidurae*, a new species of chewing louse (Insecta: Phthiraptera: Menoponidae) from South Island fantails, *Rhipidura fuliginosa fuliginosa* (Aves: Passeriformes: Dicuridae). *New Zealand Journal of Zoology* 32(2): 111–115.
- Palma, R.L.; Price, R.D. & Hellenthal, R.A. 1998: New synonymies and host records for lice of the genus *Menacanthus* (Phthiraptera: Menoponidae) from the Passeriformes (Aves). *Journal of the Royal Society of New Zealand* 28(2): 309–320.
- Palma, R.L. & Tennyson, A.J.D. 2005: Designation of a lectotype and clarification of authorship and date of publication for the New Caledonian Petrel (*Pterodroma leucoptera caledonica*). *Notornis* 52(4): 247–248.
- Paterson, A.M.; Palma, R.L. & Gray, R.D. 1999: How frequently do avian lice miss the boat? Implications for coevolutionary studies. *Systematic Biology* 48(1): 214–223.
- Paterson, A.M.; Palma, R.L. & Gray, R.D. 2003: Drowning on arrival, missing the boat, and *x*-events: how likely are sorting events? Pp. 287–309. In: Page, R.D.M. (Ed.) *Tangled trees. Phylogeny, cospeciation, and coevolution*. Chicago & London: The University of Chicago Press. x + 350 pp.
- Paulian, P. 1953: Pinnipèdes, cétacés, oiseaux des Îles Kerguelen et Amsterdam. Mission Kerguelen 1951. *Mémoires de l'Institut Scientifique de Madagascar, Série A*, 8: 111–234.
- Payne, M.R. & Prince, P.A. 1979: Identification and breeding biology of the diving petrels and at South Georgia. *New Zealand Journal of Zoology* 6: 299–318.
- Pérez D'A., V. 1985: La teoría de la colonización de las islas aplicada a las especies de *Austrogoniodes* Harrison (Mallophaga: Ischnocera: Philopteridae) en su relación con los Sphenisciformes. *Revista Chilena de Entomología* 12: 159–164.
- Pérez, J.M.; Sánchez, I. & Palma, R.L. 2013: The dilemma of conserving parasites: the case of *Felicola (Loricicola) isidoro* (Phthiraptera: Trichodectidae) and its host, the endangered Iberian lynx (*Lynx pardinus*). *Insect Conservation and Diversity* 6: 680–686.
- Pessôa, S.B. & Guimarães, L.R. 1935a: Contribuição para o conhecimento das Mallophagas das aves do Brasil. I. Novas especies do genero *Esthiopterum*. *Annaes da Faculdade de Medicina da Universidade de São Paulo* 11(3): 311–314, 2 pls.
- Pessôa, S.B. & Guimarães, L.R. 1935b: Contribuições para o conhecimento das Mallophagas das aves do Brasil. III. Especies novas ou pouco conhecidas da Fam. *Philopteridae*. *Revista de Biologia e Higiene* 6(2): 105–112.
- Pfeffer, A.; Cole, D.J.W.; Bishop, D.M.; Heath, A.C.G. & Phegan, M.D. 1996: Detection of dermatophilosis and lice (*Bovicola ovis*) on flayed pelts and cockle on the skin of live lambs. *New Zealand Veterinary Journal* 44(4): 121–125.
- Piaget, E. 1871: Description de quelques parasites du genre *Docophorus*. *Tijdschrift voor Entomologie* 14: 113–137, 2 pls.
- Piaget, E. 1878: *Acidoproctus*. *Tijdschrift voor Entomologie* 21: 178–184, pl. 12.
- Piaget, E. 1880: *Les Pédiculines. Essai Monographique*. Two volumes. Leide: E.J. Brill. xxxix + 714 pp., 56 pls.
- Piaget, E. 1883: Quelques pédiculines nouvelles ou peu connues. *Tijdschrift voor Entomologie* 26(2): 152–158, 1 pl.
- Piaget, E. 1885: *Les Pédiculines. Essai Monographique*. Supplément. Leide: E.J. Brill. xvi + 200 pp., 17 pls.
- Piaget, E. 1890a: Quelques pédiculines nouvelles. *Tijdschrift voor Entomologie* 33: 223–259, 8 pls.
- Piaget, E. 1890b: Papers read: *Nirmus assimilis*, Piaget. *The Proceedings of the Entomological Society of London*. 1890: xxii–xxiv.
- Picaglia, L. 1885: Pediculini nuovi del Museo di Zoologia ed Anatomia Comparata della R. Università di Modena. *Atti della Società Italiana de Scienze Naturali* 28: 82–90.
- Pilgrim, R.L.C. 1970: Knowledge of New Zealand Mecoptera, Mallophaga, Anoplura and Siphonaptera. *The New Zealand Entomologist* 4(3): 72–79.
- Pilgrim, R.L.C. 1974: Lice and fleas (1). *New Zealand's Nature Heritage* 3(37): 1030–1036.
- Pilgrim, R.L.C. 1976: Mallophaga on the rock pigeon (*Columba livia*) in New Zealand, with a key to their identification. *The New Zealand Entomologist* 6(2): 160–164.
- Pilgrim, R.L.C. & Palma, R.L. 1979: A redescription of *Perineus concinnus* (Mallophaga: Philopteridae). *Pacific Insects* 21(2–3): 172–178.
- Pilgrim, R.L.C. & Palma, R.L. 1982: A list of the chewing lice (Insecta: Mallophaga) from birds in New Zealand. *Notornis* 29(suppl.): 1–32 (also as *National Museum of New Zealand Miscellaneous Series* 6).
- Pilgrim, R.L.C. & Palma, R.L. 1994: New synonymy between *Philopterus antarcticus* and *Saemundssonina nivea* (Phthiraptera: Philopteridae). *New Zealand Journal of Zoology* 21(3): 239–244.

- Polack, J.S. 1838: *New Zealand: being a narrative of travels and adventures during a residence in that country between the years 1831 and 1837*. Volume I. London: Richard Bentley. xvi + 442 pp.
- Powell, J.A. 2001: *Vernon L. Kellogg Mallophaga (Phthiraptera) Collection*. Essig Museum of Entomology. <http://bnhmwp.berkeley.edu/essig/sample-page/holdings/kellogg/> (accessed March 2016).
- Price, P.W. 1980: *Evolutionary biology of parasites*. Princeton, New Jersey: Princeton University Press. xi + 237 pp.
- Price, R.D. 1967a: The *Colpocephalum* (Mallophaga: Menoponidae) of the Pelecaniformes. *The Canadian Entomologist* 99(3): 273–280.
- Price, R.D. 1967b: A new species of *Colpocephalum* (Mallophaga: Menoponidae) from the kea. *Journal of the Kansas Entomological Society* 40(1): 9–12.
- Price, R.D. 1967c: A new subgenus and new species of *Franciscoloa* Conci (Mallophaga: Menoponidae). *Pacific Insects* 9(3): 511–517.
- Price, R.D. 1969: Three new species of *Heteromenopon* (Mallophaga: Menoponidae) from Brazilian and Australian parrots. *Annals of the Entomological Society of America* 62(3): 563–566.
- Price, R.D. 1971: A review of the genus *Holomenopon* (Mallophaga: Menoponidae) from the Anseriformes. *Annals of the Entomological Society of America* 64(3): 633–646.
- Price, R.D. 1974: A review of the genus *Pseudomenopon* (Mallophaga: Menoponidae). *Annals of the Entomological Society of America* 67(1): 73–84.
- Price, R.D. 1987: Chapter 22. Order Mallophaga. Pp. 215–223. In: Stehr, F.W. (Ed.) *Immature insects*. Volume 1. Dubuque, Iowa: Kendall / Hunt Publishing Company. xiv + 754 pp.
- Price, R.D. & Beer, J.R. 1963a: Species of *Colpocephalum* (Mallophaga: Menoponidae) parasitic upon the Falconiformes. *The Canadian Entomologist* 95(7): 731–763.
- Price, R.D. 1963b: The *Kurodaia* (Mallophaga: Menoponidae) parasitic on the Strigiformes, with a key to the species of the genus. *Annals of the Entomological Society of America* 56(6): 849–857.
- Price, R.D. 1964: Species of *Colpocephalum* (Mallophaga: Menoponidae) parasitic upon the Galliformes. *Annals of the Entomological Society of America* 57(4): 391–402.
- Price, R.D. 1965a: A review of *Ciconiphilus* (Mallophaga: Menoponidae). *The Canadian Entomologist* 97(6): 657–666.
- Price, R.D. 1965b: A review of the *Colpocephalum* of the Corvidae with the description of a new species (Mallophaga: Menoponidae). *Proceedings of the Entomological Society of Washington* 67(1): 7–14.
- Price, R.D. 1965c: The *Colpocephalum* (Mallophaga: Menoponidae) of the Ciconiiformes. *Annals of the Entomological Society of America* 58(1): 111–131.
- Price, R.D. 1966: A review of the genus *Franciscoloa* Conci (Mallophaga: Menoponidae). *Pacific Insects* 8(3): 633–648.
- Price, R.D. 1967: The genus *Heteromenopon* (Mallophaga: Menoponidae), with description of a new subgenus and six new species. *Annals of the Entomological Society of America* 60(2): 328–338.
- Price, R.D. & Clay T. 1972: A review of the genus *Austromenopon* (Mallophaga: Menoponidae) from the Procellariiformes. *Annals of the Entomological Society of America* 65(2): 487–504.
- Price, R.D. & Dagleish, R.C. 2007: *Myrsidea* Waterston (Phthiraptera: Menoponidae) from the Emberizidae (Passeriformes), with descriptions of 13 new species. *Zootaxa* 1467: 1–18.
- Price, R.D. & Emerson, K.C. 1978: The *Neopsittaconirmus* (Mallophaga: Philopteridae) from *Cacatua* (Aves: Psittaciformes). *Pacific Insects* 18(1–2): 33–36.
- Price, R.D.; Hellenthal, R.A. & Palma, R.L. 2003: *World checklist of chewing lice with host associations and keys to families and genera*. Pp. 1–448. In: Price, R.D.; Hellenthal, R.A.; Palma, R.L.; Johnson, K.P. & Clayton, D.H. *The chewing lice: world checklist and biological overview*. Illinois Natural History Survey Special Publication 24. x + 501 pp.
- Price, R.D.; Johnson, K.P. & Palma, R.L. 2008: A review of the genus *Forficuloecus* Conci (Phthiraptera: Philopteridae) from parrots (Psittaciformes: Psittacidae), with descriptions of four new species. *Zootaxa* 1859: 49–62.
- Ramchurn, R. 1980: Observations on the life cycle of *Linognathus stenopsis*. *Revue Agricole et Sucrière de l'Île Maurice* 59(1): 6–8.
- Ramcke, J. 1965: Der Kopf der Schweineläus (Haematopinus suis L., Anoplura). *Zoologische Jahrbücher Abteilung für Anatomie und Ontogenie der Tiere* 82: 547–663.
- Ramli, R.; Cusack, M.; Curry, G.B. & Furness, R.W. 2000: Morphological variation of chewing lice (Insecta: Phthiraptera) from different skua taxa. *Biological Journal of the Linnean Society* 71(1): 91–101.
- Redi, F. 1668: *Esperienze intorno alla generazione degl'insetti*. Firenze: All' Insegna della Stella. 228 pp., 29 pls.
- Reed, D.L.; Allen, J.M.; Troups, M.A.; Boyd, B.M. & Ascunce, M.S. 2015: The study of primate evolution from a lousy perspective. Chapter 10: 202–214. In: Morand, S.; Krasnov, B.R. & Littlewood, D.T.J. (Eds) *Parasite diversity and diversification: Evolutionary ecology meets phylogenetics*. Cambridge University Press: Cambridge, U.K. xiv + 488 pp.



- Reed, D.L.; Light, J.E.; Allen, J.M. & Kirchman, J.J. 2007: Pair of lice lost or parasites regained: The evolutionary history of anthropoid primate lice. *BioMed Central Biology* 5: 7, 11 pp.
- Reid, B. & Williams, G.R. 1975: The kiwi. Pp. 301–330. In: Kuschel, G. (Ed.) *Biogeography and Ecology in New Zealand*. The Hague: W. Junk. xvi + 689 pp.
- Rékási, J. 1986: Magyarországi madarak tolltetvei (Mallophaga). II. *Parasitologia Hungarica* 19: 119–126, 8 pls.
- Rékási, J. & Kiss, B.J. 1984: Weitere Angaben zur Kenntnis der Federlinge (Mallophaga) der Vögel Nord-Dobrudschas Rumänien. II. *Parasitologica Hungarica* 17: 97–117.
- Retzius, A.J. 1783: 87. *Pediculus*. P. 201. In: *Caroli Lib. Bar. De Geer. Genera et Species Insectorum e generosissimi auctoris scriptis extraxit, digessit, Latine quoad partem reddidit, et terminologiam insectorum Linneanam addidit*. Lipsiae: Siegfried Lebrecht Crusium. vi + 7–220. Index Linneo-Degeerianus 32 pp.
- Rheinwald, G. 1968: Die Mallophagengattung *Ricinus* De Geer, 1778. Revision der außeramerikanischen Arten. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 65: 181–326.
- Ribbeck, R. 1972: DDR-Erstnachweis der Haustauben-Mallophage *Bonomiella columbae*. *Angewandte Parasitologie* 13(3): 129–134.
- Rivera-Parra, J.L.; Levin, I.I. & Parker, P.G. 2014: Comparative ectoparasite loads of five seabird species in the Galapagos Islands. *Journal of Parasitology* 100(5): 569–577.
- Rivera-Parra, J.L.; Levin, I.I.; Johnson, K.P. & Parker, P.G. 2015: Lineage sorting in multihost parasites: *Eidmanniella albescens* and *Fregatiella aurifasciata* on seabirds from the Galapagos Islands. *Ecology and Evolution* 5(16): 3264–3271.
- Roberts, M. 1991: Parasitological evidence for the presence of other rodent species on “kiore only” islands. *Journal of the Royal Society of New Zealand* 21(4): 349–356.
- Robertson, H.A.; Dowding, J.E.; Elliott, G.P.; Hitchmough, R.A.; Miskelly, C.M.; O’Donnell, C.F.J.; Powlesland, R.G.; Sagar, P.M.; Scofield, R.P. & Taylor, G.A. 2013: Conservation status of New Zealand birds, 2012. *Department of Conservation, New Zealand Threat Classification Series* 4: 1–22.
- Rózsa, L. & Apari, P. 2012: Why infest the loved ones – inherent human behaviour indicates former mutualism with head lice. *Parasitology* 139: 696–700.
- Rózsa, L. & Vas, Z. 2015a: Host correlates of diversification in avian lice. Chapter 11: 215–229. In: Morand, S.; Krasnov, B.R. & Littlewood, D.T.J. (Eds) *Parasite diversity and diversification: Evolutionary ecology meets phylogenetics*. Cambridge University Press: Cambridge, U.K. xiv + 488 pp.
- Rózsa, L. & Vas, Z. 2015b: Co-extinct and critically co-endangered species of parasitic lice, and conservation-induced extinction: should lice be reintroduced to their hosts? *Oryx – The International Journal of Conservation* 49(1): 107–110.
- Rudolph, D. 1983: The water-vapour uptake system of the Phthiraptera. *Journal of Insect Physiology* 29(1): 15–25.
- Rudow, F. 1866a: Sechs neue Haarlinge. *Zeitschrift für die Gesamten Naturwissenschaften* (Halle) (N.F. 2) 27(2): 109–112, 3 pls.
- Rudow, F. 1866b: Charakteristik neuer Federlinge. *Zeitschrift für die Gesamten Naturwissenschaften* (Halle) (N.F. 2) 27(6): 465–477.
- Rudow, F. 1869a: Neue Mallophagen. *Zeitschrift für die gesamnten Naturwissenschaften* (Halle) (N.F. 2) 34(11/12): 387–407.
- Rudow, F. 1869b: *Beitrag zur Kenntniss der Mallophagen oder Pelzfresser. Neue exotische Arten der Familie Philopterus*. Inaugural – Dissertation, Universität zu Leipzig. Halle: Druck von Wilh. Plötz. 47 pp.
- Rudow, F. 1870: Beobachtungen über die Lebensweise und den Bau der Mallophagen oder Pelzfresser, sowie Beschreibung neuer Arten. *Zeitschrift für die Gesamten Naturwissenschaften* (Halle) (Neue Folge 1) 35(6): 449–487.
- Ryan, S.O. & Price, R.D. 1969: A review of the genus *Eidmanniella* (Mallophaga: Menoponidae) from the Pelecaniformes. *Annals of the Entomological Society of America* 62(4): 817–823.
- Sales, J. 2005: The endangered kiwi: a review. *Folia Zoologica. International Journal of Vertebrate Zoology* 54(1): 1–20.
- Samuel, W.M. & Trainer, D.O. 1971: Seasonal fluctuations of *Tricholipeurus parallelus* (Osborn, 1896) (Mallophaga: Thichodectidae) on white-tailed deer *Odocoileus virginianus* (Zimmermann, 1780) from south Texas. *The American Midland Naturalist* 85(2): 507–513.
- Savill, R.A. 1990: Early records of the human head louse *Pediculus humanus capitis* (Phthiraptera: Pediculidae) in New Zealand. *Records of the Canterbury Museum* 10(6): 69–72.
- Schaefer, C.W. 1978: Ecological separation of the human head lice and body lice (Anoplura: Pediculidae). *Transactions of the Royal Society of Tropical Medicine and Hygiene* 72(6): 669–670.
- Schaefer, P.W. & Strandtmann, R.W. 1971: Notes on incidence and niche preference of Mallophaga and Analgoidea ectoparasitic on South polar skua (*Catharacta skua maccormicki*) on Ross Island, Antarctica. *Pacific Insects Monographs* 25: 15–16.
- Scharf, W.C. & Price, R.D. 1965: A taxonomic study of the genus *Cuculiphilus* (Mallophaga: Menoponidae). *Annals of the Entomological Society of America* 58(4): 546–555.

- Scharf, W.C. & Price, R.D. 1983: Review of the *Amyrsidea* in the subgenus *Argimenopon* (Mallophaga: Menoponidae). *Annals of the Entomological Society of America* 76(3): 441–451.
- Schrank, F. von P. 1776: *Beyträge zur Naturgeschichte*. Augsburg: Veith. 140 pp., 7 pls.
- Schrank, F. von P. 1781: *Enumeratio insectorum Austriae indigenorum*. Augustae Vindelicorum: Viduam Eberhardi Klett et Franck. ix + 548 pp., 4 pls.
- Schrank, F. von P. 1802: *Briefe naturhistorischen, physikalischen und ökonomischen Inhalts an Herrn Bernard Sebastian Nau, ehemaligen kurmainzischen Hofgerichtsrat und Professor – nebst drey vorausgeschickten naturhistorischen Abhandlungen*. Erlangen: J.C. Schubart. 384 pp.
- Schrank, F. von P. 1803: *Fauna Boica*. Landshut 3(1): 186–194.
- Scofield, R.P.; Christie, D.; Palma, R.L. & Tennyson, A.J.D. 2011: First record of streaked shearwater (*Calonectris leucomelas*) in New Zealand. *Notornis* 57(4): 212–215.
- Scopoli, J.A. 1763: *Entomologia Carniolica exhibens insecta Carnioliae indigena et distributa in ordines, genera, species, varietates, methodo Linnaeana*. Wien: Trattner. xxxvi + 421 pp.
- Séguy, E. 1924: *Les insectes parasites de l'homme et des animaux domestiques*. Paris: Lechevalier. 422 pp.
- Séguy, E. 1944: Insectes ectoparasites (Mallophages, Anoploures, Siphonaptères). *Faune de France* 43: 1–684.
- Séguy, E. 1953: Insectes mallophages, anoploures et diptères recueillis par M.P. Paulian aux Îles Kerguelen. *Mémoires de l'Institut Scientifique de Madagascar, Series E, 4*: 553–615.
- Selim, M.K.; El-Kasaby, A. & El-Refaii, A.H. 1968: External parasites of domestic pigeon in United Arabic Republic. *Angewandte Parasitologie* 9(2): 74–83.
- Selva, J.M.; González, J.; Aguirre, J.M. & Gallego, M. 1987: Parasitofauna de las palomas (*Columba livia* var.) de Barcelona. I. Mallophaga. *Revista Ibérica de Parasitología*, volumen Extraordinario: 245–250.
- Shepherd, L.D.; Worthy, T.H.; Tennyson, A.J.D.; Scofield, R.P.; Ramstad, K.M. & Lambert, D.M. 2012: Ancient DNA analyses reveal contrasting phylogeographic patterns amongst kiwi (*Apteryx* spp.) and a recently extinct lineage of spotted kiwi. *PLoS One* 7(8): 1–9 (e42384).
- Silva, H.M.; Valim, M.P. & Gama, R.A. 2014: Community of chewing lice (Phthiraptera: Amblycera and Ischnocera) parasites of resident birds at the Archipelago of São Pedro and São Paulo in northeast Brazil. *Journal of Medical Entomology* 51(5): 941–947.
- Soler-Cruz, M.D.; Benítez-Rodríguez, R. & Alcántara-Ibáñez, F. 1982: Contribution à l'étude de la taxonomie des mallophages parasites des oiseaux de la province de Grenade (Espagne). *Cahiers Office de la Recherche Scientifique et Technique Outre-mer, série Entomologie médicale et Parasitologie* 20(4): 255–264.
- Soler-Cruz, M.D. & Guevara-Benítez, D.C. 1981: *Brueelia* [sic] *densilimba* (Nitzsch, 1866) (Mallophaga), parásito de *Carduelis carduelis* L. *Revista Ibérica de Parasitología* 41(2): 275–282.
- Soler-Cruz, M.D.; Guevara-Benítez, D.C. & Alcántara-Ibáñez, F. 1985: *Brueelia merulensis* (Denny, 1842), *Brueelia marginata* (Burmeister, 1838) y *Menacanthus polonicus* Eichler y Złotorzycka, 1963 (Mallophaga, Insecta). Estudio morfológico y biométrico. *Revista Ibérica de Parasitología* 45(2): 107–117.
- Soler-Cruz, M.D.; Martín-Mateo, M.P. & Morillas-Márquez, F. 1979: *Brüelia parviguttata* (Blagoveshchensky, 1940) (Mallophaga) parásito de *Galerida cristata* L. *Revista Ibérica de Parasitología* 39: 165–173.
- Spellerberg, I.F. 1971: Mallophaga on the South Polar Skua (*Catharacta skua maccormicki*). *Pacific Insects Monographs* 25: 19–20.
- Spratt, D.M. 1983: A list of type specimens of animal parasites held in Australian institutions. *CSIRO Division of Wildlife & Rangelands Research Technical Memorandum* 19: 1–64.
- Steele, W.K.; Pilgrim, R.L.C. & Palma, R.L. 1997: Occurrence of the flea *Glaciopsyllus antarcticus* and avian lice in central Dronning Maud land. *Polar Biology* 18: 292–294.
- Stenram, H. 1956: The ecology of *Columbicola columbae* L. (Mallophaga). *Opuscula Entomologica* 21(2–3): 170–190.
- Stephenson, B.M.; Gaskin C.P.; Griffiths, R.; Jamieson, H.; Baird, K.A.; Palma, R.L. & Imber, M.J. 2009 [2008]: The New Zealand storm-petrel (*Pealeornis maoriana* Mathews, 1932): first live capture and species assessment of an enigmatic seabird. *Notornis* 55(4): 191–206.
- Storer, R.W. 2000: The metazoan parasite fauna of grebes (Aves: Podicipediformes) and its relationship to the birds' biology. *Miscellaneous Publications – Museum of Zoology, University of Michigan* 188: iv + 1–90.
- Stork, N. & Lyal, C.H.C. 1993: Extinction or 'co-extinction' rates? *Nature* 366: 307.
- Sychra, O. 2006: *Neopsittaconirmus vendulae*, a new species of louse (Phthiraptera: Philopteridae) from the cockatiel *Nymphicus hollandicus* (Psittaciformes: Cacatuidae). *Zootaxa* 1270: 57–68.
- Sychra, O.; Kolencik, S. & Palma, R.L. 2016: Three new species of *Myrsidea* (Phthiraptera: Menoponidae) from New Zealand passerines (Aves: Passeriformes). *Zootaxa* 4126(3): 397–410.

- Symmons, S. 1952: Comparative anatomy of the mallophagan head. *Transactions of the Zoological Society of London* 27(4): 349–436.
- Tandan, B.K. 1955: Mallophagan parasites from Indian birds. Part IV. Species belonging to the genera *Philopterus*, *Capraiella* and *Pectinopygus* (superfamily Ischnocera). *Annals and Magazine of Natural History* (Series 12) 8: 417–433.
- Tandan, B.K. 1958a: Mallophaga (Insecta). Part II. A new species of *Ibidoecus* Cummings, 1916. Pp. 151–154. In: Wolff, T. (Ed.) *The Natural History of Rennell Island, British Solomon Islands*. Volume 2 (Invertebrates, Pars). *Scientific Results of the Danish Rennell Expedition, 1951 and the British Museum (Natural History) Expedition, 1953*. Copenhagen & London: University of Copenhagen and the British Museum (Natural History). 234 pp., 22 pls.
- Tandan, B.K. 1958b: Mallophagan parasites from Indian birds— Part V. Species belonging to the genus *Ibidoecus* Cummings, 1916 (Ischnocera). *Transactions of the Royal Entomological Society of London* 110(14): 393–410.
- Tandan, B.K. 1972a: The species of *Apterygon* (Insecta: Phthiraptera: Amblycera) parasitic on Kiwis (*Apteryx*). *New Zealand Journal of Science* 15(1): 52–69.
- Tandan, B.K. 1972b: A new species of *Ardeicola* (Insecta: Phthiraptera: Ischnocera). *Journal of the Royal Society of New Zealand* 2(1): 49–54.
- Taschenberg, O. 1879: Über die Synonymie von *Goniocotes hologaster*. *Zeitschrift für die gesammten Naturwissenschaften* (Halle) 52: 104–107, 1 pl.
- Tandan, B.K. 1882: Die mallophagen mit besonderer Berücksichtigung der von Dr. Meyer gesammelten Arten systematisch bearbeitet. *Nova Acta der Kaiserlich Leopoldinisch-Carolinisch Deutschen Akademie der Naturforscher* 44(1): 1–244, 7 pls.
- Taylor, G.A. & Tennyson, A.J.D. 1994: Christmas Island shearwater *Puffinus nativitatis* on Curtis Island. *Notornis* 41(4): 287–291.
- Tendeiro, J. 1958: Études sur les Mallophages. Sur une nouvelle espèce du genre *Eidmanniella* von Kéler 1938 (Amblycera, Menoponidae), parasite de *Sula leucogaster leucogaster* (Boddaert) et *Sula leucogaster plotus* Forster. *Garcia de Orta, Revista da Junta das Missões Geográficas e de Investigações do Ultramar* 6(3): 443–449.
- Tendeiro, J. 1959: Études sur les mallophages. Sur quelques espèces et sous-espèces du genre *Nosopon* Hopkins (Amblycera, Menoponidae), parasites de Falconiformes. *Boletim Cultural da Guiné Portuguesa* 14(54): 193–211.
- Tendeiro, J. 1961: Mallophaga. *Memoires de l'Institut Français d'Afrique Noire* 62: 289–304, 6 pls.
- Tendeiro, J. 1963: Études sur les mallophages. Observations sur des Ischnocera africains, avec description de 12 espèces et 2 sous-espèces nouvelles (suite et fin). *Boletim Cultural da Guiné Portuguesa* 18(69): 13–106.
- Tendeiro, J. 1964: Mallophaga. *Annales du Musée Royal de l'Afrique Centrale, Tervuren* 132: 161–216, 13 pls.
- Tendeiro, J. 1965a [1962]: Estudos sobre malófagos. Revisão monográfica do género *Columbicola* Ewing (Ischnocera, Philopteridae). *Memorias da Junta de Investigações do Ultramar* (Serie 2) 32: 1–462, 175 pls
- Tendeiro, J. 1965b: Nouvelles études sur les Mallophages africains. *Revista dos Estudos Gerais Universitários de Moçambique* (Series 4) 2: 1–155.
- Tendeiro, J. 1965c: Études sur les Mallophages Parasites des Alcédinidés. I. Genres *Alcedoecus* Th. Clay et Meinertzhagen, 1939 et *Emersoniella* nov. *Revista dos Estudos Gerais Universitários de Moçambique* (Series 4) 2: 1–92.
- Tendeiro, J. 1965d: Études sur les Mallophages. Observations sur le genre *Pseudomenopon* Mjöberg, 1910, avec description de six espèces et une sous-espèce nouvelles. *Revista dos Estudos Gerais Universitários de Moçambique* (Series 4) 2: 1–88.
- Tendeiro, J. 1972a: Estudos sobre os goniodídeos. (Mallophaga, Ischnocera) dos Columbiformes. VIII— Uma nova espécie para o género *Nitzschiella* Kéler, 1939: *N. hemiphagae* n. sp., parasita da *Hemiphaga novaeseelandiae spadicea* (Latham). *Revista de Ciências Veterinárias (Lourenço Marques)* (Series A) 5: 1–15.
- Tendeiro, J. 1972b: Estudos sobre os goniodídeos. (Mallophaga, Ischnocera) dos Columbiformes. X— Descrição do género *Patellinirmus* nov., para três espécies novas parasitas de columbídeos do género *Hemiphaga* Bonaparte. *Revista de Ciências Veterinárias (Lourenço Marques)* (Series A) 5: 55–99.
- Tendeiro, J. 1973: Estudos sobre os goniodídeos. (Mallophaga, Ischnocera) dos Columbiformes. XIV— Género *Coloceras* Taschenberg, 1882. *Revista de Ciências Veterinárias (Lourenço Marques)* (Series A) 6: 199–524.
- Tendeiro, J. 1978: Estudos sobre os goniodídeos (Mallophaga, Ischnocera) dos Columbiformes. XVII— Novas observações sobre os géneros *Campanulotes* Kéler, *Saussurites* Tendeiro e *Stephanius* Tendeiro. *Garcia de Orta (Série Zoologia)* 7(1–2): 115–124, 6 pls.
- Tendeiro, J. 1980: Contributions a l'étude des mallophages des Columbiformes africains. I. Description de quatre nouvelles especes du genre *Hohorstiella* Eichler (Mallophaga, Amblycera), parasites de Columbiformes. *Annales Musee Royal de l'Afrique Centrale (Serie In-8) Sciences Zoologiques* 232: 1–21.
- Tendeiro, J. 1988: Etudes sur les *Colpocephalum* (Mallophaga, Menoponidae) parasites des Falconiformes I. Groupe *zerafae* Price & Beer. *Bonner zoologische Beiträge* 39(2/3): 77–102.
- Tennyson, A.J.D. 1986: Second record of a Manx shearwater in New Zealand. *Notornis* 33(1): 59–61.

- Tennyson, A.J.D. 1997: First record of a black-faced monarch (*Monarcha melanopsis*) in New Zealand. *Notornis* 44(4): 267–269.
- Tennyson, A.J.D. & Brackenbury, G. 1998: Channel-billed cuckoos in New Zealand in spring 1996. *Notornis* 45(3): 223–226.
- Tenorio, J.M. 1979: Catalog of entomological types in the Bishop Museum. Mallophaga. *Pacific Insects* 20(1): 5–17.
- Tenquist, J.D. 1977: Some ectoparasites of veterinary importance. *The New Zealand Entomologist* 6(3): 285–288.
- Tenquist, J.D. & Charleston, W.A.G. 1981: An annotated checklist of ectoparasites of terrestrial mammals in New Zealand. *Journal of the Royal Society of New Zealand* 11(3): 257–285.
- Tenquist, J.D. & Charleston, W.A.G. 2001: A revision of the annotated checklist of ectoparasites of terrestrial mammals in New Zealand. *Journal of the Royal Society of New Zealand* 31(3): 481–542.
- Thomas, P.L. 1958: The control of lice (*Damalinia ovis*) and keds (*Melophagus ovinus*) – A comparison of the efficiency of common insecticides. *New Zealand Journal of Agriculture Research* 1(2): 217–223.
- Thompson, G.B. 1935a: Notes on Mallophaga. II. *Annals and Magazine of Natural History* (Series 10) 15: 555–558.
- Thompson, G.B. 1935b: Preliminary descriptions of three new genera of Mallophaga (Subfamily Esthiopterinae). *Annals and Magazine of Natural History* (Series 10) 16: 148–151.
- Thompson, G.B. 1935c: A new genus of Mallophaga allied to *Pseudonirmus* Mjöberg, together with a note on the genus *Naubates* Bedford (subfamily Esthiopterinae). *Annals and Magazine of Natural History* (Series 10) 16: 485–488.
- Thompson, G.B. 1936: Three new genera of Mallophaga (Subfam. Esthiopterinae). *Annals and Magazine of Natural History* (Series 10) 18: 40–43.
- Thompson, G.B. 1937–1939: *The Piaget Collection of Mallophaga*.— Parts I–VIII. *Annals and Magazine of Natural History* (Series 10) 20: 19–27, 1 pl. *Ditto* (Series 11) 1: 268–271, 493–496; 2: 339–343, 607–611; 3: 417–421; 4: 139–144, 471–472.
- Thompson, G.B. 1937a: A list of the Denny collection of Mallophaga in the British Museum (Natural History) and their hosts. *Annals and Magazine of Natural History* (Series 10) 19: 74–81.
- Thompson, G.B. 1937b: A new genus and species of Mallophaga (Sub-family Esthiopterinae) from a petrel. *Annals and Magazine of Natural History* (Series 10) 20: 434–436.
- Thompson, G.B. 1938a: The Mallophaga (biting-lice) recorded from the Pacific Islands. *Entomologist's Monthly Magazine* 74: 184–189, 202–208.
- Thompson, G.B. 1938b: Mallophaga. *Résultats du voyage de la Belgica en 1897–99. Rapports scientifiques, Zoologie*, Mallophaga. 6 pp., 1 pl.
- Thompson, G.B. 1938c: On two species of Mallophaga from *Phaëthon rubricauda roseotincta* (Mathews). *Annals and Magazine of Natural History* (Series 11) 2: 458–464.
- Thompson, G.B. 1938d: The lice of petrels.— Part I. The elongate forms. *Annals and Magazine of Natural History* (Series 11) 2: 481–493.
- Thompson, G.B. 1939: The Mallophaga (biting-lice) recorded from the Pacific Islands. *Entomologist's Monthly Magazine* 75: 13–18, 71–76, 120–123, 209–218.
- Thompson, G.B. 1940a: Notes on species of the genus *Pectinopygus* (s. l.) (Mallophaga).— I. *Annals and Magazine of Natural History* (Series 11) 5: 372–381, 1 pl.
- Thompson, G.B. 1940b: The parasites of Australian birds and mammals. III. Mallophaga from sea-birds. *Annals and Magazine of Natural History* (Series 11) 5: 498–502, 1 pl.
- Thompson, G.B. 1940c: The lice of petrels.— Part III. The genus *Pelmatocerandra*. *Annals and Magazine of Natural History* (Series 11) 6: 103–111, 1 pl.
- Thompson, G.B. 1948a [1947]: Notes on species of the genus *Pectinopygus* (s. l.) – (Mallophaga). —IV. *Annals and Magazine of Natural History* (Series 11) 14: 317–327, 1 pl.
- Thompson, G.B. 1948b [1947]. Notes on species of the genus *Pectinopygus* (s. l.) – (Mallophaga). —V. *Annals and Magazine of Natural History* (Series 11) 14: 346–352.
- Thompson, G.B. 1948c: The lice of petrels. —Part IV. The genus *Episbates*. *Annals and Magazine of Natural History* (Series 11) 14: 661–671, 1 pl.
- Thompson, G.B. 1948d: Mallophaga collected by the Tanager Expedition. *Occasional Papers of Bernice P. Bishop Museum* 19(9): 195–200.
- Thomson, G.M. 1922: *The naturalisation of animals & plants in New Zealand*. Cambridge, U.K.: Cambridge University Press. x + 607 pp.
- Tillyard, R.J. 1926: *The Insects of Australia and New Zealand*. Sydney: Angus & Robertson, Ltd. xvi + 560 pp.
- Timmermann, G. 1936: *Saemundssonina* nov. gen., ein neues Mallophagengenus, aufgestellt für *Philopterus gonothorax* (Giebel) und verwandte Arten. *Zoologischer Anzeiger* 114(3/4): 97–100.

- Timmermann, G. 1949a: Beiträge zur Kenntnis der Ektoparasitenfauna isländischer Säugetiere und Vögel. 1. Mitteilung. Das Mallophagengenus *Saemundssonina* Timmermann, 1936. *Visindafélag Íslendinga* 2(3): 1–32.
- Timmermann, G. 1949b: Beiträge zur Kenntnis der Ektoparasitenfauna isländischer Säugetiere und Vögel. 2. Mitteilung. Das Mallophagengenus *Koeniginirmus* Eichler, 1940. *Visindafélag Íslendinga* 2(3): 83–88.
- Timmermann, G. 1950: Beiträge zur Kenntnis der Ektoparasitenfauna isländischer Säugetiere und Vögel. 4. Mitteilung. Die Gattung *Quadriceps* Clay & Meinertzhagen, 1939 und verwandte Genera Mallophagorum. *Fauna Islandica. Miscellaneous papers on Icelandic Zoology* 2: 1–8.
- Timmermann, G. 1951a: Die Mövenkneifer. Eine Revision sämtlicher bei echten Möven schmarotzenden Federlinge der Gattung *Saemundssonina* Tim., 1936. *Parasitological News, Reykjavik* 2(1): 1–12.
- Timmermann, G. 1951b: Nýjungar um lúsaættkvíslina *Saemundssonina* Tim. *Náttúrufræðingurinn* 21: 140–143.
- Timmermann, G. 1951c: Investigations on some ischnoceran bird lice (genus *Saemundssonina*) parasitic on waders. *Annals and Magazine of Natural History (Series 12)* 4: 390–401.
- Timmermann, G. 1952a: Revision der bei Seeschwalben schmarotzenden Kletterfederlinge der Gattung *Quadriceps* (Clay u. Meinertzhagen). *Zoologischer Anzeiger* 148: 71–87.
- Timmermann, G. 1952b: The species of the genus *Quadriceps* (Mallophaga) from the Larinae, with some remarks on the systematics and the phylogeny of the gulls. Part I. *Annals and Magazine of Natural History (Series 12)* 5(51): 209–222, 1 pl.
- Timmermann, G. 1952c: The species of the genus *Quadriceps* (Mallophaga) from the Larinae, with some remarks on the systematics and the phylogeny of the gulls. Part II. *Annals and Magazine of Natural History (Series 12)* 5(52): 595–600.
- Timmermann, G. 1952d: New and little-known species of *Quadriceps* (Mallophaga) from pratincoles, coursers and other plover-like birds. *Annals and Magazine of Natural History (Series 12)* 5: 1025–1037.
- Timmermann, G. 1953a: Die Federläuse des Säbelschnäblers. *Bombus – Faunistische Mitteilungen aus Nordwestdeutschland* 78/79: 329–333.
- Timmermann, G. 1953b: Die *Quadriceps*-Arten (Mallophaga) der Regenpfeifer (Unterfamilie Charadriinae). *Zoologischer Anzeiger* 150(7/8): 178–190.
- Timmermann, G. 1954a: A revision of the genus *Carduiceps* Clay & Meinertzhagen, 1939 (Mallophaga). *Annals and Magazine of Natural History (Series 12)* 7: 40–48.
- Timmermann, G. 1954b: Studies on Mallophaga from the collections of the British Museum (Nat. Hist.), London. 1. A preliminary survey of the genus *Luniceps* (Clay & Meinertzhagen), 1939. *Annals and Magazine of Natural History (Series 12)* 7: 623–637.
- Timmermann, G. 1954c: Studien über Mallophagen aus den Sammlungen des Britischen Museums (Nat. Hist.), London. II. Das Amblycerengenus *Actornithophilus* Ferris, 1916. *Annals and Magazine of Natural History (Series 12)* 7: 829–841, pl. 26.
- Timmermann, G. 1954d: Vorläufige Übersicht über das Amblyceren-Genus *Austromenopon* Bedford, 1939 (Mallophaga). *Bonner zoologische Beiträge* 5(3/4): 195–206.
- Timmermann, G. 1954e: Die *Quadriceps*-Arten (Mallophaga) der Kiebitze. *Zeitschrift für Parasitenkunde* 16: 195–208.
- Timmermann, G. 1954f: Neue and wenig bekannte Kletterfederlinge von charadriiformen Wirten. *Zoologischer Anzeiger* 152(7/8): 163–177.
- Timmermann, G. 1955: Studien über Mallophagen aus den Sammlungen des Britischen Museums (Nat. Hist.), London. 3. Mitteilung. *Annals and Magazine of Natural History (Series 12)* 8: 513–534.
- Timmermann, G. 1956: *Quadriceps niethammeri* n. sp. und andere neue Federlinge aus den Gattungen *Quadriceps*, *Saemundssonina* und *Austromenopon*. *Bonner zoologische Beiträge* 7(1/3): 186–192.
- Timmermann, G. 1957a: Studien zu einer vergleichenden Parasitologie der Charadriiformes oder Regenpfeifervögel. Teil 1: Mallophaga. *Parasitologische Schriftenreihe* 8: 1–204, 15 pls.
- Timmermann, G. 1957b: Mallophaga from Tristan da Cunha. Part II. Some remarks on the genus *Longimenopon* Thompson, 1948. *Results of the Norwegian Scientific Expedition to Tristan da Cunha 1937–1938* 41: 7–12.
- Timmermann, G. 1959a: Drei neue Sturmvogelfederlinge. *Zoologischer Anzeiger* 162: 148–153.
- Timmermann, G. 1959b: Taxonomie und hospitale Verbreitung der Mallophagengattung *Trabeculus* Rudow, 1866. *Zeitschrift für Parasitenkunde* 19: 485–502.
- Timmermann, G. 1959c: Zur Kenntnis der Gattung *Docophoroides* Giglioli, 1864 (Insecta, Mallophaga). *Mitteilungen aus dem Zoologischen Museum in Berlin* 35(1): 57–72.
- Timmermann, G. 1960: Gruppen-Revisionen bei Mallophagen. II. Genus *Halipeurus* Thompson 1936. 1. Teil: Die *Halipeurus*-Arten der “gadfly-petrels” (Genera *Pterodroma* und *Bulweria*). *Zeitschrift für Parasitenkunde* 20: 317–334.
- Timmermann, G. 1961a: Gruppen-Revisionen bei Mallophagen. II. Genus *Halipeurus* Thompson 1936. 2. Teil: Die *Halipeurus*-Arten der Wasserscherer (Puffininae), Sturmtaucher (Pelecanoididae) und Sturmschwalben (Hydrobatidae). *Zeitschrift für Parasitenkunde* 20: 401–419.

- Timmermann, G. 1961b: Gruppen-Revisionen bei Mallophagen. III. Genus *Philoceanus* Kellogg, 1903. *Zeitschrift für Parasitenkunde* 20: 525–537.
- Timmermann, G. 1961c: Gruppen-Revisionen bei Mallophagen. IV. Genera *Pseudonirmus* Mjöberg, 1910, *Bedfordiella* Thompson, 1937, und *Episbates* Harrison, 1935. *Zeitschrift für Parasitenkunde* 21: 30–45.
- Timmermann, G. 1961d: Gruppen-Revisionen bei Mallophagen. I. Genus *Naubates* Bedford, 1930. *Zoologischer Anzeiger* 166(5/6): 173–191, pl. 1.
- Timmermann, G. 1962a: Neue Zangenläuse (Mallophaga, Philopteridae) von procellariiformen und charadriiformen Wirten. *Zeitschrift für Parasitenkunde* 21: 426–436.
- Timmermann, G. 1962b: Gruppen-Revisionen bei Mallophagen. V. Zur näheren Kennzeichnung des *Ornithobius*-Komplexes (Philopteridae), parasitisch bei Entenvögeln. *Zeitschrift für Parasitenkunde* 22: 133–147.
- Timmermann, G. 1963: Gruppen-Revisionen bei Mallophagen. VI. Die *Austromenopon*-Arten der Sturmvögel. *Zeitschrift für Parasitenkunde* 22(5): 401–427.
- Timmermann, G. 1964: Gruppen-Revisionen bei Mallophagen. VII. Die *Pectinopygus*-Arten der Großkormorane (Gen. *Phalacrocorax* Brisson, 1760 s. str.). *Mitteilungen aus den Hamburgischen Zoologischen Museum und Institut (Kosswig-Festschrift)*: 271–284, 1 pl.
- Timmermann, G. 1965: Die Federlingsfauna der Sturmvögel und die Phylogenese des procellariiformen Vogelstammes. *Abhandlungen und Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg, N. F. 8 (Supplement)*: 1–249, 12 pls.
- Timmermann, G. 1966: *Haffneria* nov. gen., ein neues Mallophagengenus von Raubmöwen. *Mitteilungen aus den Hamburgischen Zoologischen Museum und Institut* 63: 85–89.
- Timmermann, G. 1969a: Gruppen-Revisionen bei Mallophagen. VIII. Die Formenkreise *Saemundssonina scolopacisphaeopodis* (Schrank), 1803, *Saemundssonina platygaster* (Denny), 1842 und *Saemundssonina africana* Timmermann, 1951. *Zoologischer Anzeiger* 183(3/4): 225–256.
- Timmermann, G. 1969b: Neue Mallophagen aus dem Bernice P. Bishop Museum, Honolulu. *Bonner zoologische Beiträge* 20(1/3): 244–252.
- Timmermann, G. 1969c: Mallophagologische Kollektaneen. *Abhandlungen und Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg, Neue Folge* 13: 195–200.
- Timmermann, G. 1971: “Regelwidrigkeiten” im Ausbreitungsbild bei Vogelmallophagen (erläutert an den Kletterfederlingen der Watvögel und Möwen). *Mitteilungen aus den Hamburgischen Zoologischen Museum und Institut* 67: 135–174.
- Timmermann, G. 1977: Mallophagologische Kollektaneen. 3. *Bonner zoologische Beiträge* 28(1/2): 135–140.
- Tombesi, M.L. & Castro, D. del C. 1995: Redescrición de los estadios ninfales de *Haematopinus suis* (Linnaeus, 1758) y su caracterización mediante un análisis multivariado (Phthiraptera, Anoplura). *Revista Brasileira de Entomologia* 39(3): 653–660.
- Toon, A. & Hughes, J.M. 2008: Are lice good proxies for host history? A comparative analysis of the Australian magpie, *Gymnorhina tibicen*, and two species of feather louse. *Heredity* 101: 127–135.
- Trainor, C.R. 2002: *A preliminary list of important bird areas in East Timor*. Bogor: BirdLife International – Asia Programme. 26 pp.
- Trouessart, E.L. & Neumann, L.G. 1888: Le pou de l’otarie (*Echinophthirius microchir*, n. sp.). *Le Naturaliste, Paris* 10: 80–81.
- Tsurumi, M. 1989: Ectoparasites (Mallophaga and Siphonaptera: Insecta) from the Streaked Shearwater *Calonectris leucomelas* collected on Toshima Island, Tokyo. *Journal of the Yamashina Institute for Ornithology* 21: 280–282.
- Tuff, D.W. 1966: Notes on *Eucolpocephalum femorale* (Mallophaga: Menoponidae). *Journal of the Kansas Entomological Society* 39(3): 379–382.
- Tuff, D.W. 1967: A review of North American *Ardeicola* (Mallophaga: Philopteridae). *Journal of the Kansas Entomological Society* 40(2): 241–263.
- Tuff, D.W. 1970: A new synonym of *Ardeicola expallida* Blagoveshtchensky (Mallophaga: Philopteridae). *Journal of Medical Entomology* 7(4): 484.
- Uchida, S. 1926: Studies on amblycerous Mallophaga of Japan. *Journal of the College of Agriculture, Tokyo* 9(1): 1–56.
- Uchida, S. 1948: Studies on the biting-lice (Mallophaga) of Japan and adjacent territories (suborder Ischnocera Pt. I). *The Japanese Medical Journal* 1(4): 303–326.
- Uchida, S. 1949: Studies on the biting-lice (Mallophaga) of Japan and adjacent territories (suborder Ischnocera Pt. II). *The Japanese Medical Journal* 1(6): 535–556.
- Valette, L.H. 1913: Apuntes descriptivos sobre algunos invertebrados encontrados en un viaje a las Islas Orcadas. *Boletín del Ministerio de Agricultura, República Argentina* 15(3): 293–318, 17 pls.
- Valim, M.P. 2009: Type specimens of lice (Insecta: Phthiraptera) held in the Museu de Zoologia da Universidade de São Paulo, Brazil. *Papéis Avulsos de Zoologia* 49(17): 197–219.


- Valim, M.P. & Palma, R.L. 2013: Three new species of the genus *Philopterooides* Mey, 2004 (Phthiraptera, Ischnocera, Philopteridae) from New Zealand. *ZooKeys* 297: 71–89.
- Valim, M.P. & Palma, R.L. 2015: A new genus and two new species of feather lice (Phthiraptera: Ischnocera: Philopteridae) from New Zealand endemic passerines (Aves: Passeriformes). *Zootaxa* 3926(4): 480–498.
- Vermeulen, E.T.; Power, M.L.; Nipperess, D.A.; Beveridge, I. & Eldridge, M.D.B. 2016: Biodiversity of parasite assemblages in the genus *Petrogale* and its relation to the phylogeny and biogeography of their hosts. *Australian Journal of Zoology* 64(1): 61–80.
- Voss, W.J. 1966: A lectotype designation for *Hoplopleura pacifica* Ewing (Anoplura: Hoplopleuridae). *Pacific Insects* 8(1): 29–32.
- Ward, R.A. 1955: Biting lice of the genus *Saemundssonina* (Mallophaga: Philopteridae) occurring on terns. *Proceedings of the United States National Museum* 105(3353): 83–100.
- Ward, R.A. & Downey, J.C. 1973: Checklist of the Mallophaga of Midway Atoll, Pacific Ocean. *Journal of Medical Entomology* 10(4): 391–396.
- Waterston, J. 1912a: On *Mackayia dimorpha*, a new genus and species of Mallophaga from the Manx shearwater. *Scottish Naturalist*. 11: 251–258.
- Waterston, J. 1912b: A new species of *Mackayia* (*M. heteracanthus*). *Scottish Naturalist*. 11: 258.
- Waterston, J. 1914: On some ectoparasites in the South African Museum, Cape Town. *Annals of the South African Museum* 10(9): 271–324, 2 pls.
- Waterston, J. 1915: On two new species of Mallophaga (Menoponidae): *Menacanthus balfouri* n. sp. and *Myrsidea victrix* n. sp. from Colombia. *Entomologist's Monthly Magazine* 51: 12–16, pl. 1.
- Waterston, J. 1917: On a new species of *Docophoroides* Gigl. (*Eurymetopus* Tasch.) from an albatross (*Diomedea melanophrys*). *Entomologist's Monthly Magazine* (Series 3) 3: 99.
- Waterston, J. 1922: A new genus of Ischnocera (Mallophaga). *Entomologist's Monthly Magazine* 58: 159.
- Watson, K.C. 1967: The terrestrial Arthropoda of Macquarie Island. *ANARE Scientific Reports, series B (I) Zoology* 99: xii + 90 pp., 2 maps, 12 pls.
- Watson, T.G.; Bishop, D.M.; Hooke, P.G.; Heath, A.C.G. & Cole, D.J.W. 1996: Efficacy of injectable doramectin against naturally acquired louse infestations on cattle. *New Zealand Journal of Agricultural Research* 39: 401–404.
- Watt, J.C. 1971: Ectoparasitic insects of birds and mammals of the Kermadec Islands. *Notornis* 18(4): 227–244.
- Webb, J.E. 1946: Spiracle structure as a guide to the phylogenetic relationships of the Anoplura (biting and sucking lice), with notes on the affinities of the mammalian hosts. *Proceedings of the Zoological Society of London* 116: 49–119.
- Webb, J.E. 1948: Siphunculata of the genus *Haematopinus* Leach infesting Equidae, with a description of a new subspecies of *Haematopinus asini* (L.) from a zebra. *Proceedings of the Zoological Society of London* 118: 578–581, pl. 1.
- Webb, J.E. 1949: The evolution and host-relationships of the sucking lice of the Ferungulata. *Proceedings of the Zoological Society of London* 119(1): 133–188.
- Wegner, Z. 1974: A morphological analysis of *Polyplax serrata* (Burmeister, 1839) (Arthropoda, Anoplura). *Acta Parasitologica Polonica* 22(18): 203–217.
- Weidner, H. 1966: Die Entomologischen Sammlungen des Zoologischen Staatsinstituts und Zoologischen Museums Hamburg. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 63: 209–264.
- Weidner, H. 1977: Die Entomologischen Sammlungen des Zoologischen Instituts und Zoologischen Museums der Universität Hamburg. XIV. (letzter) Teil. Insecta XI. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 74: 77–138.
- Werneck, F.L. 1936: Contribuição ao conhecimento dos mallophagos encontrados nos mamíferos sul-americanos. *Memórias do Instituto Oswaldo Cruz*, Rio de Janeiro 31(3): 391–590, 1 pl.
- Werneck, F.L. 1950: *Os Malófagos de Mamíferos*. Parte II: Ischnocera (continuação de Trichodectidae) e Rhyncophthirina. Rio de Janeiro: Instituto Oswaldo Cruz. 207 pp.
- Westwood, J.O. 1874: *Thesaurus entomologicus Oxoniensis*. Oxford: Clarendon Press. xxiv + 205 pp., 40 pls.
- White, A. & Doubleday, E. 1843: Annulose animals. Pp. 265–291. In: Dieffenbach, E. *Travels in New Zealand; with contributions to the geography, geology, botany, and natural history of that country*. London: John Murray. Volume 2. iv + 396 pp.
- Whiteman, N.K.; Dosanjh, V.S.; Palma, R.L.; Hull, J.M.; Kimball, R.T.; Sánchez, P.; Sarasola, J.H. & Parker, P.G. 2009: Molecular and morphological divergence in a pair of bird species and their ectoparasites. *Journal of Parasitology* 95(6): 1372–1382.
- Whitten, L.K. 1970: The control of external parasites of cattle. *New Zealand Veterinary Journal* 18(7): 146–147.
- Whitten, L.K. 1971: *Diseases of domestic animals in New Zealand*. Wellington: Editorial Services Limited. 439 pp.

- Wigglesworth, V.B. 1932: The hatching organ of *Lipeurus columbae* Linn. (Mallophaga), with a note on its phylogenetic significance. *Parasitology* 24(3): 365–367.
- Williams, R.T. 1970a: In vitro studies on the environmental biology of *Goniodes colchici* (Denny) (Mallophaga: Ischnocera). I. The effects of temperature and humidity on the bionomics of *G. colchici*. *Australian Journal of Zoology* 18: 379–389.
- Williams, R.T. 1970b: In vitro studies on the environmental biology of *Goniodes colchici* (Denny) (Mallophaga: Ischnocera). II. The effects of temperature and humidity on water loss. *Australian Journal of Zoology* 18: 391–398.
- Williams, R.T. 1971: In vitro studies on the environmental biology of *Goniodes colchici* (Denny) (Mallophaga: Ischnocera). III. The effects of temperature and humidity on the uptake of water vapour. *Journal of Experimental Biology* 55: 553–568.
- Wilson, F.H. 1933: A louse feeding on the blood of its host. *Science* 77(2003): 490.
- Wilson, J.A.; Heath, A.C.G.; Quilter, S.; McKay, C.; Litchfield, D & Nottingham, R. 1997: A preliminary investigation into resistance to synthetic pyrethroids by the sheep biting louse (*Bovicola ovis*) in New Zealand. *New Zealand Veterinary Journal* 45: 8–10.
- Windsor, D.A. 1995: Equal rights for parasites. *Conservation Biology* 9: 1–2.
- Wise, K.A.J. 1977: A synonymic checklist of the Hexapoda of the New Zealand Sub-region. The smaller orders. *Bulletin of the Auckland Institute and Museum* 11: iv + 1–76.
- Wood, L.M. 1937: in Harrison, L. Mallophaga and Siphunculata. 2(1): 1–47, pls 1–3. In: Johnston, T.H. (ed.), *Scientific reports, Australasian Antarctic Expedition 1911–14, series C — zoology and botany*. Sydney: Government Printer.
- Wundrig, G. 1936: Die Sehorgane der Mallophagen, nebst vergleichenden Untersuchungen an Liposceliden und Anopluren. *Zoologische Jahrbücher Abteilung für Anatomie und Ontogenie der Tiere* 62: 45–110.
- Yamagishi, A.; Yao, I.; Johnson, K.P. & Yoshizawa, K. 2014: Comparisons of host specificity in feather louse genera (Insecta: Phthiraptera: Philopteridae) parasitizing gulls (Aves: Laridae: *Larus*). *Zoological Science (Tokyo)* 31(6): 383–389.
- Yoshizawa, K. & Johnson, K.P. 2003: Phylogenetic position of Phthiraptera (Insecta: Paraneoptera) and elevated rate of evolution in mitochondrial 12S and 16S rDNA. *Molecular Phylogenetics and Evolution* 29: 102–114.
- Yoshizawa, K. & Johnson, K.P. 2006: Morphology of male genitalia in lice and their relatives and phylogenetic implications. *Systematic Entomology* 31: 350–361.
- Yoshizawa, K. & Johnson, K.P. 2010: How stable is the “Polyphyly of Lice” hypothesis (Insecta: Psocodea)? A comparison of phylogenetic signal in multiple genes. *Molecular Phylogenetics and Evolution* 55: 939–951.
- Zimmer, C. 2000: *Parasite rex: inside the bizarre world of Nature's most dangerous creatures*. London: Arrow Books. xxii + 298 pp.
- Złotorzycka, J. 1964a: Mallophaga parasitizing Passeriformes and Pici. II. Brueeliinae. *Acta Parasitologica Polonica* 12(24): 239–282, 4 pls.
- Złotorzycka, J. 1964b: Mallophaga parasitizing Passeriformes and Pici. III. Philopterinae. *Acta Parasitologica Polonica* 12(37): 401–430, 4 pls.
- Złotorzycka, J. 1967: Studien über *Quadriceps* s. l. (Mallophaga, Quadraceptinae). Übersicht der Arten und systematische Revision mit besonderer Berücksichtigung der synhospitalen und allohospitalen Arten. *Polskie Pismo Entomologiczne* 37(4): 705–785, 17 pls.
- Złotorzycka, J. 1970: Studien an den mitteleuropäischen Arten der Gattung *Anatoecus* Cumm. (Esthiopteridae, Mallophaga). *Polskie Pismo Entomologiczne* 40(1): 7–67, 6 pls.
- Złotorzycka, J. & Lucińska, A. 1967: Über den Federling *Bonomiella columbae* Emers. (Mallophaga, Somaphantidae) aus Polen. *Polskie Pismo Entomologiczne* 37(2): 341–348.
- Złotorzycka, J. & Modrzejewska, M. 1988: Wszczoły — Mallophaga. *Katalog Fauny Polski. Catalogus faunae Poloniae* 19(1): 1–223.
- Zonfrillo, B. 1993: Relationships of the *Pterodroma* petrels from the Madeira archipelago inferred from their feather lice. *Boletim do Museu Municipal do Funchal, Supplement* 2: 325–331.





**Figure 1:** *Boopia notafusca* Le Souëf, 1902. Male.



**Figure 2:** *Boopia notafusca*. Nymph.



**Figure 3:** *Heterodoxus ampullatus* Kéler, 1971. Male.



**Figure 4:** *Heterodoxus ampullatus*. Female.



**Figure 5:** *Gliricola (Gliricola) porcelli* (Schrank, 1781). Male.



**Figure 6:** *Gliricola (Gliricola) porcelli*. Female.



**Figure 7:** *Gyropus ovalis* Burmeister, 1838. Male.



**Figure 8:** *Gyropus ovalis*. Female.



**Figure 9:** *Laemobothrion (Laemobothrion) tinnunculi* (Linnaeus, 1758). Female.



**Figure 10:** *Actornithophilus ceruleus* (Timmermann, 1954). Male. **Figure 11:** *Actornithophilus ceruleus*. Female.



**Figure 12:** *Amyrsidea (Argimenopon) minuta* Emerson, 1961. Male. **Figure 13:** *Amyrsidea (A.) minuta* Female.




**Figure 14:** *Ancistrona vagelli* (J.C. Fabricius, 1787). Male.

**Figure 15:** *Ancistrona vagelli*. Female.



**Figure 16:** *Apterygon okarito* Palma & Price, 2004. Male.




**Figure 17:** *Apterygon okarito*. Female.



**Figure 18:** *Austromenopon navigans* (Kellogg, 1896). Male.



**Figure 19:** *Austromenopon navigans*. Female.



**Figure 20:** *Bonomiella columbae* Emerson, 1957. Male.

**Figure 21:** *Bonomiella columbae*. Female.



**Figure 22:** *Ciconiphilus decimfasciatus* (Boisduval & Lacordaire, 1835). Male. **Figure 23:** *Ciconiphilus decimfasciatus*. Female.



**Figure 24:** *Colpocephalum pilgrimi* Price, 1967. Male.




**Figure 25:** *Colpocephalum pilgrimi*. Female.



**Figure 26:** *Cuculiphilus (Cuculiphilus) fasciiventris* Carriker, 1955. Male.



**Figure 27:** *Cuculiphilus (C.) fasciiventris*. Female.




**Figure 28:** *Dennyus (Takamatsuia)* sp. Nymph.




**Figure 29:** *Eidmanniella eurygaster* (Nitzsch [in Giebel], 1866). Male. **Figure 30:** *Eidmanniella eurygaster*. Female.






**Figure 31:** *Eucolpocephalum femorale* (Piaget, 1880). Male.




**Figure 32:** *Eucolpocephalum femorale*. Female.




**Figure 33:** *Franciscocola (Franciscocola) pallida* (Piaget, 1880). Male.



**Figure 34:** *Franciscocola (Franciscocola) pallida*. Female.



**Figure 35:** *Heteromenopon (Keamenopon) kea* (Kellogg, 1907). Male. **Figure 36:** *Heteromenopon (K.) kea*. Female.



**Figure 37:** *Hohorstiella timorensis* Tendeiro, 1980. Male.

**Figure 38:** *Hohorstiella timorensis*. Female.



**Figure 39:** *Holomenopon tadornae* (Gervais, 1844). Male.



**Figure 40:** *Holomenopon tadornae*. Female.



**Figure 41:** *Kurodaia (Conciella) cryptostigmatia* (Nitzsch [in Giebel], 1861). Male.




**Figure 42:** *Kurodaia cryptostigmatia*. Female.




**Figure 43:** *Longimenopon* sp. Male.




**Figure 44:** *Longimenopon* sp. Female.



**Figure 45:** *Menacanthus pallidulus* (Neumann, 1912). Male.



**Figure 46:** *Menacanthus pallidulus*. Female.



**Figure 47:** *Menopon gallinae* (Linnaeus, 1758). Male.




**Figure 48:** *Menopon gallinae*. Female.




**Figure 49:** *Myrsidea thoracica* (Giebel, 1874). Male




**Figure 50:** *Myrsidea thoracica*. Female.




**Figure 51:** *Nosopon lucidum* (Rudow, 1869). Male.




**Figure 52:** *Nosopon lucidum*. Female.



**Figure 53:** *Plegadiphilus plegadis* (Dubinin, 1938). Male.



**Figure 54:** *Plegadiphilus plegadis*. Female.



**Figure 55:** *Pseudomenopon pilgrimi* Price, 1974. Male.




**Figure 56:** *Pseudomenopon pilgrimi*. Female.




**Figure 57:** *Trinoton nigrum* Le Souëf, 1902. Male.



**Figure 58:** *Trinoton nigrum*. Female.




**Figure 59:** *Ricinus* sp. Female.



**Figure 60:** *Acidoproctus gottwaldhirschi* Eichler, 1958. Male.

**Figure 61:** *Acidoproctus gottwaldhirschi*. Female.





**Figure 62:** *Alcedoecus alatoclypeatus* (Piaget, 1885). Male.



**Figure 63:** *Alcedoecus alatoclypeatus*. Female.



**Figure 64:** *Anaticola magnificus* Ansari, 1955. Male.




**Figure 65:** *Anaticola magnificus*. Female.




**Figure 66:** *Anatoecus icterodes oloris* Zlotorzycska, 1970. Male.

**Figure 67:** *Anatoecus icterodes oloris*. Female.




**Figure 68:** *Aquanirmus australis* Kettle, 1974. Male.

**Figure 69:** *Aquanirmus australis*. Female.



**Figure 70:** *Ardeicola neopallidus* Price, Hellenthal & Palma, 2003. Male **Figure 71:** *Ardeicola neopallidus*. Female.



**Figure 72:** *Austrogoniodes hamiltoni* Harrison, 1937. Male

**Figure 73:** *Austrogoniodes hamiltoni*. Female.



**Figure 74:** *Bedfordiella unica* Thompson, 1937. Male.




**Figure 75:** *Bedfordiella unica*. Female.



**Figure 76:** *Brueelia semiannulata* (Piaget, 1883). Male.




**Figure 77:** *Brueelia semiannulata*. Female.



**Figure 78:** *Campanulotes bidentatus compar* (Burmeister, 1838). Male. **Figure 79:** *Campanulotes bidentatus compar*. Female.



**Figure 80:** *Capraiella* sp. Female.



**Figure 81:** *Carduiceps cingulatus cingulatus* (Denny, 1842). Male. **Figure 82:** *Carduiceps cingulatus cingulatus*. Female.



**Figure 83:** *Chelopistes meleagridis* (Linnaeus, 1758). Male.

**Figure 84:** *Chelopistes meleagridis*. Female.



**Figure 85:** *Coloceras novaeseelandiae* (Tendeiro, 1972). Male.



**Figure 86:** *Coloceras novaeseelandiae*. Female.



**Figure 87:** *Columbicola columbae columbae* (Linnaeus, 1758). Male.



**Figure 88:** *Columbicola columbae columbae*. Female.



**Figure 89:** *Cuclotogaster synoicus* (Clay, 1938). Male.



**Figure 90:** *Cuclotogaster synoicus*. Female.



**Figure 91:** *Cuculicola kui* Kettle, 1980. Male.



**Figure 92:** *Cuculicola kui*. Female.





**Figure 93:** *Degeeriella fusca* (Denny, 1842). Male.




**Figure 94:** *Degeeriella fusca*. Female.



**Figure 95:** *Docophoroides simplex* (Waterston, 1914). Male.



**Figure 96:** *Docophoroides simplex*. Female.



**Figure 97:** *Emersoniella bracteata* (Nitzsch [in Giebel], 1866). Male. **Figure 98:** *Emersoniella bracteata*. Female.




**Figure 99:** *Episbates pederiformis* (Dufour, 1835). Male.

**Figure 100:** *Episbates pederiformis*. Female.




**Figure 101:** *Forficuloecus pilgrimi* Guimarães, 1985. Male **Figure 102:** *Forficuloecus pilgrimi*. Female.




**Figure 103:** *Fulicoffula stammeri* Eichler, 1958. Male.

**Figure 104:** *Fulicoffula stammeri*. Female.



**Figure 105:** *Goniocotes pusillus* (Nitzsch [in Giebel], 1866). Male. **Figure 106:** *Goniocotes pusillus*. Female.



**Figure 107:** *Goniodes retractus* Le Souëf, 1902. Male.

**Figure 108:** *Goniodes retractus*. Female.



**Figure 109:** *Haffneria grandis* (Piaget, 1880). Male.



**Figure 110:** *Haffneria grandis*. Female.



**Figure 111:** *Halipeurus confusus* Palma, 2011. Male.



**Figure 112:** *Halipeurus confusus*. Female.



**Figure 113:** *Harrisoniella ferox* (Giebel, 1867). Male.



**Figure 114:** *Harrisoniella ferox*. Female.



**Figure 115:** *Idoeacus diana* Tandan, 1958. Male.



**Figure 116:** *Idoeacus diana*. Female.




**Figure 117:** *Incidifrons porzanae* Blagoveshtchensky, 1951. Male.

**Figure 118:** *Incidifrons porzanae*. Female.




**Figure 119:** *Lagopoecus docophoroides* (Piaget, 1880). Male.

**Figure 120:** *Lagopoecus docophoroides*. Female.




**Figure 121:** *Lipeurus maculosus maculosus* Clay, 1938. Male. **Figure 122:** *Lipeurus maculosus maculosus*. Female.



**Figure 123:** *Luniceps numenii madagascariensis*. Male.  
Gustafsson & Olsson, 2012

**Figure 124:** *Luniceps numenii madagascariensis*. Female.





**Figure 125:** *Melibruelia novaeseelandiae* Valim & Palma, 2015. Male.



**Figure 126:** *Melibruelia novaeseelandiae* . Female.



**Figure 127:** *Naubates (Guenterion) lessonii* Palma & Pilgrim, 2002. Male.



**Figure 128:** *Naubates (Guenterion) lessonii*. Female.



**Figure 129:** *Naubates (Naubates) thieli* Timmermann, 1965. Male. **Figure 130:** *Naubates (Naubates) thieli*. Female.



**Figure 131:** *Neopsittaconirmus kea* (Kellogg, 1907). Male.

**Figure 132:** *Neopsittaconirmus kea*. Female.



Figure 133: *Nesiotinus demersus* Kellogg, 1903. Male.



Figure 134: *Nesiotinus demersus*. Female.



Figure 135: *Ornithobius fuscus* Le Souëf, 1902. Male.



Figure 136: *Ornithobius fuscus*. Female.



**Figure 137:** *Oxylipeurus polytrapezius polytrapezius*. Male. (Burmeister, 1838)



**Figure 138:** *Oxylipeurus polytrapezius polytrapezius*. Female.



**Figure 139:** *Paraclisis miriceps* (Kellogg & Kuwana, 1902). Male.



**Figure 140:** *Paraclisis miriceps*. Female.




**Figure 141:** *Pectinopygus varius* Timmermann, 1964. Male. **Figure 142:** *Pectinopygus varius*. Female.



**Figure 143:** *Pelmatocerandra setosa* (Giebel, 1876). Male. **Figure 144:** *Pelmatocerandra setosa*. Female.



**Figure 145:** *Penenirmus* sp. Female.



**Figure 146:** *Perineus concinnoides* Kéler, 1957. Male.

**Figure 147:** *Perineus concinnoides*. Female.



**Figure 148:** *Philoceanus fasciatus* (Carriker, 1958). Male.



**Figure 149:** *Philoceanus fasciatus*. Female.




**Figure 150:** *Philopteroides fuliginosus* Valim & Palma, 2013. Male.



**Figure 151:** *Philopteroides fuliginosus*. Female.




**Figure 152:** *Philopterus novaezealandiae* Palma & Price, 2000. Male. **Figure 153:** *Philopterus novaezealandiae*. Female.




**Figure 154:** *Pseudonirmus lugubris* (Taschenberg, 1882). Male.

**Figure 155:** *Pseudonirmus lugubris*. Female.





**Figure 156:** *Psittoecus vanzolinii* Guimarães, 1974. Male.




**Figure 157:** *Psittoecus vanzolinii*. Female.




**Figure 158:** *Quadriceps caspius* (Giebel, 1874). Male.



**Figure 159:** *Quadriceps caspius*. Female.



**Figure 160:** *Rallicola (Aptericola) rodericki* Palma, 1991. Male **Figure 161:** *Rallicola (Aptericola) rodericki*. Female.



**Figure 162:** *Rallicola (Huiacola) extinctus* (Mey, 1990). Male. **Figure 163:** *Rallicola (Huiacola) extinctus*. Female.



**Figure 164:** *Rallicola (Rallicola) tabuensis* Emerson, 1966. Male.




**Figure 165:** *Rallicola (Rallicola) tabuensis*. Female.




**Figure 166:** *Saemundssonina (Puffinoecus) enderleini* (Eichler, 1949). Male.



**Figure 167:** *Saemundssonina (P.) enderleini*. Female.




**Figure 168:** *Saemundssonina (Saemundssonina) albatrossa* Palma, 2012. Male. **Figure 169:** *Saemundssonina (S.) albatrossa*. Female.



**Figure 170:** *Strigiphilus vapidus* Clay, 1977. Male.

**Figure 171:** *Strigiphilus vapidus*. Female.



**Figure 172:** *Sturnidoecus sturni* (Schrank, 1776). Male.



**Figure 173:** *Sturnidoecus sturni*. Female.




**Figure 174:** *Trabeculus fuscoclypeatus* (Johnston & Harrison, 1912). Male.



**Figure 175:** *Trabeculus fuscoclypeatus*. Female.



**Figure 176:** *Bovicola (Bovicola) longicornis* (Nitzsch, 1818). Male.




**Figure 177:** *Bovicola (B.) longicornis*. Female.



**Figure 178:** *Bovicola (Lepikentron) breviceps* (Rudow, 1866). Female.




**Figure 179:** *Bovicola (Spinibovicola) hemitragi* (Cummings, 1916). Male. **Figure 180:** *Bovicola (S.) hemitragi*. Female.




**Figure 181:** *Felicola (Felicola) subrostratus* (Burmeister, 1838). Male.

**Figure 182:** *Felicola (F.) subrostratus*. Female.




**Figure 183:** *Trichodectes (Trichodectes) canis* (De Geer, 1778). Male. **Figure 184:** *Trichodectes (T.) canis*. Female.



**Figure 185:** *Trichodectes (Stachiella) ermineae* (Hopkins, 1941). Male. **Figure 186:** *Trichodectes (S.) ermineae*. Female.







**Figure 187:** *Tricholipeurus lipeuroides* (Mégnin, 1884). Male. **Figure 188:** *Tricholipeurus lipeuroides*. Female.



**Figure 189:** *Werneckiella ocellata* (Piaget, 1880). Female.



**Figure 190:** *Antarctophthirus microchir* (Trouessart & Neumann, 1888). Male. **Figure 191:** *Antarctophthirus microchir*. Female.



**Figure 192:** *Lepidophthirus macrorhini* Enderlein, 1904. Male.


**Figure 193:** *Lepidophthirus macrorhini*. Female.



**Figure 194:** *Haematopinus asini* (Linnaeus, 1758). Male.




**Figure 195:** *Haematopinus asini*. Female.



**Figure 196:** *Hoplopleura pacifica* Ewing, 1924. Male.




**Figure 197:** *Hoplopleura pacifica*. Female.




**Figure 198:** *Linognathus setosus* (von Olfers, 1816). Male.


**Figure 199:** *Linognathus setosus*. Female.



**Figure 200:** *Solenopotes burmeisteri* (Fahrenheit, 1919). Male. **Figure 201:** *Solenopotes burmeisteri*. Female.




**Figure 202:** *Pediculus humanus capitis* De Geer, 1778. Male.



**Figure 203:** *Pediculus humanus capitis*. Female.




**Figure 204:** *Pediculus humanus humanus* Linnaeus, 1758. Male.



**Figure 205:** *Pediculus humanus humanus*. Female.



**Figure 206:** *Pthirus pubis* (Linnaeus, 1758). Male.




**Figure 207:** *Pthirus pubis*. Female.



**Figure 208:** *Haemodipsus lyriocephalus* (Burmeister, 1838). Male.



**Figure 209:** *Haemodipsus lyriocephalus*. Female.



**Figure 210:** *Polyplax serrata* (Burmeister, 1838). Male.

**Figure 211:** *Polyplax serrata*. Female.

## TAXONOMIC INDEX

This index includes all nominal invertebrate taxa mentioned in this volume, regardless of their current taxonomic status. Entries are valid and invalid genera, species, subspecies and varieties with their different generic and subgeneric combinations. Also, taxa above generic level are given in CAPITALS. Taxa in **bold** are those regarded by the author as currently valid. Page numbers in **bold** indicate main entries. Page numbers in *italics* indicate figures.

- 10-fasciatum*, *Liotheum (Colpocephalum)* 55  
*acanthopus*, *Hoplopleura* 234  
*acanthus*, *Docophorus* 200  
*acanthus*, *Hastaephorus* 200  
*acanthus*, *Philopterus* 200  
*accentor*, *Halipeurus* 122  
*accentor*, *Halipeurus (Halipeurus)* 122, 125  
***Acidoproctus*** 11, 21, **80**  
*actophila*, *Degeeriella* 138  
***actophilus*, *Luniceps*** 13, **138**, 139, 260  
*actophilus*, *Nirmus* 138  
***Actornithophilus*** 8, 21, **32**  
*acutifrons*, *Esthiopterum* 157  
***aegialitidis sensu lato*, *Austromenopon*** 9, **41**, 262  
*aegialitidis*, *Menopon* 41  
*aepycerus*, *Tricholipeurus* 226  
***affine*, *Austromenopon*** 9, **41**, 42, 249  
*affine*, *Menopon* 41  
*affinis*, *Procellariphaga* 42  
***aitkeni*, *Strigiphilus*** 16, **211**, 265  
*alato-clypeatus sensu lato*, *Alcedoecus* 81  
***alato-clypeatus*, *Alcedoecus*** 11, **81**, 266, 321  
*alato-clypeatus*, *Docophorus* 81  
*alato-clypeatus*, *Halcyonicola* 81  
*alato-clypeatus*, *Philopterus* 81  
***albatrossa*, *Saemundssonina (Saemundssonina)*** 16, **194**, 195, 249, 250, 348  
*albemarzensis*, *Docophorus* 195  
*albemarzensis*, *Philopterus* 195  
*albemarzensis*, *Saemundssonina* 195  
***albemarzensis*, *Saemundssonina (Saemundssonina)*** 16, **195**, 253  
***albescens*, *Eidmanniella*** 10, **60**, 61, 257  
*albescens*, *Eidmanniella* 60  
*albescens*, *Menopon* 60  
*albiventris*, *Pediculus* 161  
*albiventris*, *Penenirmus* 161  
*albofasciatum*, *Menopon* 65  
*albulus*, *Ardeicola* 87  
*album*, *Esthiopterum* 147  
*albus*, *Lipeurus* 147  
***albus*, *Neopsittaconirmus*** 14, **147**, 148, 265  
***Alcedoecus*** 11, **80**, 21  
*Allobrueelia* 96  
*Allobrüelia* 96  
*alpha*, *Degeeriella* 178  
*alpha*, *Koeniginirmus (Laminonirmus) normifer* 178  
*alpha*, *Koeniginirmus normifer* 178  
*alpha*, *Quadriceps* 178



- alpha*, *Quadriceps normifer* 15, **178**, 262  
*alpinus*, *Bovicola* (*Bovicola*) 221  
*alpinus*, *Docophorus* 194, 209  
*alpinus*, *Hastaeophorus* 209  
**AMBLYCERA** 8, 19, 20, **29**  
*americana*, *Bovicola* 221  
*americana*, *Damalinia* 221  
*americanus*, *Damalinia* (*Rhabdopedilon*) 221  
**ampullatus**, *Heterodoxus* 8, **29**, 30, 269, 305  
*amsel*, *Allobrueelia* 96  
*amsel*, *Allobrueelia* 96  
*amsel*, *Brueelia* 11, **96**, 268  
*Amysidea* 9, 21, **37**  
*Anamias* 121  
*Anaticola* 11, 21, **82**  
*Anatoecus* 11, 21, **83**  
*Ancistrona* 9, 21, **38**  
*angulicollis*, *Nirmus* 113  
*angusticeps angusticeps*, *Halipeurus* (*Halipeurus*) 121  
*angusticeps fosteri*, *Halipeurus* (*Halipeurus*) 121  
*angusticeps*, *Esthiopterum* 121  
**angusticeps**, *Halipeurus* 13, **121**, 122, 130, 254  
*angusticeps*, *Halipeurus* (*Halipeurus*) 121, 122  
*angusticeps*, *Halipeurus* (*Halipeurus*) *angusticeps* 121  
*angusticeps*, *Lipeurus* 121, 130  
*annulatum*, *Esthiopterum* 155  
*annulatus*, *Lipeurus* 155  
**annulatus**, *Pectinopygus* 14, **155**, 257  
**ANOPLURA** 8, 17, 19, 20, **228**  
*Anousceps* 172  
*anserinum*, *Trinoton* 78  
*anseris*, *Anaticola* 11, **82**, 247  
*anseris*, *Esthiopterum* 82  
*anseris*, *Pediculus* 82  
*anseris*, *Ricinus* 82  
*antarctica*, *Saemundssonina* 195  
**antarctica**, *Saemundssonina* (*Saemundssonina*) 16, **195**, 251  
**antarcticus**, *Austrogoniodes* 11, **90**, 248  
*antarcticus*, *Nirmus* 170  
*antarcticus*, *Perineus* 120  
*antarcticus*, *Phlopterus* 195  
*antarcticus*, *Pseudonirmus* 120, 170, 171  
**Antarctophthirus** 18, 21, **229**  
*apophoretus*, *Quadriceps* 177  
**apophoretus**, *Quadriceps hopkinsi* 15, **177**, 263  
**Aptericola** 15, 20, 25, **186**  
**Apterygon** 9, 20, 21, **39**  
**Aquanirmus** 11, 21, **86**  
*Arctophthirus* 229  
*ardeae*, *Ardeicola* 87  
*ardeae*, *Esthiopterum* 87  
**Ardeicola** 11, 21. **87**  
**Argimenopon** 9, **37**  
*armatus*, *Phlopterus* 207  
*asini asini*, *Haematopinus* 232  
*asini macrocephalus*, *Haematopinus* 232

- asini*, *Damalinia* 228  
**asini**, *Haematopinus* 17, **232**, 271, 355  
*asini*, *Haematopinus asini* 232  
*asini*, *Pediculus* 232  
*asini*, *Werneckiella equi* 228  
*assimilis assimilis*, *Quadriceps* 172  
*assimilis cedemajori*, *Quadriceps* 174  
*assimilis*, *Degeeriella* 172  
*assimilis*, *Nirmus* 172  
**assimilis**, *Quadriceps* 15, **172**, 262  
*assimilis*, *Quadriceps assimilis* 172  
*atlanticus*, *Docophorus* 196  
*atrofulvum sensu lato*, *Austromenopon* 42  
**atrofulvum**, *Austromenopon* 9, **42**, 263, 264  
*atrofulvum*, *Menopon* 42  
*attenuatus*, *Oncophorus* 186  
*aurata*, *Degeeriella* 173  
*auratus*, *Haematophagus* 173  
*auratus*, *Phlopterus (Docophorus)* 173  
**auratus**, *Quadriceps* 15, 172, **173**, 183, 261  
**australis**, *Aquanirmus* 11, 26, **86**, 248, 322  
**australis**, *Pectinopygus* 14, **155**, 156, 257  
*australis*, *Pectinopygus (Epipelicanus)* 155  
**Austrogoniodes** 11, 21, 26, **90**  
*Austromenopon* 9, 21, **41**, 42, 49, 53, 54  
*aviator*, *Docophorus* 214  
**aviator**, *Trabeculus* 17, 213, **214**, 255  
*baculus*, *Phlopterus (Lipeurus)* 106  
**balati**, *Saemundssonina (Saemundssonina) platygaster* 16, **206**, 262  
*bassanae serrator*, *Pectinopygus (Pectinopygus)* 156  
*bassanae*, *Pectinopygus* 156  
*bassanae*, *Pectinopygus (Pectinopygus)* 156  
**bassani**, *Pectinopygus* 14, 155, **156**, 257  
*bassani*, *Pectinopygus (Pectinopygus)* 156  
*bassani*, *Pediculus* 156  
*bassani serrator*, *Pectinopygus (Pectinopygus)* 156  
*bassanus*, *Pectinopygus (Pectinopygus)* 156  
*becki sensu lato*, *Austromenopon* 43  
*becki*, *Austromenopon* 43  
*becki*, *Philoceanus* 164  
*beckii sensu lato*, *Austromenopon* 43  
**beckii**, *Austromenopon* 9, **43**, 257  
*beckii*, *Menopon* 43  
**Bedfordiella** 11, 21, **95**  
*Benatoecus* 83  
**bicolor**, *Actornithophilus* 8, **32**, 261  
*bicolor*, *Colpocephalum* 32  
*bicolor*, *Docophorus* 196  
*bicolor*, *Phlopterus* 196  
*bicolor*, *Saemundssonina* 196  
**bicolor**, *Saemundssonina (Saemundssonina)* 16, **196**, 251  
**bidentatus compar**, *Campanulotes* 12, **101**, 264, 325  
*bidentatus*, *Campanulotes* 101  
*bidentatus*, *Pediculus* 101  
*birostris*, *Degeeriella* 173  
*birostris*, *Koeniginirmus (Laminonirmus) nychthemerus* 173

- birostris*, *Koeniginirmus nychthemerus* 173  
*birostris*, *Nirmus* 173  
***birostris*, *Quadriceps*** 15, 173, 174, 263  
*biseriatim*, *Menopon* 71  
*biseriatum*, *Eomenacanthus* 69  
*biseriatum*, *Menopon* 71  
*bisignatus*, *Docophorus* 134  
***bisignatus*, *Ibidoecus*** 13, 134, 135, 259  
*bisignatus*, *Philopterus* 134  
*bituberculatus*, *Goniodes* 117  
***Bonomiella*** 9, 21, 54  
***Boopia*** 8, 21, 29  
**BOOPIIDAE** 8, 29  
*borgioli*, *Neopsittaconirmus* 147  
***Bovicola*** 17, 21, 219  
*bovis*, *Bovicola* 219  
***bovis*, *Bovicola (Bovicola)*** 17, 219, 271  
*bovis*, *Bovicola (Damalinia)* 219  
*bovis*, *Damalinia* 219  
*bovis*, *Damalinia (Bovicola)* 219  
*bovis*, *Pediculus* 219  
*bovis*, *Trichodectes* 219  
*bracteata*, *Brueelia* 112  
***bracteata*, *Emersoniella*** 12, 112, 113, 266, 330  
*bracteatus*, *Nirmus* 112  
***breueri*, *Brueelia*** 11, 96, 97, 269  
*breviceps*, *Bovicola* 222  
***breviceps*, *Bovicola (Lepikentron)*** 17, 222, 223, 271, 350  
*breviceps*, *Damalinia* 222  
*breviceps*, *Damalinia (Lepikentron)* 222  
*breviceps*, *Lepikentron* 222  
*breviceps*, *Trichodectes* 222  
*brevicornis*, *Lipeurus* 155  
*brevicornis*, *Pectinopygus* 155  
*brevifimbriata*, *Procellariphaga* 43  
***brevifimbriatum*, *Austromenopon*** 9, 43, 44, 251  
*brevifimbriatum*, *Menopon* 43  
*brevipalpe*, *Menopon* 60  
*brevipalpis*, *Eidmanniella* 62  
*brevipalpis*, *Menopon* 61  
***brevis*, *Docophoroides*** 12, 110, 249  
*brevis*, *Philopterus* 110  
***Brueelia*** 11, 21, 26, 96  
*Brueelia* 96  
*Brüelia* 96  
*bucephalus bucephalus*, *Ornithobius* 149  
*bucephalus tribulis*, *Ornithobius* 149  
*bucephalus*, *Lipeurus* 149  
***bucephalus*, *Ornithobius*** 14, 149, 246  
*bucephalus*, *Ornithobius bucephalus* 149  
*bulleri*, *Austromenopon* 48, 49  
***bulweriae*, *Austromenopon*** 9, 44, 254  
***bulweriae*, *Halipeurus*** 13, 122, 254  
*bulweriae*, *Halipeurus (Halipeurus)* 122  
*burmeisteri*, *Linognathus* 238  
*burmeisteri*, *Nitzchia* 60

- burmeisteri*, *Solenopotes* 18, **238**, 271, 356  
*cacatuae*, *Franciscolea* 63  
*callaeincola*, *Brueelia* 12, **97**, 266  
*cameratus*, *Nirmus* 137  
**Campanulotes** 12, 21, **101**  
*canis*, *Pediculus* 224  
*canis*, *Ricinus* 224  
*canis*, *Trichodectes* 224  
*canis*, *Trichodectes (Trichodectes)* 17, **224**, 225, 270, 352  
*capillatus*, *Solenopotes* 18, **238**, 239, 271  
*capistratus*, *Alcedoecus* 80  
*capistratus*, *Philopterus* 80  
*capitis*, *Pediculus* 239  
*capitis*, *Pediculus humanus* 18, **239**, 272, 357  
*caponis*, *Lipeurus* 13, **137**, 138, 246  
*caponis*, *Lipeurus (Lipeurus)* 137  
*caponis*, *Pediculus* 137  
*caponis*, *Ricinus* 137  
*caprae*, *Bovicola* 220  
*caprae*, *Bovicola (Bovicola)* 17, 219, **220**, 271  
*caprae*, *Damalinia* 220  
*caprae*, *Damalinia (Bovicola)* 220  
*caprae*, *Trichodectes* 219, 220  
**Capraiella** 12, 21, **102**  
**Carduiceps** 12, 21, **102**  
*carlinii*, *Antarctophthirus* 17, **229**, 231, 270  
*carunculatus sensu lato*, *Pectinopygus* 14, **157**, 258  
*carunculatus*, *Pectinopygus* 14, **156**, 157, 258  
*caspia*, *Degeeriella* 174  
*caspius caspius*, *Koeniginirmus (Laminonirmus)* 174  
*caspius*, *Koeniginirmus (Laminonirmus) caspius* 174  
*caspius*, *Koeniginirmus sellatus* 174  
*caspius*, *Nirmus* 174  
*caspius*, *Quadriceps* 15, **174**, 263, 345  
*Cecalymenus* 213  
*cedemajori*, *Quadriceps* 15, **174**, 262  
*cedemajori*, *Quadriceps assimilis* 174  
*cephalus*, *Philopterus* 196  
*cephalus*, *Philopterus (Docophorus)* 196  
*cephalus*, *Saemundsonia* 196  
*cephalus*, *Saemundsonia (Saemundsonia)* 16, **196**, 197, 262  
*ceruleus sensu lato*, *Actornithophilus* 32, 33  
*ceruleus*, *Actornithophilus* 8, **32**, 263, 307  
*ceruleus*, *Clypeodon* 32  
*Chadriceps* 172  
*charadrii charadrii*, *Quadriceps* 175  
*charadrii hospes*, *Quadriceps* **172**, 175  
*charadrii orarius*, *Quadriceps* 15, **175**, 261  
*charadrii punctifer*, *Quadriceps* 175  
*charadrii*, *Pediculus* 175  
*charadrii*, *Quadriceps* 175  
*charadrii*, *Quadriceps charadrii* 175  
*charcoti*, *Degeeriella* 170  
*charcoti*, *Esthiopterum* 170  
*charcoti*, *Pseudonirmus* 15, **170**, 251  
*charcoti*, *Ricinus* 170

- chathamensis*, *Saemundssonina* 197  
***chathamensis*, *Saemundssonina* (*Saemundssonina*)** 16, **197**, 262  
*chelicornis*, *Goniodes* 117  
***Chelopistes*** 12, 21, **103**  
*chilensis*, *Harrisoniella* 133  
***chinense*, *Coloceras*** 12, **104**, 105, 264  
*chinensis*, *Goniocotes* 104  
***chrysocephalus*, *Goniocotes*** 12, **116**, 246  
***Ciconiphilus*** 9, 21, **55**  
*cingulata*, *Degeeriella* 102  
***cingulatus cingulatus*, *Carduiceps*** 12, **102**, 261, 326  
***cingulatus lapponicus*, *Carduiceps*** 12, **102**, 103, 261  
***cingulatus*, *Carduiceps cingulatus*** 12, **102**, 261, 326  
*cingulatus*, *Philopterus* (*Nirmus*) 102  
*circumfasciatus* var. *kea*, *Lipeurus* 148  
***circumfasciatus*, *Perineus*** 14, **161**, 162, 163, 249, 250  
*clavatus*, *Epicolinus* 150  
*clavatus*, *Lipeurus* 150  
***clavatus*, *Oxylipeurus*** 14, **150**, 151, 246  
*climax*, *Trichodectes* 220  
*clypeatum*, *Esthiopterum* 142, 144  
*clypeatus*, *Lipeurus* 142, 144  
*clypeatus*, *Naubates* 142, 144  
***clypeatus*, *Naubates* (*Guenterion*)** 14, **142**, 143, 145, 253  
*clypeatus*, *Naubates* (*Micronaubates*) 144  
*clypeilargum*, *Holomenopon* 244  
*clypeilargum*, *Holomenopon* 66, 67  
*Clypeodon* 32  
***coenocoryphae*, *Quadriceps*** 15, **175**, 260  
***colchici*, *Goniodes*** 13, **117**, 246  
*colchici*, *Gonocephalus* 117  
*colchici*, *Philopterus* (*Goniodes*) 117  
*colchicus*, *Oxylipeurus* 151  
***colchicus*, *Oxylipeurus mesopelios*** 14, **151**, 246  
*Colinicola* 137  
***Coloceras*** 12, 21, **104**  
***Colpocephalum*** 9, 21, **56**  
***columbae columbae*, *Columbicola*** 12, **106**, 197, 264, 327  
***columbae*, *Bonomiella*** 9, **54**, 55, 264, 310  
***columbae*, *Columbicola columbae*** 12, **106**, 197, 264, 327  
*columbae*, *Esthiopterum* 106  
*columbae*, *Lipeurus* 106  
*columbae*, *Pediculus* 106  
*columbae*, *Ricinus* 106  
***Columbicola*** 12, 21, **106**  
*communis* var. *passeris*, *Docophorus* 167  
*compar*, *Campanulotes* 101  
***compar*, *Campanulotes bidentatus*** 12, **101**, 264, 325  
*compar*, *Goniocotes* 101  
*compar*, *Philopterus* (*Goniocotes*) 101  
*complexiva*, *Degeeriella* 102  
*complexivus*, *Carduiceps* 102  
*concaivifrons*, *Damalinia* (*Bovicola*) 221  
*conci*, *Austrogoniodes* 90  
***Conciella*** 10, **67**  
*concii sensu lato*, *Austrogoniodes* 90, 91

- concii*, *Austrogoniodes* 11, **90**, 91, 248, 249  
*concii*, *Cesareus* 90  
*concinnoides*, *Perineus* 14, **162**, 163, 249, 342  
*concinus*, *Perineus* 245  
*concretum*, *Menopon* 76  
*concretum*, *Pseudomenopon* 11, **76**, 77, 260  
*confusus*, *Halipeurus* 13, **122**, 126, 252, 333  
*conica conica*, *Saemundssonina* 197  
*conica conica*, *Saemundssonina* (*Saemundssonina*) 16, **197**, 198, 261  
*conica*, *Saemundssonina* 197  
*conica*, *Saemundssonina* (*Saemundssonina*) *conica* 16, **197**, 198, 261  
*conica*, *Saemundssonina conica* 197  
*conicus*, *Phlopterus* 197  
*conicus*, *Phlopterus* (*Docophorus*) 197  
*consimilis*, *Halipeurus* 13, 25, **123**, 252  
*consimilis*, *Halipeurus* (*Halipeurus*) 123  
*conspurcatum*, *Liotheum* (*Trinoton*) 78  
*constrictiventre*, *Esthiopterum* 123  
*corporis*, *Pediculus* 240  
*corporis*, *Pediculus humanus* 240  
*corporus*, *Pediculus* (*Pediculus*) *humanus* 240  
*Corvocolpocephalum* 56  
*crassicorne*, *Anaticola* 82  
*crassicorne*, *Esthiopterum* 82  
*crassicornis*, *Anaticola* 11, **82**, 83, 247, 248  
*crassicornis*, *Pediculus* 82  
*crassicornis*, sensu lato, *Anaticola* 82, 83  
*creatopae*, *Saemundssonina* 196  
*cristati*, *Austrogoniodes* 11, **91**, 248  
*crocatum*, *Austromenopon* 41  
*crocatum*, *Menopon* 41  
*cryptostigmatia*, *Kurodaia* 67  
*cryptostigmatia*, *Kurodaia* (*Conciella*) 10, **67**, 68, 265, 315  
*cryptostigmatia*, *Conciella* 67  
*cryptostigmatia*, *Menopon* 67  
*Cuclotogaster* 12, 21, **107**  
*Cuculicola* 12, 21, **108**  
*Cuculiphilus* 10, 21, **59**  
*cursitans*, *Docophorus* 211  
*cursitans*, *Phlopterus* 211  
*cursitans*, *Strigiphilus* 16, **211**, 212, 265  
*cursorium*, *Austromenopon* 9, **44**, 262  
*cursorius*, *Austromenopon* 44  
*cursorius*, *Menopon* 44  
*cuspidata*, *Rallicola* 189  
*cyclothorax*, *Bruelia* 12, **97**, 268  
*cyclothorax*, *Bruelia* 97  
*cyclothorax*, *Degeeriella* 97  
*cyclothorax*, *Nirmus* 97  
*cygni*, *Ornithobius* 149  
*damicorne*, *Coloceras* 104  
*damicornis*, *Goniodes* 104  
*damma*, *Naubates* 143  
*damma*, *Naubates* (*Guenterion*) 14, **143**, 252, 253  
*daptionis*, *Austromenopon* 43  
*daptionis*, *Procellariphaga* 43

- decimfasciatum*, *Colpocephalum* 55  
*decimfasciatus*, *Ciconiphilus* 9, **55**, 258, 259, 310  
*Degeeriella* 12, 21, **109**  
*delicata*, *Brueelia* 12, **97**, 98, 269  
*delicata*, *Bruëlia* 97  
*delicata*, *Degeeriella* 97  
*delicatus*, *Nirmus* 97  
*delphax*, *Alcedoecus* 11, **81**, 266  
*delphax*, *Docophorus* 81  
*delphax*, *Halcyonicola* 81  
*delphax*, *Philopterus* 81  
*demersa*, *Nesiotinus* 148  
*demersus*, *Nesiotinus* 14, **148**, 149, 248, 339  
*Dennyus* 10, 21, **60**  
*densilimba*, *Brueelia* 12, **98**, 269  
*densilimba*, *Bruëlia* 98  
*densilimba*, *Degeeriella* 98  
*densilimbus*, *Nirmus* 98  
*densus*, *Lipeurus* 133  
*dentatus dentatus*, *Anatoecus* (*Benatoecus*) 84  
*dentatus magnicornutus*, *Anatoecus* 11, **84**, 85, 246  
*dentatus magnicornutus*, *Anatoecus* (*Benatoecus*) 84  
*dentatus sensu lato*, *Anatoecus* 11, **83**, 84, 247, 248  
*dentatus*, *Anatoecus* 83  
*dentatus*, *Anatoecus* (*Benatoecus*) *dentatus* 84  
*dentatus*, *Pediculus* 83  
*dentatus*, *Philopterus* 83  
*desolata*, *Saemundssonina* 198  
*desolata*, *Saemundssonina* (*Saemundssonina*) 16, **198**, 253, 254  
*dianae*, *Ibidoecus* 13, **135**, 259, 334  
*dimorpha*, *Mackayia* 213  
*dimorphus*, *Trabeculus* 214  
*diomedae diomedae*, *Perineus* 152  
*diomedae enderleini*, *Perineus* 152  
*diomedae lepturus*, *Perineus* 152  
*diomedae sensu lato*, *Paraclisis* 152, 153  
*diomedae*, *Esthiopterum* 133, 152  
*diomedae*, *Harrisoniella* 134, 152  
*diomedae*, *Lipeurus* 133  
*diomedae*, *Paraclisis* 14, **152**, 153, 249, 250  
*diomedae*, *Pediculus* 133, 152  
*diomedae*, *Perineus* 134, 152  
*diomedae*, *Perineus* (s. l.) 152  
*diomedae*, *Perineus diomedae* 152  
*diomedae*, *Philopterus* 133  
*diomedae*, *Ricinus* 152  
*Diomedicola* 133  
*discocephalus*, *Degeeriella* 109  
*dispar*, *Esthiopterum* 157  
*dispar*, *Goniodes* 13, 117, **118**, 246  
*dispar*, *Nirmus* 157  
*dispar*, *Pectinopygus* 14, **157**, 257  
*dispar*, *Solenodes* 118  
*dissimilis*, *Goniodes* 13, 117, **118**, 244, 246  
*dissimilis*, *Oulocrepis* 118  
*dissimilis*, *Philopterus* (*Goniodes*) 118

- diversum*, *Esthiopterum* 123  
*diversus* var. *excavatus*, *Lipeurus* 125  
*diversus* var. *major*, *Lipeurus* 127  
***diversus*, *Halipeurus*** 13, **123**, 124, 133, 255  
*diversus*, *Halipeurus* (*Halipeurus*) 123  
*diversus*, *Lipeurus* 123  
***Docophoroides*** 12, 21, **110**  
*docophoroides*, *Colinicola* 137  
***docophoroides*, *Lagopoecus*** 13, **137**, 245, 335  
*docophoroides*, *Lipeurus* 137  
*Docophorulus* 167  
*Docophorus* 167, 168, 169  
***dominella*, *Quadriceps*** 15, **176**, 262  
*doriabagla*, *Ciconiphilus* 55  
*doriabagla*, *Pseudocolpocephalum* 55  
***drosti*, *Luniceps*** 13, **139**, 260  
*dumosum* sensu lato, *Apterygon* 39. 40  
***dumosum*, *Apterygon*** 9, **39**. 40, 245  
*eatoni*, *Lipeurus* 160  
*eatoni*, *Pelmatocerandra* 160  
**ECHINOPHTHIRIIDAE** 17, **229**  
*Echinophthirius* 229  
***Eidmanniella*** 10, 21, **60**  
*elegans*, *Bruelia* 100  
***elliotti*, *Austromenopon*** 9, **45**, 257  
*elliptica*, *Degeeriella* 176  
***ellipticus* sensu lato, *Quadriceps*** 15, 172, **176**, 262  
*ellipticus*, *Glareolites* 176  
*ellipticus*, *Epicolinus* 151  
*ellipticus*, *Nirmus* 172, 176  
***ellipticus*, *Oxylipeurus*** 14, **151**, 245  
*ellipticus*, *Quadriceps* 172  
*emarginata*, *Degeeriella* 184  
*emarginatus*, *Nirmus gloriosus* 184  
***emersoni*, *Acidoproctus*** 11, **80**, 246  
***Emersoniella*** 12, 21, **112**  
*enderleini*, *Perineus* 152  
*enderleini*, *Perineus diomedea* 152  
*enderleini*, *Puffinoecus* 192  
*enderleini*, *Saemundssonina* 192  
***enderleini*, *Saemundssonina* (*Puffinoecus*)** 16, **192**, 251, 252, 347  
***enigki*, *Austromenopon*** 9, **45**, 256  
*Epicolinus* 150  
*Epifregata* 155  
*Epipelicanus* 155  
***Episbates*** 12, 21, **113**  
*equi asini*, *Werneckiella* 228  
*equi*, *Bovicola* 227  
*equi*, *Bovicola* (*Werneckiella*) 227, 228  
*equi*, *Damalinia* 227  
*equi*, *Damalinia* (*Bovicola*) 227  
*equi*, *Damalinia* (*Werneckiella*) 227  
*equi*, *Trichodectes* 227  
***equi*, *Werneckiella*** 17, **227**, 228, 271  
*ermineae*, *Stachiella* 225  
***ermineae*, *Trichodectes* (*Stachiella*)** 17, **225**, 270, 352



- ermineae, Trichodectes* 225  
*eternitatus, Brueelia turdinulae* 100  
*eucarenum, Colpocephalum* 9, **56**, 257  
***Eucolpocephalum*** 10, 21, **62**  
*eurygaster, Eidmanniella* 10, **61**, 257, 258, 312  
*eurygaster, Menopon* 61  
*Eurymetopus* 110  
*euryrhyncha, Saemundssonina* 198, 199  
***euryrhyncha, Saemundssonina (Saemundssonina)*** 16, **198**, 199, 262  
*euryrhynchus, Docophorus* 198  
*euryrhynchus, Philopterus* 198  
*eurysternus, Haematopinus* 232  
*eurysternum, Menacanthus* 69  
***eurysternus, Haematopinus*** 17, 232, **233**, 271  
***eurysternus, Menacanthus*** 10, 19, **69**, 70, 268, 269  
*eurysternus, Menopon* 69  
*eurysternus, Pediculus* 233  
*excavatum, Esthiopterum* 125  
*exiguus, Lipeurus* 128  
*expallida, Ardeicola* 87  
***expallidus, Ardeicola*** 11, **87**, 88, 258  
*extinctus, Huiacola* 188  
***extinctus, Rallicola (Huiacola)*** 16, **188**, 189, 266, 346  
***falcinellus, Luniceps*** 13, **139**, 140, 244, 260  
*fallai, Saemundssonina* 201, 202  
*fallai, Saemundssonina lari* 201, 202  
*fallax, Oncophorus* 190  
*fallax, Rallicola* 190  
***falsus pacificus, Halipeurus*** 13, 25, **124**, 257  
*falsus pacificus, Halipeurus (Halipeurus)* 124  
***fasciiventris, Cuculiphilus (Cuculiphilus)*** 10, **59**, 265, 311  
*fasciatus, Cuculiphilus (Cuculiphilus)* 59  
*fasciatus, Naubates* 164  
*fasciatus, Pediculus* 59  
***fasciatus, Philoceanus*** 15, **164**, 256, 343  
***Felicola*** 17, 21, **223**  
***femorale, Eucolpocephalum*** 10, 62, **63**, 259, 313  
*femorale, Menopon* 63  
*ferox, Diomedicola* 133  
*ferox, Esthiopterum* 133  
*ferox, Harrisoniella* 134  
***ferox, Harrisoniella*** 13, **133**, 250, 334  
*ferox, Lipeurus* 133  
*ferox, Perineus* 133  
*Ferrisia* 56  
***flemingi, Trabeculus*** 17, 25, **214**, 255  
*forficula, Forficuloecus* 113  
*forficula, Philopterus* 113  
*forficulatus, Lipeurus* 155  
*forficulatus, Pectinopygus* 155  
***Forficuloecus*** 12, 20, 21, **113**, 115  
*fosteri, Halipeurus (Halipeurus) angusticeps* 121  
***Franciscoloa*** 10, 21, **63**  
***fregili, Colpocephalum*** 9, 56, **57**, 267  
*fregili, Liotheum (Colpocephalum)* 57  
*fringillae, Philopterus* 167

- fringillae, Ricinus* 79  
*fulicae, Incidifrons* 13, **136**, 260  
*fulicae, Pediculus* 136  
*fulicae, Philopterus (Nirmus)* 189  
*fulicae, Rallicola* 189  
*fulicae, Rallicola (Rallicola)* 16, **189**, 260  
**Fulicoffula** 12, 22, **115**  
*fuliginosum, Esthiopterum* 146  
*fuliginosus hawaiiensis, Docophorus* 197  
*fuliginosus, Lipeurus* 146  
*fuliginosus, Naubates* 144, 146  
*fuliginosus, Naubates (Naubates)* 14, **146**, 254  
*fuliginosus, Philopteroides* 15, **165**, 267, 343  
*fulvofasciatum kea, Menopon* 64  
*fulvofasciatum* var. *kea, Menopon* 64  
*fulvofasciatum* var. *minor, Menopon* 75  
*fusca, Degeeriella* 12, **109**, 259, 329  
*fuscoclypeata, Giebelia* 214  
*fuscoclypeatus sensu lato, Trabeculus* 215  
*fuscoclypeatus, Philopterus* 214  
*fuscoclypeatus, Trabeculus* 17, **214**, 215, 252, 349  
*fuscofasciatum, Austromenopon* 9, **46**, 262  
*fuscofasciatum, Menopon* 46  
*fuscolaminulata, Degeeriella* 180  
*fuscolaminulatus Quadriceps ornatus* 15, **180**, 181, 182, 263  
*fuscolaminulatus, Koeniginirmus* 180  
*fuscolaminulatus, Koeniginirmus (Laminonirmus) ornatus* 180  
*fuscolaminulatus, Koeniginirmus ornatus* 180  
*fuscolaminulatus, Quadriceps* 180  
*fuscolaminulatus, Ricinus* 180  
*fuscus, Degeeriella* 109  
*fuscus, Nirmus* 109  
*fuscus, Ornithobius* 14, **149**, 150, 247, 339  
*fuscus, Philopterus (Nirmus)* 109  
*gadowi sensu lato, Rallicola (Aptericola)* 15, **186**, 187, 245  
*gadowi, Aptericola* 186  
*gadowi, Rallicola* 186  
*gadowi, Rallicola (Aptericola)* 15, 25, **186**, 245  
*gaibagla, Ardeicola* 87  
*gaini, Philopterus* 199  
*gaini, Saemundssonina* 199  
*gaini, Saemundssonina (Saemundssonina)* 16, **199**, 251  
*galeatum, Longimenopon* 10, **68**, 253  
*Galliferrisia* 56  
*gallinae, Goniocotes* 101  
*gallinae, Goniocotes* 12, **116**, 246  
*gallinae, Menopon* 10, **72**, 246, 317  
*gallinae, Pediculus* 72  
*gallinae, Ricinus* 72, 101, 116  
*garbei, Naubates* 157  
*garbei, Naubates (Micronaubates)* 157  
*garbei, Pectinopygus* 14, **157**, 158, 257  
*garbei, Pectinopygus (Pectinopygus)* 158  
*garrodiae, Naubates* 164  
*garrodiae, Philoceanus* 15, **164**, 165, 256  
*Giebelia* 213

- giganteum*, *Menopon* 65  
*gigas*, *Ancistrona* 38  
*gigas*, *Goniocotes* 244  
*gigas*, *Goniodes* 244  
*Glareolites* 172  
***Gliricola* 8, 22, 30**  
*gloriosa*, *Degeeriella* 173  
*gloriosus emarginatus*, *Nirmus* 184  
*gloriosus*, *Nirmus* 173  
***Goniocotes* 12, 22, 116**  
***Goniodes* 13, 22, 117, 118, 119**  
*goniodicerus*, *Strigiphilus* 211  
***goniopleurus*, *Ornithobius* 14, 150, 247**  
*gonioplurus*, *Philoferus* (*Ornithobius*) 150  
*Gonocephalus* 117  
*gonothorax*, *Docophorus* 201  
*gonothorax*, *Philoferus* 192, 194, 201  
*gonothorax*, *Saemundssonina* 201  
*gonothorax*, *Saemundssonina lari* 201, 202  
***gottwaldhirschi*, *Acidoproctus* 11, 80, 247, 320**  
*gracilentus*, *Rallicola* 187  
***gracilentus*, *Rallicola* (*Aptericola*) 16, 25, 187, 245**  
*gracilicornis*, *Lipeurus* 155  
*gracilicornis*, *Pectinopygus* 155  
*gracilis*, *Docophorus* 187  
*gracilis*, *Gyropus* 30  
*gracilis*, *Rallicola* 187  
*gracilis*, *Rallicola* 187  
*gracilis*, *Rallicola* (*Aptericola*) 25, 187  
*grande*, *Esthiopterum* 120  
***grandiceps*, *Actornithophilus* 8, 33, 261**  
*grandiceps*, *Colpocephalum* 33  
*grandis sensu lato*, *Haffneria* 121  
*grandis*, *Diomedicola* 120  
***grandis*, *Haffneria* 13, 120, 121, 262, 333**  
*grandis*, *Harrisoniella* 120  
*grandis*, *Lipeurus* 120  
*grandis*, *Perineus* 120  
*gravis gravis*, *Halipeurus* 124  
***gravis priapulus*, *Halipeurus* 13, 124, 255**  
*gravis priapulus*, *Halipeurus* (*Halipeurus*) 124  
*gravis*, *Halipeurus* (*Halipeurus*) 124  
*gravis*, *Halipeurus gravis* 124  
***Guenterion* 14, 142**  
*gurlti*, *Esthiopterum* 170  
*gurlti*, *Lipeurus* 170  
***gurlti*, *Pseudonirmus* 15, 170, 171, 251**  
***gygisa*, *Saemundssonina* (*Saemundssonina*) 16, 200, 263**  
*gyricorne*, *Esthiopterum* 158  
***gyricornis*, *Pectinopygus* 14, 158, 257**  
*gyricornis*, *Pectinopygus* (*Philichthyophaga*) 158  
*gyricornis*, *Philichthyophaga* 158  
*gyricornis*, *Philoferus* (*Lipeurus*) 158  
**GYROPIDAE 8, 30**  
***Gyropus* 8, 22, 30**  
*Haematophagus* 172

- haematopi*, *Austromenopon* 9, **46**, 261  
*haematopi*, *Pediculus* 200  
*haematopi*, *Quadriceps* 172  
*haematopi*, *Ricinus* 200  
*haematopi*, *Saemundssonina* 200  
*haematopi*, *Saemundssonina* (*Saemundssonina*) 16, **200**, 261  
**HAEMATOPINIDAE** 17, **232**  
***Haematopinus*** 17, 22, **232**  
***Haemodipsus*** 18, 22, **241**  
***Haffneria*** 13, 22, **120**  
*Halcyonicola* 81  
*halcyonis*, *Emersoniella* 112  
*haliaeeti*, *Colpocephalum* 67  
*haliaeeti*, *Kurodaia* (*Kurodaia*) 67  
***Halipeurus*** 13, 22, 26, **121**  
*halobaenae*, *Saemundssonina* 206  
***hamiltoni*, *Austrogoniodes*** 11, **91**, 248, 323  
*hamiltoni*, *Cesareus* 91  
*hanáki*, *Halipeurus* 123  
*hanáki*, *Halipeurus diversus* 123  
*harrisoni* sensu lato, *Naubates* 146, 147  
*harrisoni* sensu lato, *Rallicola* 189, 190  
***harrisoni*, *Coloceras*** 12, **105**, 264  
***harrisoni*, *Docophoroides*** 12, **110**, 111, 250  
*harrisoni*, *Naubates* 146  
***harrisoni*, *Naubates* (*Naubates*)** 14, **146**, 147, 255  
*harrisoni*, *Patellinirmus* 105  
*harrisoni*, *Rallicola* 189  
***harrisoni*, *Rallicola* (*Rallicola*)** 16, 25, **189**, 190, 259  
***Harrisoniella*** 13, 22, **133**, 152  
*Hastaephorus* 194  
*hawaiiensis*, *Docophorus fuliginosus* 197  
*hawaiiensis*, *Saemundssonina* 197  
*hemichroa*, *Degeeriella* 176  
*hemichrous*, *Himantophagus* 177  
*hemichrous*, *Nirmus* 172, 176  
***hemichrous*, *Quadriceps*** 15, 172, **176**, 177, 184, 261  
***hemiphagae*, *Coloceras*** 12, **105**, 264  
*hemiphagae*, *Nitzschiella* 105  
*hemitragi*, *Bovicola* 223  
***hemitragi*, *Bovicola* (*Spinibovicola*)** 17, **223**, 271, 351  
*hemitragi*, *Damalinia* 223  
*hemitragi*, *Damalinia* (*Bovicola*) 223  
*hemitragi*, *Trichodectes* 223  
*heteracanthus*, *Mackayia* 217  
*heteracanthus*, *Trabeculus* 217, 218  
*heterocerus*, *Docophorus* 211  
***Heterodoxus*** 8, 22, **29**  
*heterographus heterographus*, *Gallipeurus* 107  
***heterographus*, *Cuclotogaster*** 12, **107**, 246  
*heterographus*, *Gallipeurus heterographus* 107  
*heterographus*, *Lipeurus* 107  
*heterographus*, *Lipeurus* (*Gallipeurus*) 107  
***Heteromenopon*** 10, 22, **64**  
*heteroproctus*, *Naubates* 142, 143, 144  
***heteroproctus*, *Naubates* (*Guenterion*)** 14, 142, **143**, 144, 145, 251

- hexacon*, *Trabeculus* 215, 216  
*hexagona*, *Saemundssonina* 200, 201  
***hexagona*, *Saemundssonina* (*Saemundssonina*)** 16, **200**, 201, 257  
*hexagonus*, *Docophorus* 200  
*hexagonus*, *Saemundssonina* 200  
*hexakon*, *Giebelia* 215, 216  
***hexakon sensu lato*, *Trabeculus*** 17, **216**, 252, 253, 255  
***hexakon*, *Trabeculus*** 17, **215**, 219, 254  
*hiaticulae*, *Pediculus* 172  
*hiaticulae*, *Quadriceps* 172  
***hihi*, *Myrsidea*** 10, **72**, 73, 266  
*Himantophagus* 172  
***himantopi*, *Austromenopon*** 9, **46**, 47, 261  
***hintoni*, *Apterygon*** 9, **40**, 245  
*hirundinis*, *Dennyus* (*Dennyus*) 60  
***Hohorstiella*** 10, 22, **64**  
*hologaster*, *Goniocotes* 116  
*hologaster*, *Philopterus* (*Goniodes*) 116  
***Holomenopon*** 10, 22, **65**  
***hopkinsi apophoretus*, *Quadriceps*** 15, **177**, 263  
***hopkinsi hopkinsi*, *Quadriceps*** 15, **177**, 263  
*hopkinsi*, *Anousceps* 177  
*hopkinsi*, *Diomedicola* 134  
***hopkinsi*, *Harrisoniella*** 13, **133**, 134, 249  
*hopkinsi*, *Quadriceps* 177  
***hopkinsi*, *Quadriceps hopkinsi*** 15, **177**, 263  
***Hoplopleura*** 18, 22, **234**  
**HOPLOPLEURIDAE** 18, **234**  
*hoplopteri*, *Colpocephalum* 33  
***hoplopteri sensu lato*, *Actornithophilus*** 9, **33**, 262  
***hospes*, *Quadriceps charadrii*** 172, 175  
*hour*, *Koeniginirmus* (*Laminonirmus*) 183  
*hour*, *Quadriceps* 183  
***hour*, *Quadriceps sellatus*** 15, **183**, 184, 264  
***Huiacola*** 16, 20, **188**  
***humanus capitis*, *Pediculus*** 18, **239**, 272, 357  
*humanus corporis*, *Pediculus* 240  
*humanus corporis*, *Pediculus* (*Pediculus*) 240  
***humanus humanus*, *Pediculus*** 18, **240**, 272, 357  
*humanus*, *Pediculus* 240  
***humanus*, *Pediculus humanus*** 18, **240**, 272, 357  
*hunteri*, *Docophoroides* 111  
*hyalina*, *Harrisoniella* 153  
***hyalina*, *Paraclisis*** 14, **153**, 249  
*hyalinum*, *Esthiopterum* 153  
*hyalinus*, *Lipeurus* 153  
*hyalinus*, *Perineus* 153  
***Ibidoecus*** 13, 22, **134**  
*icterodes icterodes*, *Anatoecus* (*Anatoecus*) 85  
*icterodes knechteli*, *Anatoecus* 85  
***icterodes oloris*, *Anatoecus*** 11, **85**, 86, 246, 322  
*icterodes oloris*, *Anatoecus* (*Anatoecus*) 85  
***icterodes sensu lato*, *Anatoecus*** 11, **85**, 247, 248  
*icterodes*, *Anatoecus* 83  
*icterodes*, *Anatoecus* (*Anatoecus*) *icterodes* 85  
*icterodes*, *Philopterus* 85

- icterodes*, *Philopterus* (*Docophorus*) 85  
*icteroides*, *A.* 85  
*inaequalis*, *Lipeurus* 150  
*inaequalis*, *Oxylipeurus* 150  
***Incidifrons*** 13, 22, **136**  
*incisa*, *Saemundssonina* 208  
*incisa*, *Saemundssonina* (*Saemundssonina*) 209  
***incisa*, *Saemundssonina* (*Saemundssonina*) *thalassidromae*** 16, **208**, 209, 256  
*incoenis* sensu lato, *Luniceps* 139  
*incoenis*, *Luniceps* 141, 244  
*incoenis*, *Nirmus* 139  
*inexpectata*, *Saemundssonina* 201  
***inexpectata*, *Saemundssonina* (*Saemundssonina*)** 16, **201**, 263  
*inguinalis*, *Phthirus* 241  
*inguinalis*, *Phthirus* 241  
*inguinalis*, *Phtirus* 241  
*inguinalis*, *Pthirus* 241  
**INSECTA** 3, **5**, 8  
*insolitunguicolata*, *Bonomiella* 54  
*intermedius*, *Halipeurus* (*Halipeurus*) 127  
*intestatus*, *Halipeurus* 130  
***irkutensis*, *Philopterus*** 15, **167**, 268, 269  
*irroratae*, *Diomedicola* 133  
*irroratae*, *Harrisoniella* 133  
**ISCHNOCERA** 8, 11, 19, 20, **80**  
***ivanliteraki*, *Myrsidea*** 10, **73**, 267  
*jacobi*, *Trichodectes* (*Stachiella*) 225  
*jamaicensis*, *Pectinopygus* 155  
*jamaicensis*, *Puffinoecus* 192  
*jamaicensis*, *Saemundssonina* 192  
***jamaicensis*, *Saemundssonina* (*Puffinoecus*)** 16, **192**, 193, 254  
*kea*, *Esthiopterum* 148  
*kea*, *Heteromenopon* 64  
***kea*, *Heteromenopon* (*Keamenopon*)** 10, **64**, 264, 314  
*kea*, *Menopon* 64  
*kea*, *Menopon fulvofasciatum* 64  
***kea*, *Neopsittaconirmus*** 14, 147, **148**, 264, 338  
*kea*, *Psittacicola* 148  
*kea*, *Psittacomenopon* 64  
***Keamenopon*** 10, **64**  
*kermadecense*, *Esthiopterum* 125  
*kermadecense*, *Halipeurus* 125  
*kermadecense*, *Halipeurus* (*Halipeurus*) 125  
*kermadecensis* sensu lato, *Halipeurus* (*Halipeurus*) 125  
***kermadecensis*, *Halipeurus*** 13, **125**, 252  
*kermadecensis*, *Halipeurus* (*Halipeurus*) 125  
*kermadecensis*, *Lipeurus* 125  
*knechteli*, *Anatoecus icterodes* 85  
*Koeniginirmus* 172  
***kui*, *Cuculicola*** 12, 26, **108**, 265, 328  
***Kurodaia*** 10, 22, **67**  
*kuwani*, *Eidmanniella* 62  
*kuwani*, *Menopon* 61  
*laculatus*, *Lipeurus* 120  
*laculatus*, *Perineus* 120  
**LAEMOBOTHRIDAE** 8, **31**

- Laemobothrion*** 8, 22, **31**  
***Lagopoecus*** 13, 22, **137**  
*Laminonirmus* 172  
*languidus, Halipeurus* 128  
*languidus, Lipeurus* 128  
*lapponicus, Carduiceps* 102  
***lapponicus, Carduiceps cingulatus*** 12, **102**, 103, 261  
*lari fallai, Saemundssonina* 201, 202  
*lari gonothorax, Saemundssonina* 201, 202  
*lari lari, Saemundssonina* 202  
*lari magna, Docophorus* 196  
***lari sensu lato, Saemundssonina (Saemundssonina)*** 16, 192, 194, **201**, 202, 263  
*lari, Actornithophilus* 35  
***lari, Actornithophilus piceus*** 9, **35**, 263  
*lari, Colpocephalum* 35  
*lari, Pediculus* 195, 201  
*lari, Philopterus* 201  
*lari, Saemundssonina* 201, 202  
*lari, Saemundssonina (Saemundssonina)* 192, 194  
*lari, Saemundssonina lari* 202  
***lata, Hohorstiella*** 10, 64, **65**, 264  
*laticorpus, Cuclotogaster* 107  
***latirostris, Cuculicola*** 12, **108**, 109, 265  
*latirostris, Degeeriella* 108  
*latirostris, Nirmus* 108  
*latum, Menopon* 64  
*latus, Trichodectes* 224  
*leontodon, Docophorus* 212  
*leontodon, Philopterus* 213  
*leontodon, Philopterus (Docophorus)* 213  
*lepida, Degeeriella* 185  
***Lepidophthirus*** 17, 22, **231**  
*lepidus, Nirmus* 185  
*lepidus, Quadriceps* 185  
***Lepikentron*** 17, **222**  
***leptopygos, Colpocephalum*** 9, 56, **57**, 259  
*leptopygos, Liothella* 57  
*lepturus, Lipeurus* 152  
*lepturus, Perineus diomedae* 152  
***lessonii, Naubates (Guenterion)*** 14, **144**, 252, 337  
***leucophryna, Halipeurus*** 13, 122, **125**, 126, 131, 132, 253  
*leucophryna, Halipeurus (Halipeurus)* 122, 125  
***leucoxanthum, Holomenopon*** 10, **65**, 66, 246, 247  
*leucoxanthum, Menopon* 65  
*limbata, Damalinia* 220  
*limbata, Damalinia (Bovicola)* 220  
*limbatus, Bovicola* 220  
***limbatus, Bovicola (Bovicola)*** 17, **220**, 221, 271  
*limbatus, Trichodectes* 220  
*limitatus, Lipeurus* 123  
***limosae, Actornithophilus*** 9, **33**, 34, 261  
***limosae, Austromenopon*** 9, **47**, 261  
***limosae, Luniceps*** 13, **140**, 261  
*limosae, Luniceps limosella* 140  
*limosae, Philopterus* 202, 209  
*limosae, Philopterus (Docophorus)* 202, 209

- limosae*, *Saemundssonina* 202, 209  
***limosae*, *Saemundssonina* (*Saemundssonina*)** 16, **202**, 261  
*limosella limosae*, *Lunaceps* 140  
***limosella*, *Lunaceps*** 13, **140**, 261  
*lingulata*, *Degeeriella* 181  
*lingulatus lingulatus*, *Koeniginirmus* (*Koeniginirmus*) 181  
*lingulatus sublingulatus*, *Koeniginirmus* (*Koeniginirmus*) 181  
*lingulatus*, *Koeniginirmus* 181  
*lingulatus*, *Koeniginirmus* (*Koeniginirmus*) *lingulatus* 181  
*lingulatus*, *Nirmus punctatus* 181  
*lingulatus*, *Quadriceps* 181  
***lingulatus*, *Quadriceps punctatus*** 15, **181**, 263  
**LINOGNATHIDAE** 18, **235**  
***Linognathus*** 18, 22, 25, **235**  
*Liothella* 56  
*Liotheum* 60  
*lipeuroides*, *Damalinia* 226  
*lipeuroides*, *Damalinia* (*Cervicola*) 226  
*lipeuroides*, *Damalinia* (*Tricholipeurus*) 226  
*lipeuroides*, *Eutrichophilus* 226  
*lipeuroides*, *Trichodectes* 226  
***lipeuroides*, *Tricholipeurus*** 17, **226**, 271, 353  
***Lipeurus*** 13, **137**  
*lituretum*, *Trinotum* 78  
*lobaticeps remota*, *Saemundssonina* 207  
*lobaticeps remota*, *Saemundssonina* (*Saemundssonina*) 207  
***lobodontis*, *Antarctophthirus*** 17, **229**, 230, 270  
*lockleyi sensu lato*, *Saemundssonina* 203  
*lockleyi*, *Saemundssonina* 203  
***lockleyi*, *Saemundssonina* (*Saemundssonina*)** 16, **203**, 204, 263, 264  
*longicornis*, *Bovicola* 221  
***longicornis*, *Bovicola* (*Bovicola*)** 17, 219, **221**, 271, 350  
*longicornis*, *Damalinia* 221  
*longicornis*, *Damalinia* (*Bovicola*) 221  
*longicornis*, *Damalinia* (*Rhabdopedilon*) 221  
*longicornis*, *Rhabdopedilon* 221  
*longicornis*, *Trichodectes* 219, 221  
***Longimenopon*** 10, 20, 22, **68**, 69  
*longithoracica*, *Procellariaphaga* 47  
***longithoracicum*, *Austromenopon*** 9, **47**, 254  
*longithoracicum*, *Menopon* 47  
*lucidum*, *Menopon* 75  
***lucidum*, *Nosopon*** 10, **75**, 259, 318  
*lugens*, *Nirmus* 190  
*lugens*, *Rallicola* 190  
***lugens*, *Rallicola* (*Rallicola*)** 16, **190**, 260  
*lugubre*, *Esthiopterum* 171  
*lugubris*, *Lipeurus* 171  
***lugubris*, *Pseudonirmus*** 15, **171**, 251, 344  
***Lunaceps*** 13, 22, **138**  
***lurida*, *Fulicoffula*** 12, **115**, 260  
*luridum*, *Esthiopterum* 115  
*luridum*, *Trinotum* 78  
*luridus*, *Lipeurus* 115  
*luridus*, *Philopterus* (*Lipeurus*) 115  
***lutescens lutescens*, *Austromenopon*** 9, **47**, 48, 260, 261



- lutescens*, *Austromenopon lutescens* 9, **47**, 48, 260, 261  
*lutescens*, *Menopon* 47  
*lyriocephalus*, *Haematopinus* 242  
**lyriocephalus**, *Haemodipsus* 18, **241**, 242, 243, 244, 269, 358  
*lyriocephalus*, *Pediculus* 242  
*lyrurus*, *Lagopoecus* 137  
*macilhennyi* *Episbates* 113  
*macilhennyi*, *Lipeurus* 113  
*Mackayia* 213  
**macquariensis sensu lato**, *Austrogoniodes* 11, **93**, 248  
**macquariensis**, *Austrogoniodes* 11, **92**, 93, 248, 249  
*macquariensis*, *Cesareus* 93  
*macquariensis*, *Pectinopygus* 159  
*macquariensis*, *Pectinopygus* (*Philichthyophaga*) 159  
*macrocephalus*, *Haematopinus* 232  
*macrocephalus*, *Haematopinus asini* 232  
*macrocephalus*, *Pediculus* 232  
**macrocephalus**, *Philopteroides* 15, **166**, 267, 268  
**macronecti**, *Perineus* 14, **163**, 251  
**macropus**, *Heterodoxus* **29**  
*macrorhini*, *Lepidophthirus* 232  
**macrorhini**, *Lepidophthirus* 17, **231**, 232, 270, 354  
**maculosus maculosus**, *Lipeurus* 13, **138**, 246, 336  
*maculosus*, *Lipeurus* (*Lipeurus*) 138  
**maculosus**, *Lipeurus maculosus* 13, **138**, 246, 336  
**madagascariensis**, *Lunaceps numenii* 13, **140**, 141, 244, 260, 336  
*magna*, *Docophorus lari* 196  
*magnicornutus*, *Anatoecus* (*Benatoecus*) *dentatus* 84  
**magnicornutus**, *Anatoecus dentatus* 11, **84**, 85, 246  
**magnificus**, *Anaticola* 11, **83**, 247, 321  
*major*, *Dennyus* (*Takamatsuia*) 60  
*major*, *Lipeurus* 127  
*major*, *Takamatsuia* 60  
*mamillatus*, *Goniodes* 120  
*mamillatus*, *Gonocephalus* 120  
*mammillatus*, *Goniodes* 120  
*marginatus*, *Acidoproctus* 80  
*marina*, *Saemundssonina* 203, 205  
**marina**, *Saemundssonina* (*Saemundssonina*) 16, **203**, 256  
*marquesanum*, *Esthiopterum* 126  
**marquesanus**, *Halipeurus* 13, **126**, 254  
*marquesanus*, *Halipeurus* (*Halipeurus*) 126  
**mawsoni**, *Antarctophthirus* 17, **230**, 270  
**mawsoni**, *Austrogoniodes* 11, **93**, 248  
*mawsoni*, *Cesareus* 93  
*maximum*, *Laemobothrion* 31  
*megalosoma*, *Amyrsidea* 37  
*megalosomum*, *Menopon* 37  
*meinertzhageni sensu lato*, *Forficuloecus* 114  
**meinertzhageni**, *Forficuloecus* 12, **113**, 114, 264  
*melanocephalus*, *Docophorus* 195, 203, 204  
*melanocephalus*, *Philopterus* 195, 203, 204  
*melanocephalus*, *Saemundssonina* 203, 204  
**melanocephalus**, *Saemundssonina* (*Saemundssonina*) 16, **204**, 263  
**meleagridis**, *Chelopistes* 12, 71, 103, **104**, 246, 326  
*meleagridis*, *Goniodes* 103, 104

- meleagridis*, *Pediculus* 71, 104  
*meleagridis*, *Ricinus* 104  
*meleagridis*, *Virgula* 104  
***Melibrueelia*** 14, 20, 22, **142**  
***Menacanthus*** 10, 22, **69**, 72  
*menadense*, *Coloceras* 104  
*menadensis*, *Goniocotes* 104  
***Menopon*** 10, 22, 31, **72**  
**MENOPONIDAE** 8, **31**  
*mergiserrati*, *Anaticola* 83  
*merulae*, *Docophorus* 169  
*merulae*, *Philopterus* 169  
*merulae*, *Philopterus* (*Docophorus*) 169  
***merulensis*, *Brueelia*** 12, 96, **98**, 99, 268  
*merulensis*, *Bruelia* 98  
*merulensis*, *Bruëlia* 98  
*merulensis*, *Degeeriella* 98  
*merulensis*, *Philopterus* (*Nirmus*) 98  
*merulensis*, *Turdinirmus* 98  
***mesopelios colchicus*, *Oxylipeurus*** 14, **151**, 246  
*mesopelios*, *Oxylipeurus* 151  
*mexicana*, *Proneptis semifissa* 184  
*mexicana*, *Quadriceps semifissa* 184  
*mexicanus*, *Quadriceps* 184  
*mexicanus*, *Quadriceps semifissus* 184  
***meyeri*, *Austromenopon*** 9, **48**, 261  
*Meyeri*, *Menopon* 48  
*micariproctus*, *Halipeurus* (*Halipeurus*) 124  
***microchir*, *Antarctophthirus*** 17, **230**, 231, 270, 354  
*microchir*, *Echinophthirus* 230  
*minus*, *Nosopon* 75  
***minuta*, *Amyrsidea* (*Argimenopon*)** 9, **37**, 246, 308  
*mirabilis*, *Giebelia* 213, 216  
***mirabilis*, *Halipeurus*** 13, **126**, 127, 255  
*mirabilis*, *Halipeurus* (*Halipeurus*) 126  
*mirabilis sensu lato*, *Trabeculus* 216, 217  
***mirabilis*, *Trabeculus*** 17, 213, **216**, 217, 255, 256  
*miriceps sensu lato*, *Paraclisis* 154  
*miriceps*, *Esthiopterum* 154  
*miriceps*, *Harrisoniella* 154  
*miriceps*, *Lipeurus* 154  
***miriceps*, *Paraclisis*** 14, **154**, 251, 340  
*miriceps*, *Perineus* 154  
***mirum*, *Apterygon*** 9, 39, **40**, 245  
*modestum*, *Esthiopterum* 120  
*modestus*, *Lipeurus* 120  
*modestus*, *Perineus* 120  
***modularis*, *Philopterus*** 15, **167**, 168, 269  
*modularis*, *Philopterus* (*Docophorus*) 167  
***mundae*, *Halipeurus*** 13, **127**, 129, 256  
*mundae*, *Halipeurus* (*Halipeurus*) 127  
***murphyi*, *Docophoroides*** 12, **111**, 251  
*murphyi*, *Eurymetopus* 111  
*mustelae*, *Pediculus* 225  
*mustelae*, *Stachiella* 226  
*mustelae*, *Trichodectes* 225

- mustelae*, *Trichodectes* (*Stachiella*) 17, **225**, 226, 270  
*mutabilis*, *Menacanthus* 69  
*Myrsidea* 10, 22, **72**, 73, 75  
*nadleri*, *Puffinoecus* 193  
*Naubates* 14, 22, **142**, **146**  
*navigans*, *Austromenopon* 9, **48**, 49, 249, 250, 309  
*navigans*, *Menopon* 48  
*navigans*, *Procellariphaga* 48  
*nebulosa*, *Brueelia* 12, **99**, 268  
*nebulosa*, *Bruëlia* 99  
*nebulosa*, *Degeeriella* 99  
*nebulosus*, *Nirmus* 99  
*Neocolpocephalum* 56  
*neopallidus*, *Ardeicola* 11, **88**, 258, 323  
*Neopsittaconirmus* 14, 22, **147**  
*nereis*, *Saemundssonina* 204  
*nereis*, *Saemundssonina* (*Saemundssonina*) 16, **204**, 256  
*Nesiotinus* 14, 22, **148**, 149  
new species, *Apterygon* 41  
*niger*, *Triniton* 78  
*nigrolimbatus*, *Esthiopterum* 163  
*nigrolimbatus*, *Lipeurus* 163  
*nigrolimbatus*, *Perineus* 14, 161, **163**, 164, 251  
*nigrum*, *Trinoton* 11, **78**, 247, 319  
*Nirmus* 109  
*Nitzschia* 60  
*Nitzschiella* 104  
*nivea*, *Saemundssonina* 195  
*noctivagus*, *Halipeurus* 13, **127**, 252  
*normifer alpha*, *Koeniginirmus* 178  
*normifer alpha*, *Koeniginirmus* (*Laminonirmus*) 178  
*normifer alpha*, *Quadriceps* 15, **178**, 262  
*normifer normifer*, *Koeniginirmus* 178  
*normifer normifer*, *Koeniginirmus* (*Laminonirmus*) 178  
*normifer normifer*, *Quadriceps* 15, **178**, 262  
*normifer parvopallidus*, *Koeniginirmus* (*Laminonirmus*) 179  
*normifer parvopallidus*, *Koeniginirmus* 179  
*normifer parvopallidus*, *Quadriceps* 15, **179**, 262  
*normifer*, *Degeeriella* 178  
*normifer*, *Koeniginirmus* 178  
*normifer*, *Koeniginirmus* (*Laminonirmus*) *normifer* 178  
*normifer*, *Koeniginirmus normifer* 178  
*normifer*, *Nirmus* 178  
*normifer*, *Quadriceps* 178, 179  
*normifer*, *Quadriceps normifer* 15, **178**, 262  
*Nosopon* 10, 22, **75**  
*notafusca*, *Boopia* 8, **29**, 269, 305  
*novaeseelandiae*, *Coloceras* 12, 104, **105**, 106, 264, 327  
*novaeseelandiae*, *Melibruelia* 14, **142**, 267, 337  
*novaeseelandiae*, *Myrsidea* 10, **73**, 267  
*novaeseelandiae*, *Patellinirmus* 104, 105  
*novaeseelandiae*, *Quadriceps* 15, **179**, 180, 262  
*novaezealandiae*, *Philopterus* 15, **168**, 266, 344  
*novaezealandiae*, *Rallicola* 186  
*novaezealandiae*, *Rallicola* (*Aptericola*) 186  
*novaezealandiae*, *Philopteroides* 15, 165, **166**, 266

- numenii madagascariensis*, *Luniceps* 13, **140**, 141, 244, 260, 336  
*numenii numenii*, *Luniceps* 140, 244  
*numenii oliveri*, *Luniceps* 13, **141**, 261  
*numenii phaeopi*, *Luniceps* 141  
*numenii*, *Luniceps* 140, 141  
*numenii*, *Luniceps numenii* 140, 244  
*numeniicola*, *Philopterus* 197  
*numeniicola*, *Saemundssonina* 197  
*numerosa*, *Procellariophaga* 43  
*numerosum*, *Menopon* 43  
*numerosus*, *Procellariophaga* 43  
*numidianus*, *Goniodes* 137  
*numidianus*, *Lagopoecus* 137  
*nychthemerus birostris*, *Koeniginirmus* 173  
*nychthemerus birostris*, *Koeniginirmus (Laminonirmus)* 173  
*nychthemerus nychthemerus*, *Koeniginirmus* 180  
*nychthemerus nychthemerus*, *Koeniginirmus (Laminonirmus)* 180  
*nychthemerus*, *Koeniginirmus (Laminonirmus) nychthemerus* 180  
*nychthemerus*, *Koeniginirmus nychthemerus* 180  
*nychthemerus*, *Quadriceps* 180  
*nycthemera*, *Degeeriella* 180  
*nycthemerus*, *Nirmus* 180  
*nycthemerus*, *Quadriceps* 15, **180**, 263  
*obscura*, *Harrisoniella* 154  
*obscura*, *Paraclisis* 14, **154**, 155, 251  
*obscurum*, *Esthiopterum* 154  
*obscurus*, *Lipeurus* 154  
*obscurus*, *Perineus* 154  
*obscurus*, *Perineus* (s. l.) 154  
*ocellata*, *Damalinia* 228  
*ocellata*, *Damalinia (Werneckiella)* 228  
*ocellata*, *Werneckiella* 17, **228**, 271, 353  
*ocellatus*, *Bovicola* 228  
*ocellatus*, *Bovicola (Werneckiella)* 228  
*ocellatus*, *Philopterus* 167  
*ocellatus*, *Philopterus (Docophorus)* 167  
*ocellatus*, *Trichodectes* 228  
*ochraceum*, *Colpocephalum* 33  
*ochraceum*, *Liotheum (Colpocephalum)* 34  
*ochraceus sensu lato*, *Actornithophilus* 9, **34**, 261, 262  
*oenomydis*, *Hoplopleura* 234  
*oestrelatae*, *Cecalymenus* 213, 217  
*ogmorhini*, *Antarctophthirus* 17, 229, **231**, 270  
*okarito*, *Apterygon* 9, **41**, 245, 309  
*oliveri*, *Degeeriella* 141  
*oliveri*, *Luniceps* 141  
*oliveri*, *Luniceps numenii* 13, **141**, 261  
*oloris*, *Anatoecus (Anatoecus) icterodes* 85  
*oloris*, *Anatoecus icterodes* 11, **85**, 86, 246, 322  
*Oncophorus* 213  
*oraria*, *Degeeriella* 175  
*orarius*, *Nirmus* 175  
*orarius*, *Quadriceps* 175  
*orarius*, *Quadriceps charadrii* 15, **175**, 261  
*orientalis*, *Puffinoecus* 193  
*orientalis*, *Saemundssonina* 193

- orientalis*, *Saemundssonina* (*Puffinoecus*) 16, **193**, 254  
*ornatus fuscolaminulatus*, *Koeniginirmus* 180  
*ornatus fuscolaminulatus*, *Koeniginirmus* (*Laminonirmus*) 180  
*ornatus fuscolaminulatus*, *Quadriceps* 15, **180**, 181, 182, 263  
*ornatus ornatus*, *Quadriceps* 172  
*ornatus*, *Koeniginirmus* 172  
***Ornithobius*** 14, 22, **149**, 150  
*ortygis*, *Goniodes* 13, **118**, 119, 246  
*ortygis*, *Philopterus* (*Goniodes*) 118  
*ortyometrae philippensis*, *Rallicola* 190  
***ortyometrae philippensis*, *Rallicola* (*Rallicola*)** 16, **190**, 259  
*ortyometrae*, *Pediculus* 190  
*ortyometrae*, *Rallicola* 190  
*ortyometrae*, *Rallicola* (*Rallicola*) 186  
*oschei*, *Austromenopon* 43  
***ossifragae*, *Austromenopon*** 9, 41, **49**, 251  
*ossifragae*, *Procellariphaga* 41, 49  
*Oulocrepis* 117  
***ovalis*, *Gyropus*** 8, **30**, 31, 269, 306  
*ovillus*, *Haematopinus* 235  
***ovillus*, *Linognathus*** 18, 25, **235**, 271  
*ovis*, *Bovicola* 222  
***ovis*, *Bovicola* (*Bovicola*)** 17, **222**, 271  
*ovis*, *Damalinia* 222  
*ovis*, *Damalinia* (*Bovicola*) 222  
*ovis*, *Pediculus* 222  
*ovis*, *Trichodectes* 222  
***Oxylipeurus*** 14, 22, **150**  
***pacifica*, *Hoplopleura*** 18, **234**, 235, 269, 355  
*pacificus*, *Halipeurus* (*Halipeurus*) *falsus* 124  
***pacificus*, *Halipeurus falsus*** 13, 25, **124**, 257  
*painei*, *Colpocephalum* 67  
*painei*, *Kurodaia* (*Kurodaia*) 67  
***pallida*, *Franciscoloa* (*Franciscoloa*)** 10, **63**, 264, 313  
*pallidulum*, *Menopon* (*Menacanthus*) 70  
***pallidulus*, *Menacanthus*** 10, **70**, 246, 316  
*pallidum*, *Colpocephalum* 63  
*pallidum*, *Esthiopterum* 88  
*pallidum*, *Liotheum* (*Menopon*) 72  
*pallidum*, *Menopon* 72  
*pallidum*, *Psittacomenopon* 63  
*pallidus*, *Ardeicola* 88  
***Paraclisis*** 14, 22, **152**  
*parallela*, *Damalinia* 227  
*parallela*, *Damalinia* (*Tricholipeurus*) 227  
*parallelus*, *Damalinia* (*Cervicola*) 227  
*parallelus*, *Trichodectes* 227  
***parallelus*, *Tricholipeurus*** 17, **227**, 271  
*parumpilosus* var. *ocellata*, *Trichodectes* 228  
*parumpilosus*, *Trichodectes* 227  
***parviguttata*, *Brueelia*** 12, **99**, 100, 268  
*parviguttata*, *Bruëlia* 99  
*parviguttata*, *Degeeriella* 99  
*parvopallidus*, *Koeniginirmus* (*Laminonirmus*) *normifer* 179  
*parvopallidus*, *Koeniginirmus normifer* 179  
*parvopallidus*, *Quadriceps* 179

- parvopallidus*, *Quadriceps normifer* 15, **179**, 262  
*Patellinirmus* 104  
*paulula*, *Procellariphaga* 49  
*paululum*, *Austromenopon* 9, **49**, 50, 254, 255, 256  
*paululum*, *Menopon* 49  
*pavonis*, *Goniodes* 13, 117, **119**, 246  
*pavonis*, *Pediculus* 119  
*pavonis*, *Ricinus* 119  
*pectinatum* *Colpocephalum* 55  
*pectinatum* *Colpocephalum* 55  
*pectinatum*, *Menopon* 55  
*pectiniventre*, *Colpocephalum* 55  
*pectiniventris*, *Ciconiphilus* 9, **55**, 56, 247  
*Pectinopygus* 14, 22, 26, **155**, 157, 160  
*pedalis*, *Haematopinus* 236  
*pedalis*, *Linognathus* 18, **236**, 271  
*pedalis*, *Trichaulus* 236  
*pederiforme*, *Esthiopterum* 113  
*pederiformis*, *Episbates* 12, **113**, 249, 330  
*pederiformis*, *Phlopterus* 113  
**PEDICULIDAE** 18, **239**  
*pediculoides*, *Actornithophilus* 9, **34**, 35, 261  
*pediculoides*, *Colpocephalum* 34  
*pediculoides*, *Longimenopon* 35  
*pediculoides*, *Rediella* 34  
**Pediculus** 18, 22, **239**  
*pelagicum*, *Esthiopterum* 128  
*pelagicus*, *Halipeurus* 13, 121, **128**, 129, 256  
*pelagicus*, *Halipeurus* (*Synnautes*) 128, 129  
*pelagicus*, *Lipeurus* 121  
*pelagicus*, *Phlopterus* (*Lipeurus*) 128  
*pelagicus sensu lato*, *Halipeurus* (*Synnautes*) 128, 129  
*pelagicus*, *Synnautes* 128  
*pelagodromae*, *Halipeurus* 13, **128**, 129, 256  
*Pelecanigogus* 56  
*pellucida*, *Eidmanniella* 10, 60, **61**, 62, 257, 258  
*pellucidum*, *Menopon* 61  
**Pelmatocerandra** 14, 22, **160**  
**Penenirmus** 14, 22, **161**  
*perdicis*, *Amyrsidea* (*Argimenopon*) 9, **37**, 38, 246  
*perdicis*, *Liotheum* (*Menopon*) 37  
**Perineus** 14, 23, **161**  
*pertusus porzanae*, *Incidifrons* 136  
*pertusus*, *Docophorus* 136  
*pertusus*, *Incidifrons* 136  
*pertusus*, *Phlopterus* 136  
*petersi*, *Saemundssonina* 195  
*peusi*, *Puffinoecus* 192  
*peusi*, *Saemundssonina* (*Puffinoecus*) 192  
*phaeopi*, *Luniceps* 141  
*phaeopi*, *Luniceps numenii* 141  
*phaeopi*, *Phlopterus* (*Nirmus*) 141  
*phaeopodis*, *Austromenopon* 9, **50**, 51, 261  
*phaeopodis*, *Pediculus* 50  
*phaetona*, *Saemundssonina* 195  
*Philichthyophaga* 155

- philippensis*, *Rallicola* (*Rallicola*) *ortygometrae* 16, **190**, 259  
*philippensis*, *Rallicola* *ortygometrae* 190  
*Philoceanus* 15, 23, 26, **164**  
**PHILOPTERIDAE** 11, **80**  
*Philopteroides* 15, 23, **165**  
*Philopterus* 15, 23, **167**, 168, 169, 170  
**PHTHIRAPTERA** 3, 5, 7, 8, 19, 20, **29**  
PHTHIRIDAE 240  
*Phthirus* 241  
*Phthirus* 241  
*Phthirus* 241  
*Physostomum* 79  
*piceus lari*, *Actornithophilus* 9, **35**, 263  
*piceus* sensu lato, *Actornithophilus* 35  
*piekarskii*, *Austromenopon* 50  
*pilgrimi*, *Ardeicola* 11, 25, **88**, 258  
*pilgrimi*, *Colpocephalum* 9, 25, **57**, 58, 264, 311  
*pilgrimi*, *Forficuloecus* 12, **114**, 265, 331  
*pilgrimi*, *Philopteroides* 15, **166**, 267  
*pilgrimi*, *Pseudomenopon* 11, 25, **77**, 259, 319  
*pilgrimi*, *Rallicola* 187  
*pilgrimi*, *Rallicola* (*Aptericola*) 16, 25, **187**, 188, 245  
*piliferous*, *Haematopinus* 236  
*piliferus*, *Haematopinus* 236  
*piliferus*, *Linognathus* 236  
*piliferus*, *Pediculus* 236  
*piliferus*, *Trichaulus* 235, 236  
*pilosum*, *Pseudomenopon* 11, 76, **77**, 260  
*pilosus*, *Pediculus* 77  
*pilosus*, *Trichodectes* 227  
*pingue*, *Austromenopon* 51  
*pingue*, *Colpocephalum* 51  
*pinguis*, *Austromenopon* 9, **51**, 249, 250  
*pinguis*, *Procellariphaga* 51  
*piratae*, *Perineus* 120  
*placodus*, *Halipeurus* 13, 127, **129**, 255, 256  
*placodus*, *Halipeurus* (*Halipeurus*) 129  
*plataleae*, *Ardeicola* 11, **88**, 89, 259  
*plataleae*, *Esthiopterum* 88  
*plataleae*, *Ibidoecus* 13, 134, **135**, 136, 259  
*plataleae*, *Pediculus* 88  
*plataleae*, *Philopterus* 135  
*plataleae*, *Philopterus* (*Docophorus*) 135  
*platycephalus*, *Docophorus* 204  
*platycephalus*, *Philopterus* 204  
*platycephalus*, *Saemundssonina* 205  
*platycephalus*, *Saemundssonina* (*Saemundssonina*) 16, **204**, 205, 256  
*platygaster balati*, *Saemundssonina* (*Saemundssonina*) 16, **206**, 262  
*platygaster* sensu lato, *Saemundssonina* (*Saemundssonina*) 16, **205**, 206, 261, 262  
*platygaster*, *Cuculiphilus* (*Cuculiphilus*) 10, **59**, 60, 265  
*platygaster*, *Cuculiphilus cuculiphilus* 59  
*platygaster*, *Menopon* 59  
*platygaster*, *Philopterus* 205  
*platygaster*, *Philopterus* (*Docophorus*) 205  
*platygaster*, *Saemundssonina* 205, 206  
**Plegadiphilus** 10, 23, **75**

- plegadis*, *Menopon* 75  
***plegadis*, *Plegadiphilus*** 10, **75**, 76, 259, 318  
*podiceps*, *Aquanirmus* 11, **87**, 248  
*podiceps*, *Philopterus* (*Nirmus*) 87  
*podicipis*, *Aquanirmus* 87  
*podicipis*, *Degeeriella* 87  
**POLYPLACIDAE** 18, **241**  
***Polyplax*** 18, 23, 241, **243**  
***polytrapezius polytrapezius*, *Oxylipeurus*** 14, **152**, 246, 340  
*polytrapezius*, *Lipeurus* 152  
*polytrapezius*, *Lipeurus* (*Oxylipeurus*) 152  
*polytrapezius*, *Oxylipeurus* 152  
***polytrapezius*, *Oxylipeurus polytrapezius*** 14, **152**, 246, 340  
*polytrichum*, *Amyrsidea* (*Argimenopon*) 37  
*polytrichum*, *Argimenopon* 37  
*popellus* sensu lato, *Austromenopon* 51, 52  
***popellus*, *Austromenopon*** 9, **51**, 52, 53, 251, 252, 254  
*popellus*, *Menopon* 51  
*popellus*, *Procellariphaga* 51  
***porcelli*, *Gliricola* (*Gliricola*)** 8, **30**, 269, 306  
***porzanae*, *Incidifrons*** 13, **136**, 137, 260, 335  
*porzanae*, *Incidifrons pertusus* 136  
*potens*, *Lipeurus* 155  
*potens*, *Pectinopygus* 155  
*priapulus*, *Halipeurus* (*Halipeurus*) *gravis* 124  
***priapulus*, *Halipeurus gravis*** 13, **124**, 255  
***pricei*, *Halipeurus*** 13, **129**, 130, 253  
*prioni*, *Lipeurus* 144  
*prioni*, *Naubates* 144  
***prioni*, *Naubates* (*Guenterion*)** 14, 133, 143, **144**, 145, 253, 254  
*procellariae* sensu lato, *Halipeurus* (*Halipeurus*) 130  
*procellariae*, *Ancistrona* 38  
*procellariae*, *Esthiopterum* 130  
***procellariae*, *Halipeurus*** 13, 122, **130**, 251, 252  
*procellariae*, *Halipeurus* (*Halipeurus*) 130  
*procellariae*, *Pediculus* 130  
*procellariae*, *Ricinus* 130  
*Procellariphaga* 41  
*Proneptis* 172  
*Pseudocolpocephalum* 55  
***Pseudomenopon*** 11, 23, **76**  
***Pseudonirmus*** 15, 23, **170**  
*Psittacicola* 147  
***Psittoecus*** 15, 23, **171**  
*pterodromae*, *Saemundssonina* 206  
***pterodromae*, *Saemundssonina* (*Saemundssonina*)** 16, **206**, 251, 253  
*pterodromi* sensu lato, *Naubates* 144  
*pterodromi*, *Naubates* 143, 145  
***pterodromi*, *Naubates* (*Guenterion*)** 14, 143, 144, **145**, 252  
**PTHIRIDAE** 18, **240**  
***Pthirus*** 18, 23, **240**  
*pubis*, *Pediculus* 240  
*pubis*, *Phthirus* 241  
*pubis*, *Phthirus* 241  
***pubis*, *Pthirus*** 18, **241**, 272, 358  
*puellula*, *Saemundssonina* 193



- puellula*, *Saemundssonina* (*Puffinoecus*) 16, **193**, 255  
*puellulus*, *Puffinoecus* 193  
***Puffinoecus*** 16, **192**, 194  
*puffinus*, *Longimenopon* 68  
*pullatus*, *Lipeurus* 155  
*punctatus lingulatus*, *Nirmus* 181  
***punctatus lingulatus*, *Quadriceps*** 15, **181**, 263  
*punctatus punctatus*, *Quadriceps* 181  
*punctatus sensu lato*, *Pectinopygus* 158, 159  
*punctatus sensu lato*, *Quadriceps* 181, 182  
***punctatus sublingulatus*, *Quadriceps*** 15, **181**, 182, 263  
*punctatus*, *Koeniginirmus* 172  
***punctatus*, *Pectinopygus*** 14, 25, **158**, 159, 258  
*punctatus*, *Quadriceps* 172, 181, 182  
*punctatus*, *Quadriceps punctatus* 181  
*punctifer*, *Quadriceps charadrii* 175  
***pusillus*, *Goniocotes*** 12, **116**, 117, 246, 332  
*pusillus*, *Goniodes* 116  
*pusillus*, *Trichodectes* 225  
***pustulosa*, *Eidmanniella*** 10, **62**, 257  
*pustulosa*, *Saemundssonina* 196  
*pustulosum*, *Menopon* 62  
*pustulosus*, *Docophorus* 196  
*pustulosus*, *Philopterus* 198  
***Quadriceps*** 15, 23, **172**, 186  
*quadripustulatus*, *Ciconiphilus* 55  
*quadripustulatus*, *Colpocephalum* 55  
*querquedulae sensu lato*, *Trinoton* 79  
*querquedulae*, *Pediculus* 78  
***querquedulae*, *Trinoton*** 11, **78**, 79, 247, 248  
***Rallicola*** 15, 16, 20, 23, 25, **186**, **189**  
*raphanus*, *Halipeurus* 121  
*remota*, *Saemundssonina* 207  
***remota*, *Saemundssonina* (*Saemundssonina*)** 16, **207**, 263  
*remota*, *Saemundssonina* (*Saemundssonina*) *lobaticeps* 207  
*remota*, *Saemundssonina lobaticeps* 207  
***renschii*, *Quadriceps*** 15, **182**, 262  
***restinctum*, *Coloceras*** 12, **106**, 264  
*restinctus*, *Coloceras* 106  
*restinctus*, *Patellinirmus* 106  
***retractus*, *Goniodes*** 13, **119**, 120, 246, 332  
*retractus*, *Gonoidea* 119  
*rhaphidium*, *Esthiopterum* 89  
***rhaphidius*, *Ardeicola*** 11, **89**, 259  
*rhaphidius*, *Lipeurus* 89  
***rhpidurae*, *Menacanthus*** 10, **70**, 71, 267  
**RHYNCHOPHTHIRINA** 19  
**RICINIDAE** 11, **79**  
***Ricinus*** 11, 23, **79**  
*ridgwayi*, *Degeeriella* 182  
*ridgwayi*, *Nirmus* 182  
***ridgwayi*, *Quadriceps*** 15, **182**, 183, 261  
*ridibundis*, *Liotheum* (*Menopon*) 53  
*ridibundum*, *Menopon* 53  
*ridibundus*, *Austromenopon* 53  
*ridibundus*, *Liotheum* (*Menopon*) 53

- robertsi*, *Naubates* 165  
***robertsi*, *Philoceanus*** 15, **165**, 256  
*robustum*, *Eucolpocephalum* 63  
*robustum*, *Menopon* 69  
*robustus*, *Menacanthus* 69  
***rodericki*, *Rallicola* (*Aptericola*)** 16, **188**, 245, 346  
*rostratus*, *Acidoproctus* 80  
*rothkoi*, *Luniceps* 139  
*rotundatus*, *Docophorus* 207  
***rufa rufa*, *Degeeriella*** 12, **109**, 259  
***rufa*, *Degeeriella rufa*** 12, **109**, 259  
*rufus*, *Nirmus* 109  
*runcinata*, *Degeeriella* 86  
*runcinatus*, *Aquanirmus* 86  
***Saemundsson*** 16, 23, **191**, **194**  
*sawadai*, *Halipeurus* 121  
*sawadai*, *Halipeurus* (*Halipeurus*) 121  
*scalaris*, *Trichodectes* 219  
***schillingi sensu lato*, *Trabeculus*** 17, 217, **218**, 251, 252  
*Schillingi*, *Docophorus* 192, 217  
***schillingi*, *Trabeculus*** 17, 192, 213, **217**, 252  
*schillingii*, *Oncophorus* 213, 217  
*scolopacis phaeopodis*, *Pediculus* 207  
*scolopacisphaeopodis scolopacisphaeopodis*, *Saemundsson* 207  
*scolopacis-phaeopodis scolopacis-phaeopodis*, *Saemundsson* 207  
***scolopacisphaeopodis scolopacisphaeopodis*, *Saemundsson* (*Saemundsson*)** 16, **207**, 208, 261  
*scolopacisphaeopodis*, *Pediculus* 207  
*scolopacisphaeopodis*, *Saemundsson* 207  
*scolopacis-phaeopodis*, *Saemundsson* 207  
***scolopacisphaeopodis*, *Saemundsson* (*Saemundsson*) *scolopacisphaeopodis*** 16, **207**, 208, 261  
*scolopacisphaeopodis*, *Saemundsson* *scolopacisphaeopodis* 207  
*scolopacis-phaeopodis*, *Saemundsson* *scolopacis-phaeopodis* 207  
*scopulacorne*, *Liotheum* (*Menopon*) 77  
***scopulacorne*, *Pseudomenopon*** 11, **77**, 78, 259  
*sc.-ph. scolopacis-phaeopodis*, *Saemundsson* 207  
***sedes*, *Actornithophilus*** 9, **35**, 36, 262  
*sellata*, *Degeeriella* 183  
*sellatus caspius*, *Koeniginirmus* 174  
***sellatus houri*, *Quadriceps*** 15, **183**, 184, 264  
*sellatus sellatus*, *Koeniginirmus* 183  
*sellatus sellatus*, *Koeniginirmus* (*Laminonirmus*) 183  
***sellatus sellatus*, *Quadriceps*** 15, **183**, 184, 263, 264  
*sellatus*, *Koeniginirmus* 183  
*sellatus*, *Koeniginirmus* (*Laminonirmus*) *sellatus* 183  
*sellatus*, *Koeniginirmus sellatus* 183  
*sellatus*, *Nirmus* 183  
*sellatus*, *Quadriceps* 183  
***sellatus*, *Quadriceps sellatus*** 15, **183**, 184, 263, 264  
***semiannulata*, *Brueelia*** 12, **100**, 267, 324  
*semiannulata*, *Bruelia* 100  
*semiannulata*, *Degeeriella* 100  
*semiannulatus*, *Nirmus* 100  
*semifissa mexicana*, *Proneptis* 184  
*semifissa mexicana*, *Quadriceps* 184  
*semifissa semifissa*, *Proneptis* 184  
*semifissa*, *Degeeriella* 184

- semifissa*, *Proneptis* 172, 184  
*semifissa*, *Proneptis semifissa* 184  
*semifissus mexicanus*, *Quadriceps* 184  
***semifissus*, *Quadriceps*** 15, 172, 177, **184**, 261  
*separata*, *Degeeriella* 184  
*separatus*, *Anousceps* 184  
*separatus*, *Nirmus* 172, 184  
***separatus*, *Quadriceps*** 15, 172, **184**, 185, 263  
*serini*, *Menopon* 73  
***serini*, *Myrsidea*** 10, **73**, 74, 269  
***serrata*, *Polyplax*** 18, 242, **243**, 244, 270, 359  
*serrator*, *Pectinopygus* (*Pectinopygus*) *bassanae* 156  
*serrator*, *Pectinopygus* (*Pectinopygus*) *bassani* 156  
*serratus*, *Haematopinus* 243  
*serratus*, *Pediculus* 243  
***setosa*, *Pelmatocerandra*** 14, **160**, 161, 257, 341  
*setosum*, *Colpocephalum* 57  
*setosum*, *Esthiopterum* 159  
***setosus*, *Linognathus*** 18, 235, **236**, 237, 270, 356  
*setosus*, *Lipeurus* 159, 160  
*setosus*, *Nirmus* 160  
***setosus*, *Pectinopygus*** 14, **159**, 258  
*setosus*, *Pediculus* 236  
*sigmoidalis*, *Menopon* 61  
*signatus* var. *pallida*, *Lipeurus* 88  
*simplex* sensu lato, *Docophoroides* 112  
***simplex*, *Docophoroides*** 12, **112**, 249, 250, 329  
*simplex*, *Eurymetopus* 112  
*simsi*, *Bedfordiella* 95  
*sincipitalis*, *Heteromenopon* 64  
***singhi*, *Anatoecus*** 11, **86**, 247  
*singularis*, *Menopon* 60  
*Solenodes* 117  
***Solenopotes*** 18, 23, **238**  
 sp. *incertae sedis*, *Apterygon* 39  
 sp., *Actornithophilus* 35, 36  
 sp., *Alcedoecus* 81  
 sp., *Anatoecus* 86  
 sp., *Ancistrona* 38  
 sp., *Antarctophthirus* 229, 230  
 sp., *Apterygon* 41  
 sp., *Aquanirmus* 87  
 sp., *Austrogoniodes* 93, 94  
 sp., *Austromenopon* 42, 44, 45, 48, 49, 50, 51, 53, 54  
 sp., *Brueelia* 96, 97, 98, 99, 100, 101, 142  
 sp., *Fulicoffula* 115  
 sp., *Goniodes* 118  
 sp., *Halipeurus* 122, 123, 129  
 sp., *Halipeurus* (*Halipeurus*) 122, 129  
 sp., *Harrisoniella* 134  
 sp., *Hohorstiella* 65  
 sp., *Holomenopon* 66, 67  
 sp., *Incidifrons* 136  
 sp., *Luniceps* 140, 141  
 sp., *Menacanthus* 69, 70, 71  
 sp., *Menopon* 69

- sp., *Myrsidea* 72, 73, 74  
 sp., *Naubates* 122, 142, 144  
 sp., *Nosopon* 75  
 sp., *Pectinopygus* 157, 160  
 sp., *Perineus* 162, 163  
 sp., *Perineus* (s. l.) 152  
 sp., *Philopterus* 165, 166, 167, 168, 169, 170  
 sp., *Quadriceps* 179  
 sp., *Rallicola* 188, 189, 191  
 sp., *Rallicola* (*Aptericola*) 188  
 sp., *Saemundssonina* 192, 194, 196, 198, 200, 201, 202, 203, 205, 209, 216, 245  
 sp., *Saemundssonina* (*Puffinoecus*) 192, 193, 194  
 sp., *Saemundssonina* (*Saemundssonina*) 194, 200, 210  
 sp., *Sturnidoecus* 245  
*spadix spadix, Halipeurus* 13, **130**, 131, 255  
*spadix subclavus, Halipeurus* 13, **131**, 255  
*spadix, Halipeurus (Halipeurus)* 130  
*spadix, Halipeurus spadix* 13, **130**, 131, 255  
 species **1, Austromenopon** 9, 45, 52, **53**, 252, 253  
 species **1, Holomenopon** 10, **66**, 248  
 species **1, Philopterus** 15, **169**, 267  
 species **1, Trabeculus** 17, 20, **218**, 252  
 species **2, Austromenopon** 9, 52, **53**, 54, 252  
 species **2, Holomenopon** 10, **67**, 248  
 species **2, Philopterus** 15, **170**, 267  
 species **2, Trabeculus** 17, **218**, 255  
 species **3, Austromenopon** 9, **54**, 260  
 species, *Anaticola* 11, **83**, 247  
 species, *Brueelia* 12, **101**, 266, 267  
 species, *Capraiella* 12, **102**, 266, 325  
 species, *Dennyus (Takamatsuia)* 10, **60**, 266, 312  
 species, *Forficuloecus* 12, 20, **114**, 265  
 species, *Heteromenopon (Keamenopon)* 10, **64**, 265  
 species, *Longimenopon* 10, 20, **68**, 251, 252, 253, 254, 256, 316  
 species, *Menacanthus* 10, **71**, 266  
 species, *Myrsidea* 10, **74**, 267, 268  
 species, *Pectinopygus* 14, **160**, 258  
 species, *Penenirmus* 14, **161**, 268, 342  
 species, *Quadriceps* 15, **185**, 263  
 species, *Ricinus* 11, **79**, 267, 320  
 species, *Saemundssonina (Puffinoecus)* 16, 192, **194**, 252, 255  
 species, *Saemundssonina (Saemundssonina)* 16, **211**, 256  
 species, *Strigiphilus* 17, 24, **212**, 265  
 species, *Sturnidoecus* 17, **213**, 268  
*sphaerocephalus, Trichodectes* 222  
*Spinibovicola* 17, **223**  
*spinulosa, Polyplax* 18, 242, **243**, 244, 269, 270  
*spinulosum, Colpocephalum* 36  
*spinulosus, Actornithophilus* 9, **36**, 261  
*spinulosus, Haematopinus* 243  
*spinulosus, Pediculus* 243, 244  
*Stachiella* 17, **225**, 226  
*stadleri, Docophorulus* 168  
*stadleri, Philopterus* 15, **168**, 169, 268  
*stammeri, Austromenopon* 9, 25, **52**, 253, 254  
*stammeri, Fulicoffula* 12, **115**, 259, 260, 331


- stammeri*, *Saemundssonina* 208  
*stammeri*, *Saemundssonina* (*Saemundssonina*) 16, **208**, 251  
*stefani*, *Goniodes* 13, **120**, 245  
*stellare*, *Esthiopterum* 89  
*stellaris* sensu lato, *Ardeicola* 89, 90  
*stellaris*, *Ardeicola* 11, **89**, 90, 259  
*stellaris*, *Philopteris* (*Lipeurus*) 89  
*stenopsis*, *Haematopinus* 237  
*stenopsis*, *Linognathus* 18, **237**, 271  
*stenopsis*, *Pediculus* 237  
*stenopsis*, *Trichaulis* 237  
*sternae*, *Pediculus* 208  
*sternae*, *Philopteris* 208  
*sternae*, *Ricinus* 208  
*sternae*, *Saemundssonina* 208  
*sternae*, *Saemundssonina* (*Saemundssonina*) 16, **208**, 263  
*stramineum*, *Liotheum* (*Menopon*) 71  
*stramineum*, *Menopon* 71  
*stramineus*, *Eomenacanthus* 71  
*stramineus*, *Menacanthus* 10, 69, **71**, 246  
*strepsilaris*, *Degeeriella* 185  
*strepsilaris*, *Quadriceps* 15, **185**, 261  
*stresemanni*, *Saemundssonina* 198, 199  
*stresemanni*, *Saemundssonina* (*Saemundssonina*) 199  
*Strigiphilus* 16, 23, **211**  
*strutheus nomen dubium*, *Austrogoniodes* 11, 20, 91, **94**, 248  
*sturni*, *Pediculus* 213  
*sturni*, *Philopteris* 213  
*sturni*, *Sturnidoecus* 17, 212, **213**, 268, 349  
*Sturnidoecus* 17, 23, **212**, 213  
*stylifer*, *Philopteris* (*Goniodes*) 104  
*stylifer*, *Rhopaloceras* 103, 104  
*styliferum*, *Rhopaloceras* 104  
*subaequale*, *Colpocephalum* 57  
*subaequale*, *Corvocolpocephalum* 57  
*subaequale*, *Liotheum* (*Colpocephalum*) 57  
*subangusticeps*, *Halipeurus* 128  
*subangusticeps*, *Lipeurus* 128  
*subclavus*, *Halipeurus* 131  
*subclavus*, *Halipeurus* (*Halipeurus*) 131  
*subclavus*, *Halipeurus spadix* 13, **131**, 255  
*subcuspidata*, *Capraiella* 102  
*subcuspidata*, *Degeeriella* 102  
*sublingulatus*, *Koeniginirmus* (*Koeniginirmus*) *lingulatus* 181  
*sublingulatus*, *Quadriceps* 181  
*sublingulatus*, *Quadriceps punctatus* 15, **181**, 182, 263  
*subrostrata*, *Felicola* 224  
*subrostratus*, *Felicola* 224  
*subrostratus*, *Felicola* (*Felicola*) 17, 223, **224**, 270, 351  
*subrostratus*, *Trichodectes* 223, 224  
*subrotunda*, *Eidmanniella* 61  
*subrotundum*, *Menopon* 61  
*subsetosus*, *Lipeurus* 157  
*subsetosus*, *Pectinopygus* 157  
*subzerafae*, *Colpocephalum* 10, **58**, 259  
*suis suis*, *Haematopinus* 234


- suis*, ***Haematopinus*** 18, 232, **234**, 271  
*suis*, *Haematopinus suis* 234  
*suis*, *Pediculus* 234  
*sula*, *Eidmanniella* 60  
***superciliosus*, *Lunaceps*** 14, **141**, 142, 260  
*Synnautes* 121  
***synoicus*, *Cuclotogaster*** 12, **108**, 246, 328  
*synoicus*, *Gallipeurus* 108  
*tabuensis*, *Rallicola* 191  
***tabuensis*, *Rallicola* (*Rallicola*)** 16, **191**, 259, 260, 347  
*tadornae* sensu lato, *Holomenopon* 66  
***tadornae*, *Holomenopon*** 10, **66**, 247, 315  
*tadornae*, *Menopon* 66  
*tadornae*, *Phlopterus* 66  
*takahe*, *Rallicola* 191  
***takahe*, *Rallicola* (*Rallicola*)** 16, 25, **191**, 260  
***Takamatsuia*** 10, **60**  
***tarsata*, *Boopia*** **29**  
*taurus*, *Lipeurus* 110  
***tausi*, *Colpocephalum*** 10, 56, **58**, 246  
*tausi*, *Galliferrisia* 58  
*taxosetus*, *Halipeurus* (*Halipeurus*) 130  
***thalassidromae incisa*, *Saemundssonina* (*Saemundssonina*)** 16, **208**, 209, 256  
***theresae*, *Halipeurus*** 13, **131**, 132, 252  
*theresae*, *Halipeurus* (*Halipeurus*) 131  
*thieli*, *Naubates* 147  
***thieli*, *Naubates* (*Naubates*)** 14, **147**, 252, 338  
***thompsoni*, *Halipeurus*** 13, **132**, 255  
*thompsoni*, *Halipeurus* (*Halipeurus*) 132  
*thompsoni*, *Harrisoniella* 134  
*thompsoni*, *Saemundssonina* 209  
***thompsoni*, *Saemundssonina* (*Saemundssonina*)** 16, **209**, 261  
***thoracica*, *Myrsidea*** 10, **74**, 268, 317  
*thoracicum*, *Menopon* 74  
***threskiornis*, *Plegadiphilus*** 11, 75, **76**, 259  
*timidum*, *Colpocephalum* 34  
*timidus*, *Actornithophilus* 34  
***timorensis*, *Hohorstiella*** 10, **65**, 254, 314  
***tinnunculi*, *Laemobothrion* (*Laemobothrion*)** 8, **31**, 259, 307  
*tinnunculi*, *Pediculus* 31  
*tinnunculi*, *Ricinus* 31  
***Trabeculus*** 17, 23, **213**, 215, 216, 218, 219  
***transversum*, *Austromenopon*** 9, **53**, 263  
*transversum*, *Liotheum* (*Menopon*) 53  
*transversum*, *Menopon* 53  
*transversus*, *Liotheum* (*Menopon*) 53  
*triangulatus* var. *alpha*, *Nirmus* 178  
*triangulatus* var., *Nirmus* 179  
*tribulis*, *Ornithobius bucephalus* 149  
*Trichaulus* 235  
*trichechi*, *Antarctophthirus* 229  
*trichechi*, *Arctophthirus* 229  
***Trichodectes*** 17, 23, 219, **224**  
**TRICHODECTIDAE** 17, **219**  
***Tricholipeurus*** 17, 23, **226**  
*tridens*, *Menopon* 76, 77

- tridens*, *Pseudomenopon* 77  
***tringae sensu lato*, *Saemundssonina* (*Saemundssonina*)** 16, 194, **209**, 210, 260  
*tringae tringae*, *Saemundssonina* (*Saemundssonina*) 210  
*tringae*, *Pediculus* 209  
*tringae*, *Saemundssonina* 209, 210  
***tringae*, *Saemundssonina* (*Saemundssonina*)** **194**, 210  
*tringae*, *Saemundssonina* (*Saemundssonina*) *tringae* 210  
***Trinoton*** 11, 23, **78**  
*Trinotum* 78  
*turbinatum sensu lato*, *Colpocephalum* 59  
***turbinatum*, *Colpocephalum*** 10, 56, **59**, 259, 264  
*turbinatum*, *Liotheum* (*Colpocephalum*) 59  
*turbinatus*, *Oncophorus* 159  
***turbinatus*, *Pectinopygus*** 14, **159**, 160, 258  
*turdi*, *Docophorus* 169  
***turdi*, *Philopterus*** 15, **169**, 268  
*turdi*, *Philopterus* (*Docophorus*) 169  
*turdinulae etemitatus*, *Brueelia* 100  
***turdinulae*, *Brueelia*** 12, **100**, 268  
***turtur*, *Halipeurus*** 13, 123, **132**, 133, 253  
*turtur*, *Halipeurus* (*Halipeurus*) 132  
*umbrinum*, *Colpocephalum* 36  
*umbrinus sensu lato*, *Actornithophilus* 36, 37  
***umbrinus*, *Actornithophilus*** 9, **36**, 37, 260  
*umbrosum*, *Colpocephalum* 36  
*umbrosus*, *Actornithophilus* 36  
***unica*, *Bedfordiella*** 11, **95**, 251, 324  
*upolensis*, *Saemundssonina* 210  
*upoluensis*, *Saemundssonina* 210  
*uppalensis*, *Docophorus* 210  
*uppalensis*, *Philopterus* 210  
*uppalensis*, *Saemundssonina* 210  
***uppalensis*, *Saemundssonina* (*Saemundssonina*)** 16, **210**, 257  
*urius*, *Haematopinus* 234  
*urius*, *Pediculus* 234  
***vagelli*, *Ancistrona*** 9, **38**, 39, 251, 252, 253, 254, 255, 256, 308  
*vagelli*, *Pediculus* 38  
*valida*, *Saemundssonina* 194  
***valida*, *Saemundssonina* (*Puffinoecus*)** 16, **194**, 255  
*validus* var. *orientalis*, *Philopterus* 193  
*validus*, *Docophorus* 194  
*validus*, *Philopterus* 194  
*validus*, *Puffinoecus* 194  
***vanalphenae*, *Austrogoniodes*** 11, **94**, 249  
*vanelli*, *Degeeriella* 172  
***vanzolinii*, *Psittoecus*** 15, **171**, 172, 265, 345  
***vapidus*, *Strigiphilus*** 17, **212**, 265, 348  
var. *alpha*, *Nirmus triangulatus* 178  
var. *excavatus*, *Lipeurus diversus* 125  
var. *kea*, *Lipeurus circumfasciatus* 148  
var. *kea*, *Menopon fulvofasciatum* 64  
var. *major*, *Lipeurus diversus* 127  
var. *minor*, *Menopon fulvofasciatum* 75  
var. *ocellata*, *Trichodectes parumpilosus* 228  
var. *orientalis*, *Philopterus validus* 193  
var. *pallida*, *Lipeurus signatus* 88

var. *passeris*, *Docophorus communis* 167  
var. *triangulatus*, *Nirmus* 179  
*variabilis*, *Phlopterus (Docophorus)* 209  
*variabilis*, *Saemundssonina* 209  
**varius**, ***Pectinopygus*** 14, 25, **160**, 257, 341  
*ventrale*, *Menopon* 37  
*ventralis*, *Amyrsidea (Amyrsidea)* 37  
*ventricosus*, *Haematopinus* 242  
**ventricosus**, ***Haemodipsus*** 18, **242**, 243, 269  
*vestimenti*, *Pediculus* 240  
*victrix*, *Myrsidea* 72  
*Virgula* 103  
*vituli*, *Haematopinus* 237  
**vituli**, ***Linognathus*** 18, **237**, 238, 271  
*vituli*, *Pediculus* 237  
*vituli*, *Trichaulus* 237  
*wallacei*, *Phlopterus* 197  
**waterstoni**, ***Austrogoniodes*** 11, 90, 91, **94**, 95, 249  
*waterstoni*, *Cesareus* 94  
*waterstoni*, *Goniocotes* 90, 94  
*waterstoni*, *Phlopterus* 171  
**waterstoni**, ***Psittoecus*** **171**  
**Werneckiella** 17, 23, **227**, 228  
*wolfherrei*, *Philoceanus* 164  
**xenicus**, ***Phlopterooides*** 15, **167**, 266  
*zerafae*, *Colpocephalum* 58  
*zonaria*, *Degeeriella* 103  
*zonarius sensu lato*, *Carduiceps* 103  
**zonarius**, ***Carduiceps*** 12, **103**, 260  
*zonarius*, *Nirmus* 103







Area codes and boundaries used to categorise specimen locality data (after Crosby *et al.* 1976)

## FAUNA OF NEW ZEALAND PUBLICATIONS

- 1 **Terebrantia** (Insecta: Thysanoptera). *Laurence A. Mound & Annette K. Walker*. ISBN 0-477-06687-9, 23 December 1982, 120 pp. .... \$29.95
- 2 **Osoriinae** (Insecta: Coleoptera: Staphylinidae). *H. Pauline McColl*. ISBN 0-477-06688-7, 23 December 1982, 96 pp. .... \$18.60
- 3 **Anthribidae** (Insecta: Coleoptera). *B. A. Holloway*. ISBN 0-477-06703-4, 23 December 1982, 272 pp. .... \$41.00
- 4 **Eriophyoidea except Eriophyinae** (Arachnida: Acari). *D. C. M. Manson*. ISBN 0-477-06745-X, 12 November 1984, 144 pp. .... \$29.95
- 5 **Eriophyinae** (Arachnida: Acari: Eriophyoidea). *D. C. M. Manson*. ISBN 0-477-06746-8, 14 November 1984, 128 pp. .... \$29.95
- 6 **Hydraenidae** (Insecta: Coleoptera). *R. G. Ordish*. ISBN 0-477-06747-6, 12 November 1984, 64 pp. .... \$18.60
- 7 **Cryptostigmata** (Arachnida: Acari) — a concise review. *M. Luxton*. ISBN 0-477-06762-X, 8 December 1985, 112 pp. .... \$29.95
- 8 **Calliphoridae** (Insecta: Diptera). *James P. Dear*. ISBN 0-477-06764-6, 24 February 1986, 88 pp. .... \$18.60
- 9 **Protura** (Insecta). *S. L. Tuxen*. ISBN 0-477-06765-4, 24 February 1986, 52 pp. .... \$18.60
- 10 **Tubulifera** (Insecta: Thysanoptera). *Laurence A. Mound & Annette K. Walker*. ISBN 0-477-06784-0, 22 September 1986, 144 pp. .... \$34.65
- 11 **Pseudococcidae** (Insecta: Hemiptera). *J. M. Cox*. ISBN 0-477-06791-3, 7 April 1987, 232 pp. .... \$49.95
- 12 **Pompilidae** (Insecta: Hymenoptera). *A. C. Harris*. ISBN 0-477-02501-3, 13 November 1987, 160 pp. .... \$39.95
- 13 **Encyrtidae** (Insecta: Hymenoptera). *J. S. Noyes*. ISBN 0-477-02517-X, 9 May 1988, 192 pp. .... \$44.95
- 14 **Lepidoptera** — annotated catalogue, and keys to family-group taxa. *J. S. Dugdale*. ISBN 0-477-02518-8, 23 September 1988, 264 pp. .... \$49.95
- 15 **Ambositrinae** (Insecta: Hymenoptera: Diapriidae). *I. D. Naumann*. ISBN 0-477-02535-8, 30 December 1988, 168 pp. .... \$39.95
- 16 **Nepticulidae** (Insecta: Lepidoptera). *Hans Donner & Christopher Wilkinson*. ISBN 0-477-02538-2, 28 April 1989, 92 pp. .... \$22.95
- 17 **Mymaridae** (Insecta: Hymenoptera) — introduction, and review of genera. *J. S. Noyes & E. W. Valentine*. ISBN 0-477-02542-0, 28 April 1989, 100 pp. .... \$24.95
- 18 **Chalcidoidea** (Insecta: Hymenoptera) — introduction, and review of genera in smaller families. *J. S. Noyes & E. W. Valentine*. ISBN 0-477-02545-5, 2 August 1989, 96 pp. .... \$24.95
- 19 **Mantodea** (Insecta), with a review of aspects of functional morphology and biology. *G. W. Ramsay*. ISBN 0-477-02581-1, 13 June 1990, 96 pp. .... \$24.95
- 20 **Bibionidae** (Insecta: Diptera). *Roy A. Harrison*. ISBN 0-477-02595-1, 13 November 1990, 28 pp. .... \$14.95
- 21 **Margarodidae** (Insecta: Hemiptera). *C. F. Morales*. ISBN 0-477-02607-9, 27 May 1991, 124 pp. .... \$34.95
- 22 **Notonemouridae** (Insecta: Plecoptera). *I. D. McLellan*, ISBN 0-477-02518-8, 27 May 1991, 64 pp. .... \$24.95
- 23 **Sciapodinae, Medeterinae** (Insecta: Diptera) with a generic review of the Dolichopodidae. *D. J. Bickel*. ISBN 0-477-02627-3, 13 January 1992, 74 pp. .... \$27.95
- 24 **Therevidae** (Insecta: Diptera). *L. Lyneborg*. ISBN 0-477-02632-X, 4 March 1992, 140 pp. .... \$34.95
- 25 **Cercopidae** (Insecta: Homoptera). *K. G. A. Hamilton & C. F. Morales*. ISBN 0-477-02636-2, 25 May 1992, 40 pp. .... \$17.95
- 26 **Tenebrionidae** (Insecta: Coleoptera): catalogue of types and keys to taxa. *J. C. Watt*. ISBN 0-477-02639-7, 13 July 1992, 70 pp. .... \$27.95
- 27 **Antarctoperlinae** (Insecta: Plecoptera). *I. D. McLellan*. ISBN 0-477-01644-8, 18 February 1993, 70 pp. .... \$27.95
- 28 **Larvae of Curculionoidea** (Insecta: Coleoptera): a systematic overview. *Brenda M. May*. ISBN 0-478-04505-0, 14 June 1993, 226 pp. .... \$55.00
- 29 **Cryptorhynchinae** (Insecta: Coleoptera: Curculionidae). *C. H. C. Lyal*. ISBN 0-478-04518-2, 2 December 1993, 308 pp. .... \$65.00
- 30 **Hepialidae** (Insecta: Lepidoptera). *J. S. Dugdale*. ISBN 0-478-04524-7, 1 March 1994, 164 pp. .... \$42.50
- 31 **Talitridae** (Crustacea: Amphipoda). *K. W. Duncan*. ISBN 0-478-04533-6, 7 October 1994, 128 pp. .... \$36.00
- 32 **Sphecidae** (Insecta: Hymenoptera). *A. C. Harris*, ISBN 0-478-04534-4, 7 October 1994, 112 pp. .... \$33.50
- 33 **Moranilini** (Insecta: Hymenoptera). *J. A. Berry*. ISBN 0-478-04538-7, 8 May 1995, 82 pp. .... \$29.95
- 34 **Anthicidae** (Insecta: Coleoptera). *F. G. Werner & D. S. Chandler*. ISBN 0-478-04547-6, 21 June 1995, 64 pp. .... \$26.50

- 35 **Cydnidae, Acanthosomatidae, and Pentatomidae** (Insecta: Heteroptera): systematics, geographical distribution, and bioecology. *M.-C. Larivière*. ISBN 0-478- 09301-2, 23 November 1995, 112 pp. ....\$42.50
- 36 **Leptophlebiidae** (Insecta: Ephemeroptera). *D. R. Towns & W. L. Peters*. ISBN 0-478-09303-9, 19 August 1996, 144 pp. ....\$39.50
- 37 **Coleoptera**: family-group review and keys to identification. *J. Klimaszewski & J. C. Watt*. ISBN 0-478- 09312-8, 13 August 1997, 199 pp. ....\$49.50
- 38 **Naturalised terrestrial Stylommatophora** (Mollusca: Gastropoda). *G. M. Barker*. ISBN 0-478-09322-5, 25 January 1999, 253 pp. ....\$72.50
- 39 **Molytini** (Insecta: Coleoptera: Curculionidae: Molytinae). *R. C. Craw*. ISBN 0-478-09325-X, 4 February 1999, 68 pp. ....\$29.50
- 40 **Cixiidae** (Insecta: Hemiptera: Auchenorrhyncha). *M.-C. Larivière*. ISBN 0-478-09334-9, 12 November 1999, 93 pp. ....\$37.50
- 41 **Coccidae** (Insecta: Hemiptera: Coccoidea). *C. J. Hodgson & R. C. Henderson*. ISBN 0-478-09335-7, 23 February 2000, 264 pp. ....\$72.50
- 42 **Aphodiinae** (Insecta: Coleoptera: Scarabaeidae). *Z. T. Stebnicka*. ISBN 0-478-09341-1, 15 June 2001, 64 pp. ....\$29.50
- 43 **Carabidae** (Insecta: Coleoptera): catalogue. *A. Larochelle & M.-C. Larivière*. ISBN 0-478-09342-X, 15 June 2001, 285 pp. ....\$72.50
- 44 **Lycosidae** (Arachnida: Araneae). *C. J. Vink*. ISBN 0-478- 09347-0, 23 December 2002, 94 pp. ....\$37.50
- 45 **Nemonychidae, Belidae, Brentidae** (Insecta: Coleoptera: Curculionoidea). *G. Kuschel*. ISBN 0-478- 09348-9, 28 April 2003, 100 pp. ....\$40.00
- 46 **Nesameletidae** (Insecta: Ephemeroptera). *Terry R. Hitchings & Arnold H. Staniczek*. ISBN 0-478-09349-7, 14 May 2003, 72 pp. ....\$32.50
- 47 **Erotylidae** (Insecta: Coleoptera: Cucujoidea): phylogeny and review. *R. A. B. Leschen*. ISBN 0-478-09350-0, 5 June 2003, 108 pp. ....\$42.50
- 48 **Scaphidiinae** (Insecta: Coleoptera: Staphylinidae). *I. Löbl & R. A. B. Leschen*. ISBN 0-478-09353-5, 18 November 2003, 94 pp. ....\$37.50
- 49 **Lithinini** (Insecta: Lepidoptera: Geometridae: Ennominae). *J. D. Weintraub & M. J. Scoble*. ISBN 0-478- 09357-8, 29 April 2004, 48 pp. ....\$24.50
- 50 **Heteroptera** (Insecta: Hemiptera): catalogue. *M.-C. Larivière & A. Larochelle*. ISBN 0-478-09358-6, 14 May 2004, 330 pp. ....\$89.00
- 51 **Coccidae** (Insecta: Hemiptera: Coccoidea): adult males, pupae and prepupae of indigenous species. *C. J. Hodgson & R. C. Henderson*. ISBN 0-478-09360-8, 22 June 2004, 228 pp. ....\$65.00
- 52 **Raphignathoidea** (Acari: Prostigmata). *Qing-Hai Fan & Zhi-Qiang Zhang*. ISBN 0-478-09371-3, 20 May 2005, 400 pp. ....\$89.00
- 53 **Harpalini** (Insecta: Coleoptera: Carabidae: Harpalinae). *A. Larochelle & M.-C. Larivière*. ISBN 0-478-09369-1, 4 July 2005, 160 pp. ....\$55.00
- 54 **Hierodoris** (Insecta: Lepidoptera: Gelechoidea: Oecophoridae), and overview of Oecophoridae. *Robert J. B. Hoare*. ISBN 0-478-09378-0, 24 December 2005, 100 pp. ....\$40.00
- 55 **Criconematina** (Nematoda: Tylenchida). *W. M. Wouts*. ISBN 0-478-09381-0, 24 March 2006, 232 pp. ....\$65.00
- 56 **Tyrophagus** (Acari: Astigmata: Acaridae). *Qing-Hai Fan & Zhi-Qiang Zhang*. ISBN 978-0-478-09386-5, 4 June 2007, 291 pp. ....\$80.00
- 57 **Apoidea** (Insecta: Hymenoptera). *B. J. Donovan*. ISBN 978-0-478-09389-6, 7 September 2007, 295 pp. ....\$89.00
- 58 **Alysiinae** (Insecta: Hymenoptera: Braconidae). *J. A. Berry*. ISBN 978-0-478-09390-2, 7 September 2007, 95 pp. . ....\$45.00
- 59 **Erotylinae** (Insecta: Coleoptera: Cucujoidea: Erotylidae): taxonomy and biogeography. *Paul E. Skelley & Richard A. B. Leschen*. ISBN 978-0-478-09391-9, 7 September 2007, 59 pp. ....\$30.00
- 60 **Carabidae** (Insecta: Coleoptera): synopsis of supraspecific taxa. *A. Larochelle & M.-C. Larivière*. ISBN 978-0-478-09394-0, 21 November 2007, 188 pp. ....\$54.00
- 61 **Lucanidae** (Insecta: Coleoptera). *B. A. Holloway*. ISBN 978-0-478-09395-7, 21 November 2007, 254 pp. . \$75.00
- 62 **Trechini** (Insecta: Coleoptera: Carabidae: Trechinae). *J. I. Townsend*. ISBN 978-0-478-34717-9 (print), 978-0-478-34716-6 (online), 16 June 2010, 101 pp. ....\$49.50
- 63 **Auchenorrhyncha** (Insecta: Hemiptera): catalogue. *M.-C. Larivière, M. J. Fletcher & A. Larochelle*. ISBN 978-0-478-34720-3 (print), 978-0-478-34721-0 (online), 16 June 2010, 232 pp. ....\$75.00

- 64 **Pisauridae** (Arachnida: Araneae). C. J. Vink & N. Dupérré. ISBN 978-0-478-34722-7 (print), 978-0-478- 34723-4 (online), 13 July 2010, 60 pp. .... \$37.50
- 65 **Izatha** (Insecta: Lepidoptera: Gelechioidea: Oecophoridae). Robert J. B. Hoare. ISBN 978-0-478-34724-1 (print), 978-0-478-34725-8 (online), 2 September 2010, 201 pp..... \$75.00
- 66 **Diaspididae** (Insecta: Hemiptera: Coccoidea). R. C. Henderson. ISBN 978-0-478-34726-5 (print), 978-0-478-34727-2 (online), 23 May 2011, 275 pp. .... \$89.00
- 67 **Peloriidiidae** (Insecta: Hemiptera: Coleorrhyncha). M.-C. Larivière, D. Burckhardt & A. Larochelle. ISBN 978-0-478- 34730-2 (print), 978-0-478-34731-9 (online), 14 November 2011, 78 pp. .... \$48.00
- 68 **Simuliidae** (Insecta: Diptera). Douglas A. Craig, Ruth E. G. Craig & Trevor K. Crosby. ISBN 978-0-478-34734- 0 (print), 978-0-478-34735-7 (online), 29 June 2012, 336 pp. .... \$95.00
- 69 **Carabidae** (Insecta: Coleoptera): synopsis of species, Cicindelinae to Trechinae (in part) . A. Larochelle & M.-C. Larivière. ISBN 978-0-478-34738-8 (print), 978-0-478- 34739-5 (online), 7 March 2013, 193 pp. .... \$75.00
- 70 **Periegopidae** (Arachnida: Araneae). C. J. Vink, N. Dupérré & J. Malumbres-Olarte. ISBN 978-0-478-34740- 1 (print), 978-0-478-34741-8 (online), 7 March 2013, 41 pp. .... \$29.00
- 71 **Fanniidae** (Insecta: Diptera). M. Cecilia Domínguez & Adrian C. Pont. ISBN 978-0-478-34745-6 (print), 978-0-478-34746-3 (online), 30 June 2014, 91 pp. .... \$49.00
- 72 **Micropterigidae** (Insecta: Lepidoptera). George W. Gibbs. ISBN 978-0-478-34759-3 (print), 978-0-478-34760- 9 (online), 30 June 2014, 127 pp..... \$59.00
- 73 **Noctuinae** (Insecta: Lepidoptera: Noctuidae) part 1: *Austramathes*, *Cosmodes*, *Proteuxoa*, *Physetica*. R.J.B. Hoare, ISBN 978-0-947525-09-5 (print), 978-0-947525-07-1 (online), 23 June 2017, 130 pp ..... \$60.00
- 74 **Ceratomerinae** (Diptera: Empidoidea: Brachystomatidae), B.J. Sinclair, ISBN 978-0-947525-10-1 (print), 978-0-947525-11-8 (online), 23 June 2017, 157 pp ..... \$66.00
- 75 **Licinini** (Insecta: Coleoptera: Carabidae: Harpalinae), R.M. Emberson. ISBN 978-0-947525-13-2 (print), 978-0-947525-12-5 (online), 16 August 2017, 80 pp ..... \$48.00
- 76 **Phthiraptera** (Insecta): a catalogue of parasitic lice from New Zealand, Ricardo L. Palma. ISBN 978-0-947525-19-4 (print), 978-0-947525-18-7 (online), 30 September 2017, 400 pp ..... \$98.00

Visit the Landcare Research Website at: <http://www.landcareresearch.co.nz/> for further information.

Orders for printed books should be sent to the Library of Landcare Research ([library@landcareresearch.co.nz](mailto:library@landcareresearch.co.nz)).

To access on-line version of this series visit: <http://fnz.landcareresearch.co.nz/> or <https://www.biotaxa.org/fnz>

## Taxonomic groups covered in the *Fauna of New Zealand* series

### Insecta

#### Coleoptera

Family-group review and keys to identification (*J. Klimaszewski & J. C. Watt*, FNZ 37, 1997)

Anthribidae (*B. A. Holloway*, FNZ 3, 1982)

Anthicidae (*F. G. Werner & D. S. Chandler*, FNZ 34, 1995)

Carabidae: catalogue (*A. Larochelle & M.-C. Larivière*, FNZ 43, 2001); synopsis of supraspecific taxa (*A. Larochelle & M.-C. Larivière*, FNZ 60, 2007); synopsis of species, Cicindelinae to Trechinae (in part) (*A. Larochelle & M.-C. Larivière*, FNZ 69, 2013)

Carabidae: Harpalinae: Harpalini (*A. Larochelle & M.-C. Larivière*, FNZ 53, 2005)

Carabidae: Trechinae: Trechini (*J. I. Townsend*, FNZ 62, 2010)

Curculionidae: Cryptorhynchinae (*C. H. C. Lyal*, FNZ 29, 1993)

Curculionidae: Molytinae: Molytini (*R. C. Craw*, FNZ 39, 1999)

Curculionoidea: Nemonychidae, Belidae, Brentidae (*G. Ku-schel*, FNZ 45, 2003)

Curculionoidea larvae: a systematic overview (*Brenda M. May*, FNZ 28, 1993)

Erotylidae: phylogeny and review (*Richard A. B. Leschen*, FNZ 47, 2003); Erotylinae: taxonomy and biogeography (*Paul E. Skelley & Richard A. B. Leschen*, FNZ 59, 2007)

Hydraenidae (*R. G. Ordish*, FNZ 6, 1984)

Lucanidae (*B. A. Holloway*, FNZ 61, 2007)

Scarabaeidae: Aphodiinae (*Z. T. Stebnicka*, FNZ 42, 2001)

Staphylinidae: Osoriinae (*H. Pauline McColl*, FNZ 2, 1982)

Staphylinidae: Scaphidiinae (*I. Löbl & Richard A. B. Leschen*, FNZ 48, 2003)

Tenebrionidae: catalogue of types and keys to taxa (*J. C. Watt*, FNZ 26, 1992)

Carabidae: Harpalinae: Licinini (*R.M. Emberson*, FNZ 75, 2017)

#### Diptera

Bibionidae (*Roy A. Harrison*, FNZ 20, 1990)

Calliphoridae (*James P. Dear*, FNZ 8, 1986)

Dolichopodidae: Sciapodinae, Medeterinae with a generic review (*D. J. Bickel*, FNZ 23, 1992)

Fanniidae (*M. Cecilia Domínguez, Adrian C. Pont*, FNZ 71, 2014)

Simuliidae (*Douglas A. Craig, Ruth E. G. Craig, Trevor K. Crosby*, FNZ 68, 2012)

Therevidae (*L. Lyneborg*, FNZ 24, 1992)

Ceratomerinae (Diptera: Empidoidea: Brachystomatidae) (*B.J. Sinclair*, FNZ 74, 2017)

#### Ephemeroptera

Leptophlebiidae (*D. R. Towns & W. L. Peters*, FNZ 36, 1996)

Nesameletidae (*Terry R. Hitchings & Arnold H. Staniczek*, FNZ 46, 2003)

#### Hemiptera

Auchenorrhyncha: catalogue (*M.-C. Larivière, M. J. Fletcher & A. Larochelle*, FNZ 63, 2010)

Cercopidae (*K. G. A. Hamilton & C. F. Morales*, FNZ 25, 1992)

Cixiidae (*M.-C. Larivière*, FNZ 40, 1999)

Coccidae (*C. J. Hodgson & R. C. Henderson*, FNZ 41, 2000); adult males, pupae and prepupae of indigenous species (*C. J. Hodgson & R. C. Henderson*, FNZ 51, 2004)

Cydnidae, Acanthosomatidae, and Pentatomidae (*M.-C. Larivière*, FNZ 35, 1995)

Diaspididae (*R. C. Henderson*, FNZ 66, 2011)

Heteroptera: catalogue (*M.-C. Larivière & A. Larochelle*, FNZ 50, 2004)

Margarodidae (*C. F. Morales*, FNZ 21, 1991)

Pseudococcidae (*J. M. Cox*, FNZ 11, 1987)

Peloriidae (*M.-C. Larivière, D. Burckhardt & A. Larochelle*, FNZ 67, 2011).

#### Hymenoptera

Apoidea (*B. J. Donovan*, FNZ 57, 2007)

Braconidae: Alysiinae (*J. A. Berry*, FNZ 58, 2007)

Chalcidoidea: introduction, and review of smaller families (*J. S. Noyes & E. W. Valentine*, FNZ 18, 1989)

Diapriidae: Ambositrinae (*I. D. Naumann*, FNZ 15, 1988)

Encyrtidae (*J. S. Noyes*, FNZ 13, 1988)

Mymaridae (*J. S. Noyes & E. W. Valentine*, FNZ 17, 1989)

Pompilidae (*A. C. Harris*, FNZ 12, 1987)

Pteromalidae: Eunotinae: Moranilini (*J. A. Berry*, FNZ 33, 1995)

Sphecidae (*A. C. Harris*, FNZ 32, 1994)

### **Lepidoptera**

Annotated catalogue, and keys to family-group taxa (*J. S. Dugdale*, FNZ 14, 1988)

Geometridae: Ennominae: Lithinini (*Jason D. Weintraub & Malcolm J. Scoble*, FNZ 49, 2004)

Hepialidae (*J. S. Dugdale*, FNZ 30, 1994)

Micropterigidae (*George W. Gibbs*, FNZ 72, 2014)

Nepticulidae (*Hans Donner & Christopher Wilkinson*, FNZ 16, 1989)

Oecophoridae: *Hierodoris* (*Robert J. B. Hoare*, FNZ 54, 2005); *Izatha* (*Robert J. B. Hoare*, FNZ 65, 2010).

Noctuinae (Insecta: Lepidoptera: Noctuidae) part 1: *Austramathes*, *Cosmodes*, *Proteuxoa*, *Physetica* (*R.J.B. Hoare*, FNZ 73, 2017)

**Mantodea**, with a review of aspects of functional morphology and biology (*G. W. Ramsay*, FNZ 19, 1990)

**Phthiraptera** (Insecta): a catalogue of parasitic lice from New Zealand (*R. L. Palma*, FNZ 76, 2017)

### **Plecoptera**

Antarctoperlinae (*I. D. McLellan*, FNZ 27, 1993)

Notonemouridae (*I. D. McLellan*, FNZ 22, 1991)

**Protura** (*S. L. Tuxen*, FNZ 9, 1986)

### **Thysanoptera**

Terebrantia (*Laurence A. Mound & Annette K. Walker*, FNZ 1, 1982)

Tubulifera (*Laurence A. Mound & Annette K. Walker*, FNZ 10, 1986)

## **Arachnida**

### **Acari**

Acaridae: *Tyrophagus* (*Qing-Hai Fan & Zhi-Qiang Zhang*, FNZ 56, 2007)

Cryptostigmata — a concise review (*M. Luxton*, FNZ 7, 1985)

Eriophyoidea except Eriophyinae (*D. C. M. Manson*, FNZ 4, 1984)

Eriophyinae (*D. C. M. Manson*, FNZ 5, 1984)

Raphignathoidea (*Qing-Hai Fan & Zhi-Qiang Zhang*, FNZ 52, 2005)

### **Araneae**

Lycosidae (*C. J. Vink*, FNZ 44, 2002)

Periegopidae (*C. J. Vink, N. Dupérré & Malumbres-Olarte*, FNZ 70, 2013)

Pisauridae (*C. J. Vink & N. Dupérré*, FNZ 64, 2010)

## **Crustacea**

### **Amphipoda**

Talitridae (*K. W. Duncan*, FNZ 31, 1994)

## **Mollusca**

### **Gastropoda**

Naturalised terrestrial Stylommatophora (*G. M. Barker*, FNZ 38, 1999)

## **Nematoda**

**Tylenchida**: Criconematina (*W. M. Wouts*, FNZ 55, 2006)

## NOTICES

This series of refereed publications has been established to encourage those with expert knowledge to publish concise yet comprehensive accounts of elements in the New Zealand fauna. The series is professional in its conception and presentation, yet every effort is made to provide resources for identification and information that are accessible to the non-specialist.

*Fauna of N.Z.* deals with non-marine invertebrates only, since the vertebrates are well documented, and marine forms are covered by the series *NIWA Biodiversity Memoirs*.

**Contributions** are invited from any person with the requisite specialist skills and resources. Material from the N.Z. Arthropod Collection is available for study.

Contributors should discuss their intentions with a member of the Editorial Board or with the Series Editor before commencing work; all necessary guidance will be given.

## NGĀ PĀNUI

Kua whakatūria tēnei huinga pukapuka hei whakahauhau i ngā tohunga whai mātauranga kia whakaputa i ngā kōrero poto, engari he whaikiko tonu, e pā ana ki ngā aitanga pepeke o Aotearoa. He tōtika tonu te āhua o ngā tuhituhi, engari ko te tino whāinga, kia mārama te marea ki ngā tohu tautuhi o ia ngārara, o ia ngārara, me te roanga atu o ngā kōrero mō tēnā, mō tēnā.

He titiro whāiti tā tēnei pukapuka ki ngā mea noho whenua, kāore he tuarā; i pēnei ai i te mea kei te mōhio whānuitia ngā mea whai tuarā, ā, ko ngā mea noho moana, koirā te tino kaupapa o te huinga pukapuka *NIWA Biodiversity Memoirs*.

Ka āhei te tangata ki te **whakauru tuhituhinga** mehemea kei a ia ngā tohungatanga me ngā rauemi e tutuki pai ai tana mahi. Heoi anō, e wātea ana te Kohinga Angawaho o Aotearoa hei āta tiro tiro mā te tangata mehemea he āwhina kei reira.

Me whāki te kaituhi i ōna whakaaro ki tētahi o te Kāhui Ārahi Whakarōpūtanga Tuarā-Kore, ki te Ētita rānei i mua i te tīmatanga, ā, mā rātou a ia e ārahi mō te wāhi ki tana tuhinga.