



Landcare Research LINK Seminar
25 August 2015

2015 International Year of the Soil

Where has all the carbon gone?

*The answer lies in
the soil.....*

David Whitehead
Louis Schipper
Miko Kirschbaum



WORKING TOGETHER




Landcare Research
Manaaki Whenua



Global carbon storage
23% atmosphere

15% vegetation

62% soil to 1 m



- Carbon input to soil is regulated by plants
- Carbon retention is regulated by physical and microbial processes
- Carbon is stored in a range of organic materials with turnover rates from days to centuries
- Disturbance can cause rapid losses and recovery is often slow



Soil carbon is essential for maintaining the productive potential of our primary industries

- soil physical structure and stability
- water retention
- nutrient cycling
- buffering and filtering

Retaining and increasing soil carbon provides opportunity to offset our greenhouse gas emissions. Research is needed to inform our international negotiations

Identify land management practices to maintain soil carbon stocks and, if possible, achieve stable, increased stocks



- Top soil carbon stocks can be high
Average for NZ's grassland soils is 100 t C/ha to a depth of 0.3 m
- Deeper in soils, carbon stocks can be much lower but have higher potential to store carbon
- Carbon stability (longevity) in soils is not well understood
- *Between 1990 and 2013 increases in NZ's methane (8%) and soil nitrous oxide emissions (23%) are equivalent to 1 Mt C*
- *This could be offset with an increase in soil carbon of 1 t C/ha over 1 Mha or 0.1 t C/ha over NZ's approx. 10 Mha grassland estate ie. 0.1% increase*
- *Increasing soil carbon stocks commits nitrogen, phosphorus, sulphur and other nutrients. This represents of order \$200 for 1 t C/ha based on today's fertiliser costs*

Improved measurements of soil carbon



Visible near infra red spectroscopy

Hedley et al(2015), Roudier et al (2015)




- rapid
- lower cost
- increased spatial and depth representation
- allows spatial scaling
- enables interpretation about carbon stability
- increased efficiency for accounting practices


Tuapaka Hill Country (Manawatu)

Soil organic carbon t C/ha
to 0.3m depth
at 50 sampling positions

Carolyn Hedley, Pierre Roudier,
Leo Valette (CSIRO)
GRA funding




Relative importance of covariates (%)



0 200 400 600 800 1000 m



What is the potential for increasing soil carbon stocks?




- From National Soils Database including long term grasslands
- Using a spatially explicit model, differences in carbon content were attributable to surface area, aluminium and pH
- Potential carbon saturation deficit was estimated from the difference between the upper (90th percentile) and current level 50th percentile)
- 0 – 0.15 m average potential saturation deficit 32%
- 0.15 – 0.3 m average potential saturation deficit 83%
- At 40 mg C/g (0 - 0.15 m) filling the deficit equivalent to 30% increase carbon stocks

We need to use management practices that maintain and increase soil carbon

It's changes in carbon stocks that are important



Pasture renewal



Irrigation

Fertiliser addition



Exotic worms



Mixed swards



Stocking, supplemental feed




Biochar addition

We need to use management practices that maintain and increase soil carbon

It's changes in carbon stocks that are important





Pasture renewal



Irrigation

Fertiliser addition

Stocking, supplemental feed



Exotic worms



Mixed swards



Biochar addition

Change in grassland carbon stocks

National Soils Database resampling




- Resampling sites to 1 m previously sampled 20-40 years previously
- Analysed archived soil samples

Schipper et al. (2014)

Change in grassland carbon stocks

National Soils Database resampling after 20-40 years



Change in grassland carbon stocks

National Soils Database resampling

2014 (148 sites)

- Allophanic (-0.5 t/ha/y) and Gley (-0.3 t/ha/y) losing carbon
- Other mineral soils no significant change
- Hill country gaining carbon (0.6 t/ha/y)

- No apparent effect of grazing type

Change in grassland carbon stocks

500 soils resampling

Approach

- 158 sites resampled after 7 years to 0.1 m depth
- Range of people collected samples
- No reanalysis of archived soils

Findings

- Gains on dairy 0.32 t C/ha/y and drystock 0.57 t C/ha/y
- Not significant from zero
- Combined was significant 0.42 t C/ha/y for flat land
- Gains on hill country 1.33 t C/ha/y

Carbon changes in organic soils

- Loss of 2.9 t C/ha/y
- Size of error ? Only one site
- Many peats many metres deep and losses will continue as long as they are drained for farming: many centuries
 - at about 0.02 m/y




Excessive drainage, Moanatuatua

Campbell et al. (2015)

Management effects: P fertiliser

Winchmore, South Canterbury
Whatawhata, Waikato


No benefit of adding P on
soil carbon recovery



Note: shallow sampling

Management effects: irrigation

Winchmore, South Canterbury



Also confirmed to 1 m depth (Condrón et al. 2014)

Management effects: irrigation


Preliminary data

South Island

0 – 0.3 m depth


10 farms

Similar but less significant effects at North Island sites



Management effects: pasture renewal

- Occurs every 5 to 10 years
- Sprayed off and can involve cultivation
- Total carbon losses of between 0.8 and 4.1 t C/ha (2-3% of carbon stock to 0.3 m)
- Losses and gains dependent on soil water availability
- Likely recovered between renewals
- 2 farms only





Rutledge et al. (2014)

Can diverse pastures capture more carbon?





Lucerne
Chicory
Plantain
Ryegrass
Clover






Troughton Farm, Waikato
Established late 2011 on 3 ryegrass/clover areas
Treatments imposed early 2013

Can diverse pastures capture more carbon?




Carbon balance

Net carbon exchange



Carbon imports
(feed, effluent)




Carbon exports
(milk/silage)



Farm scale carbon balance

Scott Farm, Waikato, 4 years



Carbon sink overall
average $\sim 600 \pm 320$ kg C/ha/y

Both weather and
management impact
the annual carbon balance

But how stable is this carbon and how much more can NZ soil store?

Summary from historical observations

Time trends in soil carbon stocks

Flat land

- losses up to 0.5 t C/ha/y from Allophanic and Gley soils to 0.3 m depth over 30 years
- large ongoing losses up to 2.9 t C/ha/y from organic soils (1 site!)
- no change other soil orders
- some evidence of recent increases in top 0.1 m depth (method?)

Hill country

- increases up to 0.6 t C/ha/y observed both short and long term

Summary from historical observations

Management effects on soil carbon stocks

- P fertiliser: no detectable change
- N fertiliser: no information available
- Irrigation: decrease BUT size of loss to be determined shortly
- Pasture renewal: small decrease probably recovers if infrequent
- Diverse swards: short-term increases but no long-term data yet

How to estimate changes at national scale?

- Limited historical observations do not provide clarity
- Trends depend on soil type, slope and management
- Complexities of multiple variables interacting
 - eg. soil type, climate, irrigation, fertiliser, animal stocking
- Currently no regular soil carbon monitoring in New Zealand

How to forecast future soil carbon changes?

- Continue re-sampling and analysis at historically sampled sites
- Need process-based studies to understand and predict
- **Future progress depends on the use of models to interpret and forecast management effects and best practices**

National scale soil carbon trends

	Per area change (t C ha ⁻¹ yr ⁻¹)	Area (ha)	Total change (MtCO ₂ C yr ⁻¹)
Tussocks/ low-producing	0.0 ± 0.26	4 116 750	0.0 (-3.92 to 3.92)
Allophanic soils/ flat land	-0.54 ± 0.32	454 182	-0.9 (-1.43 to -0.37)
Gley soils/ flat land	-0.32 ± 0.31	655 411	-0.77 (-1.51 to -0.02)
Organic soils	-2.9 ± 1.3	140 589	-1.49 (-2.17 to -0.82)
Other soils/ flat land	0.0 ± 0.19	3 492 757	0.0 (-2.43 to 2.43)
Hill-country soils (mid-slope)	0.6 ± 0.31	1 047 042	2.3 (1.11 to 3.49)
Hill-country soils (other slopes)	no data	2 330 473	no data
National total			-0.86 (-5.76 to 4.04)




30-year analyses of carbon stocks
in upper 0.3 m
Schipper et al. (2014)

National scale soil carbon trends


	Per area change (t C ha ⁻¹ yr ⁻¹)	Area (ha)	Total change (Mt C yr ⁻¹)
Tussocks/ low-producing	0.0 ± 0.26	4 116 750	0.0 (-3.92 to 3.92)
All flat land	0.4 ± 0.33	4 602 350	7.09 (1.52 to 12.66)
Organic soils	-2.9 ± 1.3	140 589	-1.49 (-2.17 to -0.82)
Hill-country soils (mid-slope)	1.33 ± 1.02	1 047 042	5.11 (1.19 to 9.02)
Hill-country soils (other slopes)	no data	2 330 473	no data
National total			10.7 (2.81 to 18.59)

7-year analyses of soil quality in
upper 0.1 m
Parfitt et al. (2014)

Soil carbon trends - extrapolation




Carbon accounting rules (Net-Net)




at land?

at land?






CenW model






Modelling results are consistent with observations





Supplemental feed



Fertiliser addition



Changing rainfall, irrigation



Conclusions

- National-scale estimates of carbon stocks changes rely on a small number of measurements
- Changing trends could be real or not! Many questions remain
- There is potential to increase carbon in New Zealand soils
- Changes depend on carbon gain, grazing off-take, carbon stabilisation and turn-over
- Carbon can increase with supplemental feeding, fertiliser addition, and irrigation on very dry sites
- Carbon increases can be achieved at the cost of reduced milk production
- Management practices most likely to achieve increase are:
 - optimising nitrogen addition and irrigation
 - increasing carbon inputs from roots eg. mixed swards

Landcare Research

Carolyn Hedley
Miko Kirschbaum
Andrew Manderson
Stephen McNeill
Pete Millard
Gabriel Moinet
Paul Mudge
Roger Parfitt
Beckie Phillips
Nicolas Puche
Pierre Roudier
David Whitehead

AgResearch

Mike Dodd
Frank Kelliher
Alec Mackay
Nicole Schon

Massey University

Marta Camps Arbestain
Roberto Calvelo Pereira

Plant & Food Research

Mike Beare
Denis Curtin

University Waikato

Dave Campbell
Sam McNally
Jack Pronger
Susanna Rutledge
Louis Schipper
Aaron Wall

Scion

Simeon Smail

GNS Science

Troy Baisden

NZAGRC PGgRc

Harry Clark
Andy Reisinger

CSIRO


Jeff Baldock

MPI

Gerald Rys




Soil C trends – temporal variability



Case study from Te Whanga catchment (de Rose, 2013)

Model-data comparison




Simulations with CenW (Kirschbaum et al., 2015)
Data from Waikato University (Rutledge, Mudge,
Schipper et al.)

Understanding Biochar




Biochar and Carbon Stability

BIOCHAR



Biochar mineralizes more slowly than the biomass it was produced from

UNPYROLYZED BIOMASS



Biochar C storage capacity differs widely!

Biochars produced from ash rich material (e.g., manure) at low temperature

C storage value

Biochars produced from woody material at high temperature

Class 1 (< 300 g C kg⁻¹ biochar will remain stable for > 100 years)

Class 5 (> 600 g C kg⁻¹ biochar will remain stable for > 100 years)

Biochar and Fertiliser Value



Biochars
produced from
pine

Fertiliser value

Biochars
produced from
poultry litter,
tomato waste

Class 0 (no fertiliser value for hypothetical
Corn needs at doses $\leq 10 \text{ t ha}^{-1}$)

Class 4 (fertiliser class 4;
e.g., $\text{K}_{2\text{t}}$, $\text{P}_{2\text{t}}$, $\text{S}_{5\text{t}}$, $\text{Mg}_{3\text{t}}$)

Biochar and Liming Value



Biochars
produced from
pine at low
temperature

Liming value

Biochars
produced from
tomato waste,
Paper sludge

Class 0 (liming eq < 1%)

Class 3 (liming eq > 20%)